

# Bicolorings and Equitable Bicolorings of Matrices

Michele Conforti\*    Gérard Cornuéjols †    Giacomo Zambelli †

dedicated to Manfred Padberg

## Abstract

Two classical theorems of Ghouila-Houri and Berge characterize total unimodularity and balancedness in terms of equitable bicolorings and bicolorings, respectively. In this paper, we prove a bicoloring result that provides a common generalization of these two theorems.

A  $0/\pm 1$  matrix is *balanced* if it does not contain a square submatrix with exactly two nonzero entries per row and per column such that the sum of all the entries is congruent to 2 modulo 4. This notion was introduced by Berge [1] for  $0/1$  matrices and generalized by Truemper [15] to  $0/\pm 1$  matrices.

A  $0/\pm 1$  matrix is *bicolorable* if its columns can be partitioned into blue columns and red columns so that every row with at least two nonzero entries contains either two nonzero entries of opposite sign in columns of the same color or two nonzero entries of the same sign in columns of different colors. Berge [1] showed that a  $0/1$  matrix  $A$  is balanced if and only if every submatrix of  $A$  is bicolorable. Conforti and Cornuéjols [6] extended this result to  $0/\pm 1$  matrices. Cameron and Edmonds [3] gave a simple greedy algorithm to find a bicoloring of a balanced matrix. In fact, given any  $0/\pm 1$  matrix  $A$ , their algorithm finds either a bicoloring of  $A$  or a square submatrix of  $A$  with exactly two nonzero entries per row and per column such that the sum of all the entries is congruent to 2 modulo 4. Does this algorithm provide an easy test for balancedness? The answer is no, because the algorithm may find a bicoloring of  $A$  even when  $A$  is not balanced.

A real matrix is *totally unimodular* (t.u.) if every nonsingular square submatrix has determinant  $\pm 1$  (note that every t.u. matrix must be a  $0/\pm 1$  matrix).

A  $0/\pm 1$  matrix  $A$  has an *equitable bicoloring* if its columns can be partitioned into red and blue columns so that, for every row of  $A$ , the sum of the entries in the red columns differs by at most one from the sum of the entries in the blue columns. Ghouila-Houri

---

\*Dipartimento di Matematica Pura ed Applicata, Università di Padova, Via Belzoni 7, 35131, Padova, Italy

†Graduate School of Industrial Administration, Carnegie Mellon University, Schenley Park, Pittsburgh, Pennsylvania 15213-3890

This work was supported in part by NSF grant DMI-0098427 and ONR grant N00014-97-1-0196.

[9] showed that a  $0/\pm 1$  matrix is totally unimodular if and only if every submatrix of  $A$  has an equitable bicoloring.

A  $0/\pm 1$  matrix which is not totally unimodular but whose submatrices are all totally unimodular is said *almost totally unimodular*. Camion [4] proved the following:

**Theorem 1 (Camion [4] and Gomory [cited in [4]])** *Let  $A$  be an almost totally unimodular  $0/\pm 1$  matrix. Then  $A$  is square,  $\det A = \pm 2$  and  $A^{-1}$  has only  $\pm \frac{1}{2}$  entries. Furthermore, each row and each column of  $A$  has an even number of nonzero entries and the sum of all entries in  $A$  equals 2 modulo 4.*

A nice proof of this result can be found in Padberg [12], [13]. Note that a matrix is balanced if and only if it does not contain any almost totally unimodular matrix with two nonzero entries in each row. For any positive integer  $k$ , we say that a  $0/\pm 1$  matrix  $A$  is  *$k$ -balanced* if it does not contain any almost totally unimodular submatrix with at most  $2k$  nonzero entries in each row. Obviously, an  $m \times n$   $0/\pm 1$  matrix  $A$  is balanced if and only if it is 1-balanced, while  $A$  is totally unimodular if and only if  $A$  is  $k$ -balanced for some  $k \geq \lfloor n/2 \rfloor$ . The class of  $k$ -balanced matrices was introduced by Conforti, Cornuéjols and Truemper in [7].

For any integer  $k$ , we denote by  $\mathbf{k}$  a vector with all entries equal to  $k$ . For any  $m \times n$   $0/\pm 1$  matrix  $A$ , we denote by  $n(A)$  the vector with  $m$  components whose  $i$ th component is the number of  $-1$ 's in the  $i$ th row of  $A$ .

**Theorem 2 (Conforti, Cornuéjols and Truemper [7])** *Let  $A$  be an  $m \times n$   $k$ -balanced  $0/\pm 1$  matrix with rows  $a^i$ ,  $i \in [m]$ ,  $b$  be a vector with entries  $b_i$ ,  $i \in [m]$ , and let  $S_1, S_2, S_3$  be a partition of  $[m]$ . Then*

$$P(A, b) = \{x \in \mathbb{R}^n : \begin{aligned} &a^i x \leq b_i \text{ for } i \in S_1 \\ &a^i x = b_i \text{ for } i \in S_2 \\ &a^i x \geq b_i \text{ for } i \in S_3 \\ &\mathbf{0} \leq x \leq \mathbf{1} \end{aligned}\}$$

*is an integral polytope for all integral vectors  $b$  such that  $-n(A) \leq b \leq \mathbf{k} - n(A)$ .*

This theorem generalizes previous results by Hoffman and Kruskal [10] for totally unimodular matrices, Berge [2] for 0/1 balanced matrices, Conforti and Cornuéjols [6] for  $0/\pm 1$  balanced matrices, and Truemper and Chandrasekaran [16] for  $k$ -balanced 0/1 matrices. As an application of Theorem 2, consider the SAT problem where, in each clause of a set of CNF clauses, at least  $k$  literals must evaluate to True. This SAT problem can be formulated as  $Ax \geq \mathbf{k} - n(A)$ ,  $x \in \{0, 1\}^n$ . If the matrix  $A$  is  $k$ -balanced, it follows from Theorem 2 that the polytope  $Ax \geq \mathbf{k} - n(A)$ ,  $\mathbf{0} \leq x \leq \mathbf{1}$  is integral and therefore the SAT problem can be solved by linear programming.

A  $0/\pm 1$  matrix  $A$  has a  *$k$ -equitable bicoloring* if its columns can be partitioned into blue columns and red columns so that:

- the bicoloring is equitable for the row submatrix  $A'$  determined by the rows of  $A$  with at most  $2k$  nonzero entries,
- every row with more than  $2k$  nonzero entries contains  $k$  pairwise disjoint pairs of nonzero entries such that each pair contains either entries of opposite sign in columns of the same color or entries of the same sign in columns of different colors.

Obviously, an  $m \times n$   $0/\pm 1$  matrix  $A$  is bicolored if and only if  $A$  has a 1-equitable bicoloring, while  $A$  has an equitable bicoloring if and only if  $A$  has a  $k$ -equitable bicoloring for  $k \geq \lfloor n/2 \rfloor$ . The following theorem provides a new characterization of the class of  $k$ -balanced matrices, which generalizes the bicoloring results mentioned above for balanced and totally unimodular matrices.

**Theorem 3** *A  $0/\pm 1$  matrix  $A$  is  $k$ -balanced if and only if every submatrix of  $A$  has a  $k$ -equitable bicoloring.*

**Proof.** Assume first that  $A$  is  $k$ -balanced and let  $B$  be any submatrix of  $A$ . Assume, up to row permutation, that

$$B = \begin{pmatrix} B' \\ B'' \end{pmatrix}$$

where  $B'$  is the row submatrix of  $B$  determined by the rows of  $B$  with  $2k$  or fewer nonzero entries. Consider the system

$$\begin{aligned} B'x &\geq \left\lfloor \frac{B'\mathbf{1}}{2} \right\rfloor \\ -B'x &\geq -\left\lceil \frac{B'\mathbf{1}}{2} \right\rceil \\ B''x &\geq \mathbf{k} - n(B'') \\ -B''x &\geq \mathbf{k} - n(-B'') \\ \mathbf{0} &\leq x \leq \mathbf{1} \end{aligned} \tag{1}$$

Since  $B$  is  $k$ -balanced, also  $\begin{pmatrix} B \\ -B \end{pmatrix}$  is  $k$ -balanced. Therefore the constraint matrix of system (1) above is  $k$ -balanced. One can readily verify that  $-n(B') \leq \lfloor \frac{B'\mathbf{1}}{2} \rfloor \leq \mathbf{k} - n(B')$  and  $-n(-B') \leq -\lceil \frac{B'\mathbf{1}}{2} \rceil \leq \mathbf{k} - n(-B')$ . Therefore, by Theorem 2 applied with  $S_1 = S_2 = \emptyset$ , system (1) defines an integral polytope. Since the vector  $(\frac{1}{2}, \dots, \frac{1}{2})$  is a solution for (1), the polytope is nonempty and contains a 0/1 point  $\bar{x}$ . Color a column  $i$  of  $B$  blue if  $\bar{x}_i = 1$ , red otherwise. It can be easily verified that such a bicoloring is, in fact,  $k$ -equitable.

Conversely, assume that  $A$  is not  $k$ -balanced. Then  $A$  contains an almost totally unimodular matrix  $B$  with at most  $2k$  nonzero elements per row. Suppose that  $B$  has a  $k$ -equitable bicoloring, then such a bicoloring must be equitable since each row has, at most,  $2k$  nonzero elements. By Theorem 1,  $B$  has an even number of nonzero elements in

each row. Therefore the sum of the columns colored blue equals the sum of the columns colored red, therefore  $B$  is a singular matrix, a contradiction.  $\square$

Given a  $0/\pm 1$  matrix  $A$  and positive integer  $k$ , one can find in polynomial time a  $k$ -equitable bicoloring of  $A$  or a certificate that  $A$  is not  $k$ -balanced as follows:

Find a basic feasible solution of (1). If the solution is not integral,  $A$  is not  $k$ -balanced by Theorem 2. If the solution is a 0/1 vector, it yields a  $k$ -equitable bicoloring as in the proof of Theorem 3.

Note that, as with the algorithm of Cameron and Edmonds [3], a 0/1 vector may be found even when the matrix  $A$  is not  $k$ -balanced.

Using the fact that the vector  $(\frac{1}{2}, \dots, \frac{1}{2})$  is a feasible solution of (1), a basic feasible solution of (1) can actually be derived in strongly polynomial time using an algorithm of Megiddo [11].

## References

- [1] C. Berge, Sur Certain Hypergraphes Généralisant les Graphes Bipartis, in *Combinatorial Theory and Its Applications I* (P. Erdős, A. Rényi, and V. Sós, eds.), *Colloquia Mathematica Societatis János Bolyai* 4, North Holland, Amsterdam (1970) 119-133.
- [2] C. Berge, Balanced Matrices, *Mathematical Programming*, 2 (1972) 19-31.
- [3] K. Cameron and J. Edmonds, Existentially Polytime Theorems, *DIMACS Series in Discrete Mathematics and Theoretical Computer Science* 1, American Mathematical Society, Providence, RI (1990) 83-100.
- [4] P. Camion, Characterization of Totally Unimodular Matrices, *Proceedings of the American Mathematical Society*, 16 (1965), 1068-1073.
- [5] P. Camion, Caractérisation des Matrices Unimodulaires, *Cahier du Centre d'Etudes de Recherche Opérationnelle*, 5 (1963) 181-190.
- [6] M. Conforti and G. Cornuéjols, Balanced  $0, \pm 1$  Matrices, Bicoloring and Total Dual Integrality, *Mathematical Programming*, 71 (1995) 249-258.
- [7] M. Conforti, G. Cornuéjols and K. Truemper, From Totally Unimodular to Balanced  $0, \pm 1$  Matrices: A Family of Integer Polytopes, *Mathematics of Operation Research*, 19 (1994) 21-23.
- [8] D.R. Fulkerson, A.J. Hoffman and R. Oppenheim, On Balanced Matrices, *Mathematical Programming Study*, 1 (1974) 120-132.
- [9] A. Ghouila-Houri, Caractérisations des Matrices Totalement Unimodulaires, *Comptes Rendus de l'Académie des Sciences*, 254 (1962) 1192-1193.

- [10] A.J. Hoffman and J.B. Kruskal, Integral Boundary Points of Convex Polyhedra, in *Linear Inequalities and Related Systems* (H.W. Kuhn and A.W. Tucker, eds.), Princeton University Press, Princeton, NJ (1956) 223-246.
- [11] N. Megiddo, On Finding Primal- and Dual-Optimal Bases, *Journal of Computing*, 3 (1991) 63-65.
- [12] M. Padberg, Characterization of Totally Unimodular, Balanced and Perfect Matrices, in *Combinatorial Programming: Methods and Applications* ( B. Roy, ed.), Reidel, Dordrecht (1975) 275-284.
- [13] M. Padberg, Total Unimodularity and the Euler Subgraph Problem, *Operation Research Letters*, 7 (1988) 173-179.
- [14] M. Padberg, *Linear Optimization and Extensions*, Springer, Berlin (1995).
- [15] K. Truemper, Alpha Balanced Graphs and Matrices and  $GF(3)$ -Representability of Matroids, *Journal of Combinatorial Theory Series B*, 32 (1982) 112-139.
- [16] K. Truemper and R. Chandrasekaran, Local Unimodularity of Matrix-Vector Pairs, *Linear Algebra and its Applications*, 22 (1978) 65-78.