

Seafood Watch Seafood Report:

Shrimp Volume III



Photo© South Carolina Institute of Scientific Inquiry

Farm-Raised Shrimp Worldwide Overview

**Alice Cascorbi
Fisheries Research Analyst
Monterey Bay Aquarium**

**final
5 April 2004**

About Seafood Watch® and the Seafood Reports

Monterey Bay Aquarium's Seafood Watch® program evaluates the ecological sustainability of wild-caught and farmed seafood commonly found in the United States marketplace. Seafood Watch defines sustainable seafood as species, whether fished or farmed, that can exist into the long-term by maintaining or increasing stock abundance and conserving the structure, function, biodiversity and productivity of the surrounding ecosystem. Seafood Watch® makes its science-based recommendations available to the public in the form of regional pocket guides that can be downloaded from the web (www.montereybayaquarium.org) or obtained from the program by emailing seafoodwatch@mbayaq.org. The program's goals are to raise awareness of important ocean conservation issues and to shift the buying habits of consumers, restaurateurs and other seafood purveyors to support sustainable fishing and aquaculture practices.

Each sustainability recommendation on the regional pocket guides is supported by a Seafood Report. Each report synthesizes and analyzes the most current ecological, fisheries and ecosystem science on a species then evaluates this information against the program's conservation ethic to arrive at a recommendation of "Best Choices", "Proceed with Caution" or "Avoid". In producing the Seafood Reports, Seafood Watch seeks out research published in academic, peer-reviewed journals whenever possible. Other sources of information include government technical publications, fishery management plans and supporting documents, and other scientific reviews of ecological sustainability. Seafood Watch Fishery Analysts also communicate regularly with ecologists, fisheries and aquaculture scientists, and members of industry and conservation organizations when evaluating fisheries and aquaculture practices. Capture fisheries and aquaculture practices are highly dynamic; as the scientific information on each species changes, Seafood Watch's sustainability recommendations and the underlying Seafood Reports will be updated to reflect these changes.

Parties interested in capture fisheries, aquaculture practices and the sustainability of ocean ecosystems are welcome to use Seafood Reports in any way they find useful. For more information about Seafood Watch® and Seafood Reports, please contact the Seafood Watch program at Monterey Bay Aquarium by calling 831-647-6873 or sending an email to seafoodwatch@mbayaq.org.

Disclaimer

Seafood Watch® strives to have all Seafood Reports reviewed for accuracy and completeness by external scientists with expertise in ecology, fishery science and aquaculture. Scientific review, however, does not constitute an endorsement of the Seafood Watch® program or its recommendations on the part of the reviewing scientists. Seafood Watch® is solely responsible for the conclusions reached in this report.

Seafood Watch® and Seafood Reports are made possible through a grant from the David and Lucile Packard Foundation.

Introduction to Series on Shrimps:

This is Volume III in a series of three seafood reports covering the shrimps most frequently found in West Coast markets and restaurants.

Volume Index:

- I Wild-Caught Coldwater Shrimp (Caridea; the pandalids and genus *Crangon*)
- II Wild-Caught Warmwater Shrimp (the penaeids)
- III Farmed Shrimp

General Shrimp Biology


Shrimps belong to the order Decapoda, a crustacean order which also includes the lobsters, true crabs, and hermit crabs. All decapods possess a full carapace or "head shield", and, eponomously, five pairs of walking legs [1]. Their first three pairs of thoracic appendages are modified into "maxillipeds", or feeding legs [2].

Shrimps are distinguished from the other decapods by having the front-most section of the abdomen about the same size as the rest of the sections, and by having five pairs of abdominal appendages, or pleopods, adapted for swimming [1].

There are more than 3,000 living species of shrimp worldwide [2]. Many are tiny, or inhabit niches unsuited to mass harvest [1]. Those harvested on a commercial basis share two characteristics: they are relatively large, ranging roughly from 2-10 cm. carapace length, and they school, shoal, migrate toward baited traps, or otherwise aggregate so that they are amenable to capture. Worldwide, about 40 species of shrimp meet these criteria and are harvested commercially [3]. About ten species have been raised in captivity; for some species, such as the Pacific white shrimp *Penaeus vannamei*, selective breeding is developing truly "domesticated" breeds of shrimp.

Scientific Names and Shrimp vs. Prawn

While there is no hard and fast rule about applying the names "shrimp" and "prawn" [Watling, 2004; Shumway, 2003], certain scientific references state that "shrimp" refers to the infraorder Caridea, which includes the widely-harvested coldwater genera *Pandalus* and *Crangon* [2]. With more than 2,000 species, these so-called "true shrimp" [2] are the largest group of shrimplike decapods [2]. They are distinguished by the fact that small side flaps of the exoskeleton overlap on their first, second, and third abdominal segments [2]. Under this definition, "prawn" refers to members of the infraorder Penaeidea, which includes the penaeids or tropical shrimp [2]. Also known as the "primitive shrimp", prawns are recognizable because their first and second anterior segments are about the same size [2].



A typical prawn

A typical shrimp

Image after MuseumVictoria, Australia

However, there is vast confusion among the common names of these animals. The "spot prawn" (*Pandalus platyceros*) of the U.S. West Coast is in fact a shrimp [2], [3]; while in British usage only the genus *Palaemon*, with its prominent head spine or rostrum, can be called a prawn [1].

In U.S. markets, "shrimp" is the default name for all these animals. "Prawn" often refers to freshwater shrimp or large saltwater shrimp. The term "scampi" refers not to a species but to a cooking method: any large shrimp cooked in butter and garlic [4]. Commercially-harvested shrimp may be divided into three categories, based upon their habitat: coldwater or northern species; warmwater, tropical, or southern species; and freshwater species [4].

Overview of Market Information

The market for shrimp continues to expand, and farmed shrimp supply an ever-increasing share of that market. About three-quarters of world shrimp production is wild-caught, further broken down into 70% warmwater and 30% coldwater shrimp. The remaining quarter of total production is farm-raised shrimp [5, 6]. However, wild shrimp tend to be consumed in the country of catch, while farmed shrimp are more likely to be traded internationally [Clay, 2003]. Perhaps 50% of shrimp traded internationally are farm-raised [Clay, 2003].

With worldwide shrimp fisheries at or near maximum sustainable yield, any growth in shrimp production must come from farm-raised shrimp. Many nations are turning to farm-raised shrimp as an attractive source of international trade revenue. In 2003, one study found that shrimp farming is growing at 12% to 15% per year [Clay, 2003; Rosenberg, 2003].

According to the latest figures available in 2003, about 87% of the U.S. shrimp supply was imported. This includes farmed and wild, warm- and coldwater- shrimp, and U.S. national import statistics do not distinguish between these categories [NMFS Statistics, 2003].

Executive Summary:

Several species of saltwater shrimp and one freshwater prawn are being cultivated in captivity. Farming methods range from simple ponds located in coastal areas to high-tech inland systems that filter and recirculate their water. The vast majority of farmed shrimp comes from economically-disadvantaged tropical nations, including India, Thailand, Indonesia, Ecuador, China, Bangladesh, and Vietnam. In many areas, shrimp farming has had an adverse effect on coastal habitat, water quality and local village economies. Shrimp are also farmed in the southern United States, with more regulatory oversight than is common in most other nations. In the United States and elsewhere, experimental farms are in the process of developing methods for low-impact shrimp culture. As of the latest figures available in 2003, about 87% of the U.S. shrimp supply was imported. This includes farmed and wild, warm- and coldwater- shrimp, with very little way for the consumer to discern them. It is hoped that the advent of country-of-origin labelling (COOL) in autumn 2004 will help U.S. consumers discern the sources of their shrimp.

*Seafood Watch is indebted to the outside experts who graciously volunteered their time to review the facts presented in this report for completeness and scientific accuracy: Dr. Jason Clay, World Wildlife Fund-U.S.; Dr. Todd Steiner, Sea Turtle Restoration Project. Thanks are also due Dr. Albert Tacon, Dr. Sandy Shumway, and Dr. Les Watling, whose reviews of other Seafood Watch shrimp reports contributed to the final form of this report. **It is important to note that scientific review does not constitute an endorsement of Seafood Watch on the part of the reviewing scientists; the Seafood Watch staff is solely responsible for the conclusions reached in this report.***

Overall Seafood Ranks:

International Farmed Shrimp (general recommendation): Avoid

List of Component Ranks	Low Conservation Concern	Medium Conservation Concern	High Conservation Concern	Critical Conservation Concern
Feed Efficiency		√		
Risks to Wild Stocks		√		
Disease Transfer		√		
Habitat Effects			√	
Pollution		√		
Chemical Use		√		
Management Effectiveness			√	


U.S. Farmed Shrimp (general recommendation): Proceed with Caution

List of Component Ranks	Low Conservation Concern	Medium Conservation Concern	High Conservation Concern	Critical Conservation Concern
Feed Efficiency		√		
Risks to Wild Stocks	√			
Disease Transfer	√			
Habitat Effects		√		
Pollution		√		
Chemical Use	√			
Management Effectiveness		√		

FARM-RAISED SHRIMP Worldwide Overview



Top left: Black tiger prawn, *Penaeus monodon*. Illustration ©Monterey Bay Aquarium. Top right: Pacific white shrimp, *Penaeus vannamei*. Photo© South Carolina Institute of Scientific Inquiry. Below: Freshwater prawn, *Macrobrachium rosenbergii*. Illustration courtesy UN FAO FIGIS database.



Shrimp is a hugely popular seafood in the developed nations of the world, including the United States, the European Union, Japan, and Australia. In the United States, shrimp recently surpassed canned tuna as the seafood most consumed per capita [5]. Unlike canned tuna, however, shrimp remains a luxury product in the developed world, commanding prices of six to eight dollars per pound for the farmer [7]. In Asia, a shrimp farm can offer 50 times the annual return of rice farming [Clay, 2003]. Tropical nations can raise three or more crops of warmwater shrimp per year [8; Clay, 2003], making shrimp farming an attractive source of foreign exchange for developing nations. Even within developed nations, farmers searching for a profitable product respond to the lure of shrimp. In some communities in the southern U.S., displaced fishermen have turned to shrimp farming as a way to continue making a living on the coast [8].

Several species of shrimp are now being farm-raised (Table 1). Because of their quick growth and large size, warmwater penaeid shrimp have been the focus of aquacultural production [4, 9]. The Asian black tiger shrimp *Penaeus monodon* reaches harvest weight of 35 grams after 120 days and is widely cultivated in the Eastern Hemisphere [9]. In the Western Hemisphere, including the United States, farmers focus on the Pacific white shrimp, *Penaeus vannamei*[9]. In Asia, other penaeids are favored. There has also been interest in farming shrimp in fresh water--this would open the possibility of shrimp farming to many inland areas that do not enjoy a ready supply of sea water. The long-clawed freshwater prawn *Macrobrachium rosenbergii* has been farmed experimentally in the Pacific basin and the U.S. [7], although these shrimp are aggressive with each other and require a culture method that allows each adult its own hiding place [7]. Some

penaeid species, among them the Pacific white shrimp *Penaeus vannamei*, the Chinese white shrimp *P. chinensis*, and the black tiger prawn *P. monodon*, can adapt to water much less saline than seawater [7, 10, 11]. This allows raising of shrimps in non-seawater systems---often in groundwater. These penaeid species are not aggressive, can be cultured at high densities, and have become the mainstay of shrimp farming worldwide [9].

Life History

Most shrimps are omnivorous, catching or scavenging whatever plant or animal material is readily available. The intestine runs the dorsal length of the abdomen; it is the brown line sometimes called the "mud vein" on cooked shrimp. Like other arthropods, shrimps have no internal skeleton, being protected instead by a chitinous exoskeleton which must be repeatedly shed as the animal grows [1]. The sexes are separate, and females tend to be larger than males. Some species release their eggs into the water column, while others brood the fertilized eggs on the female's abdomen until hatching. Newly-hatched shrimp larvae bear little resemblance to their elders; each must undergo up to 12 molts to attain final form as a juvenile shrimp. The tiny shrimp larvae drift with the plankton, where they are important food for many fishes and invertebrates [1]; those that escape predators and find favorable currents may live long enough to reproduce.

Pandalid shrimps such as the spot prawn may live for three to seven years [12, 13]; in contrast, many of the warm-water penaeid shrimps complete their life cycles in one to three years[14].

This diagram of penaeid life history is typical of the natural life cycle of many farm-raised shrimp species:



Image after Benfield and Downer, 2001 [15].

Table 1: Shrimp species in aquaculture

Scientific Name	common name(s)	where farmed	Percentage of total world farmed shrimp production [9]
saltwater shrimp--major species			
<i>Penaeus monodon</i>	black tiger shrimp, tiger prawn	Asia, Indonesia, Australia	47%
<i>Penaeus vannamei</i>	white shrimp, Pacific white shrimp, Vanna White	North and South America, Caribbean	16%
<i>Peneus chinensis</i>	white shrimp, Chinese white shrimp, fleshy prawn	Asia (tolerates cooler water than other peneids)	14%
<i>Penaeus stylirostris</i>		North and South America, Caribbean	**
<i>Penaeus japonicus</i>		Asia; some South America	**
<i>Penaeus penicillatus</i>		Asia, Indonesia, Australia	**
saltwater shrimp--minor species			
<i>Penaeus schmittii</i> <i>P. semiculcatus</i> <i>P. brasiliensis</i> <i>P. paulensis</i> <i>P. setiferus</i> <i>P. subtilis</i> <i>P. duorarum</i> <i>P. occidentalis</i> <i>P. californiensis</i>			**
			** all together total 23%
fresh water species			
<i>Macrobrachium rosenbergii</i>	freshwater prawn, long-clawed prawn	Asia, U.S. (experimental)	
<i>P. vannamei</i>	Pacific white shrimp	U.S. (experimental)	
<i>P. monodon</i>	black tiger shrimp	Thailand (experimental)	

Statement on the Availability of Science:

Because shrimp farming can be highly lucrative, many studies of farming methods are available in peer-reviewed journals and from universities and agricultural extension programs. Studies of the environmental impacts of shrimp farming have been produced since the early 1990's. In 1999, the Global Aquaculture Alliance, an international coalition of shrimp producers and marketers, published a monograph entitled Codes of Practice for Responsible Shrimp Farming (GAA, 1999). In 2001, at least two peer-reviewed journals published extensive reviews of shrimp farming practices and their environmental impacts worldwide [16, 17], and a monograph was published on the same subject [18]. Extensive work on shrimp farming has been published by the Shrimp Farming and the Environment Consortium, a group made up of the World Bank, U.N. Food and Agriculture Organization, World Wildlife Fund, and NACA [Clay, 2003].

Market Information:

It is important to note that, in the United States, the various species of shrimp are generally sold interchangeably, traded not by species, but by *size*. It matters little to most restaurateurs whether their breaded shrimp is *Penaeus setiferus* or *Pandalus jordani*, as long as it's the right size. In the United States, farmed and wild-caught shrimp enter the same market and are traded interchangeably [5].

The Shrimp Count

Because shrimp are so small, they are sold by a count (number) per pound rather than by individual weight [19]. This is expressed as a range. For example, a 16/20 count means it takes 16 to 20 shrimp of that size to make up a pound [19]. The smaller the count, the larger the shrimp.

Size Name	Count per Pound		
	Green headless	Peeled	Cooked
Extra Colossal	Under 10	Under 15	16/20
Colossal	Under 15	16/20	21/25
Extra jumbo	16/20	21/25	26/30
Jumbo	21/25	26/30	31/35
Extra large	26/30	31/35	36/40
Large	31/40	36/45	41/50
Medium large	36/40	41/45	46/50
Medium	41/50	46/55	51/60
Small	51/60	56/65	61/70
Extra small	61/70	66/75	71/80
Tiny	Over 70		

From Seafood Business Seafood Handbook, 1999

Market Names

Perhaps more than any other seafood commodity, the market names of shrimp are seldom standardized. Several different species are commonly called "white shrimp", and the situation is the same for "pink shrimp", "rock shrimp", and "tiger shrimp" [4], [20], [21]. Moreover, widely-distributed species have many common names. As one example, the circumpolar species *Pandalus borealis* may be marketed as pink shrimp, northern shrimp, Alaska pink shrimp, northern pink shrimp, Pacific pink shrimp, or salad shrimp [22;3].

Seasonal Availability:

In the United States, demand for shrimp is greatest during the winter holiday season, which, in terms of shrimp consumption, means Thanksgiving in late November, Christmas in December, and Superbowl Sunday in early January [Clay, 2003]. The availability of farmed shrimp is greatest in the autumn, as farms in many temperate areas harvest a single yearly crop September or October and then close down for the winter [8]. Tropical shrimp farms may harvest more than one crop per year, but even these tend to bring a crop to market in time for the winter holiday season, further increasing the shrimp supply from November through January [5; Clay, 2003]. This abundant supply tends to push wholesale shrimp prices down [5].

Shrimp is scarcest in the early spring (late January through early March), when many northern fisheries are closed for the winter, temperate farms have yet to re-open, and tropical farms are still growing out their first new crop [8, 23]. Spring is when shrimp wholesalers expect the highest prices for their products [5].

Product Forms

Not only are there many varieties of shrimp for sale worldwide, but there is a great diversity in product forms. Product can be divided into two basic types: raw and cooked [19]. It can then be further divided into fresh and frozen [19]. Within these broad categories, almost all shrimp sold in the U.S. market is sold as head-off tails, and the bulk of that is sold frozen [19]. Primary product forms for frozen shrimp are:

Green Headless: The standard market form. Includes the six tail segments, with vein, shell and tail fin. "Green" does not refer to shell color but to the uncooked, raw state of the shrimp. Also called "shell-on" or "headless" [19].

Peeled: Green headless shrimp without the shell [19].

PUD: Peeled, un-deveined, tail fin on or off; raw or cooked. The vein, running the length of the tail, is the intestine, also called the mud vein or sand vein [19].

Tail-on Round: Undeveined shrimp with tail fin on [19].


P&D: Peeled, deveined, tail fin on or off; raw or cooked. Another name for IQF P&D shrimp is PDI (peeled, deveined, individually frozen) [19].

Cleaned: Shrimp that is peeled and washed, a process that removes some or all of the vein but is not thorough enough to warrant the P&D label [19].

Shell-on Cooked: Cooked tail, with vein, shell and tail fin [19].

Split, Butterfly, Fantail: Tail-on shrimp that are cut deeply when being deveined [19].

Pieces: Shrimp with fewer than four or five whole segments [19].



Images from Seafood Business Seafood Handbook, 1999.

Frozen Products: Frozen shrimp generally comes in two forms: blocks (shrimp frozen en masse) and individually quick-frozen (IQF) packs [19]. Both shrimp blocks and IQF shrimp are glazed with a protective ice coating to prevent dehydration [19].

Breaded Shrimp: Shrimp, whether tail-on or tail-off, is the most-common breaded shellfish on the market [19].

Import and Export Sources and Statistics:

About 3/4 of world shrimp production is wild-caught. The remaining 1/4 of total production is farm-raised shrimp [5, 6]. In 2003, one study found that shrimp farming is growing at 12% to 15% per year [Clay, 2003; Rosenberg, 2003]. Perhaps 50% of shrimp traded internationally are farm-raised [Clay, 2003].


According to the latest figures available in 2003, about 87% of the U.S. shrimp supply was imported. This includes farmed and wild, warm- and coldwater- shrimp, and U.S. national import statistics do not distinguish between these categories [NMFS Statistics, 2003].

On the world market, farmed shrimp is typically traded in boxes of 1 to 20 kilograms [Clay, 2003]. Although the pre-consumer market for farmed shrimp (from farm to wholesaler) tracks information about the species, size, and country of origin of each box of farmed shrimp [Clay, 2003], marketers often remove this information before shrimp is sold at retail outlets, leaving the consumer with no easy way to discern the origins of their shrimp [Cutland & Cherry, 2002]. It is hoped that the advent of country-of-origin labelling (COOL) in autumn 2004 will help U.S. consumers discern the sources of their shrimp.


The U.S. imports more seafood than it exports; for several years, this trade deficit has been driven by what seafood market analyst H.M. Johnson calls "the tidal wave of shrimp imports" [5]. In 2000, imports of shrimp to the U.S. topped one billion pounds (heads-off weight) for the first time in history. That translates to 343,418 metric tons, with a wholesale value of about \$3.7 billion. As of the latest figures available in 2003, about 87% of the U.S. shrimp supply was imported. This includes farmed and wild, warm- and coldwater- shrimp [Cutland & Cherry, 2002]. So-called "value-added" or processed shrimp products are an important part of the import picture. Between 1997 and 2000, imports of breaded shrimp increased by 85% and cooked shrimp by 23% [5].

In 2001, the total U.S. shrimp supply was about 1.3 billion pounds (589,670 metric tons), including domestic landings of about 180 million pounds (81646.6 metric tons) [Cutland and Cherry, 2002]. Domestic landings (of *wild* shrimp) thus accounted for about 13% of the U.S. shrimp market in 2001. The exact percentage of the market that is domestic *farmed* shrimp is unavailable in NMFS or FDA statistics [NMFS Stats, 2003].

Shrimp has the unusual distinction of being the one seafood preferred equally in all regions of the United States [AquaNIC, 2001]. In 2001, for the first time in history, shrimp became the most-consumed seafood in the United States, as per-capita consumption of shrimp surpassed consumption of canned tuna [Johnson, 2003]. In 2001, Americans ate 3.4 pounds of shrimp per capita—an increase of 0.2 pounds from 2000 and 0.4 pounds from 1999 [Johnson, 2003].



Source: National Marine Fisheries Service



Source: National Marine Fisheries Service/National Fisheries Institute

Figure 1 (left): Origin of U.S. shrimp supply. (Right): domestic shrimp production vs. consumption. Source: NMFS



According to the latest figures available in 2003, about 87% of the U.S. shrimp supply was imported. For many years [Johnson, 2003; GLOBEFISH 2003], Thailand has been the leading import source (Figure 1). In 2001, Thai shrimp accounted for 29% of the U.S. market [Cutland and Cherry, 2002]. Mexico, India, and Vietnam tied for second place, with 7% each [Cutland and Cherry, 2002]. China, Indonesia, and Bangladesh also make substantial contributions to U.S. shrimp imports [Johnson, 2003; NMFS 2003; Globefish, 2003; Clay, 2003]. In all, Asian nations account for 66% of America's imported shrimp [Johnson, 2003; NMFS 2003].

Consumption Information and Trends:



Worldwide, shrimp consumption has been on the rise for more than a decade [25]. Demand for this luxury item shows no sign of slowing in the major markets of Japan, Europe and the United States. In the United States, shrimp has the unusual distinction of being the one seafood preferred equally in all regions of the country [9]. U.S. shrimp consumption rose in 2000 to 3.2 pounds per capita, up .2 pounds from 1999 [5]. These statistics, like the global shrimp market itself, make no distinction between warmwater and coldwater shrimp, farm-raised or wild-caught. However, data show that about three-quarters of world shrimp production is wild-caught. The remaining quarter of total production is farm-raised shrimp [5, 6].

Trends in Farmed Shrimp Production:

Worldwide, more than a million metric tons of farm-raised shrimp are produced each year; in 2003, one study found that shrimp farming is growing at 12% to 15% per year [Clay, 2003; Rosenberg, 2003]. In 2000, farmed shrimp production topped 700,000 metric tons [9]; FAO reported approximately 1.1 million mt in 2001 and 1.6 mt in 2002 [Clay, 2003]. In total, about one-quarter of world shrimp production is farm-raised [5]. However, wild shrimp tend to be consumed in the country of catch, while farmed shrimp are more likely to be traded internationally [Clay, 2003]. Perhaps 50% of the shrimp traded internationally are farm-raised [Clay, 2003].



L. Total capture of *Penaeus vannamei* (wild-caught only). R. Total production of *P.vannamei* (wild plus farmed).



Left: Total capture of *Machrobrachium rosenbergii* (wild-caught only). Right: Total production of *M. rosenbergii* (wild plus farmed). Graphs courtesy UN FAO FIGIS database.

Shrimp Aquaculture Information: International Shrimp Farming

Operation Types: International

Shrimp are most commonly cultivated in open-air ponds. At their simplest, these are enclosures of tidal estuary, ringed with earthen levees. Small postlarvae, often netted from the wild, are introduced into the ponds. In the simplest systems, the shrimp feed on algae and other ecosystem products [Clay, 2003]. In slightly more technological operations, supplemental food is introduced in the form of feed pellets. When the shrimp have grown to market size, a levee on the seaward side of the pond is opened. A net or other collector is installed over the outlet, releasing the water but trapping the shrimp [18].

More advanced systems line the bottoms of the ponds with clay to prevent wastes from seeping into the underlying soil and groundwater [18]. In Brazil, pond bottoms are often lined with plastic [Clay, 2003]. Further advances include the use of mechanical aeration, which enables more intensive stocking of the ponds; mechanical feeders, which standardize feedings; and various types of mechanical or biological filters to treat effluent when the ponds are drained [18].

Some facilities now specialize in breeding shrimp in captivity and supplying postlarvae to other farms. While this takes some of the pressure off wild shrimp populations, there is not enough broodstock cultivated to supply the worldwide demand [8].

In Thailand, the United States and Israel, a few shrimp farms employ fully-enclosed systems which filter and recirculate their water. For the most part, these are still experimental efforts that do not yet supply much shrimp to the market [7, 11, 26].

Intensity of Cultivation: International

Shrimp can be cultivated in several different ways. Coastal ponds stock their shrimp at medium densities and often aerate the water mechanically; this is considered a “semi-intensive” form of aquaculture [27]. Recirculating inland systems stock their shrimp at high densities; this method requires more equipment and is considered “very intensive” [9]. In Latin America, most shrimp farms stock their ponds at low densities and manage without aeration [28]. In India, 80% of farmed shrimp come from small, marginal holdings which use a variety of culture techniques, ranging from traditional paddy culture to specially-built ponds utilizing aerators, pellet feed dispensers, and other mechanical methods [29].

The limiting factor on shrimp density is often disease. In the early 1990's, two viral diseases (white spot and Taura syndrome) swept through Asia's shrimp farming industry [30]. Spread by infected broodstock, these diseases were soon detected in the Americas [30] and continue to threaten shrimp farms worldwide [26]. The highest densities of shrimp are achieved in high-tech recirculating freshwater systems, which start with pathogen-free postlarvae [26].

Type of Feed and Feed Conversion Ratio: International

As omnivorous scavengers, shrimp can feed on a wide range of protein sources, but large-scale shrimp aquaculture typically depends upon commercially-formulated shrimp feeds [11]. These always contain fish meal as a source of protein, and often contain fish oil as a source of lipids [11; Clay, 2003]. Although marine and freshwater shrimp can be successfully cultivated on diets containing no fishmeal or fish oil [SEAFeds Workshop Report, 2003], commercial shrimp typically contains fish derivatives. According to one authority, it currently takes between 1.7 and

2.21 kilograms of wild fish to produce one kilogram of farmed shrimp [Tacon, SEAFeds Workshop Report, 2003]. However, Clay (2003) suggests that current ratios range from 1.4 :1 to 2.0 :1, and reports that super-intensive shrimp farms in Belize now use only 0.7 kg of wild fish to produce 1 kg of shrimp.

Because fish meal and oil are typically the most expensive ingredients in a compound feed [Tacon, 2003], the fish-feeds industry is working to decrease the percentage of fishmeal and fish oil in compound feeds [SEAFeds Workshop Report, 2003]; this may explain why the 2001 review by Naylor et al. states that the average ratio of wild fish to farmed shrimp is 2.25 to1 [31]. As can be seen from the tables below, in 2001, marine shrimp consumed about 12.6% of the total production of compound aquacultural feeds, into which went 19.3% of the world's fish meal and 7% of the world's fish oil [SEAFeds Workshop, 2003].

Table 2: Conversion efficiency of capture fish to farmed fish (kg per kg fresh weight)
(from Tacon, SEAFeds Workshop Report, 2003)


<i>Species group</i>	<i>pelagic input per unit of production</i>
eels	3.4-4.2
marine fish	2.9-3.7
salmonids	2.6-3.3
marine shrimp	1.7-2.1
freshwater crustaceans	1.0-1.3
milkfish	0.3-0.4
catfish	0.3-0.4
tilapia	0.2-0.3
feeding carp	0.1-0.2

Fish meal and oil is derived mostly from directed fisheries for small, sardine-like species; the major fishing nations are Peru, Chile, and the Scandinavian countries [SEAFeds Workshop Report, 2003]. These are fast-breeding, short-lived, planktivorous fishes, probably well suited by their life history to support heavy fishing [SEAFeds Workshop Report, 2003]. With the exception of one overfished stock of blue whiting, the fish stocks targeted for reduction fisheries are believed to be within safe biological limits [Pike, 2003]. The stocks should be considered fully fished, as total landings remain relatively constant (including predictable declines caused by El Nino) [Pike, 2003].

The directed reduction fisheries are the world's largest source of fishmeal, but not the only source. Fish meal may also be derived from fish taken as bycatch in other fisheries [Steiner, 2003], or from fish trimmings from the processing of wild or farmed fish [SEAFeds Workshop Report, 2003]. These sources have the potential to expand the availability of fishmeal without increasing pressure on the sardine-like fishes [SEAFeds Workshop Report, 2003]. In some areas of the world, including Canada, fish trimmings now supply a significant fraction of the fishmeal used in aquacultural feeds [SEAFeds Workshop Report, 2003]. However, public-health laws enacted in the wake of the "mad cow" disaster forbid the use of fish trimmings in fish feeds within the European Union [SEAFeds Workshop Report, 2003].

**ESTIMATED GLOBAL COMPOUND AQUAFEED PRODUCTION IN 2001
FOR MAJOR FARMED SPECIES**

(values expressed as % total aquafeed production, dry as-fed basis)



TOTAL ESTIMATED COMPOUND AQUAFEED PRODUCTION IN 2001 – 16.70 MMT

In 2001 the aquafeed sector consumed

<u>Fish meal</u>	<u>2.62 mmt</u>	<u>Fish oil</u>	<u>0.59 mmt</u>
Salmonids	29.4%		64.5%
Marine fish	22.6%		20.3%
Shrimp	19.3%		7.0%
Carp	15.3%		-
Eels	6.9%		2.5%
Tilapia	2.7%		1.9%
Milkfish	1.4%		0.7%
Catfish	0.9%		1.0%

**Total fish meal and fish oil used in 2001 3.21 mmt
43.1% of total 6.08 mmt, 53.6% of total 1.10 mmt**

Figure 2: Graphs from SEAFeds Workshop Report, 2003 detailing world production of compound feeds for aquaculture (above) and percentages of fish meal and oil used in the feeding of various farmed species (below).

Management and Monitoring: International

Shrimp farms are subject to the various laws and monitoring protocols of each country where shrimp are raised (GAA, 1999). In many nations, laws for environmental protection may not exist, or, if they exist, may or may not be enforced [32]. In Bangladesh, illegal environmental destruction by large shrimp farming concerns has led to mass protests by displaced villagers, who clash, sometimes violently, with local authorities [33]. In Thailand, government managers recently recommended a moratorium on inland shrimp farming, partly because of a lack of confidence that the government could enforce environmental regulations [10].

Environmental Impacts: International

A factor in favor of farm-raised shrimp is that their production involves no bycatch of sea turtles or mature fish. However, most farms in Asia and Central and South America rely upon wild-caught broodstock, as the supply of captive-bred broodstock is very limited [17; Clay, 2003]. And perhaps 5% of shrimp farms in developing nations still depend on wild-caught postlarvae [18; Clay, 2003]. While the capture of larvae has become a cottage industry in some economically-disadvantaged coastal communities [18], larva fishing takes a heavy toll of bycatch of the youngest stages of many fish and invertebrates [6, 18].

On the issue of habitat damage, the story varies depending on where and how the shrimp is cultured [27]. To date, approximately 1.5 million hectares of coastal habitat (mangrove forests, marshes, salt flats and agricultural lands) have been converted to shrimp ponds [16]. Worldwide, about 10% of the loss of mangrove forests can be attributed to shrimp farming; that figure rises to 20% in certain areas [Lassen, 2004]. Hernandez-Cornejo and Ruiz-Luna (2000) reported little environmental impact for shrimp cultivated on barren salt flats of Mexico's Sinaloa coastline [27]. Trott and Alongi (2000) found that nutrients decreased to pre-cultivation levels within three years of the cessation of farming in certain Australian mangrove swamps where shrimp cultivation had not been intensive and the site enjoyed robust tidal flushing [34]. The rate of conversion of virgin wetlands to shrimp farms has slowed in recent years [Clay, 2003]. However, broadly speaking, in tropical nations, shrimp farming is associated with continuing reports of coastal habitat destruction [18, 27, 33]; displacement of small-scale sustainable fisheries [18, 33]; harmful discharges of nutrient-enriched wastewater [11, 35]; and unregulated use of antibiotics [36].

In early 2002, the European Union banned imports of farmed shrimp from China, Indonesia and Vietnam because of antibiotic residues in the shrimp [38]. A 2002 analysis of the social and ecological sustainability of shrimp aquaculture in Thailand and Vietnam, conducted by the Royal Swedish Academy of Sciences, notes that, in Thailand, “years of experience with intensified systems...have not led to sustainable solutions” and concludes that “current pathways in both countries are unlikely to lead to a sustainable industry. A complete transformation of the way shrimp are grown, fed, processed, distributed, and regulated is needed” (Ambio, 2002).

Experiments continue with inland shrimp farming in Thailand, Israel and elsewhere [26, 30]. Inland farms avoid any impact on coastal habitat, and may be sited in existing agricultural regions with no more impact to the environment than conventional farms [37]. Because of the strict need to protect the crop from disease in a recirculating system, these farms use pathogen-free captive-bred larvae as a matter of course [7, 11]. Recirculating systems entail relatively high production costs, but offer benefits of disease control, year-round production and wastewater containment.

Nation-By-Nation Overview of International Shrimp Farming

Clay [2003] notes that the environmental soundness of shrimp farms varies considerably from farm to farm, even more than it varies from nation to nation. Individual, progressive shrimp farms in Australia, Belize, Brazil, Madagascar, Mexico, and all Asian nations may equal or surpass the environmental friendliness of United States shrimp farms [Clay, 2003]. However, as U.S. shrimp markets typically preclude tracing shrimp back to its source farm, Seafood Watch is forced to make blanket recommendations based on generalized information about shrimp farming practices in various nations.

Australia: While Australia supplies only a small fraction of farmed shrimp imported into the U.S., environmental standards and enforcement in Australia are comparable to those in the United States [Clay, 2003].

Bangladesh: In coastal Bangladesh, the black tiger prawn *P. monodon* is cultured in mixed-species ponds along with other shrimps (*P. indicus*, *Metapeneaus monoceros*, *M. brevicornis*) and several species of finfish [Alam and Phillips, 2004]. While the tiger prawn is the main product of these farms, accounting for 52% of crop biomass and about 89% of crop value, the raising of finfish along with shrimps has been a hedge for farmers when shrimp crops are lost due to viral disease [Alam and Phillips, 2004].

Belize: Clay (2003) reports that a few super-intensive shrimp farms in Belize now use only 0.7 kg of wild fish to produce 1 kg of shrimp.

Brazil: Brazil is the largest producer of farmed shrimp in the Western Hemisphere, harvesting 60,128 mt in 2002 [Nunes, 2004]. While most tropical shrimp farms can produce two or three crops per year, shrimp farms in Brazil average 3.5 crops per year [Clay, 2003] and some report 3.8 crops per year [Jost, 2004]. Brazil's shrimp farming industry began in the early 1980s, but suffered setbacks due to disease outbreaks in the early 1990s. Currently, there is a trend away from low-intensity ponds where shrimp spend their entire life cycle to high-intensity operations with specialized nursery tanks for postlarvae and lined ponds for adults [Nunes, 2004]. These intensified operations depend upon paddlewheel aerators [Nunes, 2004]. Such modifications help minimize the spread of disease. In order to maximize feed-conversion ratios and minimize nutrient pollution of ponds and groundwater, Brazilian shrimp farmers are moving away from broadcast feeding to the use of feeding trays [Nunes, 2004].

China: Total production of 380,000 mt in 2002 [Wang et al., 2004]. In 2002, shrimp farms were estimated to cover 240,000 hectares, with average yield per hectare about 1.6 tons [Wang et al., 2004]. Traditionally, semi-intensive farming along coastline using *Penaeus chinensis*. Farming began in the late 1970s, burgeoned along the coast until severe outbreak of white-spot disease in 1993. Farming declined for some years, but has recovered with introduction of disease-resistant *P. vannamei*. There has been a recent expansion of intensive industrial indoor ponds, especially in Shandong Province. Indoor ponds are now estimated to cover 10,000-15,000 hectares and to produce 90,000-100,000 mt annually [Wang et al., 2004]. Currently, the species profile of Chinese-produced farmed shrimp is: *P. vannamei* 60%; *P. chinensis* 15%; *M. japonicus* 15%; *P. monodon* 8%, and other species 2% [Wang et al., 2004].

Ecuador: A major outbreak of viral disease decimated Ecuador's coastal shrimp farms in 1999. By 2001, shrimp production had dropped 70% from its 1998 level [Alava, 2004]. Since 1999, shrimp farming has been expanding inland as an alternative to the coastal zone. About 90% of Ecuador's inland shrimp farms are in Guayas province, and many are located on the sites of old freshwater red-claw lobster pond farms [Alava, 2004]. In these inland areas, *P. vannamei* are raised in very-low-salinity conditions (i.e., hard groundwater). Ecuador has an environmental permitting process in place to address impacts of these inland farms on groundwater, surface water, and agricultural lands [Alava, 2004]. One study found relatively major impacts on vegetation and soil erodability during construction and operation of these inland shrimp farms, and moderate to low impacts on groundwater and surface water [Alava, 2004].

Honduras: The leading shrimp producer in Central America, Honduras shrimp farming began in 1972 and expanded throughout the 1980s [Valderama and Engle, 2002]. In the early 1990s, problems with Taura virus, introduced from Asia (probably with infected broodstock) eroded the profitability of the operations [Valderama and Engle, 2002]. Introduction of Taura-resistant lines and use of antibiotics restored productivity, and production peaked in 1998 at 12,000 mt of shrimp from 14,000 hectares of pond [Valderama and Engle, 2002]. However, introduction of the white-spot virus in 1999 has curbed production once again [Valderama and Engle, 2002]. The heart of Honduras' shrimp farming industry is the Gulf of Fonesca, near Choluteca [Valderama and Engle, 2002; Marquez, 2001]. The rapid expansion of shrimp farming in this area is associated with destruction of mangrove forests, despite legal protection for mangrove habitat under Honduran law [Marquez, 2001]. Heavy seasonal discharges of wastewater from shrimp ponds into the ocean, as well as ongoing discharges of wastes from shrimp packing plants, also violate Honduran law, but no enforcement efforts are made [Marquez, 2001]. One very large Honduran farm, Sea Farms International, is responsible for 50% of the national production; this operation reports feed-conversion ratios of 1.5 : 1 [Gautier et al., 2003] and claims to replant mangroves for their biofiltering benefits [Gautier et al, 2003]. Sea Farms International operates over 7,000 hectares of shrimp ponds in low-density, non-aerated production. The ponds are lightly fertilized with nitrogen and phosphorus; a water-quality monitoring program run by the shrimp farming industry found no change in the nutrient content of the Gulf of Fonesca between 1993 and 2003 [Gautier et al, 2003].

Mexico: Mexico's shrimp farming industry, once based on the simplest technology of unlined ponds in coastal areas, has been hard-hit by disease and by the worldwide glut of inexpensive shrimp from Asia [Reyna, 2004; Zendejas-Hernandez, 2004]. Mexico exports most of its shrimp production, both farmed and wild, to the United States [Gutierrez and Zendejas, 2004]. Recently, farmed production surpassed wild-caught production, due in part to the fact that the increasing cost of boat fuel has encouraged shrimp fishermen to get out of the business [Gutierrez and Zendejas, 2004]. Much of Mexican shrimp aquaculture has developed in the northwestern states of Sinaloa and Sonora [Reyna, 2004; Gutierrez and Zendejas, 2004]. In North Sonora, the most productive region, all farms are semi-intensive [Gutierrez and Zendejas, 2004]. There is increased interest in farming practices that minimize the chance of disease outbreaks; these include the use of disease-free captive-bred postlarvae, development of a disease-testing infrastructure, and treatment of outbreaks with antibiotics [Reyna, 2004]. A few coastal farms are now using mangrove forests as natural biofilters for shrimp farm effluent. The shrimp ponds are constructed behind (inland from) the mangrove belt, which processes nutrients and sediments released from the ponds [Lassen, 2004].

Philippines: Disease has had a severe impact upon Philippine shrimp farming: from a high of 90,426 mt in 1994, production had declined to 35,493 mt in 2003 [Yap and Villaluz, 2004]. Surviving, successful farms tend to have the following characteristics and use the following culture practices: access to clean seawater, either by being located close to shorelines that drop off into deep water, or by having access to a saltwater well, or by treating new seawater in a reservoir; a clean and compact pond bottom, kept clean and compact either by being located above the tide line or by being heavily backfilled with crushed limestone; use of health-tested postlarvae; adequate aeration of the ponds; “good feed management”; addition of bio-remediators to the ponds, such as fish and/or probiotic microbes; letting ponds sites dry out thoroughly between shrimp crops or else alternating shrimp with crops of tilapia or milkfish; “adequate production-based incentives to technicians and workers” [Yap and Villaluz, 2004].

Peru: Peru’s cold climate limits shrimp farming; perhaps 3,000 hectares are available for shrimp farming [Moya et al., 2004]. The disease epidemics of the late 1990s have hit the Peruvian industry particularly hard, and perhaps 50% of the shrimp ponds constructed in the past decade are now idle [Moya et al., 2004]. Farmers pursue one of two strategies to defeat disease: stocking extensive areas at low densities, or enclosing small lined ponds under greenhouses and conducting intensive, high-density aquaculture with disease-free postlarvae [Moya et al., 2004]. In these intensive indoor systems, which are always intensively aerated, water exchange varies from zero to 12% per day [Moya et al., 2004].

Thailand: Thailand has been the world’s largest producer of farmed shrimp since 1991 [Szuster et al., 2004]. Unlike the large shrimp-farming industries of China, Taiwan, and Ecuador, which collapsed in the early 1990s from a combination of habitat degradation and disease, Thailand’s industry has continued to expand [Szuster et al., 2004]. Most farms are small-scale, family-operated [Fegan, 2004], and most ponds are aerated [Clay, 2003]. Widespread shrimp farming began only about 1986 [Fegan, 2004]. When Taura virus began to decimate coastal farms in the early 1990s, producers moved inland and brought in virus-resistant *P. vannamei* [Fegan, 2004]. All the *P. vannamei* raised in Thailand are from captive-bred, pathogen-free postlarvae, although farmers still depend on wild broodstock for *P. monodon* [Fegan, 2004]. Many farms now use intensive culture methods that include high stocking density and heavy aeration in closed systems with little water exchange and with settling ponds for wastes [Fegan, 2004]. While Thai shrimp farmers often receive technical support from feed companies, there is still only limited vertical integration in the industry [Fegan, 2004].

Vietnam: Shrimp farms currently cover 400,000 hectares of Vietnamese land, mostly in South Vietnam [Phuong, 2004]. Seventy percent of this land is employed in “extensive” farming; i.e., low-density open pond culture [Phuong, 2004]. A report by the Environmental Justice Foundation asserts that, since 1975, Vietnam’s Mekong Delta has lost 70% of its mangrove habitat due to the expansion of shrimp farming [EJF, 2001]. This assertion is questioned by Clay [2003], who states that the wartime use of the defoliant Agent Orange has been the single largest cause of mangrove loss in Vietnam. Vietnam produces only 30% of the shrimp postlarvae used by its industry [Phuong, 2004]. Most of the “extensive” farms rely on wild postlarvae [Phuong, 2004]. In extensive farming, the shrimp are stocked at low densities; shrimp tend to feed on algae and microorganisms that develop naturally in the ponds, and inputs of feed or chemicals to the ponds are very low [Phuong, 2004]. Some farmers practice so-called “improved extensive” farming, where shrimp crops alternate with rice in flooded fields [Phuong, 2004]. However, the use of semi-intensive and intensive farming methods is expanding rapidly in Vietnam. These entail high production costs, but can maximize profits, if disease does not wipe out the crop [Phuong, 2004].

Shrimp farming is still expanding rapidly in Vietnam's coastal areas, but the infrastructure for shrimp health monitoring is still underdeveloped [Phuong, 2004]. Seed stock is in short supply, and viral infections are on the rise [Phuong, 2004]. In some areas, 40% of each shrimp crop is lost to viral disease [Phuong, 2004]. Use of antibiotics is widespread [Phuong, 2004].

Synthesis and Analysis of Criteria: International Farmed Shrimp

Effects on Wild Fisheries

Does the aquaculture activity adversely impact wild fisheries?

a) Introduction of non-indigenous species and/or pathogens

Shrimp farming has spread viral shrimp diseases around the globe [25, 29]. In some areas, stocks of non-native shrimp have become established—e.g., as a result of escapes from shrimp farms, Pacific white shrimp are now found in the Gulf of Mexico [Steiner, 2003]. Their impact upon the ecosystem is unknown. In other areas, such as around the Hawaiian archipelago, establishment of feral shrimp populations is unlikely because suitable estuary breeding habitat for penaeids does not exist [Dalzell, 2003].

b) Collection of wild broodstock and postlarvae

Most farms in Asia and Central and South America rely upon wild-caught broodstock, as the supply of captive-bred broodstock is very limited [17; Clay, 2003]. And perhaps 5% of shrimp farms in developing nations still depend on wild-caught postlarvae [18; Clay, 2003]. While the capture of larvae has become a cottage industry in some economically-disadvantaged coastal communities [18], larva fishing takes a heavy toll of bycatch of the youngest stages of many fish and invertebrates [6, 18].

c) Wild-caught Food

Many small-scale shrimp farms in developing nations do not use commercially-formulated shrimp feeds. This means that their reliance on fishmeal from wild fish is less than that of large-scale shrimp farming operations [14]. However, large-scale shrimp farming operations in Asia, Central America and South America do rely upon commercially-produced feeds [14]. Fishmeal is always a component of these rations [Clay, 2003]. Naylor et al. [2001] reported a typical ratio of 2.25 kilos wild fish (wet weight) to produce each kilo of shrimp (wet weight) [31]. But industry may have improved upon this ratio in the past few years. According to one authority, it currently takes between 1.7 and 2.21 kilograms of wild fish to produce one kilogram of farmed shrimp [Tacon, SEAFeds Workshop Report, 2003]. However, Clay (2003) suggests that current ratios range from 1.4 :1 to 2.0 :1, and reports that super-intensive shrimp farms in Belize now use only 0.7 kg of wild fish to produce 1 kg of shrimp. The actual environmental impact of reduction (fishmeal-directed) fisheries remains unquantified. With the exception of one overfished stock of blue whiting near Scotland, world stocks of reduction fishes are considered fully fished and at sustainable levels of abundance [Pike, 2003]. Ironically, in Asia, some of the fishmeal fed to farmed shrimp is made from the incidental catch (bycatch) taken in local trawl fisheries for wild shrimp [Steiner, 2003]. This connects shrimp farming in these regions to the trawl fisheries implicated in the decline of endangered sea turtles [Steiner, 2003].

Environmental and Ecological Impacts

Does the aquaculture activity impact environmental health and productivity?

a) Habitat modification

Broadly speaking, international shrimp farming is associated with continuing reports of habitat degradation, particularly in coastal regions of southeast Asia and India [17,26,32]. Coastal shrimp

farming in Asia and South America has been associated with the displacement of small-scale sustainable fisheries that rely upon intact coastal habitat [17, 32].

b) Effluent discharge

Broadly speaking, international shrimp farming is associated with continuing reports of nutrient effluent discharge [14, 32] as well as the unregulated use of antibiotics banned in the United States and European Union [35, 38]. However, there has been progress in recent years in many nations in the use of settlement ponds and “biofilters”(crop plants or artificial wetlands) to reduce sediment discharge and nutrient pollution from shrimp farms [Clay, 2003].

c) Food web interactions

The direct effect upon ocean food webs of the capture of fish for shrimp feed is unknown. Many developing-world shrimp farms depend on wild-caught postlarvae to stock their ponds. While the capture of larvae has become a cottage industry in some economically-disadvantaged coastal communities, larva fishing takes a heavy toll of bycatch of the youngest stages of many fish and invertebrates [17, 24].

Consumer Information

International farmed shrimp account for about 25% of all shrimp produced [9, 24] and for 45-50% of shrimp traded internationally [Clay, 2003]. The largest producers are Thailand, China, and India [13], and the largest suppliers to the U.S. are Thailand, India, and Mexico [9]. Because shrimp is traditionally sold in the U.S. without regard to species, nation of origin, or catch/culture method, it is currently nearly impossible for the consumer to discern international farmed shrimp in the market [9]. It is hoped that the advent of country-of-origin labelling (COOL) in autumn 2004 will help U.S. consumers discern the source of their shrimp.

List of Component Ranks	Low Conservation Concern	Medium Conservation Concern	High Conservation Concern	Critical Conservation Concern
Feed Efficiency		√		
Risks to Wild Stocks		√		
Disease Transfer		√		
Habitat Effects			√	
Pollution		√		
Chemical Use		√		
Management Effectiveness			√	

Overall Seafood Rank: Avoid

Seafood Watch Recommendation for International Farmed Shrimp

Avoid

Shrimp Aquaculture Information: United States

Operation Types: United States

Shrimp farming in the United States is comparatively small-scale. All told, shrimp farms cover only about 600 acres in the U.S. [Clay, 2003]. In their 1999 monograph “Codes of Practice for Responsible Shrimp Farming”, the Global Aquaculture Alliance concludes that, compared to Asia and South and Central America, the United States will always be a minor producer of farmed shrimp because of high production costs, few suitable sites, and its relatively cool climate (GAA, 1999). Nonetheless, the enormous value of shrimp leads to ongoing attempts to expand U.S. shrimp aquaculture.

U.S. shrimp farmers have access to advanced technology and tend to farm with more machinery than farmers in Asia or South America (GAA, 1999). U.S. shrimp farms are usually seawater coastal pond culture or low-salinity inland pond culture (Stickney 2002), with a few high-tech enclosed systems in use inland [7, 11, 26].

Some penaeid species, among them the Pacific white shrimp *Penaeus vannamei*, the Chinese white shrimp *P. chinensis*, and the black tiger prawn *P. monodon*, can adapt to water much less saline than seawater [7, 10, 11]. This allows raising of shrimps in non-seawater systems---often in groundwater. In certain areas, groundwater contains all the minerals needed by the shrimp. In other areas (such as the southern United States) the mineral profile of groundwater is very different than seawater of the same salinity [Green, 2004]. In the U.S. states of Arkansas and Alabama, shrimp farmers have learned to supplement groundwater with trace minerals to allow *P. vannamei* to thrive [Green, 2004; Davis et al., 2004].

Intensity of Cultivation: United States

Coastal ponds stock their shrimp at medium densities and often aerate the water mechanically; this is considered a “semi-intensive” form of aquaculture [27]. Recirculating inland systems stock their shrimp at high densities; this method requires more equipment and is considered “very intensive” [9] or “super-intensive” [Clay, 2003].

The limiting factor on shrimp density is often disease. In the early 1990s, two viral diseases (white spot disease and Taura syndrome) swept through Asia's shrimp farming industry [30]. Spread by infected broodstock, these diseases were soon detected in the Americas [30]. These viral diseases put several Texas coastal farms out of business (Stickney, 2002) and continue to threaten shrimp farms worldwide [26]. The highest densities of shrimp are achieved in high-tech enclosed freshwater systems, which start with pathogen-free postlarvae [26].

Type of Feed and Feed Conversion Ratio: United States

All U.S. shrimp farms use commercially-formulated shrimp feeds [11] [Stickney 2002]. Fishmeal is always a component of these rations [Clay, 2003]. Naylor et al. [1999] reported a typical ratio of 2.25 kilos wild fish (wet weight) to produce each kilo of shrimp (wet weight) [31]. But this value may have been improved upon by industry. According to one authority, it currently takes between 1.7 and 2.21 kilograms of wild fish to produce one kilogram of farmed shrimp [Tacon, SEAFeds Workshop Report, 2003]. However, Clay (2003) suggests that current ratios range from 1.4 :1 to 2.0 :1, and reports that super-intensive shrimp farms in Belize now use only 0.7 kg of wild fish to produce 1 kg of shrimp. There is no reason to suspect that feed efficiencies in U.S.

practice are very different from those found in international surveys. Please see the feed conversions discussion in the International section (above) for a fuller discussion.

It cannot be overemphasized: the feed conversion ratios above are “wet fish weight to wet shrimp weight”, not FEED weight to shrimp weight.

CEATECH’s Kauai Shrimp Farm, based on Kauai, Hawaii, states an institutional objective of producing shrimp in the most ecologically sustainable way possible (please see Seafood Watch Kauai Shrimp Evaluation for references). CEATECH’s techniques can reasonably be considered best practices among shrimp farms *currently in commercial production* in the United States. Kauai reports FEED conversion ratios of about 2.1 to 1. Since Kauai uses a shrimp feed containing about 35% “protein meal”, which is itself a mixture of fish and soy proteins, this indicates a dry-weight protein-conversion ratio that cannot be “worse” than 0.735 to 1 (if all the protein were derived from fish meal). At a standard wet-weight to dry-weight ratio of 4:1 [Clay,2003], this worst-case scenario would mean a wet fish—wet shrimp feed conversion ratio of 2.94 to 1. Again, such would be the case *if* all the protein in Kauai’s feed were derived from fishmeal. As we know, some fraction of the formula is soy protein; as fishmeal is the most expensive component of shrimp feeds, feed makers are always trying to minimize fishmeal content [Clay, 2003; Stickney, 2002; Tacon, 2003].

Management and Monitoring: United States

In the United States, shrimp farms are subject to EPA water-quality regulations, which are enforced by state fisheries agencies and/or state pollution control agencies [30]; (Stickney 2002; Treece 2002). Overall, regulations are more sophisticated and compliance is better in the United States than in many other nations [Clay, 2003].

Environmental Impacts, United States:

U.S. pond culture of saltwater shrimp started in the late 1960s and expanded in the early 1970s [9]. U.S. coastal shrimp farms are more closely regulated (or more strictly enforced) than the farms of many other nations [37, 38; Clay, 2003]. Because foreign farmed shrimp is often much cheaper than domestically-raised shrimp, the U.S. shrimp farming industry faces constant economic challenges.

U.S. farmed shrimp avoid the worst environmental excesses of the tropical farms. Besides tighter regulation and better enforcement of environmental law, U.S. shrimp farmers are subject to pressure from citizens' groups about environmental impacts [38]. In their own economic interest, U.S. shrimp farmers continue to experiment with new technologies to increase efficiency and minimize the environmental impact of their operations [37; Clay, 2003]. In addition, U.S. farms use captive-bred shrimp larvae [7, 9].

Experiments continue with inland shrimp farming in the United States and elsewhere [26, 30]. Several U.S. producers are experimenting with low-emission, enclosed fresh-water culture systems [7, 11, 26]. Inland farms avoid any impact on coastal habitat, and may be sited in existing agricultural regions with no more impact to the environment than conventional farms [37]. Because of the strict need to protect the crop from disease in a recirculating system, these farms use pathogen-free captive-bred larvae as a matter of course [7, 11]. Recirculating systems entail relatively high production costs, but offer benefits of disease control, year-round production and wastewater containment. Recirculating systems are drawing more and more interest from U.S. shrimp farmers [37].

State-By-State Overview of U.S. Shrimp Farming Operations

Most U.S. farmed shrimp is raised in Texas [Stickney, 2002; Treece, 2002]. There are a few operations in the states of Hawaii, Florida, South Carolina, and Kentucky [Stickney, 2002; Treece, 2002; Clay, 2003]. Very recently, shrimp farming has been introduced to the inland delta regions of Mississippi and Alabama [Stickney, 2002; Treece 2002; Clay 2003]. A few experimental efforts are underway in Arizona and other states.

Alabama: Very recently, shrimp farming has been introduced to the inland delta regions of Mississippi and Alabama, traditional strongholds of catfish farming [Stickney, 2002; Treece 2002; Clay 2003]. The lime-rich water of this area provides enough mineral content to support shrimp species, such as *P. vannamei*, that can adapt to low-salinity waters [Stickney, 2002]. The ecosystems of these areas are able to support nutrient-rich pond culture of catfish, and no additional impact has been noted from the shrimp farms [Stickney, 2002]. Inland shrimp farming began in Alabama in 1999 with the conversion of a catfish fingerling pond to production of *P. vannamei* [Davis et al., 2004]. In 2001, there were five producers covering a total of about 25 hectares, all in west-central Alabama [Davis et al., 2004]. Alabama shrimp producers find that they must supplement their pond water with minerals, especially potassium, to obtain commercially-viable harvest levels. Alabama shrimp farmers have an advantage in that they can piggyback on feed, equipment and loan infrastructures already in place for catfish farming [Davis et al., 2004].

Arizona: Four farms produced 175 metric tons of shrimp from 123 hectares of ponds in 2003 [Wilkenfield et al., 2004]. Most ponds are dirt-lined and stocked at low densities [Wilkenfield et al., 2004]. Shrimp farming began in Arizona only in 1998; producers hoped to take advantage of cheap land and abundant sunshine. Production has fallen in recent years because of low prices for shrimp and high operating costs. Arizona shrimp farmers are struggling with algal blooms, seasonal temperature fluctuations, and shrimp mortality caused by improper mineral balance in the water [Wilkenfield et al., 2004]. Each Arizona producer is struggling to find a niche market to sustain itself [Wilkenfield et al., 2004].

Arkansas: Small amounts of shrimp are being pond-raised in Arkansas. The mineral content of Arkansas groundwater is different from seawater of the same salinity, requiring farmers to supplement their ponds with minerals [Green, 2004].

Florida: In 2001, the last year for which figures are available, Florida had 13 shrimp farms covering a total area of 36 acres [Adams, Sweat and Martinez, 2004]. Total acreage was expected to rise to 170 acres in 2002. Florida farms produced both postlarvae and table-sized shrimp. Both outdoor ponds and enclosed raceway systems were in use [Adams, Sweat and Martinez, 2004].

Hawaii: Two large commercial shrimp operations producing shrimp for the national market, several smaller farms producing pathogen-free broodstock, and a few local low-density ponds producing shrimp for local consumption make up Hawaii's shrimp farming industry. Please see the Seafood Watch Hawaii Farmed Shrimp Report for a fuller discussion. State water quality agencies report little or no problem with pollution or habitat degradation from Hawaiian shrimp farms [Tomomitsu, 2002; Young, 2003]. One of the two large commercial operations is Ceatech's Kauai Shrimp Farm, which employs environmentally-friendly best practices including an inland location, antibiotic-free organic culture methods, and settling ponds for wastes. Seafood Watch

has rated Ceatech's Kauai shrimp a "Best Choice". Please see the Seafood Watch Kauai Shrimp Farm Evaluation Report for a fuller discussion.

Mississippi: Very recently, shrimp farming has been introduced to the inland delta regions of Mississippi and Alabama, traditional strongholds of catfish farming (Stickney, 2002; Treece 2002; Clay 2003). The lime-rich water of this area provides enough mineral content to support shrimp species, such as *P. vannamei*, that can adapt to low-salinity waters (Stickney, 2002). The ecosystems of these areas are able to support nutrient-rich pond culture of catfish, and no additional impact has been noted from the shrimp farms (Stickney, 2002).

Puerto Rico: There are few sites suitable for shrimp farming in Puerto Rico. Production has declined steadily from 200 mt in 1992 to 47 mt in 1997. This decline is partly due to the failure of a large prawn-raising operation in the early 1990s. Currently, just one large operation, Eureka Marine, is in place in Puerto Rico. In 1997, the last year for which figures are available, this farm produced 47 mt of shrimp from 30 hectares of lined ponds. Species raised include *Macrobrachium* prawns and other non-native shrimps [Alston & Cabarcas-Nunez, 2004].

South Carolina: The climate of South Carolina restricts shrimp farmers to one crop per year [Stokes, Browdy & Atwood, 2004]. Unable to compete with inexpensive imported shrimp, many South Carolina shrimp farms have now converted to crayfish [Clay, 2003]. In 2003, there were ten shrimp farming operations in the state, covering 75 hectares of ponds, 20 hectares of impoundments, and two (indoor) greenhouse raceway intensive-production facilities [Stokes, Browdy & Atwood, 2004]. Seven of the ten growers raise captive-bred *P. vannamei* as table shrimp, while three raise wild-caught Atlantic white shrimp (*P. setiferus*) to sell as bait [Stokes, Browdy & Atwood, 2004]. The pond and impoundment farms are located at the coast and draw salt water from coastal creeks [Stokes, Browdy & Atwood, 2004]. Because of the expense of coastal land for siting open-pond shrimp farms and the glut of inexpensive foreign farmed shrimp, plus the short growing season and the risk of losing crops to disease, the only segment of the shrimp farming industry expanding in South Carolina is indoor intensive greenhouse culture. Raceways can be sited on inexpensive inland property. Properly run with high-health postlarvae, raceways can produce three or four crops per year [Stokes, Browdy & Atwood, 2004].

Texas: Currently, most U.S. shrimp is farmed along the south coast of Texas (Stickney, 2002; Treece, 2002). In the mid-1990s, there was some concern about environmental degradation associated with wastewater discharges from shrimp farms along the South Texas coast [8, 38]. Farms along the Arroyo Colorado coast were granted unlimited water-use rights by the state of Texas (Stickney, 2002) and used this water to produce shrimp at high densities, pumping large amounts of seawater from the bay and discharging equal amounts of wastewater laden with sediment and nutrients (Stickney, 2002). Local homeowners and dock owners soon complained; after investigation by Texas pollution control authorities, the farms were cited and their water ration was cut (Stickney, 2002). These farms elected to stay in business by switching to recirculating systems, retrofitting their water intake canals as settling ponds and lowering their stocking density of shrimp. In 2002, these farms were still producing shrimp, with little impact on the local environment (Stickney, 2002). In 2004, a company based in the inland, west-Texas town of Imperial, began nationwide marketing of shrimp farmed in water from an underground saltwater aquifer [Permian Sea Shrimp, 2004]. This inland, closed-system farm appears to be one of the innovative shrimp farms applying the best available technology to produce an environmentally-friendly product.

Synthesis and Analysis of Criteria: U.S. Farmed Shrimp

Effects on Wild Fisheries

Does the aquaculture activity adversely impact wild fisheries?

a) Introduction of non-indigenous species and/or pathogens

In some areas, stocks of non-native shrimp have become established—e.g., as a result of escapes from shrimp farms, Pacific white shrimp are now found in the Gulf of Mexico [Steiner, 2003]. Their impact upon the ecosystem is unknown. In other areas, such as Hawaii, establishment of feral shrimp populations is unlikely because suitable estuary breeding habitat for penaeids does not exist [Dalzell, 2003]. Shrimp farming has spread viral shrimp diseases around the globe [25, 29]. The white spot and Taura virus have been detected in wild Gulf of Mexico shrimp [25]. So far, researchers have noted no effect of these viruses upon the overall health of the wild populations [25]. It is not known with certainty that these viruses escaped from captive shrimp, but some consider it likely, as the viruses seem to have originated in Asia [29].

b) Collection of wild seed stock

U.S. shrimp farmers do not use wild seed stock [10, 12].

c) Use of wild-caught food

U.S. shrimp farmers rely on commercially-formulated shrimp feeds [14]. While fishmeal is often a component of these rations, the impact on wild fisheries remains unquantified. The typical conversion ratio is between 1.7 and 2.1 kg of wild fish (wet weight) per kg farmed shrimp produced (wet weight) [Tacon, SEAFeds Workshop Report, 2003]. The actual environmental impact of reduction (fishmeal-directed) fisheries remains unquantified, although, with the exception of one overfished stock of blue whiting near Scotland, world stocks of reduction fishes are considered fully fished and at sustainable levels of abundance [Pike, 2003].

Environmental and ecological impacts

Does the aquaculture activity impact environmental health and productivity?

a) Habitat modification

In the United States, habitat modification for shrimp farming has been very minor (Stickney, 2002; Tomomitsu, 2002; Young, 2003). The climate and costs prohibit widescale development of shrimp farms in the U.S. (GAA, 1999), with the result that this industry has little environmental impact.

b) Effluent discharge

There has been little problem with effluent discharge or nutrient pollution from shrimp farms in Hawaii [Tomomitsu, 2002; Young, 2003]. Problems with effluent discharge along the Texas coast, the U.S.'s largest shrimp-producing area, have been largely solved by the introduction of recirculating techniques (Stickney, 2002).

c) Food web interactions

The direct effect upon ocean food webs of the capture of fish for shrimp feed is unknown.

Consumer Information

In the United States, shrimp is traditionally marketed without regard to species, nation of origin, or catch/culture method. Because of this, it is nearly impossible for the consumer to select U.S. farmed shrimp in the market [9]. Even Texas shrimp packers mix wild-caught with farmed, and

international with domestic, when processing shrimp in the United States' most important shrimp-farming region (Stickney, 2002).

U.S. Farmed Shrimp (general recommendation): Proceed with Caution

List of Component Ranks	Low Conservation Concern	Medium Conservation Concern	High Conservation Concern	Critical Conservation Concern
Feed Efficiency		√		
Risks to Wild Stocks	√			
Disease Transfer	√			
Habitat Effects		√		
Pollution		√		
Chemical Use	√			
Management Effectiveness		√		

Seafood Watch Recommendation for U.S. Farmed Shrimp

Coastal farmed U.S. shrimp should go on our Proceed with Caution list. U.S. shrimp produced by enclosed inland systems could be a Best Choice.

References:

1. Fenner A. Chase, J.a.D.P.A., *Chapter 23--Caridea: The Shrimps*, in *Intertidal Invertebrates of California*, D.P.A. Robert H. Morris, Eugene C. Haderlie, Editor. 1980, Stanford University Press: Stanford, California.
2. Watanabe, J., *Advanced Guide Class on Invertebrates*. 2001, Stanford University, Hopkins Marine Station, and Monterey Bay Aquarium: Monterey, CA.
3. NMFS, *Our Living Oceans*. 1998.
4. Batten, T., *Shrimp*. 2001, University of Delaware, Sea Grant.
5. Johnson, H.M., *Annual Report on the U.S. Seafood Industry*, ninth, tenth, and eleventh editions. 2001; 2002; 2003, H.M. Johnson & Associates: Jacksonville, OR. p. 98.
6. Clucas, I., *A Study of the Options for Utilization of Bycatch and Discards from Marine Capture Fisheries*. 1997, United Nations Food and Agriculture Organization: Rome. p. 230.
7. Demarco, R., *Aquafuture Farm and freshwater shrimp farming*, A. Cascorbi, Editor. 2001: Watsonville, CA.
8. Baker, P., *Coastal Shrimp Farming in Texas: A Case Study*, in *Murky Waters: Environmental Effects of Aquaculture in the United States*, R. Goldberg, Editor. 1997, EDF Publications: Washington, DC.
9. (AquaNIC), N.C.f.A.E., *Saltwater Shrimp Aquaculture Curriculum Guide, Species-Specific Module*. 1995, NCAE, affiliated with Purdue University and Iowa State University: Alexandria, Virginia.
10. Flaherty, M., B. Szuster, and P. Miller, *Low salinity shrimp farming in Thailand*. *Ambio*. Stockholm, 2000. **29**(3): p. 174-179.
11. Peter Van Wyk et al, H., *Farming marine shrimp in recirculating freshwater systems*. 1999, Florida Department of Agriculture and Consumer Services. p. 251.
12. Idoine, J., *Northern Shrimp*, in *Status of fishery resources off the northeastern United States.*, S.H. Clask, Editor. 2001, NOAA.
13. Schlining, K., *The Spot Prawn Resource in Carmel Submarine Canyon, California: Aspects of Fisheries and Habitat Associations*, in *Marine Biology, Moss Landing Marine Labs*. 1999, University of California Stanislaus: Moss Landing, CA.
14. Fisheries, L.D.o.W.a., *Early Life Cycle of the White Shrimp: A Review of the Literature*. 2000, Louisiana Department of Wildlife and Fisheries: Baton Rouge, LA.
15. Benfield, M.C.a.R.G.D., *Spatial and temporal variability in the nearshore distributions of postlarval *Farfantepenaeus aztecus* along Galveston Island, Texas*. *Estuarine Coastal Shelf Sci*, 2001. **52**(4): p. 445-456.
16. Paez Osuna, F., *The Environmental Impact of Shrimp Aquaculture: Causes, Effects, and Mitigating Alternatives*. *Environmental Management*, 2001. **28**(1): p. 131-140.
17. Paez-Osuna, F., *The environmental impact of shrimp aquaculture: a global perspective*. *Environmental Pollution*, 2001. **112**(2): p. 229-231.
18. Clay, J.W., *Environmental and social BMPS for shrimp aquaculture*. 2001, 143 J.M Parker Coliseum Louisiana State University Baton Rouge LA 70803: World Aquaculture Society.
19. Handbook, S., *Shellfish Species section; Shrimp Product Forms section*. 1999, Seafood Business, Inc.
20. NOAA, *Gulf of Mexico Shrimp Statistics*. 2001, National Marine Fisheries Service.
21. Victoria, M., *Gourmet's guide to the names of edible Crustacea*. 2000, Museum Victoria: Victoria, Australia.
22. Industry, M.S., *About Shrimp*. 2001.
23. McGovern, D., *WorldCatch Market Report: Shrimp buyers take action on bargains in shrimp market*. 2001, WorldCatch.com.
24. NMFS, *Commercial fisheries, annual landings. Annual trade by product for all countries. U.S. import and export statistics*. 2001.
25. FAO, *The State of World Fisheries and Aquaculture, 2000*. 2000, United Nations Food and Agriculture Organization: Geneva. p. 142.
26. Samocha, T.M., et al., *Intensive and super-intensive production of the Pacific white *Litopenaeus vannamei* in greenhouse-enclosed raceway systems*. 2001, 143 J.M Parker Coliseum Louisiana State University Baton Rouge LA 70803: World Aquaculture Society.
27. Hernandez-Cornejo, R. and A. Ruiz-Luna, *Development of shrimp farming in the coastal zone of southern Sinaloa (Mexico): operating characteristics, environmental issues, and perspectives*. *Ocean & Coastal Management*, 2000. **43**(7): p. 597-607.
28. Teichert-Coddington, D.R., D. Martinez, and E. Ramirez, *Partial nutrient budgets for semi-intensive shrimp farms in Honduras*. *Aquaculture*, 2000. **190**(1-2): p. 139-154.
29. Srinath, K., et al., *Group farming for sustainable aquaculture*. *Ocean & Coastal Management*, 2000. **43**(7): p. 557-571.

30. Treece, G.D. and L. Hamper, *Achieving environmentally friendly shrimp farming in Texas, USA*. World Aquaculture, 2000. **31**(2): p. 49-53.
 31. Naylor, Rosamund, et al. 2001. *Is aquaculture a net loss?* Conservation Biology in Practice, 2001. **2**(4): p. 26,27.
 32. Rao, G.R.M., *Present status of coastal aquaculture in India*. Appl. Fish. Aquac., 2001. **1**(1): p. 143-144.
 33. Wistrand, A., *Shrimp cultivation puts environment in danger*. 2001, Nijera Kori (NGO): Bangladesh.
 34. Trott, L.A. and D.M. Alongi, *The Impact of Shrimp Pond Effluent on Water Quality and Phytoplankton Biomass in a Tropical Mangrove Estuary*. Marine Pollution Bulletin, 2000. **40**(11): p. 947-951.
 35. Jones, A.B., et al., *Assessing Ecological Impacts of Shrimp and Sewage Effluent: Biological Indicators with Standard Water Quality Analyses*. Estuarine, Coastal and Shelf Science, 2001. **52**(1): p. 91-109.
 36. Salayakanond, V., *Farmers warned of contaminated prawns*, in *Business Day Thailand*. 2001.
 37. Rebecca Goldberg, M.E.a.R.N., *Marine Aquaculture in the United States: Environmental Impacts and Policy Options*. 2001, Pew Oceans Commission: Arlington, VA.
 38. McKinney, L. and B. Harvey, *Recreational fishing and aquaculture: Throwing a line in the pond*. 2001, 143 J.M Parker Coliseum Louisiana State University Baton Rouge LA 70803: World Aquaculture Society.
 38. Nation, 2002. Thai shrimp company pushes for chloramphenicol ban. News wire report, 16 January 2002, distributed by WorldSources, Inc.
- Adams, Chuck, Don Sweat, and Carlos Martinez. 2004. Status of shrimp farming activities in Florida. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Alam, S.M. Nazmul and Michael J. Phillips. 2004. Economic impact of marine finfish in shrimp farming systems in Bangladesh. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Alava, Juan Jose. 2004. Environmental impact assessment on inland shrimp farming in Ecuador. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Alston, Dallas & Alexis Cabarcas-Nunez, 2004. Caribbean shrimp culture with emphasis on Puerto Rico. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Ambio, 2002. Industrial transformation and shrimp aquaculture in Thailand and Vietnam: pathways to ecological, social, and economic sustainability? Royal Swedish Academy of Sciences, Ambio, v.31 #4, June 2002.
- Cutland, Laura and Drew Cherry. 2002. Seafood Consumption Trends in the United States: The Present and Future. Industry report from IntraFish, 31 pp. Available at www.intrafish.com
- Clay, Jason. 2003. Personal communications with Seafood Watch program, by mail and email, July 2003. Review of Seafood Watch Hawaiian Farmed Shrimp report. Email address jason.clay@wwfus.org
- Clay, Jason. 2004. Senior Fellow, World Wildlife Fund-U.S. Comments received in the review of this report.
- Dalzell, Paul. 2003. Crustacean fisheries specialist, Western Pacific Fisheries Management Council. Personal communications 3/19/03 by email and 4/14/03 by phone. Email paul.dalzell@noaa.gov; phone 808-522-6042
- Davis, D. Allen, Gregory Whitis, G. Teichert-Coddington, David B. Rouse and C.E. Boyd, 2004. Inland penaeid shrimp farming in Alabama. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- EJF, 2001. Habitat loss associated with shrimp farming. Environmental Justice Foundation.
- Fegan, Dan. 2004. Shrimp farming in Thailand. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- GAA, 1999. Codes of practice for responsible shrimp farming. Global Aquaculture Alliance, St. Louis, Missouri
- Gautier, Dominic, Hector L. Corrales, and John M. Wigglesworth. 2004. Success with a low-impact production system in Honduras. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Globefish, 2003. Commodity Update: Shrimp. Extract from Globefish databank—prices, exports, imports, catches, consumption. Fish Information Network of U.N. FAO, FAO/Globefish, Rome, Italy.

- Green, Bartholomew W. 2004. Shrimp culture in low-salinity water in Arkansas. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Gutierrez, Daniel and Jesus Zendejas. 2004. *Litopenaeus vannamei* aquaculture in North Sonora, Mexico: challenges and opportunities for a fast-growing sustainable industry. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Jost, Werner. 2004. Camenor as an example of a shrimp farm in the Brazilian shrimp industry. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Lassen, Thor. 2004. Mangrove conservation and shrimp aquaculture. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Marquez, Jorge Vareia. 2001. The human rights consequences of inequitable trade and development expansion: the abuse of law and community rights in the Gulf of Fomesca, Honduras. Mangrove Action Project, Earth Island Institute. Available online at www.earthisland.org/map/ineq-trd.htm
- Moya, Marcos, Walter Carlson, and Torres Tulio. 2004. Pacific white shrimp *Peneaus vannamei* production in covered ponds in northern Peru. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- NMFS 2002 Report to Congress, 2003. Report to Congress: Status of Fisheries of the United States for 2002. Published April 2003. National Marine Fisheries Service, Silver Spring, MD.
- NMFS Stats, 2003. U.S. import and export statistics for fisheries, international and domestic. Available online at <http://www.st.nmfs.gov/commercial/index.html> and <http://www.st.nmfs.gov/st1/trade/index.html>
- Nunes, Alberto J. 2004. Competitive production strategies in the grow-out culture of *Litopenaeus vannamei* in Northeast Brazil. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Permian Sea Shrimp, 2004. Press release detailing operations of Permian Sea Shrimp company, Imperial, Texas www.permianshrimp.com
- Phuong, N.T. 2004. Shrimp farming in Vietnam: an overview. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- Pike, Ian. 2003. Feed grade fisheries: management and sustainability. Keynote address, SEAFeds Workshop, Stirling Management Centre, April 8-9, 2003. Abstract published in Workshop Report.
- Rosenberg, Robert (ed). 2003. World Shrimp Farming.
- Reyna, Jose Jesus. 2004. Driving factors for shrimp farming success in Mexico. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.
- SEAFeds Workshop Report, 2003. Report summarizing results of the workshop on developing sustainable environmental aquaculture feeds, held at Stirling Management Centre, April 8-9, 2003. 30 pp.
- Shumway, Sandy. 2003. University of Connecticut. Comments received in the review of Seafood Watch coldwater shrimp report.
- Steiner, Todd. 2003. Sea Turtle Restoration Project. Comments received in the review of this report.
- Stickney, Robert. 2002. Director, Texas Sea Grant Aquaculture Program. Personal communication, by phone, 8/26/02.
- Stokes, Al D., Craig Browdy & Heidi Atwood, 2004. The status of marine shrimp farming in South Carolina, 2003. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.

Shrimp_farmed_SFW_ac_final.doc

Szuster, Brian, Mark Flaherty, and Marut Diparos. 2004. Marine shrimp farming in Thailand: an industry in transition. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.

Tacon, Albert. 2003. Personal communication with Seafood Watch program, July 18, 2003. Review of Seafood Watch Farmed Hawaiian Shrimp document. Email address AGJTACON@aol.com

Tacon, Albert. 2003. Sustainable aquaculture feeds: an overview and global perspective. Keynote address, SEAFeds Workshop, Stirling Management Centre, April 8-9, 2003. Abstract published in Workshop Report.

Treece, Gramble. 2002. Extension specialist, shrimp aquaculture, Texas Sea Grant. Personal communication, 8/26/02

Tomomitsu, Mark, 2002. Water quality specialist, Hawaii Department of Health, Clean Water Branch, Honolulu, HI. Personal communication, by phone, 12/09/02. tel: (808) 586-4309

Valderama, Diego, and Carole R. Engle. 2002. Economic optimization of shrimp farming in Honduras. Journal of the World Aquaculture Society, v.33 no. 4

Wang, Quingyin, et al. 2004. Development of the marine shrimp farming industry in China: opportunities and Challenges. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.

Watling, Les. 2004. University of Maine. Comments received in the review of Seafood Watch coldwater shrimp report.

Wilkenfield, Joshua, Craig Collins, Roger Drudge, and Tark Rush. 2004. Inland culture of the white shrimp *Litopenaeus vannamei* in Arizona: going forward or backward? Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.

Yap, Wilfredo, and Antonio c. Villaluz. 2004. Practices and economics of successful shrimp farms in the Philippines. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.

Young, Leonard, 2003. Aquaculture specialist, USDA/Hawaii Aquaculture Development Program, Honolulu, HI. Personal communication, by phone, 4/14/03. Phone number 808-587-0030

Zendejas-Hernandez, Jesus. 2004. Shrimp farming in Mexico: a description and production cost analysis. Presentation to World Aquaculture Society, March 1-5 2004, Honolulu, Hawaii.