

Fenchel-Lagrange versus Geometric Duality in Convex Optimization^{1, 2}

R. I. BOT₃³, S. M. GRAD⁴, AND G. WANKA⁵

Communicated by V. F. Demyanov

¹The first author was supported in part by Gottlieb Daimler and Karl Benz Stiftung 02-48/99.

²The second author was supported in part by Karl und Ruth Mayer-Stiftung.

³Assistant Professor, Faculty of Mathematics, Chemnitz University of Technology, Chemnitz, Germany.

⁴Ph. D. Student, Faculty of Mathematics, Chemnitz University of Technology, Germany.

⁵Professor, Faculty of Mathematics, Chemnitz University of Technology, Chemnitz, Germany.

Abstract. We present a new duality theory in order to treat convex optimization problems and we prove that the geometric duality used by C.H. Scott and T.R. Jefferson in different papers during the last quarter of century is a special case of it. Moreover, weaker sufficient conditions in order to achieve strong duality are considered and optimality conditions are derived. Next we apply our approach to some problems considered by Scott and Jefferson, determining their duals. We give weaker sufficient conditions in order to achieve strong duality and the corresponding optimality conditions. Finally, posynomial geometric programming is viewed also as a particular case of the duality approach we present.

Key Words. Geometric programming, convex optimization, perturbation theory, Lagrange and Fenchel duality, conjugate functions.

1. Introduction

A quarter of century after the inception of their series of papers on geometric programming (see Ref. 1, among other papers) Scott and Jefferson have published recently the article cited as Ref. 2 where they treat a class of optimization problems by means of geometric duality. This fact denotes that the approach often used by these authors enjoys a continuous attention. Our main purpose is to introduce a duality theory for convex optimization problems which has many advantages in competition to the classical generalized geometric duality established by Peterson (Ref. 3) and used later in a simplified version by Scott and Jefferson (sometimes together with S. Jorjani) in many papers (see Refs. 1-2, 4-14).

We have proved in Ref. 15 that Peterson's geometric dual problems can be obtained using the perturbation theory presented in Refs. 16-17. Here we show that the geometric dual problem used by Scott and Jefferson is actually the Fenchel-Lagrange dual of the primal geometric problem. Moreover, we provide much weaker sufficient conditions in order to achieve strong duality than the ones considered by them. Later we review some problems treated by means of geometric duality by these authors and we prove that their dual problems can be determined easier with our approach, where

artificial constructions of the implicit, explicit and cone constraints are not required.

Unlike the cited authors, that do not bother about the sufficient conditions for strong duality, mentioning just that they come from Peterson's work Ref. 3, we resort to weaker conditions than the ones obtained there. Using them, we display the strong duality assertion and the optimality conditions for these problems.

The Fenchel-Lagrange dual problem has been developed by Boş and Wanka in some papers (Refs. 15, 18-22) and it is constructed by means of the perturbation theory in Refs. 16-17. Its name, introduced for the first time in Ref. 18, reveals its origin, as it is a combination between the well-known dual problems due to W. Fenchel and J.L. Lagrange. So far it has been considered for problems with constraint functions defined over the whole space \mathbb{R}^n , while here they are defined over a subset of \mathbb{R}^n .

We consider a convex optimization problem (P), the so-called primal problem, that consists in minimizing a function defined over the space \mathbb{R}^n whose values are not necessarily all finite when its variable is required to belong to a subset of \mathbb{R}^n and to satisfy the non-positivity of some constraint real-valued functions defined over the same subset. To it we determine the Fenchel-Lagrange dual problem, describing its construction in detail. Then we introduce a constraint qualification whose fulfillment

is sufficient in order to have strong duality for the two problems mentioned above.

After proving the strong duality assertion we formulate and prove also the optimality conditions for these problems and this ends the second part of the present work.

Further we consider a special case of the general convex optimization problem (P) denoted (P_K) where the variable is forced to belong also to a closed convex cone in \mathbb{R}^n . The generalized geometric problem used by Scott and Jefferson turns out to be a special case of this problem for a suitable choice of the functions and sets involved. Strong duality and optimality conditions for (P_K) follow and from them are derived the ones regarding the mentioned geometric problem. The sufficient conditions required for strong duality are weaker than the ones used before by the cited authors, as we do not ask the functions and the sets involved to be also closed alongside their convexity.

The fourth part of the paper deals with some problems Scott and Jefferson have treated by means of geometric programming duality. We prove that their artificial constructions in order to bring the problems into the form of the primal geometric problem are not necessary, as the same duals can be obtained simpler by applying our Fenchel-Lagrange duality. A large variety of problems is presented: minmax program (Ref. 11), entropy constrained program (Ref. 14), facility location problem (Ref. 12),

quadratic concave fractional program (Ref. 9), problem of minimizing a sum of convex ratios (Ref. 10) and quasiconcave multiplicative program (Ref. 2). Strong duality, obtained under weaker conditions than in the original papers, and optimality conditions assertions are delivered for these problems. We also show that the posynomial geometric programming (Ref. 23) is a special case of the Fenchel-Lagrange duality.

2. The Fenchel-Lagrange dual problem

Let us consider a convex subset X of \mathbb{R}^n and a convex function $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, $\overline{\mathbb{R}} = \mathbb{R} \cup \{\pm\infty\}$ and $g = (g_1, \dots, g_k)^T : X \rightarrow \mathbb{R}^k$, with $g_i, i = 1, \dots, k$, convex functions. We introduce the following convex optimization problem, further called primal problem,

$$(P) \quad \inf_{x \in X, g(x) \leq 0} f(x).$$

As usual, $g(x) \leq 0$ means $g_i(x) \leq 0$ for all $i = 1, \dots, k$. We denote by $\text{dom}(f) = \{x \in \mathbb{R}^n : f(x) < +\infty\}$ the effective domain of the function $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$.

A short observation is necessary: in the previous works of two of the authors (Refs. 18-22) the constraint functions $g_i, i = 1, \dots, k$, are defined over the whole space, unlike here. By defining the constraint functions over a subset of \mathbb{R}^n we are able to treat a larger class of convex optimization problems. In order to determine the so-called Fenchel-Lagrange dual problem to (P) we need to introduce the perturbation

function (cf. Refs. 18-22) $\Phi : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \overline{\mathbb{R}}$, $\Phi(x, y, z) = f(x + y)$ if $x \in X$ and $g(x) \leq z$, otherwise taking the value $+\infty$, with the perturbation variables y and z . Following the perturbation method described in Refs. 16-17 the next step is to calculate the conjugate function of Φ . When a function k is defined over a subset U of \mathbb{R}^n , we define for it the so-called conjugate relative to the set U by $k_U^* : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, $k_U^*(x^*) = \sup_{x \in U} \{ \langle x^*, x \rangle - k(x) \}$. One may notice that when $U = \mathbb{R}^n$ the conjugate of k relative to the set U becomes the classical (Fenchel-Rockafellar) conjugate function of k . This will be denoted by k^* .

Let us proceed with the definition for $\Phi^* : \mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^k \rightarrow \overline{\mathbb{R}}$,

$$\begin{aligned} \Phi^*(x^*, p, q) &= \sup_{\substack{x, y \in \mathbb{R}^n, \\ z \in \mathbb{R}^k}} \left\{ \langle x^*, x \rangle + \langle p, y \rangle + \langle q, z \rangle - \Phi(x, y, z) \right\} \\ &= \sup_{\substack{x \in X, y \in \mathbb{R}^n, \\ g(x) \leq z}} \left\{ \langle x^*, x \rangle + \langle p, y \rangle + \langle q, z \rangle - f(x + y) \right\} \end{aligned}$$

We have used the denotations $\langle x^*, x \rangle := x^{*T}x$, $\langle p, y \rangle := p^T y$ and $\langle q, z \rangle := q^T z$ for the Euclidean scalar products in the corresponding spaces \mathbb{R}^n and \mathbb{R}^k , respectively. In order to calculate Φ^* let us introduce new variables r instead of y and s instead of z by $r := x + y$ and $s := z - g(x)$. The supremum above is computed with respect to three independent variables, so it can be separated into a sum of three suprema,

$$\Phi^*(x^*, p, q) = \sup_{s \in \mathbb{R}_+^k} \langle q, s \rangle + \sup_{r \in \mathbb{R}^n} \left\{ \langle p, r \rangle - f(r) \right\} + \sup_{x \in X} \left\{ \langle x^* - p, x \rangle + \langle q, g(x) \rangle \right\},$$

which is actually $f^*(p) - \inf_{x \in X} \left\{ \langle p - x^*, x \rangle - \langle q, g(x) \rangle \right\}$ if $-q \in \mathbb{R}_+^k = \{z : z \in \mathbb{R}^k, 0 \leq z\}$

and $+\infty$ otherwise. According to Ref. 16 the dual problem to the problem (P) is

$$(D) \quad \sup_{p \in \mathbb{R}^n, q \in \mathbb{R}^k} \left\{ -\Phi^*(0, p, q) \right\},$$

that becomes in our case after changing the sign of the variable q

$$(D) \quad \sup_{p \in \mathbb{R}^n, q \in \mathbb{R}_+^k} \left\{ -f^*(p) + \inf_{x \in X} [\langle p, x \rangle + \langle q, g(x) \rangle] \right\}.$$

It is obvious from the construction of the dual that the weak duality assertion between (P) and (D), i. e. the value of the primal objective function at any primal feasible point is greater than or equal to the value of the dual objective function at any dual feasible point, always stands. We will not mention further for each pair of dual problems we treat that the weak duality is true, as it is valid in the most general case without any additional assumption. By strong duality we understand the situation in which the optimal objective values of the primal and dual are equal and the dual has an optimal solution. Unlike weak duality, strong duality can fail in the general case. To avoid any undesired event of this kind, we introduce a constraint qualification that guarantees the validity of strong duality in case it is fulfilled. First let us divide the

index set $\{1, \dots, k\}$ into two subsets,

$$L := \left\{ i \in \{1, \dots, k\} : g_i : X \rightarrow \mathbb{R} \text{ is the restriction to } X \text{ of an affine function on } \mathbb{R}^n \right\},$$

and $N := \{1, \dots, k\} \setminus L$. The constraint qualification follows

$$(CQ) \quad \exists x' \in \text{ri}(X) \cap \text{ri}(\text{dom}(f)) : g_i(x') \leq 0 \text{ if } i \in L \text{ and } g_i(x') < 0 \text{ if } i \in N,$$

where $\text{ri}(X)$ denotes the relative interior of the set X . We are ready now to formulate the strong duality assertion. Before that let us denote by $\inf(P)$ and $\sup(D)$ the optimal objective values of the primal and dual problem, respectively.

Theorem 2.1. Provided that the constraint qualification (CQ) is fulfilled, there is strong duality between problems (P) and (D) , i. e. their optimal objective values are equal and the dual problem has an optimal solution.

Proof. We can write the problem (P) equivalently

$$(P) \quad \inf_{x \in X \cap \text{dom}(f), g(x) \leq 0} f(x).$$

By Theorem 6.5 in Ref. 17, (CQ) yields $x' \in \text{ri}(X \cap \text{dom}(f)) = \text{ri}(X) \cap \text{ri}(\text{dom}(f))$.

Theorem 5.7 in Ref. 24 states under the present hypotheses the existence of a $\bar{q} \geq 0$

such that $\inf(P) = \max_{q \geq 0} \inf_{x \in X \cap \text{dom}(f)} [f(x) + \langle q, g(x) \rangle] = \inf_{x \in X \cap \text{dom}(f)} [f(x) + \langle \bar{q}, g(x) \rangle]$.

Defining $h : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, $h(x) = \langle \bar{q}, g(x) \rangle$ if $x \in X$ and $+\infty$ otherwise, we can rewrite the

right-hand side term as $\inf(P) = \inf_{x \in \mathbb{R}^n} [f(x) + h(x)]$. Because $\text{ri}(\text{dom}(f)) \cap \text{ri}(\text{dom}(h))$

$= \text{ri}(\text{dom}(f)) \cap \text{ri}(X) \neq \emptyset$, by Theorem 31.1 (Fenchel's Duality Theorem) in Ref. 17,

there exists a $\bar{p} \in \mathbb{R}^n$ such that this infimum is

$$\begin{aligned} \inf(P) &= \max_{p \in \mathbb{R}^n} [-f^*(p) - h^*(-p)] = -f^*(\bar{p}) - h^*(-\bar{p}) = -f^*(\bar{p}) - \sup_{x \in \mathbb{R}^n} \{\langle -\bar{p}, x \rangle\} \\ -h(x) &= -f^*(\bar{p}) - \sup_{x \in X} \{\langle -\bar{p}, x \rangle - \langle \bar{q}, g(x) \rangle\} = -f^*(\bar{p}) + \inf_{x \in X} \{\langle \bar{p}, x \rangle + \langle \bar{q}, g(x) \rangle\}. \quad (1) \end{aligned}$$

In the right-hand term of (1) we see the objective function of the dual problem (D)

at (\bar{p}, \bar{q}) . From weak duality follows that the supremum of (D) is attained, becoming

maximum, at (\bar{p}, \bar{q}) , which turns out to be an optimal solution of the dual. \square

Remark 2.1. In the proof above we proved first that under (CQ) there holds strong duality between the primal problem and its Lagrange dual problem. Then we proved strong duality between the Lagrange dual and its Fenchel dual problem, the last one proving to be exactly the Fenchel-Lagrange dual problem introduced earlier.

Next we derive necessary and sufficient optimality conditions regarding the problems (P) and (D).

Theorem 2.2. (a) If the constraint qualification (CQ) is fulfilled and the primal problem (P) has an optimal solution \bar{x} , then the dual problem has an optimal solution (\bar{p}, \bar{q}) and the following optimality conditions are satisfied

(i) $f(\bar{x}) + f^*(\bar{p}) = \langle \bar{p}, \bar{x} \rangle,$

$$(ii) \inf_{x \in X} [\langle \bar{p}, x \rangle + \langle \bar{q}, g(x) \rangle] = \langle \bar{p}, \bar{x} \rangle,$$

$$(iii) \langle \bar{q}, g(\bar{x}) \rangle = 0.$$

(b) If \bar{x} is a feasible point to (P) and (\bar{p}, \bar{q}) is feasible to the dual problem (D) fulfilling the optimality conditions (i)-(iii), then there is strong duality between (P) and (D) and the mentioned feasible points turn out to be optimal solutions.

Proof. (a) Theorem 2.1 guarantees strong duality between (P) and (D) . So the dual problem has an optimal solution. Let us denote it by (\bar{p}, \bar{q}) . The equality of the optimal objective values of (P) and (D) implies

$$f(\bar{x}) + f^*(\bar{p}) - \inf_{x \in X} [\langle \bar{p}, x \rangle + \langle \bar{q}, g(x) \rangle] = 0. \quad (2)$$

It is obvious that $\inf_{x \in X} [\langle \bar{p}, x \rangle + \langle \bar{q}, g(x) \rangle] \leq \langle \bar{p}, \bar{x} \rangle + \langle \bar{q}, g(\bar{x}) \rangle$, while Young's inequality states $f(\bar{x}) + f^*(\bar{p}) \geq \langle \bar{p}, \bar{x} \rangle$. Combining (2) with these relations leads to $0 \geq \langle \bar{p}, \bar{x} \rangle - \langle \bar{p}, \bar{x} \rangle + \langle \bar{q}, g(\bar{x}) \rangle = \langle \bar{q}, g(\bar{x}) \rangle$, but $\langle \bar{q}, g(\bar{x}) \rangle \leq 0$ because of the feasibility of \bar{x} to (P) and \bar{q} to (D) , respectively. Therefore (iii) is true, while (i) and (ii) follow immediately by adding and subtracting $\langle \bar{p}, \bar{x} \rangle$ to (2) which yields

$$[f(\bar{x}) + f^*(\bar{p}) - \langle \bar{p}, \bar{x} \rangle] + [\langle \bar{p}, \bar{x} \rangle - \inf_{x \in X} [\langle \bar{p}, x \rangle + \langle \bar{q}, g(x) \rangle]] = 0.$$

(b) The calculations presented above can be carried out in reverse order. \square

Remark 2.2. The optimality conditions (i)-(iii) are sufficient in the most general case as (b) applies without any convexity assumption or constraint qualification.

3. A particular case: the geometric programming duality

Our main stimulus in writing this paper was to prove that the simplified generalized geometric duality is nothing but a particular case of Fenchel-Lagrange duality. So far we have presented the basic facts and results regarding the Fenchel-Lagrange duality and now we consider first a special case of the primal problem (P) which is still more general than the geometric primal problem. So we consider the problem

$$(P_K) \quad \inf_{x \in X, g(x) \leq 0, x \in K} f(x),$$

where K is a closed convex cone in \mathbb{R}^n . Its Fenchel-Lagrange dual problem is

$$(D_K) \quad \sup_{p \in \mathbb{R}^n, q \in \mathbb{R}_+^k} \left\{ -f^*(p) + \inf_{x \in X \cap K} [\langle p, x \rangle + \langle q, g(x) \rangle] \right\}.$$

The constraint qualification that is sufficient for the existence of strong duality in this case is, with the notations introduced before,

$$(CQ_K) \quad \exists x' \in \text{ri}(X) \cap \text{ri}(\text{dom}(f)) : g_i(x') \leq 0 \text{ if } i \in L, g_i(x') < 0 \text{ if } i \in N \text{ and } x' \in \text{ri}(K).$$

The strong duality assertion regarding (P_K) and an equivalent form of its dual follows.

Theorem 3.1. If the constraint qualification (CQ_K) is satisfied, there is strong

duality between the primal problem (P_K) and the equivalent formulation of its dual,

$$(D'_K) \quad \sup_{p \in \mathbb{R}^n, q \in \mathbb{R}_+^k, t \in K^*} \left\{ -f^*(p) + \inf_{x \in X} [\langle p - t, x \rangle + \langle q, g(x) \rangle] \right\}.$$

Proof. (P_K) being a special case of the problem (P) , like (D_K) of its dual (D) and because $\text{ri}(X \cap K) = \text{ri}(X) \cap \text{ri}(K)$, strong duality is valid for (P_K) and (D_K) .

But the presence of the cone K in the formula of (D_K) is not so desired, so we need to find an alternative formulation to this dual problem. Let us rewrite the infimum

$$\text{contained in its formulation as } \inf_{x \in X \cap K} [\langle p, x \rangle + \langle q, g(x) \rangle] = \inf_{x \in \mathbb{R}^n} [\langle p, x \rangle + h(x) + \delta_K(x)],$$

where the function h is defined like in the proof of Theorem 2.1 and we use also the indicator function $\delta_K : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}, \delta_K(x) = 0, \text{ if } x \in K \text{ and } \delta_K(x) = +\infty, \text{ if } x \notin K.$

As a conjugate function, the right-hand side of the relation above is equal to

$-\left(\langle p, \cdot \rangle + h + \delta_K\right)^*(0)$, which, by Theorem 20.1 in Ref. 17, can be written as

$$\begin{aligned} & - \inf_{t \in \mathbb{R}^n} \left[(\langle p, \cdot \rangle + h)^*(t) + \delta_K^*(-t) \right] = - \inf_{t \in K^*} \sup_{x \in X} \{ \langle t, x \rangle - \langle p, x \rangle - \langle q, g(x) \rangle \} \\ & = \sup_{t \in K^*} \inf_{x \in X} \{ \langle p - t, x \rangle + \langle q, g(x) \rangle \} \end{aligned}$$

and the existence of a $\bar{t} \in K^*$ where this supremum is attained is granted. Here K^*

$= \{y \in \mathbb{R}^n : \langle y, x \rangle \geq 0, \forall x \in K\}$ is the dual cone of K . Obviously, the dual problem

(D_K) is equivalent to (D'_K) and strong duality between (P_K) and (D'_K) holds. \square

Remark 3.1. One can obtain the dual problem (D'_K) also by perturbations, in a

similar way we obtained (D) in the previous section by including the cone constraint in the constraint function (cf. Refs. 18-20). We refer further to (D'_K) as the dual problem of (P_K) .

Necessary and sufficient optimality conditions are derived from the general case.

Theorem 3.2. (a) If the constraint qualification (CQ_K) is fulfilled and the primal problem (P_K) has an optimal solution \bar{x} , then the dual problem (D'_K) has an optimal solution $(\bar{p}, \bar{q}, \bar{t})$ and the following optimality conditions are satisfied

$$(i) \quad f(\bar{x}) + f^*(\bar{p}) = \langle \bar{p}, \bar{x} \rangle,$$

$$(ii) \quad \inf_{x \in X} [\langle \bar{p} - \bar{t}, x \rangle + \langle \bar{q}, g(x) \rangle] = \langle \bar{p}, \bar{x} \rangle,$$

$$(iii) \quad \langle \bar{q}, g(\bar{x}) \rangle = 0,$$

$$(iv) \quad \langle \bar{t}, \bar{x} \rangle = 0.$$

(b) If \bar{x} is a feasible point to (P_K) and $(\bar{p}, \bar{q}, \bar{t})$ is feasible to the dual problem (D'_K) fulfilling the optimality conditions (i)-(iv), then there is strong duality between (P_K) and (D'_K) and the mentioned feasible points turn out to be optimal solutions.

With a suitable choice of the functions and the sets involved in the problem (P_K) one obtains a convex optimization problem equivalent to the primal geometric problem used by Scott and Jefferson in many papers (Refs. 1-2, 4-14),

$$(P_g) \quad \inf_{\substack{x=(x^0, x^1, \dots, x^k) \in C_0 \times C_1 \times \dots \times C_k, \\ g^i(x^i) \leq 0, i=1, \dots, k, x \in K}} g^0(x^0),$$

where $C_i \subseteq \mathbb{R}^{l_i}$, $i = 0, \dots, k$, $\sum_{i=0}^k l_i = n$, are convex sets, $g^i : C_i \rightarrow \mathbb{R}$, $i = 0, \dots, k$, convex

functions and $K \subseteq \mathbb{R}^n$ is a closed convex cone. The proper selection follows

$X = \mathbb{R}^{l_0} \times C_1 \times \dots \times C_k$, $f : \mathbb{R}^n \rightarrow \bar{\mathbb{R}}$, $f(x) = g^0(x^0)$ if $x \in C_0 \times \mathbb{R}^{n-l_0}$ and $+\infty$

otherwise, $g_i : X \rightarrow \mathbb{R}$, $i = 1, \dots, k$, $g_i(x) = g^i(x^i)$, $i = 1, \dots, k$, $x = (x^0, x^1, \dots, x^k) \in X$.

Now we can write the Fenchel-Lagrange dual problem to (P_g) (cf. (D'_K))

$$(D_g) \quad \sup_{p \in \mathbb{R}^n, q \in \mathbb{R}_+^k, t \in K^*} \left\{ -f^*(p) + \inf_{x \in \mathbb{R}^{l_0} \times C_1 \times \dots \times C_k} \left[\langle p - t, x \rangle + \sum_{i=1}^k q_i g^i(x^i) \right] \right\}.$$

The conjugate of f at $p = (p^0, p^1, \dots, p^k) \in \mathbb{R}^{l_0} \times \mathbb{R}^{l_1} \times \dots \times \mathbb{R}^{l_k}$ is

$$f^*(p) = \sup_{x=(x^0, x^1, \dots, x^k) \in C_0 \times \mathbb{R}^{l_1} \times \dots \times \mathbb{R}^{l_k}} \left\{ \sum_{i=0}^k \langle p^i, x^i \rangle - g^0(x^0) \right\},$$

which is equal to $g_{C_0}^{0*}(p^0)$ when $p^i = 0$ for all $i = 1, \dots, k$, otherwise being $+\infty$.

As the infimum that appears is separable into a sum of infima, the dual becomes

$$(D_g) \quad \sup_{\substack{p^0 \in \mathbb{R}^{l_0}, \\ q \in \mathbb{R}_+^k, t \in K^*}} \left\{ -g_{C_0}^{0*}(p^0) + \inf_{x^0 \in \mathbb{R}^{l_0}} \langle p^0 - t^0, x^0 \rangle + \sum_{i=1}^k \inf_{x^i \in C_i} [\langle -t^i, x^i \rangle + q_i g^i(x^i)] \right\},$$

if we consider $t = (t^0, t^1, \dots, t^k) \in \mathbb{R}^{l_0} \times \mathbb{R}^{l_1} \times \dots \times \mathbb{R}^{l_k}$. As $\inf_{x^0 \in \mathbb{R}^{l_0}} \langle p^0 - t^0, x^0 \rangle$ is equal to

0 when $p^0 = t^0$, otherwise being $-\infty$, the dual problem to (P_g) turns into

$$(D_g) \quad \sup_{q \in \mathbb{R}_+^k, t \in K^*} \left\{ -g_{C_0}^{0*}(t^0) - \sum_{i=1}^k \sup_{x^i \in C_i} [\langle t^i, x^i \rangle - q_i g^i(x^i)] \right\}$$

This is exactly the geometric dual problem encountered in all cited papers by Scott

and Jefferson, written without resorting to the homogenous extension of the conjugate

functions that can replace the inner suprema in (D_g) . The constraint qualification sufficient to guarantee strong duality for this pair of problems is derived from (CQ_K) , $(CQ_g) \exists x' \in \text{ri}(K): g^i(x') \leq 0, i \in L, g^i(x') < 0, i \in N$ and $x'^i \in \text{ri}(C_i), i = 0, \dots, k$.

Theorem 3.3. The satisfaction of the constraint qualification (CQ_g) is sufficient to guarantee strong duality regarding (P_g) and (D_g) .

Remark 3.2. The cited papers of the mentioned authors do not assert trenchantly any strong duality allegation, containing just the optimality conditions, while for the background of their achievement the reader is referred to Ref. 3. There all the functions and the sets involved are postulated as being closed, alongside their convexity assumptions that proved to be sufficient in our proofs when the constraint qualification is fulfilled. Moreover, the possibility to impose a milder constraint qualification regarding the affine functions whose restrictions to the given set are among the constraint functions is not taken into consideration at all.

The optimality conditions concerning (P_g) and (D_g) spring from Theorem 3.2.

Theorem 3.4. (a) If the constraint qualification (CQ_g) is fulfilled and the primal problem (P_g) has an optimal solution $\bar{x} = (\bar{x}^0, \bar{x}^1, \dots, \bar{x}^k)$, then the dual problem (D_g) has an optimal solution (\bar{q}, \bar{t}) and the following optimality conditions are satisfied

$$(i) \ g^0(\bar{x}^0) + g_{C_0}^{0*}(\bar{t}^0) = \langle \bar{t}^0, \bar{x}^0 \rangle,$$

$$(ii) \ (\bar{q}_i g_i)_{C_i}^*(\bar{t}_i) = \langle \bar{t}_i, \bar{x}^i \rangle, \ i = 1, \dots, k,$$

$$(iii) \ \bar{q}_i g^i(\bar{x}^i) = 0, \ i = 0, \dots, k,$$

$$(iv) \ \langle \bar{t}, \bar{x} \rangle = 0.$$

(b) If \bar{x} is a feasible point to the problem (P_g) and (\bar{q}, \bar{t}) is feasible to the dual problem (D_g) fulfilling the optimality conditions (i)-(iv), then there is strong duality between (P_g) and (D_g) and the mentioned feasible points turn out to be optimal solutions.

Remark 3.3. The optimality conditions we derived are equivalent to the ones displayed by Scott and Jefferson in the cited papers, but have a different form.

4. An overview of some special cases

This last section reviews some of the problems treated during the last quarter of century by Scott and Jefferson, sometimes together with Jorjani, by means of simplified generalized geometric programming. All the problems were artificially transformed into the framework required by geometric programming by introducing new variables in order to separate implicit and explicit constraints and building some cone where the new vector-variable is forced to lie. Then their dual problems arose from the general theory as well as the optimality conditions. We determine the Fenchel-Lagrange dual

problem for each problem, then we specialize the adequate constraint qualification and state the strong duality assertion followed by the optimality conditions all without proofs as they are direct consequences of Theorems 2.1 and 2.2. One may notice that even if the functions and the sets are considered closed in the original papers we remove this redundant property, as we demonstrate that strong duality and optimality conditions stand out even without its presence when the sufficient constraint qualification is valid. We have chosen six problems that we have considered more interesting, but also the problems in Refs. 1, 4-8 or 13 may benefit from the same treatment. The last subsection is dedicated to the well-known posynomial geometric programming which is undertaken into our duality theory, too. Other papers of Scott and Jefferson treat some problems by means of posynomial geometric programming, so we might have included some of these problems here.

4.1. Minmax programs (Ref. 11). The first problem we deal with is

$$(P_1) \quad \inf_{x \in C, Ax \geq b, g(x) \leq 0} \max_{i=1, \dots, I} \{f_i(x)\},$$

with $f_i : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$, $\text{dom}(f_i) = C$, $i = 1, \dots, I$, and $g = (g_1, \dots, g_J)^T : C \rightarrow \mathbb{R}^J$ convex functions, $C \subseteq \mathbb{R}^n$ a convex set, $A \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. In the original paper the functions $f_i, i = 1, \dots, I$, $g_j, j = 1, \dots, J$, and the set C are required to be also closed,

but strong duality is valid without this additional assumption. To treat the problem

(P_1) with the method presented in the second section, it is rewritten as

$$(P_1) \quad \inf_{\substack{x \in C, s \in \mathbb{R}, b - Ax \leq 0, \\ g(x) \leq 0, f_i(x) - s \leq 0, i=1, \dots, I}} s.$$

The Fenchel-Lagrange dual problem to (P_1) is, considering the objective function

$$u : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}, u(x, s) = s \text{ and } q^f = (q_1^f, \dots, q_I^f)^T,$$

$$(D_1) \quad \sup_{\substack{p^x \in \mathbb{R}^n, p^s \in \mathbb{R}, \\ q^l \in \mathbb{R}_+^m, q^f \in \mathbb{R}_+^I, q^g \in \mathbb{R}_+^J}} \left\{ -u^*(p^x, p^s) + \inf_{\substack{x \in C, \\ s \in \mathbb{R}}} [\langle p^x, x \rangle + \langle p^s, s \rangle] \right. \\ \left. + \sum_{i \in I} q_i^f (f_i(x) - s) + \langle q^g, g(x) \rangle + \langle q^l, b - Ax \rangle \right\}.$$

The conjugate of the objective function is $u^*(p^x, p^s) = 0$ if $p^s = 1$ and $p^x = 0$ and $+\infty$

otherwise. Noticing that the infimum in (D_1) is separable into a sum of two infima,

one concerning $s \in \mathbb{R}$, the other $x \in C$, the dual problem turns into

$$(D_1) \quad \sup_{\substack{q^l \in \mathbb{R}_+^m, q^f \in \mathbb{R}_+^I, \\ q^g \in \mathbb{R}_+^J}} \left\{ \inf_{x \in C} \left[\sum_{i \in I} q_i^f f_i(x) + \langle q^g, g(x) \rangle - \langle A^T q^l, x \rangle \right] + \inf_{s \in \mathbb{R}} \left[s - s \sum_{i \in I} q_i^f \right] + \langle q^l, b \rangle \right\}.$$

The second infimum is equal to 0 when $\sum_{i \in I} q_i^f = 1$, otherwise having the value $-\infty$,

while the first, transformed into a supremum, can be viewed as a conjugate function

relative to the set C . Applying Theorem 20.1 in Ref. 17 and denoting $q^g = (q_1^g, \dots, q_J^g)^T$,

the dual problem becomes

$$(D_1) \quad \sup_{\substack{q^l \in \mathbb{R}_+^m, q^g \in \mathbb{R}_+^J, q^f \in \mathbb{R}_+^I, \\ \sum_{i=1}^I q_i^f = 1, \sum_{i=1}^I u_i + \sum_{j=1}^J v_j = A^T q^l}} \left\{ \langle q^l, b \rangle - \sum_{i=1}^I (q_i^f f_i)^*(u_i) - \sum_{j=1}^J (q_j^g g_j)_C^*(v_j) \right\},$$

identical to the dual problem found in Ref. 11. A sufficient circumstance to be able to formulate the strong duality assertion is the following constraint qualification, where the sets L and N are considered as before,

(CQ_1) $\exists x' \in \text{ri}(C)$: $Ax' \geq b$ and $g_j(x') \leq 0$ if $j \in L$, respectively $g_j(x') < 0$ for $j \in N$.

Theorem 4.1. If the constraint qualification (CQ_1) is satisfied, then the strong duality between (P_1) and (D_1) is assured.

Since the optimality conditions are not delivered in Ref. 11, here they are, determined via our method.

Theorem 4.2. (a) If the constraint qualification (CQ_1) is fulfilled and \bar{x} is an optimal solution to (P_1) , then strong duality between the problems (P_1) and (D_1) is attained and the dual problem has an optimal solution $(\bar{q}^l, \bar{q}^f, \bar{q}^g, \bar{u}, \bar{v})$, where $\bar{u} = (\bar{u}_1, \dots, \bar{u}_I)^T$ and $\bar{v} = (\bar{v}_1, \dots, \bar{v}_J)^T$, satisfying the following optimality conditions

$$(i) f_i(\bar{x}) - \max_{i=1, \dots, I} \{f_i(\bar{x})\} = 0 \text{ if } \bar{q}_i^f > 0, i = 1, \dots, I,$$

$$(ii) \langle \bar{q}^l, b - A\bar{x} \rangle = 0,$$

$$(iii) \langle \bar{q}^g, g(\bar{x}) \rangle = 0,$$

$$(iv) \left(\bar{q}_i^f f_i \right)^* (\bar{u}_i) + \bar{q}_i^f f_i(\bar{x}) = \langle \bar{u}_i, \bar{x} \rangle, i = 1, \dots, I,$$

$$(v) \left(\bar{q}_j^g g_j \right)_C^* (\bar{v}_j) + \bar{q}_j^g g_j(\bar{x}) = \langle \bar{v}_j, \bar{x} \rangle, j = 1, \dots, J.$$

(b) Having a feasible solution \bar{x} to the primal and one $(\bar{q}^l, \bar{q}^f, \bar{q}^g, \bar{u}, \bar{v})$ to the dual satisfying the optimality conditions (i)-(v), the mentioned feasible solutions turn out to be optimal solutions to the corresponding problems and strong duality stands.

4.2. Entropy constrained programs (Ref. 14). A minute exposition of the way how the Fenchel-Lagrange duality is applicable to the problem treated in Ref. 14 is available in Ref. 15. Let us summarize it here. The problem

$$(P_2) \quad \inf_{\substack{Ax \geq b, -\sum_{i=1}^n x_i \ln x_i \geq H, \\ \sum_{i=1}^n x_i = 1, x_i \geq 0}} \langle c, x \rangle,$$

where $x = (x_1, \dots, x_n)^T \in \mathbb{R}^n$, $c = (c_1, \dots, c_n)^T \in \mathbb{R}^n$, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$, prompts the following Fenchel-Lagrange dual problem

$$(D_2) \quad \sup_{q^l \in \mathbb{R}_+^m, q^H > 0} \left\{ \langle b, q^l \rangle - q^H \ln \sum_{i=1}^n \exp((A^T q^l - c)_i / q^H) + q^H H \right\},$$

almost identical to the dual problem found in Ref. 14. The difference is that the interval variable q^H lies in $\mathbb{R}_+ \setminus \{0\}$ instead of \mathbb{R}_+ . This does not affect the optimal objective value of the dual problem. We have denoted the i -th entry of the vector $A^T q^l - c$ by $(A^T q^l - c)_i$. With the help of the constraint qualification

$$(CQ_2) \quad \exists x' \in \text{int}(\mathbb{R}_+^n) : H + \sum_{i=1}^n x'_i \ln x'_i < 0, b - Ax' \leq 0 \text{ and } \sum_{i=1}^n x'_i = 1,$$

the strong duality affirmation is ready to be formulated.

Theorem 4.3. If the constraint qualification (CQ_2) is satisfied, then the strong

duality between (P_2) and (D_2) is assured.

The optimality conditions are available in Ref. 15.

4.3. Facility location problem (Ref. 12). In the original paper the authors calculate the geometric duals for some problems involving norms. We have chosen one of them to be presented here, namely

$$(P_3) \quad \inf_{\|x-a_j\| \leq d_j, j=1, \dots, m} \left\{ \sum_{j=1}^m w_j \|x - a_j\| \right\},$$

where $a_j \in \mathbb{R}^n$, $w_j > 0$, $d_j > 0$, for $j = 1, \dots, m$. The Fenchel-Lagrange dual problem is (applying Theorem 20.1 in Ref. 17)

$$(D_3) \quad \sup_{\substack{p^j \in \mathbb{R}^n, \\ \sum_{j=1}^m p^j = p, \\ q \in \mathbb{R}_+^m}} \left\{ - \sum_{j=1}^m \left(w_j \| \cdot - a_j \| \right)^*(p^j) + \inf_{x \in \mathbb{R}^n} \left[\langle p, x \rangle + \sum_{j=1}^m q_j (\|x - a_j\| - d_j) \right] \right\}.$$

Knowing that, for $j = 1, \dots, m$, $(w_j \| \cdot - a_j \|)^*(p^j) = \langle a_j, p^j \rangle$, if $\|p^j\| \leq w_j$,

being $+\infty$ otherwise and turning the infimum into supremum, we get the following equivalent formulation of the dual problem, given also in Ref. 12,

$$(D_3) \quad \sup_{\substack{p^j \in \mathbb{R}^n, r^j \in \mathbb{R}^n, \|p^j\| \leq w_j, \|r^j\| \leq q_j, j=1, \dots, m, \\ \sum_{j=1}^m (p^j + r^j) = 0, q \in \mathbb{R}_+^m}} \left\{ - \langle q, d \rangle - \sum_{j=1}^m \langle a_j, p^j + r^j \rangle \right\}.$$

Of course, we have set here $d = (d_1, \dots, d_m)^T \in \mathbb{R}^m$ and $q = (q_1, \dots, q_m)^T \in \mathbb{R}^m$. A

sufficient background for the existence of strong duality is in this case

$$(CQ_3) \quad \exists x' \in \mathbb{R}^n : \|x' - a_j\| < d_j, j = 1, \dots, m.$$

Theorem 4.4. If the constraint qualification (CQ_3) is satisfied, then the strong duality between (P_3) and (D_3) is assured.

Although there is no mention of the optimality conditions in Ref. 12 for this pair of dual problems we have derived the following result.

Theorem 4.5. (a) If the constraint qualification (CQ_3) is fulfilled and \bar{x} is an optimal solution to (P_3), then strong duality between the problems (P_3) and (D_3) is attained and the dual problem has an optimal solution $(\bar{p}^1, \dots, \bar{p}^m, \bar{r}^1, \dots, \bar{r}^m, \bar{q}_1, \dots, \bar{q}_m)$ satisfying the following optimality conditions

(i) $w_j \|\bar{x} - a_j\| = \langle \bar{p}^j, \bar{x} - a_j \rangle$ and $\|\bar{p}^j\| = w_j$ when $\bar{x} \neq a_j$, $j = 1, \dots, m$.

(ii) $\bar{q}_j \|\bar{x} - a_j\| = \langle \bar{r}^j, \bar{x} - a_j \rangle$, $\bar{q}_j \geq 0$, $j = 1, \dots, m$, and if $\bar{q}_j > 0$ so is $\|\bar{r}^j\| = \bar{q}_j$. For $\bar{q}_j = 0$ there is also $\bar{r}^j = 0$. If $\bar{x} = a_j$ for any $j \in \{1, \dots, m\}$, then $\bar{q}_j = 0$ and $\bar{r}^j = 0$.

(iii) $\|\bar{x} - a_j\| = d_j$, for $j \in \{1, \dots, m\}$ such that $\bar{q}_j > 0$.

(iv) $\sum_{j=1}^m (\bar{p}^j + \bar{r}^j) = 0$.

(b) Having a feasible solution \bar{x} to the primal problem and one to the dual $(\bar{p}^1, \dots, \bar{p}^m, \bar{r}^1, \dots, \bar{r}^m, \bar{q}_1, \dots, \bar{q}_m)$ satisfying the optimality conditions (i)-(iv), then the mentioned feasible solutions turn out to be optimal solutions to the corresponding problems and strong duality holds.

4.4. **Quadratic concave fractional programs (Ref. 9).** Another problem

artificially pressed into the selective framework of geometric programming is

$$(P_4) \quad \inf_{Cx \leq b} (Q(x)/f(x)),$$

where $Q(x) = (1/2)x^T Ax$, $x \in \mathbb{R}^n$, $A \in \mathbb{R}^{n \times n}$ is a symmetric positive definite matrix,

$C \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$ and $f : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ a concave function, strictly positive over the

feasible set of the problem. Because no analytic representation of the conjugate of the

objective function is available, the problem is rewritten as

$$(P_4) \quad \inf_{\substack{sQ((1/s)x) - f(x) \leq 0, \\ Cx \leq b, s \in \mathbb{R}_+ \setminus \{0\}}} s.$$

To compute the Fenchel-Lagrange dual problem, we need first the conjugate of the

objective function $u : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$, $u(x, s) = s$. Using the result presented for the

same objective function in section 4.1, the dual problem becomes

$$(D_4) \quad \sup_{\substack{q^x \in \mathbb{R}_+^m, \\ q^s \in \mathbb{R}_+}} \left\{ \inf_{\substack{x \in \mathbb{R}^n, \\ s > 0}} \left[s + q^s \left(sQ((1/s)x) - f(x) \right) + \langle q^x, Cx - b \rangle \right] \right\}.$$

The infimum over (x, s) , transformed into a supremum, can be viewed as a conju-

gate function that is determined after some standard calculations. The formula that

results for the dual problem is identical to the geometric dual obtained in Ref. 9,

$$(D_4) \quad \sup_{\substack{q^x \in \mathbb{R}_+^m, q^s \in \mathbb{R}_+, \\ (1/2)u^T A^{-1}u \leq q^s, \\ u+v = -C^T q^x}} \left\{ -b^T q^x - (-q^s f)^*(v) \right\},$$

moreover simplifiable even to

$$(D_4) \quad \sup_{\substack{q^x \in \mathbb{R}_+^m, q^s \in \mathbb{R}_+, \\ (1/2)(-v - C^T q^x)^T A^{-1}(-v - C^T q^x) \leq q^s}} \left\{ -b^T q^x - (-q^s f)^*(v) \right\}.$$

Of course we have removed the assumption of closedness that has been imposed on the function f before. Because of the linearity of the constraints of (P_4) no constraint qualification is required in this case.

Theorem 4.6. Provided that the primal problem has at least a feasible point, strong duality between problems (P_4) and (D_4) is assured.

The corresponding optimality conditions are the same as in Ref. 9, therefore we omit them.

4.5. Sum of convex ratios (Ref. 10). An extension to vector optimization of the problem treated here can be found in Ref. 21. Here we consider as primal problem

$$(P_5) \quad \min_{Cx \leq b} \left\{ h(x) + \sum_{i=1}^N (f_i^2(x)/g_i(x)) \right\},$$

where $f_i, h : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ are convex functions, $g_i : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ concave, $f_i(x) \geq 0, g_i(x) > 0, i = 1, \dots, N$, for all x feasible to (P_5) , $b \in \mathbb{R}^m, C \in \mathbb{R}^{m \times n}$. Following the steps in Ref.

21 the Fenchel-Lagrange dual to (P_5) is

$$(D_5) \quad \sup_{\substack{q^x \in \mathbb{R}_+^m, q^s, q^t \in \mathbb{R}_+^N, \\ q_i^t \geq (q_i^s/2)^2, a_i, d_i, p^x \in \mathbb{R}^n, i=1, \dots, N, \\ \sum_{i=1}^N (a_i + d_i) = -p^x - C^T q^x}} \left\{ -h^*(p^x) - \sum_{i=1}^N (q_i^s f_i)^*(a_i) - \sum_{i=1}^N (-q_i^t g_i)^*(d_i) - \langle q^x, b \rangle \right\}.$$

Writing the homogenous extensions of the conjugate functions one gets the dual prob-

lem obtained in the original paper. Let us stress that we have ignored the hypotheses of closedness associated to the functions f , g and h in Ref. 10, as strong duality is valid even in their absence.

Theorem 4.7. Provided that the primal problem (P_5) has a finite optimal objective value, strong duality between problems (P_5) and (D_5) is assured.

The optimality conditions may be derived as in Ref. 21.

4.6. Quasiconcave multiplicative programs (Ref. 2). Despite its intricateness, geometric programming seems to be still very popular, as its direct applications still get published. The latest we could find is on a class of quasiconcave multiplicative programs that originally look like

$$(P_6) \quad \sup_{Ax \leq b} \left\{ \prod_{i=1}^k [f_i(x)]^{a_i} \right\},$$

with $f_i : \mathbb{R}^n \rightarrow \overline{\mathbb{R}}$ concave functions, positive over the feasible set of the problem, $a_i > 0$, $i = 1, \dots, k$, $A \in \mathbb{R}^{m \times n}$, $b \in \mathbb{R}^m$. The problem is brought into another layout in order to be properly treated,

$$(\tilde{P}_6) \quad \inf_{f_i(x) \geq s_i, i=1, \dots, k, Ax \leq b, s \in \text{int}(\mathbb{R}_+^k)} \left\{ - \sum_{i=1}^k a_i \ln s_i \right\}.$$

Denoting $f = (f_1, \dots, f_k)^T$, $s = (s_1, \dots, s_k)^T$ and introducing $u : \mathbb{R}^n \times \mathbb{R}^k \rightarrow \overline{\mathbb{R}}$ defined by

$$u(x, s) = - \sum_{i=1}^k a_i \ln s_i, \text{ if } (x, s) \in \mathbb{R}^n \times \text{int}(\mathbb{R}_+^k), \text{ otherwise being } +\infty, \text{ the raw formula}$$

of the Fenchel-Lagrange dual to (\tilde{P}_6) is

$$(\tilde{D}_6) \quad \sup_{\substack{p^x \in \mathbb{R}^n, p^s \in \mathbb{R}^k, \\ q^l \in \mathbb{R}_+^m, q^f \in \mathbb{R}_+^k}} \left\{ -u^*(p^x, p^s) + \inf_{\substack{x \in \mathbb{R}^n, \\ s \in \text{int}(\mathbb{R}_+^k)}} \left[\langle p^x, x \rangle + \langle p^s, s \rangle + \langle q^l, Ax - b \rangle + \langle q^f, s - f(x) \rangle \right] \right\}.$$

The conjugate of the objective function at $p^s = (p_1^s, \dots, p_k^s)^T$ is $u^*(p^x, p^s) = -\sum_{i=1}^k a_i \left(1 - \ln(a_i / (-p_i^s)) \right)$, if $p^x = 0, p^s < 0$ and $+\infty$, otherwise. The infimum in the dual problem

can also be separated into a sum of two infima, one concerning $s \in \text{int}(\mathbb{R}_+^k)$, the other $x \in \mathbb{R}^n$. Let us write again the dual using the last observations and denoting $-p_i^s$ by

$p_i^s, i = 1, \dots, k,$

$$(\tilde{D}_6) \quad \sup_{\substack{p^s \in \text{int}(\mathbb{R}_+^k), \\ q^l \in \mathbb{R}_+^m, q^f \in \mathbb{R}_+^k}} \left\{ \sum_{i=1}^k a_i \left(1 - \ln(a_i / p_i^s) \right) + \inf_{s \in \text{int}(\mathbb{R}_+^k)} \langle q^f - p^s, s \rangle \right. \\ \left. + \inf_{x \in \mathbb{R}^n} \left[\langle q^l, Ax \rangle - \langle q^f, f(x) \rangle \right] - \langle q^l, b \rangle \right\}.$$

The infimum regarding s is equal to 0 when $q^f - p^s \geq 0$, otherwise being $-\infty$, while the one over $x \in \mathbb{R}^n$ can be rewritten as a supremum and viewed as a conjugate of a sum of functions. The dual problem yields

$$(\tilde{D}_6) \quad \sup_{\substack{p^s \in \text{int}(\mathbb{R}_+^k), q^f \geq p^s, \\ q^l \in \mathbb{R}_+^m, \sum_{i=1}^k v_i = -A^T q^l}} \left\{ \sum_{i=1}^k \left(a_i - a_i \ln(a_i / p_i^s) \right) - \sum_{i=1}^k q_i^f (-f_i)^* \left((1/q_i^f) v_i \right) - \langle b, q^l \rangle \right\},$$

where $q^f = (q_1^f, \dots, q_k^f)^T$, and as the supremum regarding the variable p^s can be easily computed, being attained for $p^s = -q^f$, we get the following final version of the dual, equivalent to the one found in Ref. 2,

$$(\tilde{D}_6) \quad \sup_{\substack{q^l \in \mathbb{R}_+^m, q^f \in \text{int}(\mathbb{R}_+^k), \\ \sum_{i=1}^k v_i = -A^T q^l}} \left\{ \sum_{i=1}^k \left(a_i - a_i \ln(a_i/q_i^f) \right) - \sum_{i=1}^k q_i^f (-f_i)^* \left((1/q_i^f) v_i \right) - \langle b, q^l \rangle \right\}.$$

For strong duality a constraint qualification would normally be required because within the constraints of (\tilde{P}_6) there are affine as well as non-affine functions. But when the problem (P_6) has a non-empty feasible set, the existence of a feasible point x' so that $f_i(x') > 0, i = 1, \dots, k$, is assured and there is also an $s' > 0$ so that $f_i(x') > s' > 0, i = 1, \dots, k$. So the constraint qualification that comes from the general case is automatically fulfilled for (\tilde{P}_6) . Without any additional assumption, such as closedness, required for the functions $f_i, i = 1, \dots, k$, in the cited paper, one may formulate the strong duality statement.

Theorem 4.8. Provided that the primal problem has a feasible point, strong duality between problems (\tilde{P}_6) and (\tilde{D}_6) is assured.

The optimality conditions concerning this pair of dual problems follow.

Theorem 4.9. (a) If the problem (\tilde{P}_6) has an optimal solution \bar{x} , then strong duality between the problems (\tilde{P}_6) and (\tilde{D}_6) is attained and the dual problem has an optimal solution $(\bar{v}_1, \dots, \bar{v}_k, \bar{q}^l, \bar{q}^f)$ satisfying the following optimality conditions

$$(i) \quad (-f_i)^* \left((1/\bar{q}_i^f) \bar{v}_i \right) - f_i(\bar{x}) = \left\langle (1/\bar{q}_i^f) \bar{v}_i, \bar{x} \right\rangle, i = 1, \dots, k,$$

$$(ii) \quad \langle A^T \bar{x} - b, \bar{q}^l \rangle = 0,$$

$$(iii) \sum_{i=1}^k \bar{v}_i = -\bar{A}^T q^l,$$

$$(iv) \ln(f_i(\bar{x})) + \left(\bar{q}_i^f/a_i\right)f_i(\bar{x}) = \ln\left(a_i/\bar{q}_i^f\right) - 1, i = 1, \dots, k.$$

(b) Having a feasible solution \bar{x} to the primal problem and one to the dual $(\bar{v}_1, \dots, \bar{v}_k, \bar{q}^l, \bar{q}^f)$ satisfying the optimality conditions (i)-(iv), they turn out to be optimal solutions to the corresponding problems and strong duality holds.

4.7. Posynomial geometric programming (Ref. 23).

We are going to prove now that also the posynomial geometric programming duality can be viewed as a special case of the Fenchel-Lagrange duality. As it has been already proved (cf. Ref. 1) that the generalized geometric programming includes the posynomial instance as special case, our result is not so surprising within the framework of this paper. The primal-dual pair of posynomial geometric problems is composed by

$$(P_7) \quad \inf_{t \in \mathbb{R}_+, g_j(t) \leq 1, j=1, \dots, s} g_0(t),$$

$$\text{where } g_k(t) = \sum_{i \in J[k]} c_i \prod_{j=1}^m t^{a_{ij}}, k = 0, \dots, s, a_{ij} \in \mathbb{R}, j = 1, \dots, m, c_i > 0, i = 1, \dots, n, J[k]$$

$$= \{m_k, m_k + 1, \dots, n_k\}, k = 0, \dots, s, m_0 = 1, m_1 = n_0 + 1, \dots, m_k = n_{k-1} + 1, \dots, n_s = n,$$

and

$$(D_7) \quad \sup_{\delta \in \mathbb{R}_+, \sum_{i \in J[0]} \delta_i = 1, \sum_{i=1}^n \delta_i a_{ij} = 0, j=1, \dots, m} \left[\prod_{i=1}^n (c_i/\delta_i)^{\delta_i} \right] \prod_{k=1}^s \lambda_k(\delta)^{\lambda_k(\delta)},$$

where $\lambda_k(\delta) = \sum_{i \in J[k]} \delta_i, k = 1, \dots, s$. The primal posynomial problem is equivalent to

$$(\tilde{P}_7) \quad \inf_{\substack{\ln \left(\sum_{i \in J[k]} c_i \exp(x_i) \right) \leq 0, k=1, \dots, s, x \in \mathcal{A}}} \left\{ \ln \left(\sum_{i \in J[0]} c_i \exp(x_i) \right) \right\},$$

where \mathcal{A} denotes the linear subspace generated by the columns of the exponent matrix

(a_{ij}) (cf. Refs. 1 and 23). Let us name also the primal objective function $u(x)$. We

determine the Fenchel-Lagrange dual problem to (\tilde{P}_7) from the formula of (D'_K) , with

\mathcal{A}^\perp indicating the orthogonal subspace of \mathcal{A} , for $q = (q_1, \dots, q_s)^T$ and $t = (t_1, \dots, t_n)^T$

$$(\tilde{D}_7) \quad \sup_{p \in \mathbb{R}^n, t \in \mathcal{A}^\perp, q \in \mathbb{R}_+^s} \left\{ -u^*(p) + \inf_{x \in \mathcal{A}} \left[\langle p - t, x \rangle + \sum_{k=1}^s q_k \ln \left(\sum_{i \in J[k]} c_i \exp(x_i) \right) \right] \right\}.$$

For the conjugate of the objective function we have $u^*(p) = \sum_{i \in J[0]} p_i \ln(p_i/c_i)$, if

$p_j = 0, j \in J[k], k = 1, \dots, s, p = (p_1, \dots, p_n)^T \in \mathbb{R}_+^n$ and $\sum_{i \in J[0]} p_i = 1$. Otherwise its

value is $+\infty$. Similar results can be derived if we write the infimum within the dual

as a sum of suprema over $(x_i)_{i \in J[k]}, k = 1, \dots, s$, just with the changed constraints

$\sum_{i \in J[k]} t_i = q_k$. Also there follows $p_i = t_i, i \in J[0]$. Like in entropy optimization we

consider $0 \ln(0/c_i) = 0, c_i > 0, i = 1, \dots, n$. After these, the dual problem becomes

$$(\tilde{D}_7) \quad \sup_{t \in \mathcal{A}^\perp, q \in \mathbb{R}_+^s, t \geq 0, \sum_{i \in J[0]} t_i = 1, \sum_{i \in J[k]} t_i = q_k, k=1, \dots, s} \left\{ \sum_{i=1}^n t_i \ln(c_i/t_i) + \sum_{k=1}^s q_k \ln q_k \right\},$$

that is equivalent to the problem (D_7) , namely they have the same optimal solution

and the objective function of (D_7) is the exponential of the objective function of (\tilde{D}_7) .

Finally, the condition that guarantees strong duality, derived from the constraint

qualification (CQ) , is the so-called superconsistency introduced in Ref. 23, i.e.

$$(CQ_7) \quad \exists t' > 0 : g_k(t') < 1, k = 1, \dots, s.$$

Theorem 4.10. If the constraint qualification (CQ_7) is satisfied, then the strong duality between (P_7) and (D_7) is assured.

We present also the optimality conditions concerning this pair of problems.

Theorem 4.11. If the constraint qualification (CQ_7) is valid and \bar{x} is an optimal solution to (P_7) , then strong duality holds between the problems (P_7) and (D_7) and the dual has an optimal solution (\bar{t}, \bar{q}) satisfying the following optimality conditions

- (i) $\ln \left\{ \sum_{i \in J[0]} c_i \exp(\bar{x}_i) \right\} + \sum_{i \in J[0]} \bar{t}_i \ln(\bar{t}_i/c_i) = \langle \bar{t}^{J[0]}, \bar{x}^{J[0]} \rangle,$
- (ii) $\bar{q}_k \ln \left\{ \sum_{i \in J[k]} c_i \exp(\bar{x}_i) \right\} + \sum_{i \in J[k]} \bar{t}_i \ln(\bar{t}_i/c_i) - \bar{q}_k \ln \bar{q}_k = \langle \bar{t}^{J[k]}, \bar{x}^{J[k]} \rangle, k = 1, \dots, s,$
- (iii) $\sum_{i \in J[k]} c_i \exp(\bar{x}_i) = 1$ when $\bar{q}_k > 0, k = 1, \dots, s,$
- (iv) $\langle \bar{x}, \bar{t} \rangle = 0,$

where $x^{J[k]} = (x_{m_k}, \dots, x_{n_k})^T$ and $t^{J[k]} = (t_{m_k}, \dots, t_{n_k})^T, k = 0, \dots, s.$

(b) Having a feasible solution \bar{x} to the primal problem and one to the dual (\bar{t}, \bar{q}) satisfying the optimality conditions (i)-(iv), they turn out to be optimal solutions to the corresponding problems and strong duality holds.

5. Conclusions

Let us sum up now the main results in this paper. First we have introduced

a new duality approach, named Fenchel-Lagrange duality because of the way it is established and of the properties it has. Strong duality and optimality conditions for a general convex optimization primal problem and its Fenchel-Lagrange dual are presented, provided that a constraint qualification is valid. Then we prove that the geometric duality used by Scott and Jefferson in many of their papers turns out to be a special case of our Fenchel-Lagrange duality. In these papers the mentioned authors present the geometric dual problem to the primal under assumptions of convexity and closedness concerning the functions and the sets involved there. We established the same dual problem to the primal exploiting the Fenchel-Lagrange duality we presented earlier. Strong duality and optimality conditions are revealed to stand in much weaker circumstances, i. e. the closedness can be cancelled from the initial assumptions, while the constraint qualification used by the mentioned authors can be generalized and weakened, respectively. We review some convex optimization problems treated by the cited authors by means of geometric programming duality, showing how their duals can be obtained easier and more directly via Fenchel-Lagrange duality. Strong duality is accomplished under weaker sufficient conditions than in the original papers, namely the closedness assumptions for the functions and sets involved can be removed.

References

1. SCOTT, C. H., and JEFFERSON, T. R., *Avenues of Geometric Programming I. Theory*, New Zealand Operational Research, Vol. 6, No. 2, pp. 109–136, 1978.
2. SCOTT, C. H., and JEFFERSON, T. R., *On Duality for a Class of Quasiconcave Multiplicative Programs*, Journal of Optimization Theory and Applications, Vol. 117, No. 3, pp. 575–583, 2003.
3. PETERSON, E. L., *Geometric Programming*, SIAM Review, Vol. 18, pp. 1–51, 1976.
4. JEFFERSON, T. R., and SCOTT, C. H., *Quadratic Geometric Programming with Application to Machining Economics*, Mathematical Programming, Vol. 31, pp. 137–152, 1985.
5. JEFFERSON, T. R., and SCOTT, C. H., *Quality Tolerancing and Conjugate Duality*, Annals of Operations Research, Vol. 105, pp. 185–200, 2001.
6. JEFFERSON, T. R., WANG, Y. P., and SCOTT, C. H., *Composite Geometric Programming*, Journal of Optimization Theory and Applications, Vol. 64, No. 1, pp. 101–118, 1990.

7. SCOTT, C. H., and JEFFERSON, T. R., *Avenues of Geometric Programming II. Applications*, New Zealand Operational Research, Vol.7, No. 1, pp. 51–68, 1979.
8. SCOTT, C. H., and JEFFERSON, T. R., *Conjugate Duality in Generalized Fractional Programming*, Journal of Optimization Theory and Applications, Vol. 60, No. 3, pp. 475–483, 1989.
9. SCOTT, C. H., and JEFFERSON, T. R., *Convex Dual for Quadratic Concave Fractional Programs*, Journal of Optimization Theory and Applications, Vol. 91, No. 1, pp. 115–122, 1996.
10. SCOTT, C. H., and JEFFERSON, T. R., *Duality for a Sum of Convex Ratios*, Optimization, Vol. 40, pp. 303–312, 1997.
11. SCOTT, C. H., and JEFFERSON, T. R., *Duality for Minmax Programs*, Journal of Mathematical Analysis and Applications, Vol. 100, pp. 385–393, 1984.
12. SCOTT, C. H., JEFFERSON, T. R., and JORJANI, S., *Conjugate Duality in Facility Location*, Facility Location: A Survey of Applications and Methods, Edited by Z. Drezner, Springer Verlag, Heidelberg, Germany, pp. 89–101, 1995.

13. SCOTT, C. H., JEFFERSON, T. R., and JORJANI, S., *Duality for Programs Involving Non-Separable Objective Functions*, International Journal of Systems Science, Vol. 19, No. 9, pp. 1911–1918, 1988.
14. SCOTT, C. H., JEFFERSON, T. R., and JORJANI, S., *On Duality for Entropy Constrained Programs*, Information Sciences, Vol. 35, pp. 211–221, 1985.
15. BOŢ, R. I., GRAD, S. M., and WANKA, G., *Entropy Constrained Programs and Geometric Duality obtained via Fenchel-Lagrange Duality Approach*, Nonlinear Analysis Forum, Vol. 9, No. 1, pp. 65–85, 2004.
16. EKELAND, I., and TEMAM, R., *Convex Analysis and Variational Problems*, North-Holland Publishing Company, Amsterdam, The Netherlands, 1976.
17. ROCKAFELLAR, R. T., *Convex Analysis*, Princeton University Press, Princeton, New Jersey, 1970.
18. BOŢ, R. I., KASSAY, G., and WANKA, G., *Strong Duality for Generalized Convex Optimization Problems*, to appear in Journal of Optimization Theory and Applications, 2005.
19. WANKA, G., and BOŢ, R. I., *A New Duality Approach for Multiobjective Con-*

- Convex Optimization Problems*, Journal of Nonlinear and Convex Analysis, Vol. 3, No. 1, pp. 41–57, 2002.
20. WANKA, G., and BOŢ, R. I., *Multiobjective Duality for Convex-Linear Problems II*, Mathematical Methods of Operations Research, Vol. 53, No. 3, pp. 419–433, 2001.
21. WANKA, G., and BOŢ, R. I., *Multiobjective Duality for Convex Ratios*, Journal of Mathematical Analysis and Applications, Vol. 275, No. 1, pp. 354–368, 2002.
22. WANKA, G., and BOŢ, R. I., *On the Relations Between Different Dual Problems in Convex Mathematical Programming*, Operations Research Proceedings 2001, Edited by P. Chamoni, R. Leisten, A. Martin, J. Minnemann and H. Stadtler, Springer-Verlag, Heidelberg, Germany, pp. 255–262, 2002.
23. DUFFIN, R. J., PETERSON, E. L., and ZENER, C., *Geometric Programming - Theory and Applications*, John Wiley and Sons, New York, New York, 1967.
24. ELSTER, K. -H., REINHARDT, R., SCHÄUBLE, M., and DONATH, G., *Einführung in die nichtlineare Optimierung*, Mathematisch - Naturwissenschaftliche Bibliothek, BSB B.G. Teubner Verlagsgesellschaft, Leipzig, Germany, 1977.