

ON THE NUMERICAL EVALUATION OF FREDHOLM DETERMINANTS

FOLKMAR BORNEMANN*

Abstract. Some significant quantities in mathematics and physics are most naturally expressed as the Fredholm determinant of an integral operator, most notably many of the distribution functions in random matrix theory. Though their numerical values are of interest, there is no systematic numerical treatment of Fredholm determinants to be found in the literature. Instead, the few numerical evaluations that are available rely on eigenfunction expansions of the operator, if expressible in terms of special functions, or on alternative, numerically more straightforwardly accessible analytic expressions, e.g., in terms of Painlevé transcendents, that have masterfully been derived in some cases. In this paper we close the gap in the literature by studying projection methods and, above all, a simple, easily implementable, general method for the numerical evaluation of Fredholm determinants that is derived from the classical Nyström method for the solution of Fredholm equations of the second kind. Using Gauss–Legendre or Clenshaw–Curtis as the underlying quadrature rule, we prove that the approximation error essentially behaves like the quadrature error for the sections of the kernel. In particular, we get exponential convergence for analytic kernels, which are typical in random matrix theory. The application of the method to the distribution functions of the Gaussian unitary ensemble (GUE), in the bulk and the edge scaling limit, is discussed in detail. After extending the method to systems of integral operators, we evaluate the two-point correlation functions of the more recently studied Airy and Airy₁ processes.

Key words. Fredholm determinant, Nyström’s method, projection method, trace class operators, random matrix theory, Tracy–Widom distribution, Airy and Airy₁ processes

AMS subject classifications. 65R20, 65F40, 47G10, 15A52

1. Introduction. Fredholm’s (1903) landmark paper¹ on linear integral equations is generally considered to be the forefather of those mathematical concepts that finally led to modern functional analysis and operator theory—see the historical accounts in Kline (1972, pp. 1058–1075) and Dieudonné (1981, Chap. V). Fredholm was interested in the solvability of what is now called a Fredholm equation of the second kind,

$$u(x) + z \int_a^b K(x, y)u(y) dy = f(x) \quad (x \in (a, b)), \quad (1.1)$$

and explicit formulas for the solution thereof, for a right hand side f and a kernel K , both assumed to be continuous functions. He introduced his now famous determinant

$$d(z) = \sum_{k=0}^{\infty} \frac{z^k}{k!} \int_a^b \cdots \int_a^b \det (K(t_p, t_q))_{p, q=1}^n dt_1 \cdots dt_n, \quad (1.2)$$

which is an entire function of $z \in \mathbb{C}$, and succeeded in showing that the integral equation is uniquely solvable if and only if $d(z) \neq 0$.

Realizing the tremendous potential of Fredholm’s theory, Hilbert started working on integral equations in a flurry and, in a series of six papers from 1904 to 1910,²

*Zentrum Mathematik, Technische Universität München, Boltzmannstr. 3, 85747 Garching, Germany (bornemann@ma.tum.de). Manuscript as of June 24, 2008.

¹Simon (2005, p. VII) writes: “This deep paper is extremely readable and I recommend it to those wishing a pleasurable afternoon.” An English translation of the paper can be found in Birkhoff (1973, pp. 449–465).

²Later reproduced as one of the first books on linear integral equations (Hilbert 1912).

transformed the determinantal framing to the beginnings of what later, in the hands of Schmidt, Carleman, Riesz, and others, would become the theory of compact operators in Hilbert spaces. Consequently, over the years Fredholm determinants have faded from the core of general accounts on integral equations to the historical remarks section—if they are mentioned at all.³

So, given this state of affairs, why then study the numerical evaluation of the Fredholm determinant $d(z)$? The reason is, simply enough, that the Fredholm determinant and the more general notions, generalizing (1.1) and (1.2) to

$$u + zAu = f, \quad d(z) = \det(I + zA),$$

for certain classes of compact operators A on Hilbert spaces, have always remained important tools in operator theory and mathematical physics (Gohberg, Goldberg and Krupnik 2000, Simon 2005). In turn, they have found many significant applications: e.g., in atomic collision theory (Jost and Pais 1951, Moiseiwitsch 1977), inverse scattering (Dyson 1976), in Floquet theory of periodic differential equations (Eastham 1973), in the infinite-dimensional method of stationary phase and Feynman path integrals (Albeverio and Høegh-Krohn 1977, Rezende 1994), as the two-point correlation function of the two-dimensional Ising model (Wilkinson 1978), in renormalization in quantum field theory (Simon 2005), as distribution functions in random matrix theory (Mehta 2004, Deift 1999, Katz and Sarnak 1999) and combinatorial growth processes (Johansson 2000, Prähofer and Spohn 2002, Sasamoto 2005, Borodin, Ferrari, Prähofer and Sasamoto 2007). As Lax (2002, p. 260) puts it most aptly upon including Fredholm’s theory as a chapter of its own in his recent textbook on functional analysis: “Since this determinant appears in some modern theories, it is time to resurrect it.”

In view of this renewed interest in operator determinants, what numerical methods are available for their evaluation? Interestingly, this question has apparently never—at least to our knowledge—been systematically addressed in the numerical analysis literature.⁴ Even experts in the applications of Fredholm determinants commonly seem to have been thinking (Spohn 2008) that an evaluation is only possible

³For example, Hilbert (1912), Webster (1927), and Whittaker and Watson (1927) start with the Fredholm determinant right from the beginning; yet already Courant and Hilbert (1953, pp.142–147), the translation of the German edition from 1931, give it just a short mention (“since we shall not make any use of the Fredholm formulas later on”); while Smithies (1958, Chap. V) and Hochstadt (1973, Chap. 7), acknowledging the fact that “classical” Fredholm theory yields a number of results that functional analytic techniques do not, postpone Fredholm’s theory to a later chapter; whereas Baker (1977), Porter and Stirling (1990), Prössdorf and Silbermann (1991), Hackbusch (1995), and Kress (1999) ignore the Fredholm determinant altogether. Among the newer books on linear integral equations, the monumental four volume work of Fenyő and Stolle (1982–1984) is one of the few we know of that give Fredholm determinants a balanced treatment.

⁴Though we can only speculate about the reasons, there is something like a disapproving attitude towards determinants in general that seems to be quite common among people working in “continuous” applied mathematics. Here are a few scattered examples: Meyer (2000) writes at the beginning of the chapter on determinants (p. 460): “Today matrix and linear algebra are in the main stream of applied mathematics, while the role of determinants has been relegated to a minor backwater position.” Axler (1995) has a paper with the provocative title “Down with Determinants!” and a correspondingly worked out textbook on linear algebra (1997). The quintessential book of Golub and Van Loan (1996) on “Matrix Computations” does not explicitly address the computation of determinants at all, it is only implicitly stated as part of Theorem 3.2.1. Higham (2002) writes at the beginning of Section 14.6: “Like the matrix inverse, the determinant is a quantity that rarely needs to be computed.” He then continues with the argument, well known to every numerical analyst, that the determinant cannot be used as a measure of ill conditioning since it scales as $\det(\alpha A) = \alpha^m \det(A)$ for a $m \times m$ -matrix A , $\alpha \in \mathbb{R}$. Certainly there

if either the eigenvalues of the integral operator are, more or less, explicitly known or if an alternative analytic expression has been found that is numerically more accessible—in each specific case anew, lacking a general procedure.

The Nyström-type method advocated in this paper. In contrast, we study a simple general numerical method for Fredholm determinants which is exceptionally efficient for smooth kernels, yielding small *absolute* errors (i.e., errors that are small with respect to the scale $\det(I) = 1$ inherently given by the operator determinant). To this end we follow the line of thought of Nyström's (1930) classical quadrature method for the numerical solution of the Fredholm equation (1.1). Namely, given a quadrature rule

$$Q(f) = \sum_{j=1}^m w_j f(x_j) \approx \int_a^b f(x) dx,$$

Nyström discretized (1.1) as the linear system

$$u_i + z \sum_{j=1}^m w_j K(x_i, x_j) u_j = f(x_i) \quad (i = 1, \dots, m), \quad (1.3)$$

which has to be solved for $u_i \approx u(x_i)$ ($i = 1, \dots, m$). Nyström's method is extremely simple and, yet, extremely effective for *smooth* kernels. So much so that Delves and Mohamed (1985, p. 245), in a chapter comparing different numerical methods for Fredholm equations of the second kind, write:

Despite the theoretical and practical care lavished on the more complicated algorithms, the clear winner of this contest has been the Nyström routine with the m -point Gauss–Legendre rule. This routine is extremely simple; it includes a call to a routine which provides the points and weights for the quadrature rule, about twelve lines of code to set up the Nyström equations and a call to the routine which solves these equations. Such results are enough to make a numerical analyst weep.

By keeping this conceptual and algorithmic simplicity, the method studied in this paper approximates the Fredholm determinant $d(z)$ simply by the determinant of the $m \times m$ -matrix that is applied to the vector (u_i) in the Nyström equation (1.3):

$$d_Q(z) = \det (\delta_{ij} + z w_i K(x_i, x_j))_{i,j=1}^m. \quad (1.4)$$

If the weights w_j of the quadrature rule are positive (which is always the better choice), we will use the *equivalent* symmetric variant

$$d_Q(z) = \det (\delta_{ij} + z w_i^{1/2} K(x_i, x_j) w_j^{1/2})_{i,j=1}^m. \quad (1.5)$$

is much truth in all of their theses, and Cramer's rule and the characteristic polynomial, which were the most common reasons for a call to the numerical evaluation of determinants (Stewart 1998, p. 176), have most righteously been banned from the toolbox of a numerical analyst for reasons of efficiency. However, with respect to the infinite dimensional case, the elimination of determinants from the thinking of numerical analysts as a subject of computations might have been all too successful. For instance, the scaling argument does not apply in the infinite dimensional case: operator determinants $\det(I + A)$ are defined for compact perturbations of the identity, which perfectly determines the scaling since, for $\alpha \neq 1$, $\alpha(I + A)$ cannot be written in the form $I + \tilde{A}$ with another compact operator \tilde{A} . (This is because the identity operator is *not* compact then.)

Using Gauss–Legendre or Curtis–Clenshaw quadrature rules, the computational cost⁵ of the method is of order $O(m^3)$. The implementation in Matlab, or Mathematica, is straightforward and takes just a few lines of code.⁶ In Matlab:

```
function d = DetNyström(K,z,a,b,m)
[w,x] = QuadratureRule(a,b,m);
w = sqrt(w);
[xj,xi] = meshgrid(x,x);
d = det(eye(m)+z*(w'*w).*K(xi,xj));
```

In Mathematica:

```
DetNyström[K_, z_, a_, b_, m_] :=
Module[{w, x}, {w, x} = QuadratureRule[a, b, m]; w = Sqrt[w];
Det[IdentityMatrix[m] + z Outer[Times, w, w] Outer[K, x, x]]];
```

Strictly speaking we are not the first to suggest this simple method, though. In fact, it was Hilbert himself (1904, pp. 52–60) in his very first paper on integral equations, who motivated⁷ the expression (1.2) of the Fredholm determinant by essentially using this method with the *rectangular rule* for quadrature, proving locally uniform convergence; see also Hilbert (1912, pp. 4–10) and, for the motivational argument given just heuristically, without a proof of convergence, Whittaker and Watson (1927, Sect. 11.2), Tricomi (1957, pp. 66–68) (who speaks of a “poetic license” to be applied “without too many scruples”), Smithies (1958, pp. 65–68), Hochstadt (1973, pp. 243–239), and Fenyő and Stolle (1982–1984, Vol. II, pp. 82–84)—to name just a few but influential cases. Quite astonishingly, despite of all its presence as a motivational tool in the expositions of the classical theory, we have found just one example of the use of this method (with Gauss–Legendre quadrature) in an actual numerical calculation: a paper by the physicists Reinhardt and Szabo (1970) on low-energy elastic scattering of electrons from hydrogen atoms. However, the error estimates (Theorem 6.2) that we will give in this paper seem to be new at least; we will prove that the approximation error essentially behaves like the quadrature error for the sections $x \mapsto K(x, y)$ and $y \mapsto K(x, y)$ of the kernel. In particular, we will get exponential convergence rates for analytic kernels.

Examples. Perhaps the generality and efficiency offered by our direct numerical approach to Fredholm determinants, as compared to analytic methods if they are

⁵The computational cost of $O(m^3)$ for the matrix determinant using either Gaussian elimination with partial pivoting (Stewart 1998, p. 176) or Hyman’s method (Higham 2002, Sect. 14.6) clearly dominates the cost of $O(m \log m)$ for the weights and points of Clenshaw–Curtis quadrature using the FFT, as well as the cost of $O(m^2)$ for Gauss–Legendre quadrature using the Golub–Welsh algorithm; for implementation details of these quadrature rules see Waldvogel (2006), Trefethen (2008), and the appendix of this paper. The latter paper carefully compares Clenshaw–Curtis with Gauss–Legendre and concludes (p. 84): “Gauss quadrature is a beautiful and powerful idea. Yet the Clenshaw–Curtis formula has essentially the same performance for most integrands.”

⁶The command `[w, x] = QuadratureRule(a, b, m)` is supposed to supply the weights and points of a m -point quadrature rule on the interval $[a, b]$ as a $1 \times m$ vector w and a $m \times 1$ -vector x , respectively. For Gauss–Legendre and Clenshaw–Curtis, such a Matlab code can be found in the appendix.

⁷Fredholm himself does not give the slightest hint of a motivation in his early papers (1900, 1903). He apparently conceived his determinant in *analogy* to similar expressions that his fellow countryman von Koch (1892) had obtained for infinite matrices a few years earlier; see Fredholm (1909, p. 95), Dieudonné (1981, p. 99), or Pietsch (2007, p. 409).

available at all, is best appreciated by an example. The probability $E_2(0; s)$ that an interval of length s does not contain, in the bulk scaling limit of level spacing 1, an eigenvalue of the Gaussian unitary ensemble (GUE) is given by the Fredholm determinant of the sine kernel,

$$E_2(0; s) = \det \left(I - A_s \downharpoonright_{L^2(0, s)} \right), \quad A_s u(x) = \int_0^s \frac{\sin(\pi(x-y))}{\pi(x-y)} u(y) dy;$$

see Gaudin (1961) and Mehta (2004, Sect. 6.3). Gaudin has further shown that the eigenfunctions of this selfadjoint integral operator are exactly given by a particular family of special functions, namely the radial prolate spheroidal wave functions with certain parameters. Using tables (Stratton, Morse, Chu, Little and Corbató 1956) of these functions he was finally able to evaluate $E_2(0; s)$ numerically. On the other hand, in an admirably intricate analytic tour de force Jimbo, Miwa, Mōri and Sato (1980) expressed the Fredholm determinant of the sine kernel as

$$E_s(0; s) = \exp \left(\int_0^{\pi s} \frac{\sigma(x)}{x} dx \right) \quad (1.6)$$

in terms of the sigma, or Hirota, representation of the fifth Painlevé equation, namely

$$(x\sigma'')^2 + 4(x\sigma' - \sigma)(x\sigma' - \sigma + \sigma'^2) = 0, \quad \sigma(x) \sim -\frac{x}{\pi} - \frac{x^2}{\pi^2} \quad (x \rightarrow 0),$$

see also Mehta (2004, Chap. 21) and Tracy and Widom (2000, Sect. 4.1). With respect to the numerical evaluation, the latter two authors conclude in a footnote, most probably by comparing to Gaudin's method: "Without the Painlevé representations, the numerical evaluation of the Fredholm determinants is quite involved." However, one does not need to know more than the smooth kernel $\sin(\pi(x-y))/(\pi(x-y))$ itself to approximate $E_2(0; s)$ with the method of this paper. For instance, the Gauss–Legendre rule with just $m = 5$ quadrature points already gives, in 0.2 ms computing time, 15 accurate digits of the value

$$E_2(0, 0.1) = 0.900027271798259 \dots,$$

that is, by calculating the determinant of a 5×5 -matrix easily built from the kernel.

Even though it is satisfying to have an alternative and simpler way of calculating already known quantities, it is far more exciting to be able to calculate quantities that otherwise have defeated numerical evaluations so far. For instance, the joint distribution functions of the Airy and the Airy_1 processes are given as determinants of systems of integral operators, see Prähofer and Spohn (2002), Johansson (2003), Sasamoto (2005) and Borodin et al. (2007). Even though a nonlinear partial differential equation of third order in three variables has been found by Adler and van Moerbeke (2005, Eq. (4.12)) for the logarithm of the joint distribution function of the Airy process at two different times, this masterful analytic result is probably of next to no numerical use. And in any case, no such analytic results are yet known for the Airy_1 process. However, the Nyström-type method studied in this paper can easily be extended to systems of integral operators. In this way, we have succeeded in evaluating the two-point correlation functions of both stochastic processes, see Section 8.

Outline of this paper. For the proper functional analytic setting, in Section 2 we review some basic facts about trace class and Hilbert–Schmidt operators. In Section 3 we review the concept of the determinant $\det(I + A)$ for trace class operators A and its relation to the Fredholm determinant. In Section 4 we study perturbation bounds implying that numerical calculations of determinants can only be expected to be accurate with respect to *absolute* errors in general. In Section 5 we use the functional analytic formulation of the problem to obtain convergence estimates for projection methods of Galerkin and Ritz–Galerkin type. The convergence rate is shown to depend on a proper balance between the decay of the singular values of the operator and the growth of bounds on the derivatives of the corresponding singular functions. This is in sharp contrast with Section 6, where we study the convergence of the Nyström-type method (1.5) by directly addressing the original definition of the Fredholm determinant. Here, only the smoothness properties of the kernel enter the convergence estimates. It turns out that, for kernels of low regularity, the order of convergence of the Nyström-type method can be even higher than that of a Ritz–Galerkin method. In Section 7 we give examples for the exponential convergence rates enjoyed by analytic kernels. To this end we discuss the details of the numerical evaluation of the determinants of the sine and Airy kernels, which express the probability distributions $E_2(0; s)$ and $F_2(s)$ (the Tracy–Widom distribution) of random matrix theory. Finally, after extending the Nyström-type method to systems of integral operators we report in Section 8 on the numerical evaluation of the two-point correlation functions of the Airy and Airy₁ processes.

2. Trace Class and Hilbert–Schmidt Operators. We begin by recalling some basic material about the spectral theory of nonselfadjoint compact operators, which can be found, e.g., in Gohberg, Goldberg and Kaashoek (1990), Lax (2002) and Simon (2005). We consider a complex, separable Hilbert space \mathcal{H} with an inner product $\langle \cdot, \cdot \rangle$ that is linear in the *second* factor and conjugate linear in the first. The set of bounded linear operators will be denoted by $\mathcal{B}(\mathcal{H})$, the compact operators by $\mathcal{J}_\infty(\mathcal{H})$. The spectrum of a compact operator $A \in \mathcal{J}_\infty(\mathcal{H})$ has no non-zero limit point; its non-zero points are eigenvalues of finite algebraic multiplicity. We list these eigenvalues as $(\lambda_n(A))_{n=1}^{N(A)}$, counting multiplicity, where $N(A)$ is either a finite non-negative integer or infinity, and order them by

$$|\lambda_1(A)| \geq |\lambda_2(A)| \geq \dots$$

The positive eigenvalues

$$s_1(A) \geq s_2(A) \geq \dots > 0$$

of the associated positive-semidefinite, selfadjoint operator

$$|A| = (A^* A)^{1/2}$$

are called the singular values of A . Correspondingly, there is the Schmidt or singular-value representation of A , that is, the norm convergent expansion

$$A = \sum_{n=1}^{N(|A|)} s_n(A) \langle u_n, \cdot \rangle v_n, \quad (2.1)$$

where the u_n and v_n are certain (not necessarily complete) orthonormal sets in \mathcal{H} . Note that $s_n(A) = |\lambda_n(A)|$ if A is selfadjoint. In general we have Weyl's inequality

$$\sum_{n=1}^N |\lambda_n(A)|^p \leq \sum_{n=1}^N s_n(A)^p \quad (N \leq N(A), 1 \leq p < \infty). \quad (2.2)$$

The Schatten–von Neumann classes of compact operators are defined as

$$\mathcal{J}_p(\mathcal{H}) = \{A \in \mathcal{J}_\infty(\mathcal{H}) : \sum_{n=1}^{N(|A|)} s_n(A)^p < \infty\} \quad (1 \leq p < \infty)$$

with the corresponding norm⁸

$$\|A\|_{\mathcal{J}_p} = \left(\sum_{n=1}^{N(|A|)} s_n(A)^p \right)^{1/p}.$$

The operator norm on $\mathcal{J}_\infty(\mathcal{H})$ perfectly fits into this setting if we realize that

$$\|A\| = s_1(A) = \max_{n=1, \dots, N(|A|)} s_n(A) = \|A\|_{\mathcal{J}_\infty}.$$

There are the continuous embeddings $\mathcal{J}_p(\mathcal{H}) \subset \mathcal{J}_q(\mathcal{H})$ for $1 \leq p \leq q \leq \infty$ with

$$\|A\|_{\mathcal{J}_q} \leq \|A\|_{\mathcal{J}_p}.$$

The classes $\mathcal{J}_p(\mathcal{H})$ are two-sided operator ideals in $\mathcal{B}(\mathcal{H})$, that is, for $A \in \mathcal{J}_p(\mathcal{H})$ ($1 \leq p \leq \infty$) and $B \in \mathcal{B}(\mathcal{H})$ we have $AB, BA \in \mathcal{J}_p(\mathcal{H})$ with

$$\|AB\|_{\mathcal{J}_p} \leq \|A\|_{\mathcal{J}_p} \|B\|, \quad \|BA\|_{\mathcal{J}_p} \leq \|B\| \|A\|_{\mathcal{J}_p}. \quad (2.3)$$

Of special interest to us are the *trace class operators* $\mathcal{J}_1(\mathcal{H})$ and the *Hilbert–Schmidt operators* $\mathcal{J}_2(\mathcal{H})$. The product of two Hilbert–Schmidt operators is of trace class:

$$\|AB\|_{\mathcal{J}_1} \leq \|A\|_{\mathcal{J}_2} \|B\|_{\mathcal{J}_2} \quad (A, B \in \mathcal{J}_2(\mathcal{H})).$$

The *trace* of a trace class operator A is defined by

$$\mathrm{tr}(A) = \sum_{n=1}^{\infty} \langle u_n, Au_n \rangle$$

for any orthonormal basis $(u_n)_n$. A deep theorem of Lidskii's (Simon 2005, Chap. 3) tells us that

$$\mathrm{tr}(A) = \sum_{n=1}^{N(A)} \lambda_n(A), \quad (2.4)$$

which implies by Weyl's inequality (2.2) that

$$|\mathrm{tr}(A)| \leq \sum_{n=1}^{N(A)} |\lambda_n(A)| \leq \mathrm{tr}(|A|) = \|A\|_{\mathcal{J}_1}. \quad (2.5)$$

⁸In matrix theory these norms are not commonly used—with the exception of $p = 2$: $\|A\|_{\mathcal{J}_2}$ is then the Schur or Frobenius norm of the matrix A .

Likewise, for a Hilbert–Schmidt operator $A \in \mathcal{J}_2(\mathcal{H})$ we have

$$\operatorname{tr}(A^2) = \sum_{n=1}^{N(A)} \lambda_n(A)^2, \quad |\operatorname{tr}(A^2)| \leq \sum_{n=1}^{N(A)} |\lambda_n(A)|^2 \leq \|A\|_{\mathcal{J}_2}^2. \quad (2.6)$$

Integral operators with L^2 -kernel. In the Hilbert space $\mathcal{H} = L^2(a, b)$ of square-integrable functions on a finite interval (a, b) the Hilbert–Schmidt operators are exactly given by the integral operators with L^2 -kernel. That is, there is a one-to-one correspondence (Simon 2005, Thm. 2.11) between $A \in \mathcal{J}_2(\mathcal{H})$ and $K \in L^2((a, b)^2)$ mediated through

$$Au(x) = \int_a^b K(x, y)u(y) dy \quad (2.7)$$

with equality of norms $\|A\|_{\mathcal{J}_2} = \|K\|_{L^2}$: the spaces $\mathcal{J}_2(\mathcal{H})$ and $L^2((a, b)^2)$ are thus isometrically isomorphic. In particular, by (2.6) and a well known basic result on infinite products (Knopp 1964, p. 232), we get for such operators that

$$\prod_{n=1}^{N(A)} (1 + \lambda_n(A)) \text{ converges (absolutely)} \Leftrightarrow \sum_{n=1}^{N(A)} \lambda_n(A) \text{ converges (absolutely)}.$$

Since the product is a natural candidate for the definition of $\det(I + A)$ it makes sense requiring A to be of trace class; for then, by (2.5), the absolute convergence of the sum can be guaranteed.

Integral operators with a continuous kernel. A continuous kernel $K \in C([a, b]^2)$ is certainly square-integrable. Therefore, the induced integral operator (2.7) defines a Hilbert–Schmidt operator A on the Hilbert space $\mathcal{H} = L^2(a, b)$. Moreover, other than for L^2 kernels in general, the integral

$$\int_a^b K(x, x) dx$$

over the diagonal of $(a, b)^2$ is now well defined and constitutes, in analogy to the matrix case, a “natural” candidate for the trace of the integral operator. Indeed, if an integral operator A with continuous kernel is of trace class, one can prove (Gohberg et al. 2000, Thm. 8.1)

$$\operatorname{tr}(A) = \int_a^b K(x, x) dx. \quad (2.8)$$

Unfortunately, however, just the continuity of the kernel K does not guarantee the induced integral operator A to be of trace class.⁹ Yet, there is some encouraging positive experience stated by Simon (2005, p. 25):

However, the counter-examples which prevent nice theorems from holding are generally rather contrived so that I have found the following to be true: If an integral operator with kernel K occurs in some ‘natural’ way and $\int |K(x, x)| dx < \infty$, then the operator can (almost always) be proven to be trace class (although sometimes only after some considerable effort).

⁹A counter-example was discovered by Carleman (1918), see also Gohberg et al. (2000, p. 71).

Nevertheless, we will state some simple criteria that often work well:

1. If the continuous kernel K can be represented in the form

$$K(x, y) = \int_c^d K_1(x, y)K_2(z, y) dz \quad (x, y \in [a, b])$$

with $K_1 \in L^2((a, b) \times (c, d))$, $K_2 \in L^2((c, d) \times (a, b))$, then the induced integral operator A is trace class on $L^2(a, b)$. This is, because A can then be written as the product of two Hilbert–Schmidt operators.

2. If $K(x, y)$ and $\partial_y K(x, y)$ are continuous on $[a, b]^2$, then the induced integral operator A is trace class on $L^2(a, b)$. This is, because we can write A by partial integration in the form

$$Au(x) = K(x, b) \int_a^b u(y) dy - \int_a^b \left(\int_y^b \partial_z K(x, z) dz \right) u(y) dy$$

as a sum of a rank one operator and an integral operator that is trace class by the first criterion. In particular, integral operators with smooth kernels are trace class (Lax 2002, p. 345).

3. A continuous Hermitian¹⁰ kernel $K(x, y)$ on $[a, b]$ that satisfies a Hölder condition in the second argument with exponent $\alpha > 1/2$, namely

$$|K(x, y_1) - K(x, y_2)| \leq C|y_1 - y_2|^\alpha \quad (x, y_1, y_2 \in [a, b]),$$

induces an integral operator A that is trace class on $L^2(a, b)$; see Gohberg et al. (2000, Thm. IV.8.2).

4. If the continuous kernel K induces a selfadjoint, positive-semidefinite integral operator A , then A is trace class (Gohberg et al. 2000, Thm. IV.8.3). The hypothesis on A is fulfilled for positive-semidefinite kernels K , that is, if

$$\sum_{j,k=1}^n \bar{z}_j z_k K(x_j, x_k) \geq 0 \quad (2.9)$$

for any $x_1, \dots, x_n \in (a, b)$, $z \in \mathbb{C}^n$ and any $n \in \mathbb{N}$ (Simon 2005, p. 24).

3. Definition and Properties of Fredholm and Operator Determinants. In this section we give a general operator theoretical definition of infinite dimensional determinants and study their relation to the Fredholm determinant. For a trace class operator $A \in \mathcal{J}_1(\mathcal{H})$ there are several equivalent constructions that all define one and the same *entire* function

$$d(z) = \det(I + zA) \quad (z \in \mathbb{C});$$

in fact, each construction has been chosen at least once, in different places of the literature, as the basic definition of the operator determinant:

¹⁰An L^2 -kernel K is Hermitian if $K(x, y) = \overline{K(y, x)}$. This property is equivalent to the fact that the induced Hilbert–Schmidt integral operator A is selfadjoint, $A^* = A$.

1. Gohberg and Kreĭn (1969, p. 157) define the determinant by the locally uniformly convergent (infinite) product

$$\det(I + zA) = \prod_{n=1}^{N(A)} (1 + z\lambda_n(A)), \quad (3.1)$$

which possesses zeros exactly at $z_n = -1/\lambda_n(A)$, counting multiplicity.

2. Gohberg et al. (1990, p. 115) define the determinant as follows. Given any sequence of finite rank operators A_n with $A_n \rightarrow A$ converging in trace class norm, the sequence of finite dimensional determinants¹¹

$$\det\left(I + zA_n \upharpoonright_{\text{ran}(A_n)}\right) \quad (3.2)$$

(which are polynomials in z) converges locally uniform to $\det(I + zA)$, independently of the choice of the sequence A_n . The existence of at least one such sequence follows from the singular value representation (2.1).

3. Dunford and Schwartz (1963, p. 1029) define the determinant by what is often called Plemelj's formula¹²

$$\det(I + zA) = \exp(\text{tr} \log(I + zA)) = \exp\left(-\sum_{n=1}^{\infty} \frac{(-z)^n}{n} \text{tr} A^n\right), \quad (3.3)$$

which converges for $|z| < 1/|\lambda_1(A)|$ and can analytically be continued as an entire function to all $z \in \mathbb{C}$.

4. Grothendieck (1956, p. 347) and Simon (1977, p. 254) define the determinant most elegantly with a little exterior algebra (Greub 1967). With $\wedge^n(A) \in \mathcal{J}_1(\wedge^n(\mathcal{H}))$ being the n^{th} exterior product of A , the power series

$$\det(I + zA) = \sum_{n=0}^{\infty} z^n \text{tr} \wedge^n(A) \quad (3.4)$$

converges for all $z \in \mathbb{C}$. Note that $\text{tr} \wedge^n(A) = \sum_{i_1 < \dots < i_n} \lambda_{i_1}(A) \cdots \lambda_{i_n}(A)$ is just the n^{th} symmetric function of the eigenvalues of A .

Proofs of the equivalence can be found in (Gohberg et al. 2000, Chap. 2) and (Simon 2005, Chap. 3). We will make use of all of them in the course of this paper. We state two important properties (Simon 2005, Thm. 3.5) of the operator determinant: First its multiplication formula,

$$\det(I + A + B + AB) = \det(I + B) \det(I + A) \quad (A, B \in \mathcal{J}_1(\mathcal{H})), \quad (3.5)$$

then the characterization of invertibility: $\det(I + A) \neq 0$ if and only if the inverse operator $(I + A)^{-1}$ exists.

¹¹Gohberg et al. (2000) have later extended this idea to generally define traces and determinants on embedded algebras of compact operators by a continuous extension from the finite dimensional case. Even within this general theory the trace class operators enjoy a most unique position: it is only for them that the values of trace and determinant are *independent* of the algebra chosen for their definition. On the contrary, if A is Hilbert–Schmidt but not trace class, by varying the embedded algebra, the values of the trace $\text{tr}(A)$ can be given *any* complex number and the values of the determinant $\det(I + A)$ are either always zero or can be made to take *any* value in the set $\mathbb{C} \setminus \{0\}$ (Gohberg et al. 2000, Chap. VII).

¹²Plemelj (1904, Eq. (62)) had given a corresponding form of the Fredholm determinant for integral operators. However, it can already be found in Fredholm (1903, p. 384).

The matrix case. In Section 6 we will study the convergence of finite dimensional determinants to operator determinants in terms of the power series (3.4). Therefore, we give this series a more common look and feel for the case of a matrix $A \in \mathbb{C}^{m \times m}$. By evaluating the traces with respect to a Schur basis of A one gets

$$\mathrm{tr} \bigwedge^n(A) = \sum_{i_1 < \dots < i_n} \det(A_{i_p, i_q})_{p, q=1}^n = \frac{1}{n!} \sum_{i_1, \dots, i_n=1}^m \det(A_{i_p, i_q})_{p, q=1}^n,$$

that is, the sum of all $n \times n$ principal minors of A . This yields the von Koch (1892) form of the matrix determinant

$$\det(I + zA) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^m \det(A_{i_p, i_q})_{p, q=1}^n \quad (A \in \mathbb{C}^{m \times m}). \quad (3.6)$$

(In fact, the series must terminate at $n = m$ since $\det(I + zA)$ is a polynomial of degree m in z .) A more elementary proof of this classical formula, by a Taylor expansion of the polynomial $\det(I + zA)$, can be found, e.g., in Meyer (2000, p. 495).

The Fredholm determinant for integral operators with continuous kernel. Suppose that the continuous kernel $K \in C([a, b]^2)$ induces an integral operator A that is trace class on the Hilbert space $\mathcal{H} = L^2(a, b)$. Then, the traces of $\bigwedge^n(A)$ evaluate to (Simon 2005, Thm. 3.10)

$$\mathrm{tr} \bigwedge^n(A) = \frac{1}{n!} \int_{(a, b)^n} \det(K(t_p, t_q))_{p, q=1}^n dt_1 \cdots dt_n \quad (n = 0, 1, 2, \dots).$$

The power series representation (3.4) of the operator determinant is therefore exactly Fredholm's expression (1.2), that is,

$$\det(I + zA) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \int_{(a, b)^n} \det(K(t_p, t_q))_{p, q=1}^n dt_1 \cdots dt_n. \quad (3.7)$$

The similarity with von Koch's formula (3.6) is striking and, in fact, it was just an analogy in form that had led Fredholm to conceive his expression for the determinant. It is important to note, however, that the right hand side of (3.7) perfectly makes sense for any continuous kernel, independently of whether the corresponding integral operator is trace class or not.

The regularized determinant for Hilbert–Schmidt operators. For a general Hilbert–Schmidt operator we only know the convergence of $\sum_n \lambda(A)^2$ but not of $\sum_n \lambda_n(A)$. Therefore, the product (3.1), which is meant to define $\det(I + zA)$, is not known to converge in general. Instead, Hilbert (1904) and Carleman (1921) introduced the entire function¹³

$$\det_2(I + zA) = \prod_{n=1}^{N(A)} (1 + z\lambda_n(A)) e^{-z\lambda_n(A)} \quad (A \in \mathcal{J}_2(\mathcal{H})),$$

¹³In fact, using such exponential “convergence factors” is a classical technique in complex analysis to construct, by means of infinite products, entire functions from their intended sets of zeros, see Ablowitz and Fokas (2003, Sect. 3.6). A famous example is

$$\frac{1}{\Gamma(z)} = ze^{\gamma z} \prod_{n=1}^{\infty} \left(1 + \frac{z}{n}\right) e^{-z/n},$$

which corresponds to the eigenvalues $\lambda_n(A) = 1/n$ of a Hilbert–Schmidt operator that is not trace class.

which also has the property to possess zeros exactly at $z_n = -1/\lambda_n(A)$, counting multiplicity. Plemelj's formula (3.3) becomes (Gohberg et al. 2000, p. 167)

$$\det_2(I + zA) = \exp\left(-\sum_{n=2}^{\infty} \frac{(-z)^n}{n} \operatorname{tr} A^n\right)$$

for $|z| < 1/|\lambda_1(A)|$, which perfectly makes sense since A^2, A^3, \dots are trace class for A being Hilbert–Schmidt.¹⁴ Note that for trace class operators we have

$$\det(I + zA) = \det_2(I + zA) \exp(z \cdot \operatorname{tr} A) \quad (A \in \mathcal{J}_1(\mathcal{H})).$$

For integral operators A of the form (2.7) with a continuous kernel K on $\mathcal{H} = L^2(a, b)$ the Fredholm determinant (1.2) is related to the Hilbert–Carleman determinant by the equation (Hochstadt 1973, p. 250)

$$d(z) = \det_2(I + zA) \exp\left(z \int_a^b K(x, x) dx\right)$$

in general, even if A is not of trace class. It is important to note, though, that if $A \notin \mathcal{J}_1(\mathcal{H})$ with such a kernel, we have $\int_a^b K(x, x) dx \neq \operatorname{tr}(A)$ simply because $\operatorname{tr}(A)$ is not well defined by (2.4) anymore then.

4. Perturbation Bounds. In studying the conditioning of operator and matrix determinants we rely on the fundamental perturbation estimate

$$|\det(I + A) - \det(I + B)| \leq \|A - B\|_{\mathcal{J}_1} \exp(1 + \max(\|A\|_{\mathcal{J}_1}, \|B\|_{\mathcal{J}_1})) \quad (4.1)$$

for trace class operators, which can beautifully be proven by means of complex analysis (Simon 2005, p. 45). This estimate can be put to the form

$$|\det(I + (A + E)) - \det(I + A)| \leq e^{1+\|A\|_{\mathcal{J}_1}} \cdot \|E\|_{\mathcal{J}_1} + O(\|E\|_{\mathcal{J}_1}^2), \quad (4.2)$$

showing that the condition number κ_{abs} of the operator determinant $\det(I + A)$, with respect to absolute errors measured in trace class norm, is bounded by

$$\kappa_{\text{abs}} \leq e^{1+\|A\|_{\mathcal{J}_1}}.$$

This bound can considerably be improved for certain selfadjoint operators that will play an important role in Section 7.

¹⁴Equivalently one can define (Simon 2005, p. 50) the regularized determinant by

$$\det_2(I + zA) = \det(I + ((I + zA)e^{-zA} - I)) \quad (A \in \mathcal{J}_2(\mathcal{H})).$$

This is because one can then show $(I + zA)e^{-zA} - I \in \mathcal{J}_1(\mathcal{H})$. For integral operators A on $L^2(a, b)$ with an L^2 kernel K , Hilbert (1904, p. 82) had found the equivalent expression

$$\det_2(I + zA) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \int_{(a,b)^n} \begin{vmatrix} 0 & K(t_1, t_2) & \cdots & K(t_1, t_n) \\ K(t_2, t_1) & 0 & \cdots & K(t_2, t_n) \\ \vdots & \vdots & \ddots & \vdots \\ K(t_n, t_1) & K(t_n, t_2) & \cdots & 0 \end{vmatrix} dt_1 \cdots dt_n,$$

simply replacing the problematic “diagonal” entries $K(t_j, t_j)$ in Fredholm's determinant (3.7) by zero. Simon (2005, Thm. 9.4) gives an elegant proof of Hilbert's formula.

LEMMA 4.1. Let $A \in \mathcal{J}_1(\mathcal{H})$ be selfadjoint, positive-semidefinite with $\lambda_1(A) < 1$. If $\|E\|_{\mathcal{J}_1} < \|(I - A)^{-1}\|^{-1}$ then

$$|\det(I - (A + E)) - \det(I - A)| \leq \|E\|_{\mathcal{J}_1}. \quad (4.3)$$

That is, the condition number κ_{abs} of the determinant $\det(I - A)$, with respect to absolute errors measured in trace class norm, is bounded by $\kappa_{abs} \leq 1$.

Proof. Because of $1 > \lambda_1(A) \geq \lambda_2(A) \geq \dots \geq 0$ there exists the inverse operator $(I - A)^{-1}$. The product formula (3.1) implies $\det(I - A) > 0$, the multiplicativity (3.5) of the determinant gives

$$\det(I - (A + E)) = \det(I - A) \det(I - (I - A)^{-1}E).$$

Upon applying Plemelj's formula (3.3) and the estimates (2.3) and (2.5) we get

$$\begin{aligned} |\log \det(I - (I - A)^{-1}E)| &= \left| \operatorname{tr} \left(\sum_{n=1}^{\infty} \frac{((I - A)^{-1}E)^n}{n} \right) \right| \\ &\leq \sum_{n=1}^{\infty} \frac{\|(I - A)^{-1}\|^n \cdot \|E\|_{\mathcal{J}_1}^n}{n} = \log \left(\frac{1}{1 - \|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1}} \right) \end{aligned}$$

if $\|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1} < 1$. Hence, exponentiation yields

$$\begin{aligned} 1 - \|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1} &\leq \det(I - (I - A)^{-1}E) \\ &\leq \frac{1}{1 - \|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1}} \leq 1 + \|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1}, \end{aligned}$$

that is

$$|\det(I - (A + E)) - \det(I - A)| \leq \det(I - A) \cdot \|(I - A)^{-1}\| \cdot \|E\|_{\mathcal{J}_1}.$$

Now, by the spectral theorem for bounded selfadjoint operators we have

$$\|(I - A)^{-1}\| = \frac{1}{1 - \lambda_1(A)} \leq \prod_{n=1}^{N(A)} \frac{1}{1 - \lambda_n(A)} = \frac{1}{\det(I - A)}$$

and therefore $\det(I - A) \cdot \|(I - A)^{-1}\| \leq 1$, which finally proves the assertion. \square

Thus, for the operators that satisfy the assumptions of this lemma the determinant is a really *well* conditioned quantity—with respect to absolute errors, like the eigenvalues of a Hermitian matrix (Golub and Van Loan 1996, p. 396).

Implications on the accuracy of numerical methods. The Nyström-type method of Section 6 requires the calculation of the determinant $\det(I + A)$ of some matrix $A \in \mathbb{C}^{m \times m}$. In the presence of roundoff errors, a backward stable method like Gaussian elimination with partial pivoting (Higham 2002, Sect. 14.6) gives a result that is *exact* for some matrix $\tilde{A} = A + E$ with

$$|E_{j,k}| \leq \epsilon |A_{j,k}| \quad (j, k = 1, \dots, m) \quad (4.4)$$

where ϵ is a small multiple of the unit roundoff error of the floating-point arithmetic used. We now use the perturbation bounds of this section to estimate the resulting error in the value of determinant. Since the trace class norm is not a monotone matrix norm (Higham 2002, Def. 6.1), we cannot make direct use of the componentwise estimate (4.4). Instead, we majorize the trace class norm of $m \times m$ matrices A by the Hilbert–Schmidt (Frobenius) norm, which is monotone, using

$$\|A\|_{\mathcal{J}_1} \leq \sqrt{m}\|A\|_{\mathcal{J}_2}, \quad \|A\|_{\mathcal{J}_2} = \left(\sum_{j,k=1}^m |A_{j,k}|^2 \right)^{1/2}.$$

Thus, the general perturbation bound (4.2) yields the following a priori estimate of the roundoff error affecting the value of the determinant:

$$|\det(I + \tilde{A}) - \det(I + A)| \leq \sqrt{m}\|A\|_{\mathcal{J}_2} \exp(1 + \|A\|_{\mathcal{J}_1}) \cdot \epsilon + O(\epsilon^2).$$

If the matrix A satisfies the assumptions of Lemma 4.1, the perturbation bound (4.3) gives the even sharper estimate

$$|\det(I - \tilde{A}) - \det(I - A)| \leq \sqrt{m}\|A\|_{\mathcal{J}_2} \cdot \epsilon. \quad (4.5)$$

Therefore, if $\det(I - A) \ll \|A\|_{\mathcal{J}_2}$, we have to be prepared that we probably cannot compute the determinant to the full precision given by the computer arithmetic used. Some digits will be lost. A conservative estimate stemming from (4.5) predicts the loss of at most

$$\log_{10} \left(\frac{\sqrt{m} \cdot \|A\|_{\mathcal{J}_2}}{\det(I - A)} \right)$$

decimal places. For instance, this will affect the *tails* of the probability distributions to be calculated in Section 7.

5. Projection Methods. The general idea (3.2) of defining the infinite dimensional determinant $\det(I + A)$ for a trace class operator A by a continuous extension from the finite dimensional case immediately leads to the concept of a projection method of Galerkin type. We consider a sequence of m -dimensional subspaces $V_m \subset \mathcal{H}$ together with their corresponding orthonormal projections

$$P_m : \mathcal{H} \rightarrow V_m.$$

The Galerkin projection $P_m A P_m$ of the trace class operator A is of finite rank. Given an orthonormal basis ϕ_1, \dots, ϕ_m of V_m , its determinant can be effectively calculated as the finite dimensional expression

$$\det(I + z P_m A P_m) = \det(I + z P_m A P_m \upharpoonright_{V_m}) = \det(\delta_{ij} + z \langle \phi_i, A \phi_j \rangle)_{i,j=1}^m \quad (5.1)$$

if the matrix elements $\langle \phi_i, A \phi_j \rangle$ are numerically accessible.

Because of $\|P_m\| \leq 1$, and thus $\|P_m A P_m\|_{\mathcal{J}_1} \leq \|A\|_{\mathcal{J}_1} \leq 1$, the perturbation bound (4.1) gives the simple error estimate

$$|\det(I + z P_m A P_m) - \det(I + z A)| \leq \|P_m A P_m - A\|_{\mathcal{J}_1} \cdot |z| e^{1+|z| \cdot \|A\|_{\mathcal{J}_1}}. \quad (5.2)$$

For the method to be convergent we therefore have to show that $P_m A P_m \rightarrow A$ in *trace class norm*. By a general result about the approximation of trace class operators (Gohberg et al. 1990, Thm. 4.3) all we need to know is that P_m converges *pointwise*¹⁵

¹⁵In infinite dimensions P_m cannot converge *in norm* since the identity operator is not compact.

to the identity operator I . This pointwise convergence is obviously equivalent to the *consistency* of the family of subspaces V_m , that is,

$$\bigcup_{m=1}^{\infty} V_m \text{ is a dense subspace of } \mathcal{H}. \quad (5.3)$$

In summary, we have proven the following theorem.

THEOREM 5.1. *Let A be a trace class operator. If the sequence of subspaces satisfies the consistency condition (5.3), the corresponding Galerkin approximation (5.1) of the operator determinant converges,*

$$\det(I + z P_m A P_m) \rightarrow \det(I + z A) \quad (m \rightarrow \infty),$$

uniformly for bounded z .

A quantitative estimate of the error, that is, in view of (5.2), of the projection error $\|P_m A P_m - A\|_{\mathcal{J}_1}$ in trace class norm, can be based on the singular value representation (2.1) of A and its finite-rank truncation A_N : (We assume that A is non-degenerate, that is, $N(|A|) = \infty$; since otherwise we could simplify the following by putting $A_N = A$.)

$$A = \sum_{n=1}^{\infty} s_n(A) \langle u_n, \cdot \rangle v_n, \quad A_N = \sum_{n=1}^N s_n(A) \langle u_n, \cdot \rangle v_n.$$

We obtain, by using $\|P_m\| \leq 1$ once more,

$$\begin{aligned} \|P_m A P_m - A\|_{\mathcal{J}_1} &\leq \|P_m A P_m - P_m A_N P_m\|_{\mathcal{J}_1} + \|P_m A_N P_m - A_N\|_{\mathcal{J}_1} + \|A_N - A\|_{\mathcal{J}_1} \\ &\leq 2\|A_N - A\|_{\mathcal{J}_1} + \|P_m A_N P_m - A_N\|_{\mathcal{J}_1} \\ &\leq 2 \sum_{n=N+1}^{\infty} s_n(A) + \sum_{n=1}^N s_n(A) \|\langle P_m u_n, \cdot \rangle P_m v_n - \langle u_n, \cdot \rangle v_n\|_{\mathcal{J}_1} \\ &\leq 2 \sum_{n=N+1}^{\infty} s_n(A) + \sum_{n=1}^N s_n(A) (\|u_n - P_m u_n\| + \|v_n - P_m v_n\|). \end{aligned} \quad (5.4)$$

There are two competing effects that contribute to making this error bound small: First, there is the convergence of the truncated series of singular values,

$$\sum_{n=N+1}^{\infty} s_n(A) \rightarrow 0 \quad (N \rightarrow \infty),$$

which is *independent* of m . Second, there is, for *fixed* N , the collective approximation

$$P_m u_n \rightarrow u_n, \quad P_m v_n \rightarrow v_n \quad (m \rightarrow \infty)$$

of the first N singular functions u_n, v_n ($n = 1, \dots, N$). For instance, given $\epsilon > 0$, we can first choose N large enough to push the first error term in (5.4) below $\epsilon/2$. After fixing such an N , the second error term can be pushed below $\epsilon/2$ for m large enough. This way we have proven Theorem 5.1 once more. However, in general the convergence of the second term might considerably slow down for growing N . Therefore,

a good quantitative bound requires a carefully balanced choice of N (depending on m), which in turn requires some detailed knowledge about the decay of the singular values on the one hand and of the growth of the derivatives of the singular functions on the other hand (see the example at the end of this section). While some general results are available in the literature for the singular values—e.g., for integral operators A induced by a kernel $K \in C^{k-1,1}([a, b]^2)$ the bound

$$s_n(A) = O(n^{-k-\frac{1}{2}}) \quad (n \rightarrow \infty) \quad (5.5)$$

obtained by Smithies (1937, Thm. 12)—the results are sparse for the singular functions (Fenyő and Stolle 1982–1984, §8.10). Since the quadrature method in Section 6 does not require any such knowledge, we refrain from stating a general result and content ourselves with the case that there is no projection error in the singular functions; that is, we consider projection methods of Ritz–Galerkin type for selfadjoint operators.

THEOREM 5.2. *Let A be a selfadjoint integral operator that is induced by a continuous Hermitian kernel K and that is trace class on the Hilbert space $\mathcal{H} = L^2(a, b)$. Assume that A is not of finite rank and let (u_n) be an orthonormal basis of eigenfunctions of A . We consider the associated Ritz projection P_m , that is, the orthonormal projection*

$$P_m : \mathcal{H} \rightarrow V_m = \text{span}\{u_1, \dots, u_m\}.$$

Note that in this case

$$\det(I + z P_m A P_m) = \prod_{n=1}^m (1 + z \lambda_n(A)).$$

If $K \in C^{k-1,1}([a, b]^2)$, then there holds the error estimate (5.2) with

$$\|P_m A P_m - A\|_{\mathcal{J}_1} = o(m^{\frac{1}{2}-k}) \quad (m \rightarrow \infty).$$

If K is bounded analytic on $\mathcal{E}_\rho \times \mathcal{E}_\rho$ (with the ellipse \mathcal{E}_ρ defined in Theorem A.2), then the error estimate improves to

$$\|P_m A P_m - A\|_{\mathcal{J}_1} = O(\rho^{-m(1-\epsilon)/4}) \quad (m \rightarrow \infty),$$

for any fixed choice of $\epsilon > 0$.

Proof. With the spectral decompositions

$$A = \sum_{n=1}^{\infty} \lambda_n(A) \langle u_n, \cdot \rangle u_n, \quad P_m A P_m = A_m = \sum_{n=1}^m \lambda_n(A) \langle u_n, \cdot \rangle u_n,$$

at hand the bound (5.4) simplifies, for $N = m$, to

$$\|P_m A P_m - A\|_{\mathcal{J}_1} = \sum_{n=m+1}^{\infty} |\lambda_n(A)|.$$

Now, by some results of Hille and Tamarkin (1931, Thm. 7.2 and 10.1), we have, for $K \in C^{k-1,1}([a, b]^2)$, the eigenvalue decay

$$\lambda_n(A) = o(n^{-k-\frac{1}{2}}) \quad (n \rightarrow \infty) \quad (5.6)$$

(which is just slightly stronger than Smithies' singular value bound (5.5)) and, for K bounded analytic on $\mathcal{E}_\rho \times \mathcal{E}_\rho$,

$$\lambda_n(A) = O(\rho^{-n(1-\epsilon)/4}) \quad (n \rightarrow \infty);$$

which proves both assertions. \square

However, for kernels of low regularity, by taking into account the specifics of a particular example one often gets better results than stated in this theorem. (An example with an analytic kernel, enjoying the excellent convergence rates of the second part this theorem, can be found in Section 7.)

An example: Poisson's equation. For a given $f \in L^2(0, 1)$ the Poisson equation

$$-u''(x) = f(x), \quad u(0) = u(1) = 0,$$

with Dirichlet boundary conditions is solved (Hochstadt 1973, p. 5) by the application of the integral operator A ,

$$u(x) = Af(x) = \int_0^1 K(x, y)f(y) dy, \quad (5.7)$$

which is induced by the Green's kernel

$$K(x, y) = \begin{cases} x(1-y) & x \leq y, \\ y(1-x) & \text{otherwise.} \end{cases} \quad (5.8)$$

Since K is Lipschitz continuous, Hermitian, and positive definite, we know from the results of Section 2 that A is a selfadjoint trace class operator on $\mathcal{H} = L^2(0, 1)$. The eigenvalues and normalized eigenfunctions of A are those of the Poisson equation which are known to be

$$\lambda_n(A) = \frac{1}{\pi^2 n^2}, \quad u_n(x) = \sqrt{2} \sin(n\pi x) \quad (n = 1, 2, \dots).$$

Note that the actual decay of the eigenvalues is better than the general Hille–Tarmarkin bound (5.6) which would, because of $K \in C^{0,1}([0, 1]^2)$, just give $\lambda_n(A) = o(n^{-3/2})$. The trace formulas (2.4) and (2.8) reproduce a classical formula of Euler's, namely

$$\sum_{n=1}^{\infty} \frac{1}{\pi^2 n^2} = \text{tr}(A) = \int_0^1 K(x, x) dx = \int_0^1 x(1-x) dx = \frac{1}{6};$$

whereas, by (3.1) and the product representation of the sine function, the Fredholm determinant explicitly evaluates to the entire function

$$\det(I - zA) = \prod_{n=1}^{\infty} \left(1 - \frac{z}{\pi^2 n^2}\right) = \frac{\sin(\sqrt{z})}{\sqrt{z}}. \quad (5.9)$$

The sharper perturbation bound of Lemma 4.1 applies and we get, for each finite dimensional subspace $V_m \subset \mathcal{H}$ and the corresponding orthonormal projection $P_m : \mathcal{H} \rightarrow V_m$, the error estimate

$$|\det(I - P_m A P_m) - \det(I - A)| \leq \|P_m A P_m - A\|_{\mathcal{J}_1}. \quad (5.10)$$

Now, we study two particular families of subspaces.




FIG. 1. Convergence of Ritz–Galerkin (circles) and Galerkin (dots) for approximating the Fredholm determinant of the integral operator induced by Green’s kernel of Poisson’s equation. The solid line shows the upper bound $1/\pi^2 m$ of Ritz–Galerkin as given in (5.11).

Trigonometric polynomials. Here, we consider the subspaces

$$V_m = \text{span}\{\sin(n\pi \cdot) : n = 1, \dots, m\} = \text{span}\{u_n : n = 1, \dots, m\},$$

which are exactly those spanned by the eigenfunctions of A . In this case, the projection method is of Ritz–Galerkin type; the estimates become particularly simple since we have the explicit spectral decomposition

$$P_m A P_m - A = \sum_{n=m+1}^{\infty} \lambda_n(A) \langle u_n, \cdot \rangle u_n$$

of the error operator. Hence, the error bound (5.10) directly evaluates to

$$\begin{aligned} & |\det(I - P_m A P_m) - \det(I - A)| \\ & \leq \|P_m A P_m - A\|_{\mathcal{J}_1} = \sum_{n=m+1}^{\infty} \lambda_n(A) = \frac{1}{\pi^2} \sum_{n=m+1}^{\infty} \frac{1}{n^2} \leq \frac{1}{\pi^2 m}. \end{aligned} \quad (5.11)$$

Figure 1 shows that this upper bound overestimates the error in the Fredholm determinant by just about 20%.

Algebraic polynomials. Here, we consider the subspaces of algebraic polynomials of order m , that is,

$$V_m = \{u \in L^2(0, 1) : u \text{ is a polynomial of degree at most } m - 1\}.$$

We apply the bound given in (5.4) and obtain (keeping in mind that A is selfadjoint)

$$\|P_m A P_m - A\|_{\mathcal{J}_1} \leq 2 \sum_{n=N+1}^{\infty} \lambda_n(A) + 2 \sum_{n=1}^N \lambda_n(A) \|u_n - P_m u_n\|$$

with a truncation index N yet to be skilfully chosen. As before in (5.11), the first term of this error bound can be estimated by $2/\pi^2 N$. For the second term we recall that the projection error $\|u_n - P_m u_n\|$ is, in fact, just the error of polynomial best approximation of the eigenfunction u_n with respect to the L^2 -norm. A standard Jackson-type

inequality (DeVore and Lorentz 1993, p. 219) from approximation theory teaches us that

$$\|u_n - P_m u_n\| \leq \frac{c_k}{m^k} \|u_n^{(k)}\| = c_k \frac{(\pi n)^k}{m^k},$$

where c_k denotes a constant that depends on the smoothness level k . A *fixed* eigenfunction u_n (being an entire function in fact) is therefore approximated beyond every algebraic order in the dimension m ; but with increasingly larger constants for higher “wave numbers” n . We thus get, with some further constant \tilde{c}_k depending on $k \geq 2$,

$$\|P_m A P_m - A\|_{\mathcal{J}_1} \leq \tilde{c}_k \left(\frac{1}{N} + \frac{N^{k-1}}{(k-1)m^k} \right).$$

We now balance the two error terms by minimizing this bound: the optimal truncation index N turns out to be exactly $N = m$, which finally yields the estimate

$$|\det(I - P_m A P_m) - \det(I - A)| \leq \|P_m A P_m - A\|_{\mathcal{J}_1} \leq \frac{\tilde{c}_k}{1 - k^{-1}} m^{-1}.$$

Thus, in contrast to the approximation of a *single* eigenfunction, for the Fredholm determinant the *order* of the convergence estimate does finally not depend on the choice of k anymore; we obtain the same $O(m^{-1})$ behavior as for the Ritz–Galerkin method. In fact, a concrete numerical calculation¹⁶ shows that this error estimate really reflects the actual order of the error decay of the Galerkin method, see Figure 1.

Remark. The analysis of this section has shown that the error decay of the projection methods is essentially determined by the decay

$$\sum_{k=m+1}^{\infty} s_k(A) \rightarrow 0$$

of the singular values of A , which in turn is related to the smoothness of the kernel K of the integral operator A . In the next section, the error analysis of Nyström-type quadrature methods will relate in a much more direct fashion to the smoothness of

¹⁶By (5.1) all we need to know for implementing the Galerkin method are the matrix elements $\langle \phi_i, A \phi_j \rangle$ for the normalized orthogonal polynomials ϕ_n (that is, properly rescaled Legendre polynomials) on the interval $[0, 1]$. A somewhat lengthy but straightforward calculation shows that these elements are given by

$$(\langle \phi_i, A \phi_j \rangle)_{i,j} = \begin{pmatrix} \frac{1}{12} & 0 & b_0 & & \\ 0 & \frac{1}{60} & 0 & b_1 & \\ b_0 & 0 & a_1 & \ddots & \ddots \\ & b_1 & \ddots & a_2 & \\ & & \ddots & & \ddots \end{pmatrix}$$

with the coefficients

$$a_n = \frac{1}{2(2n+1)(2n+5)}, \quad b_n = -\frac{1}{4(2n+3)\sqrt{(2n+1)(2n+5)}}.$$

the kernel K , giving even much better error bounds, a priori and in actual numerical computations. For instance, the Green's kernel (5.7) of low regularity can be treated by an m -dimensional approximation of the determinant with an actual convergence rate of $O(m^{-2})$ instead of $O(m^{-1})$ as for the projection methods. Moreover, these methods are much simpler and straightforwardly implemented.

6. Quadrature Methods. In this section we directly approximate the Fredholm determinant (1.2) using the Nyström-type method (1.4) that we have motivated at length in Section 1. We assume throughout that the kernel K is a continuous function on $[a, b]^2$. The notation simplifies considerably by introducing the n -dimensional functions K_n defined by

$$K_n(t_1, \dots, t_n) = \det (K(t_p, t_q))_{p,q=1}^n.$$

Their properties are given in Lemma A.4 of the appendix. We then write the Fredholm determinant shortly as

$$d(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n.$$

For a given quadrature formula

$$Q(f) = \sum_{j=1}^m w_j f(x_j) \approx \int_a^b f(x) dx$$

we define the associated Nyström-type approximation of $d(z)$ by the expression

$$d_Q(z) = \det (\delta_{ij} + z w_i K(x_i, x_j))_{i,j=1}^m. \quad (6.1)$$

The key to error estimates and a convergence proof is the observation that we can rewrite $d_Q(z)$ in a form that closely resembles the Fredholm determinant. Namely, by using the von Koch form (3.6) of matrix determinants, the multi-linearity of minors, and by introducing the n -dimensional product rule (A.3) associated with Q (see the appendix), we get

$$\begin{aligned} d_Q(z) &= 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \sum_{j_1, \dots, j_n=1}^m \det (w_{j_p} K(x_{j_p}, x_{j_q}))_{p,q=1}^n \\ &= 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \sum_{j_1, \dots, j_n=1}^m w_{j_1} \cdots w_{j_n} \det (K(x_{j_p}, x_{j_q}))_{p,q=1}^n \\ &= 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \sum_{j_1, \dots, j_n=1}^m w_{j_1} \cdots w_{j_n} K_n(x_{j_1}, \dots, x_{j_n}) \\ &= 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} Q^n(K_n). \end{aligned}$$

Thus, alternatively to the motivation given in the introductory Section 1, we could have introduced the Nyström-type method by approximating each of the multi-dimensional integrals in the power series defining the Fredholm determinant with a

product quadrature rule. Using this form, we observe that the error is given by

$$d_Q(z) - d(z) = \sum_{n=1}^{\infty} \frac{z^n}{n!} \left(Q^n(K_n) - \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right), \quad (6.2)$$

that is, by the exponentially generating function of the quadrature errors for the functions K_n . The following theorem generalizes a result that Hilbert (1904, p. 59) had proven for a specific class of quadrature formulae, namely, the rectangular rule.

THEOREM 6.1. *If a family Q_m of quadrature rules converges for continuous functions, then the corresponding Nyström-type approximation of the Fredholm determinant converges,*

$$d_{Q_m}(z) \rightarrow d(z) \quad (m \rightarrow \infty),$$

uniformly for bounded z .

Proof. Let z be bounded by M and choose any $\epsilon > 0$. We split the series (6.2) at an index N yet to be chosen, getting

$$\begin{aligned} |d_{Q_m}(z) - d(z)| &\leq \sum_{n=1}^N \frac{M^n}{n!} \left| Q_m^n(K_n) - \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \\ &\quad + \sum_{n=N+1}^{\infty} \frac{M^n}{n!} \left| Q_m^n(K_n) - \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \end{aligned}$$

Let Λ be the stability bound of the convergent family Q_m of quadrature rules (see Theorem A.1 of the appendix) and put $\Lambda_1 = \max(\Lambda, b - a)$. Then, by Lemma A.4, the second part of the splitting can be bounded by

$$\begin{aligned} &\sum_{n=N+1}^{\infty} \frac{M^n}{n!} \left| Q_m^n(K_n) - \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \\ &\leq \sum_{n=N+1}^{\infty} \frac{M^n}{n!} \left(|Q_m^n(K_n)| + \left| \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \right) \\ &\leq \sum_{n=N+1}^{\infty} \frac{M^n}{n!} (\Lambda^n + (b - a)^n) \|K_n\|_{L^\infty} \leq 2 \sum_{n=N+1}^{\infty} \frac{n^{n/2}}{n!} (M\Lambda_1 \|K\|_{L^\infty})^n. \end{aligned}$$

The last series converges by Lemma A.5 and the bound can, therefore, be pushed below $\epsilon/2$ by choosing N large enough. After fixing such an N , we can certainly also push the first part of the splitting, that is,

$$\sum_{n=1}^N \frac{M^n}{n!} \left| Q_m^n(K_n) - \int_{[a,b]^n} K_n(t_1, \dots, t_n) dt_1 \cdots dt_n \right|,$$

below $\epsilon/2$, now for m large enough, say $m \geq m_0$, using the convergence of the product rules Q_m^n induced by Q_m (see Theorem A.3). In summary we get

$$|d_{Q_m}(z) - d(z)| \leq \epsilon$$

for all $|z| \leq M$ and $m \geq m_0$, which proves the asserted convergence of the Nyström-type method. \square

If the kernel K enjoys, additionally, some smoothness, we can prove error estimates that exhibit, essentially, the same rates of convergence as for the quadrature of the sections $x \mapsto K(x, y)$ and $y \mapsto K(x, y)$.

THEOREM 6.2. *If $K \in C^{k-1,1}([a, b]^2)$, then for each quadrature rule Q of order $\nu \geq k$ with positive weights there holds the error estimate*

$$|d_Q(z) - d(z)| \leq c_k 2^k (b-a)^k \nu^{-k} \Phi(|z|(b-a) \|K\|_k),$$

where c_k is the constant (depending only on k) from Theorem A.2, and $\|K\|_k$ and Φ are the norm and function defined in (A.6) and (A.10), respectively.

If K is bounded analytic on $\mathcal{E}_\rho \times \mathcal{E}_\rho$ (with the ellipse \mathcal{E}_ρ defined in Theorem A.2), then for each quadrature rule Q of order ν with positive weights there holds the error estimate

$$|d_Q(z) - d(z)| \leq \frac{4\rho^{-\nu}}{1-\rho^{-1}} \Phi(|z|(b-a) \|K\|_{L^\infty(\mathcal{E}_\rho \times \mathcal{E}_\rho)}).$$

Proof. By Theorem A.3 and Lemma A.4 we can estimate the error (6.2) in both cases in the form

$$|d_Q(z) - d(z)| \leq \alpha \sum_{n=1}^{\infty} \frac{n^{(n+2)/2}}{n!} (|z|\beta)^n = \alpha \Phi(|z|\beta);$$

with the particular values $\alpha = c_k 2^k (b-a)^k \nu^{-k}$ and $\beta = (b-a) \|K\|_k$ in the first case, and $\alpha = 4\rho^{-\nu}/(1-\rho^{-1})$ and $\beta = (b-a) \|K\|_{L^\infty(\mathcal{E}_\rho \times \mathcal{E}_\rho)}$ in the second case. This proves both assertions. \square

An example with an analytic kernel, enjoying the excellent convergence rates of the second part this theorem, can be found in Section 7.

Note that Theorem 6.2 is based on a general result (Theorem A.2) about quadrature errors that stems from the convergence rates of polynomial best approximation. There are cases (typically of low regularity), however, for which certain quadrature formulae enjoy convergence rates that are actually *better* than best approximation. The Nyström-type method inherits this behavior; one would just have to repeat the proof of Theorem 6.2 then. We refrain from stating a general theorem, since this would involve bounds on the highest derivatives involving weights¹⁷ that take into account the boundary of the interval $[a, b]$. Instead, we content ourselves with the detailed discussion of a particular example.

An example: Poisson's equation. We revisit the example of Section 5, that is the integral operator (5.7) belonging to the Green's kernel K (defined in (5.8)) of Poisson's equation. Recall from (5.9) that

$$d(-1) = \det(I - A) = \sin(1).$$

The kernel K is just Lipschitz continuous, that is, $K \in C^{0,1}([0, 1]^2)$. If we apply the Nyström-type method with the m -point Gauss–Legendre (order $\nu = 2m$) or the Curtis–Clenshaw (order $\nu = m$) formulae as the underlying quadrature rule Q_m , Theorem 6.2 proves an error bound of the form

$$d_{Q_m}(-1) - d(-1) = O(m^{-1}),$$

¹⁷For the interval $[-1, 1]$ this weight would be $(1-x^2)^{1/2}$, see Davis and Rabinowitz (1984, §4.8.1).




FIG. 2. Convergence of the Nyström-type method for approximating the Fredholm determinant of the integral operator induced by Green’s kernel (5.8) of Poisson’s equation; the underlying quadrature rules Q_m are the m -point Gauss–Legendre (dots) and Clenshaw–Curtis (circles) rules. Note that, in accordance with Trefethen (2008), both behave essentially the same. The solid line shows the function $1/25m^2$, just to indicate the rate of convergence. For comparison we have included the results of the Ritz–Galerkin method (stars) from Figure 1.

which superficially indicates the same convergence rate as for the m -dimensional Galerkin methods of Section 5. However, the actual numerical computation shown in Figure 2 exhibits the far better convergence rate of $O(m^{-2})$. This deviation can be understood in detail as follows:

On the one hand, by inverse theorems of approximation theory (DeVore and Lorentz 1993, p. 220), valid for *proper* subintervals of $[a, b]$, the polynomial best approximation (of degree m) of sections of the Green’s kernel K cannot give a better rate than $O(m^{-1})$; since otherwise those sections could not show jumps in the first derivative. Given the line of arguments leading from polynomial best approximation to Theorem 6.2, the error estimate of $O(m^{-1})$ was therefore the *best* that could be established *this way*.

On the other hand, the sections of the Green’s kernel look like piecewise linear hat functions. Therefore, the coefficients a_m of their Chebyshev expansions decay as $O(m^{-2})$ (Davis and Rabinowitz 1984, Eq. (4.8.1.3)). Given this decay rate, one can then prove—see, for Gauss–Legendre, Davis and Rabinowitz (1984, Eq. (4.8.1.7)) and, for Clenshaw–Curtis, Riess and Johnson (1972, Eq. (9))—that the quadrature error is of rate $O(m^{-2})$, too. Now, one can lift this estimate to the Nyström-like method essentially as in Theorem 6.2; thus *proving* in fact that

$$d_{Q_m}(-1) - d(-1) = O(m^{-2}),$$

as numerically observed.

Remark. This “superconvergence” property of certain quadrature rules, as opposed to best approximation, for kernels with jumps in a higher derivative is therefore also the deeper reason that the Nyström-type method then outperforms the projection methods of Section 5 (see Figure 2): Best approximation, by direct (Jackson) and inverse (Bernstein) theorems of approximation theory, is strongly tied with the regularity of K . And this, in turn, is tied to the decay of the singular values of the induced integral operator A , which governs the convergence rates of projection methods.




FIG. 3. The probability $E_2(0; s)$ that an interval of length s does not contain, in the bulk scaling limit of level spacing 1, an eigenvalue of the Gaussian unitary ensemble (GUE). The result shown was calculated with the Nyström-like method based on Gauss–Legendre with $m = 30$; and cross-checked against the asymptotic expansion $\log E_2(0; s) = -\pi^2 s^2/8 - \log(s)/4 + \log(2)/3 - \log(\pi)/4 + 3\zeta'(-1) + O(s^{-1})$ for $s \rightarrow \infty$ (Deift et al. 1997).

A note on implementation. If the quadrature weights are positive (which in view of Theorem A.1 is anyway the better choice), as is the case for Gauss–Legendre and Clenshaw–Curtis, we recommend to implement the Nyström-type method (6.1) in the equivalent, symmetric form

$$d_Q(z) = \det(I + zA_Q), \quad A_Q = \left(w_i^{1/2} K(x_i, x_j) w_j^{1/2} \right)_{i,j=1}^m. \quad (6.3)$$

(Accordingly short Matlab and Mathematica code is given in the introductory Section 1.) The reason is that the $m \times m$ -matrix A_Q inherits some important structural properties from the integral operator A :

- If A is selfadjoint, then A_Q is Hermitian (see Footnote 10).
- If A is positive semidefinite, then, by (2.9), A_Q is positive semidefinite, too.

This way, for instance, the computational cost for calculating the finite-dimensional determinant is cut to half, if by structural inheritance $I + zA_Q$ is Hermitian positive definite; the Cholesky decomposition can then be employed instead of Gaussian elimination with partial pivoting.

7. Application to Some Entire Kernels of Random Matrix Theory. In this section we study two important examples, stemming from random matrix theory, with entire kernels. By Theorem 6.2, the Nyström-type method based on Gauss–Legendre or Curtis–Clenshaw quadrature has to exhibit exponential convergence.

7.1. The sine kernel. The probability $E_2(0; s)$ (shown in Figure 3) that an interval of length s does not contain, in the bulk scaling limit of level spacing 1, an eigenvalue of the Gaussian unitary ensemble (GUE) is given (Mehta 2004, Sect. 6.3) by the Fredholm determinant

$$E_2(0; s) = \det(I - A_s)$$

of the integral operator A_s on $L^2(0, s)$ that is induced by the sine kernel K :

$$A_s u(x) = \int_0^s K(x, y) u(y) dy, \quad K(x, y) = \frac{\sin(\pi(x - y))}{\pi(x - y)}.$$

Note that $K(x, y)$ is Hermitian and entire on $\mathbb{C} \times \mathbb{C}$; thus A_s is a selfadjoint operator of trace class on $L^2(0, s)$. (This is already much more than we would need to know for successfully applying and understanding the Nyström-type method. However, to facilitate a comparison with the Ritz–Galerkin method, we analyze the operator A_s in some more detail.) The factorization

$$K(x, y) = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{i(x-y)\xi} d\xi = \frac{1}{2\pi} \int_{-\pi}^{\pi} e^{ix\xi} e^{-iy\xi} d\xi \quad (7.1)$$

of the kernel implies that A_s is positive *definite* with maximal eigenvalue $\lambda_1(A_s) < 1$; since, for $0 \neq u \in L^2(0, s)$, we obtain

$$\begin{aligned} 0 < \langle u, A_s u \rangle &= \int_{-\pi}^{\pi} \left| \frac{1}{\sqrt{2\pi}} \int_0^s e^{-ix\xi} u(x) dx \right|^2 d\xi \\ &= \int_{-\pi}^{\pi} |\hat{u}(\xi)|^2 d\xi < \int_{-\infty}^{\infty} |\hat{u}(\xi)|^2 d\xi = \|\hat{u}\|_{L^2}^2 = \|u\|_{L^2}^2. \end{aligned}$$

Here, in stating that the inequalities are *strict*, we have used the fact that the Fourier transform \hat{u} of the function $u \in L^2(0, s)$, which has compact support, is an *entire* function. Therefore, the perturbation bound of Lemma 4.1 applies and we obtain, for Ritz–Galerkin as for any Galerkin method, like in the example of Section 5, the basic error estimate

$$|\det(I - P_m A_s P_m) - \det(I - A_s)| \leq \|P_m A_s P_m - A_s\|_{\mathcal{J}_1}.$$

Now, Theorems 5.2 and 6.2 predict a rapid, exponentially fast convergence of the Ritz–Galerkin and the Nyström-type methods: In fact, an m -dimensional approximation will give an error that decays like $O(e^{-cm})$, for any fixed $c > 0$ since, for entire kernels, the parameter $\rho > 1$ can be chosen arbitrarily in these theorems.

Details of the implementation of the Ritz–Galerkin method. There is certainly no general recipe on how to actually construct the Ritz–Galerkin method for a specific example, since one would have to know, more or less exactly, the eigenvalues of A . In the case of the sine kernel, however, Gaudin (1961) had succeeded in doing so. (See also Katz and Sarnak (1999, p. 411) and Mehta (2004, pp. 124–126).) He had observed that the integral operator \tilde{A}_t on $L^2(-1, 1)$, defined by

$$\tilde{A}_t u(x) = \int_{-1}^1 e^{i\pi t xy} u(y) dy$$

(which is, by (7.1), basically a rescaled “square-root” of A_{2t}), is commuting with the selfadjoint, second-order differential operator

$$Lu(x) = \frac{d}{dx} \left((x^2 - 1)u'(x) \right) + \pi^2 t^2 x^2 u(x)$$

with boundary conditions

$$(1 - x^2)u(x)|_{x=\pm 1} = (1 - x^2)u'(x)|_{x=\pm 1} = 0.$$

Thus, both operators share the same set of eigenfunctions u_n , namely the radial prolate spheroidal wave functions (using the notation of Mathematica 6.0)

$$u_n(x) = S_{n,0}^{(1)}(\pi t, x) \quad (n = 0, 1, 2, \dots).$$




FIG. 4. Convergence of various m -dimensional approximations of the Fredholm determinants $E_2(0;1)$ (left) and $E_2(0;2)$ (right): the Nyström-type quadrature methods based on Gauss–Legendre (dots) and Curtis–Clenshaw (circles), as well as Gaudin’s (1961) Ritz–Galerkin method based on spheroidal wave functions (stars). The dashed line shows the amount, according to (4.5), of roundoff error due to the numerical evaluation of the finite-dimensional determinants; all calculations were done in IEEE double arithmetic with machine precision $\epsilon = 2^{-53}$.

These special functions are even for n even, and odd for n odd. By plugging them into the integral operator \tilde{A}_t Gaudin had obtained, after evaluating at $x = 0$, the eigenvalues

$$\lambda_{2k}(\tilde{A}_t) = \frac{1}{u_{2k}(0)} \int_{-1}^1 u_{2k}(y) dy, \quad \lambda_{2k+1}(\tilde{A}_t) = \frac{i\pi t}{u'_{2k+1}(0)} \int_{-1}^1 u_{2k+1}(y) y dy.$$

Finally, we have (starting with the index $n = 0$ here)

$$\lambda_n(A_s) = \frac{s}{4} |\lambda_n(\tilde{A}_{s/2})|^2 \quad (n = 0, 1, 2, \dots).$$

Hence, the m -dimensional Ritz–Galerkin approximation of $\det(I - A_s)$ is given by

$$\det(I - P_m A_s P_m) = \prod_{n=0}^{m-1} (1 - \lambda_n(A_s)).$$

While Gaudin himself had to rely on tables of the spheroidal wave functions (Stratton et al. 1956), we can use the fairly recent implementation of these special functions by Falloon, Abbott and Wang (2003), which now comes with Mathematica 6.0. This way, we get the following implementation:

```

u[t_, n_, x_] := SpheroidalS1[n, 0, π t, x]
μ[t_, n_?EvenQ] :=  $\frac{\text{NIntegrate}[u[t, n, y], \{y, -1, 1\}]}{u[t, n, 0]}$ 
μ[t_, n_?OddQ] :=  $\frac{i \pi t \text{NIntegrate}[y u[t, n, y], \{y, -1, 1\}]}{u^{(0,0,1)}[t, n, 0]}$ 
λ[s_, n_] := λ[s, n] =  $\frac{s}{4} \text{Abs}\left[\mu\left[\frac{s}{2}, n\right]\right]^2$ 
DetRitzGalerkin[s_, m_] :=  $\prod_{n=0}^{m-1} (1 - \lambda[s, n])$ 

```

Given all this, one can understand that Jimbo et al.’s (1980) beautiful discovery of expressing $E_2(0; s)$ by formula (1.6) in terms of the fifth Painlevé transcendent was generally considered to be a major break-through even for its numerical evaluation




FIG. 5. The Tracy–Widom distribution $F_2(s)$; that is, in the edge scaling limit, the probability of the maximal eigenvalue of the Gaussian unitary ensemble (GUE) being not larger than s . The result shown was calculated with the Nyström-like method based on Gauss–Legendre with $m = 50$; and cross-checked against the asymptotic expansion $\log F_2(-s) = -s^3/12 - \log(s)/8 + \log(2)/24 + \zeta'(-1) + O(s^{-3/2})$ for $s \rightarrow \infty$ (Deift et al. 2008).

(Tracy and Widom 2000, Footnote 10). However, note how much less knowledge suffices for the application of the far more general Nyström-type method: continuity of K makes it applicable, and K being entire guarantees rapid, exponentially fast convergence. That is all.

An actual numerical experiment. Figure 4 shows the convergence (in IEEE machine arithmetic) of an actual calculation of the numerical values $E_2(0;1)$ and $E_2(0;2)$. We observe that the Nyström-type method based on Gauss–Legendre has an exponentially fast convergence rate comparable to the Ritz–Galerkin method. Clenshaw–Curtis needs a dimension m that is about twice as large as for Gauss–Legendre to achieve the same accuracy. This matches the fact that Clenshaw–Curtis has the order $\nu = m$, which is half the order $\nu = 2m$ of Gauss–Legendre, and shows that the bounds of Theorem 6.2 are rather sharp with respect to ν (there is no “kink” phenomenon here, cf. Trefethen (2008, p. 84)). The dashed line shows the amount, as estimated in (4.5), of roundoff error that stems from the numerical evaluation of the finite dimensional $m \times m$ -determinant itself. Note that this bound is essentially the same for all the three methods and can easily be calculated in course of the numerical evaluation. We observe that this bound is explaining the actual onset of numerical “noise” in all the three methods reasonably well.

Remark. Note that the Nyström-type method *outperforms* the Ritz–Galerkin method by far. First, the Nyström-type method is general, simple, and straightforwardly implemented (see the code given in Section 1); in contrast, the Ritz–Galerkin depends on some detailed knowledge about the eigenvalues and requires numerical access to the spheroidal wave functions. Second, there is no substantial gain, as compared to the Gauss–Legendre based method, in the convergence rate from knowing the eigenvalues exactly. Third, and most important, the computing time for the Ritz–Galerkin runs well into several minutes, whereas both versions of the Nyström-type method require just a few *milliseconds*.

7.2. The Airy kernel. The Tracy–Widom distribution $F_2(s)$ (shown in Figure 5), that is, in the edge scaling limit, the probability of the maximal eigenvalue of the Gaussian unitary ensemble (GUE) being not larger than s , is given (Mehta 2004,

§24.2) by the Fredholm determinant

$$F_2(s) = \det(I - A_s) \quad (7.2)$$

of the integral operator A_s on $L^2(s, \infty)$ that is induced by the Airy kernel K :

$$A_s u(x) = \int_s^\infty K(x, y) u(y) dy, \quad K(x, y) = \frac{\text{Ai}(x)\text{Ai}'(y) - \text{Ai}(y)\text{Ai}'(x)}{x - y}. \quad (7.3)$$

Note that K is Hermitian and entire on $\mathbb{C} \times \mathbb{C}$; thus A_s is selfadjoint. (Again, this is already about all we would need to know for successfully applying and understanding the Nyström-type method. However, we would like to show that, as for the sine kernel, the strong perturbation bound of Lemma 4.1 applies to the Airy kernel, too.) There is the factorization (Tracy and Widom 1994, Eq. (4.5))

$$K(x, y) = \int_0^\infty \text{Ai}(x + \xi)\text{Ai}(y + \xi) d\xi,$$

which relates the Airy kernel with the Airy transform (Vallée and Soares 2004, §4.2) in a similar way as the sine kernel is related by (7.1) with the Fourier transform. This proves, because of the super-exponential decay of $\text{Ai}(x) \rightarrow 0$ as $x \rightarrow 0$, that A_s is the product of two Hilbert–Schmidt operators on $L^2(s, \infty)$ and therefore of trace class. Moreover, A_s is positive semi-definite with maximal eigenvalue $\lambda_1(A) \leq 1$; since by the Parseval–Plancherel equality (Vallée and Soares 2004, Eq. (4.27)) of the Airy transform we obtain, for $u \in L^2(s, \infty)$,

$$\begin{aligned} 0 \leq \langle u, A_s u \rangle &= \int_0^\infty \left| \int_0^\infty \text{Ai}(x + \xi) u(x) dx \right|^2 d\xi \\ &\leq \int_{-\infty}^\infty \left| \int_0^\infty \text{Ai}(x + \xi) u(x) dx \right|^2 d\xi = \|u\|_{L^2}^2. \end{aligned}$$

More refined analytic arguments, or a carefully controlled numerical approximation, show the strict inequality $\lambda_1(A) < 1$; the perturbation bound of Lemma 4.1 applies.

Modification of the Nyström-type method for infinite intervals. The quadrature methods of Section 6 are not directly applicable here, since the integral operator A_s is defined by an integral over the infinite interval (s, ∞) . We have the following three options:

1. Using a Gauss-type quadrature formula on (s, ∞) that is tailor-made for the super-exponential decay of the Airy function. Such formulae have recently been constructed by Gautschi (2002).
2. Truncating the integral in (7.3) at some point $T > s$. That is, before using the Nyström-type method with a quadrature formula on the finite interval $[s, T]$ (for which the second part of Theorem 6.2 is then applicable, showing exponential convergence), we approximate the Fredholm determinant (7.2) by

$$\det(I - P_T A_s P_T) = \det\left(I - A_s \upharpoonright_{L^2(s, T)}\right),$$

where the orthonormal projection $P_T : L^2(s, \infty) \rightarrow L^2(s, T)$, $Pu = u \cdot \chi_{[s, T]}$, denotes the multiplication operator by the characteristic function of $[s, T]$.




FIG. 6. Values of the expression $\|P_T A_s P_T - A_s\|_{J_2}$, which bounds, by (7.4), the error in $F_2(s)$ committed by truncating the integral in (7.3) at a point $T > s$.

This way we commit an additional truncation error, which has, by passing through the perturbation bound of Lemma 4.1, the computable bound

$$|\det(I - P_T A_s P_T) - \det(I - A_s)| \leq \|P_T A_s P_T - A_s\|_{J_1} \leq \|P_T A_s P_T - A_s\|_{J_2} = \left(\int_T^\infty \int_T^\infty |K(x, y)|^2 dx dy \right)^{1/2}. \quad (7.4)$$

Figure 6 shows this bound as a function of the truncation point T . We observe that, for the purpose of calculating (within IEEE machine arithmetic) $F_2(s)$ for $s \in [-8, 2]$ —as shown in Figure 5—a truncation point at $T = 16$ would be more than sufficiently safe.

3. Transforming the infinite intervals to finite ones. By using a monotone and smooth transformation $\phi_s : (0, 1) \rightarrow (s, \infty)$, defining the transformed integral operator \tilde{A}_s on $L^2(0, 1)$ by

$$\tilde{A}_s u(\xi) = \int_0^1 \tilde{K}_s(\xi, \eta) u(\eta) d\eta, \quad \tilde{K}_s(\xi, \eta) = \sqrt{\phi_s'(\xi)\phi_s'(\eta)} K(\phi_s(\xi), \phi_s(\eta)),$$

gives the identity

$$F_s(s) = \det(I - A_s \upharpoonright_{L^2(s, \infty)}) = \det(I - \tilde{A}_s \upharpoonright_{L^2(0, 1)}).$$

For the super-exponentially decaying Airy kernel K we suggest the transformation

$$\phi_s(\xi) = s + 10 \tan(\pi\xi/2) \quad (\xi \in (0, 1)). \quad (7.5)$$

Note that though $\tilde{K}(\xi, \eta)$ is a smooth function on $[0, 1]$ it possesses, as a function on $\mathbb{C} \times \mathbb{C}$, essential singularities on the lines $\xi = 1$ or $\eta = 1$. Hence, we can only apply the first part of Theorem 6.2 here, which then shows, for Gauss–Legendre and Clenshaw–Curtis, a super-algebraic convergence rate, that is, $O(m^{-k})$ for arbitrarily high algebraic order k . The actual numerical experiments reported in Figure 7 show, in fact, even exponential convergence.

From the general-purpose point of view, we recommend the third option. It is straightforward and does not require any specific knowledge, or construction, as would the first and second option.




FIG. 7. Convergence of the m -dimensional Nyström-type approximation—using the transformation (7.5)—of the Fredholm determinants $F_2(-2)$ (left) and $F_2(-4)$ (right), based on Gauss–Legendre (dots) and Curtis–Clenshaw (circles). The dashed line shows the amount, according to (4.5), of roundoff error due to the numerical evaluation of the finite-dimensional determinants; all calculations were done in IEEE double arithmetic ($\epsilon = 2^{-53}$).

Remarks on other numerical methods to evaluate $F_2(s)$. As for the sine kernel, there is a selfadjoint second-order ordinary differential operator commuting with A_s (Tracy and Widom 1994, p. 166). Though this has been used to derive some asymptotic formulas, nothing is known in terms of special functions that would enable us to base a Ritz–Galerkin method on it. As Mehta (2004, p. 453) puts it: “In the case of the Airy kernel ... the differential equation did not receive much attention and its solutions are not known.”

Prior to our work of calculating $F_2(s)$ directly from its determinantal expression, all the published numerical calculations started with Tracy and Widom’s (1994) remarkable discovery of expressing $F_2(s)$ in terms of the second Painlevé transcendent; namely

$$F_2(s) = \exp\left(-\int_s^\infty (z-s)q(z)^2 dz\right)$$

with $q(z)$ being the Hastings–McLeod (1980) solution of Painlevé II,

$$q''(z) = 2q(z)^3 + zq(z), \quad q(z) \sim \text{Ai}(z) \text{ as } z \rightarrow \infty. \quad (7.6)$$

Initial value methods for the numerical integration of (7.6) suffer from severe stability problems (Prähofer and Spohn 2004). Instead, the numerically stable way of solving (7.6) goes by considering $q(z)$ as a connecting orbit, the other asymptotic state being

$$q(z) \sim \sqrt{\frac{-z}{2}} \text{ as } z \rightarrow -\infty,$$

and using numerical two-point boundary value solvers (Dieng 2005).

8. Extension to Systems of Integral Operators. We now consider an $N \times N$ system of integrals operators that is induced by continuous kernels $K_{ij} \in C(I_i \times I_j)$ ($i, j = 1, \dots, N$), where the $I_i \subset \mathbb{R}$ denote some finite intervals. The corresponding system of integral equations

$$u_i(x) + z \sum_{j=1}^N \int_{I_j} K_{ij}(x, y) u_j(y) dy = f_i(x) \quad (x \in I_i, i, j = 1, \dots, N) \quad (8.1)$$

defines, with $u = (u_1, \dots, u_N)$ and $f = (f_1, \dots, f_N)$, an operator equation

$$u + zAu = f$$

on the Hilbert space $\mathcal{H} = L^2(I_1) \oplus \dots \oplus L^2(I_N)$.

8.1. The Fredholm determinant for systems. Assuming A to be trace class, let us express $\det(I + zA)$ in terms of the system (K_{ij}) of kernels. To this end we show that the system (8.1) is equivalent to a *single* integral equation; an idea that, essentially, can already be found in the early work of Fredholm (1903, p. 388). To simplify notation, we assume that the I_k are *disjoint* (a simple transformation of the system of integral equations by a set of translations will arrange for this). We then have¹⁸

$$\mathcal{H} = \bigoplus_{k=1}^N L^2(I_k) \cong L^2(I), \quad I = I_1 \cup \dots \cup I_n.$$

by means of the natural isometric isomorphism

$$(u_1, \dots, u_N) \mapsto u = \sum_{k=1}^N \chi_k u_k$$

where χ_k denotes the characteristic function of the interval I_k . Given this picture, the operator A can be viewed being the integral operator on $L^2(I)$ that is induced by the kernel

$$K(x, y) = \sum_{i,j=1}^N \chi_i(x) K_{ij}(x, y) \chi_j(y).$$

By (3.7) we finally get (cf. Gohberg et al. (2000, Thm 6.1))

$$\begin{aligned} \det(I + zA) &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \int_{I^n} \det(K(t_p, t_q))_{p,q=1}^n dt_1 \cdots dt_n \\ &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \int_{I^n} \underbrace{\left(\sum_{i_1, \dots, i_n=1}^N \chi_{i_1}(t_1) \cdots \chi_{i_n}(t_n) \right)}_{=1} \det(K(t_p, t_q))_{p,q=1}^n dt_1 \cdots dt_n \\ &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \int_{I_{i_1} \times \dots \times I_{i_n}} \det(K(t_p, t_q))_{p,q=1}^n dt_1 \cdots dt_n \\ &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \int_{I_{i_1} \times \dots \times I_{i_n}} \det(K_{i_p i_q}(t_p, t_q))_{p,q=1}^n dt_1 \cdots dt_n. \end{aligned}$$

¹⁸The general case could be dealt with by the topological sum, or coproduct, of the intervals I_k ,

$$\prod_{k=1}^N I_k = \bigcup_{k=1}^N I_k \times \{k\}.$$

One would then use (Johansson 2003) the natural isometric isomorphism

$$\mathcal{H} = \bigoplus_{k=1}^N L^2(I_k) \cong L^2\left(\prod_{k=1}^N I_k\right).$$

By eventually transforming back to the originally given, non-disjoint intervals I_k , the last expression is the general formula that we have sought for: $\det(I + zA) = d(z)$ with

$$d(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \int_{I_{i_1} \times \dots \times I_{i_n}} \det \left(K_{i_p i_q}(t_p, t_q) \right)_{p,q=1}^n dt_1 \cdots dt_n. \quad (8.2)$$

This is a perfectly well defined entire function for *any* system K_{ij} of continuous kernels, independently of whether A is a trace class operator or not. We call it the Fredholm determinant of the system.

The determinant of block matrices. In preparation of our discussion of Nyström-type methods for approximating (8.2) we shortly discuss the determinant of $N \times N$ -block matrices

$$A = \begin{pmatrix} A_{11} & \cdots & A_{1N} \\ \vdots & & \vdots \\ A_{N1} & \cdots & A_{NN} \end{pmatrix} \in \mathbb{C}^{M \times M}, \quad A_{ij} \in \mathbb{C}^{m_i \times m_j}, \quad M = m_1 + \cdots + m_N.$$

Starting with von Koch's formula (3.6), an argument¹⁹ that is similar to the one that has led us to (8.2) yields

$$\det(I + zA) = \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \sum_{k_1=1}^{m_{i_1}} \cdots \sum_{k_n=1}^{m_{i_n}} \det \left((A_{i_p i_q})_{k_p, k_q} \right)_{p,q=1}^n. \quad (8.3)$$

8.2. Quadrature methods for systems. Given a quadrature formula for each of the intervals I_i , namely

$$Q_i(f) = \sum_{j=1}^{m_i} w_{ij} f(x_{ij}) \approx \int_{I_i} f(x) dx, \quad (8.4)$$

we aim at generalizing the Nyström-type method of Section 6. We restrict ourselves to the case of positive weights, $w_{ij} > 0$, and generalize the method from the single operator case as given in (6.3) to the system case in the following form:

$$d_Q(z) = \det(I + zA_Q), \quad A_Q = \begin{pmatrix} A_{11} & \cdots & A_{1N} \\ \vdots & & \vdots \\ A_{N1} & \cdots & A_{NN} \end{pmatrix} \quad (8.5)$$

with the sub-matrices A_{ij} defined by the entries

$$(A_{ij})_{p,q} = w_{ip}^{1/2} K_{ij}(x_{ip}, x_{jq}) w_{jq}^{1/2} \quad (p = 1, \dots, m_i, q = 1, \dots, m_j).$$

¹⁹Alternatively, we can use (3.4) and, recursively, the "binomial" formula (Greub 1967, p. 121)

$$\bigwedge^k (V_0 \oplus V_1) = \bigoplus_{j=0}^k \left(\bigwedge^j V_0 \right) \otimes \left(\bigwedge^{k-j} V_1 \right)$$

of exterior algebra, which is valid for general vector spaces V_0 and V_1 .

This can be as straightforwardly implemented as in the case of a single operator. Now, a convergence theory can be built on a representation of the error $d_Q(z) - d(z)$ that is analogous to (6.2). To this end we simplify the notation by introducing the following functions on $I_{i_1} \times \cdots \times I_{i_n}$,

$$K_{i_1, \dots, i_n}(t_1, \dots, t_n) = \det \left(K_{i_p i_q}(t_p, t_q) \right)_{p, q=1}^n,$$

and by defining, for functions f on $I_{i_1} \times \cdots \times I_{i_n}$, the product quadrature formula

$$\begin{aligned} \left(\prod_{k=1}^n Q_{i_k} \right) (f) &= \sum_{j_1=1}^{m_{i_1}} \cdots \sum_{j_n=1}^{m_{i_n}} w_{i_1 j_1} \cdots w_{i_n j_n} f(x_{i_1 j_1}, \dots, x_{i_n j_n}) \\ &\approx \int_{I_{i_1} \times \cdots \times I_{i_n}} f(t_1, \dots, t_n) dt_1 \cdots dt_n. \end{aligned}$$

Thus, we can rewrite the Fredholm determinant (8.2) in the form

$$d(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \int_{I_{i_1} \times \cdots \times I_{i_n}} K_{i_1, \dots, i_n}(t_1, \dots, t_n) dt_1 \cdots dt_n.$$

Likewise, by observing the generalized von Koch formula (8.3), we put the definition (8.5) of $d_Q(z)$ to the form

$$d_Q(z) = 1 + \sum_{n=1}^{\infty} \frac{z^n}{n!} \sum_{i_1, \dots, i_n=1}^N \left(\prod_{k=1}^n Q_{i_k} \right) (K_{i_1, \dots, i_n}).$$

Thus, once again, the Nyström-type method amounts for approximating each multidimensional integral of the power series of the Fredholm determinant by using a product quadrature rule. Given this representation, Theorem 6.2 can straightforwardly be generalized to the system case:

THEOREM 8.1. *If $K_{ij} \in C^{k-1,1}(I_i \times I_j)$, then for each set (8.4) of quadrature formulae of a common order $v \geq k$ with positive weights there holds the error estimate*

$$d_Q(z) - d(z) = O(v^{-k}) \quad (v \rightarrow \infty),$$

uniformly for bounded z .

If the K_{ij} are bounded analytic on $\mathcal{E}_\rho(I_i) \times \mathcal{E}_\rho(I_j)$ (with the ellipse $\mathcal{E}_\rho(I_i)$ defined, with respect to I_i , as in Theorem A.2), then for each set (8.4) of quadrature formulae of a common order v with positive weights there holds the error estimate

$$d_Q(z) - d(z) = O(\rho^{-v}) \quad (v \rightarrow \infty),$$

uniformly for bounded z .

8.3. Examples from random matrix theory. Here, we apply the Nyström-type method (8.5) to two 2×2 -systems of integral operators that have recently been studied in random matrix theory.




FIG. 8. Values of the two-point correlation function $\text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0))$ of the Airy process $\mathcal{A}_2(t)$ (solid line). The dashed line shows the first term of the asymptotic expansion $\text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0)) \sim t^{-2}$ as $t \rightarrow \infty$.

Two-point correlation of the Airy process. The Airy process $\mathcal{A}_2(t)$ describes, in a properly rescaled limit of infinite dimension, the maximum eigenvalue of Hermitian matrix ensemble whose entries develop according to the Ornstein–Uhlenbeck process. This stationary stochastic process was introduced by Prähofer and Spohn (2002) and further studied by Johansson (2003). These authors have shown that the joint probability function is given by a Fredholm determinant; namely

$$\mathbb{P}(\mathcal{A}_2(t) \leq s_1, \mathcal{A}_2(0) \leq s_2) = \det \left(I - \begin{pmatrix} A_0 & A_t \\ A_{-t} & A_0 \end{pmatrix} \Big|_{L^2(s_1, \infty) \oplus L^2(s_2, \infty)} \right) \quad (8.6)$$

with integral operators A_t that are induced by the kernel functions

$$K_t(x, y) = \begin{cases} \int_0^\infty e^{-\xi t} \text{Ai}(x + \xi) \text{Ai}(y + \xi) d\xi, & t > 0, \\ -\int_{-\infty}^0 e^{-\xi t} \text{Ai}(x + \xi) \text{Ai}(y + \xi) d\xi, & \text{otherwise.} \end{cases} \quad (8.7)$$

Of particular interest is the two-point correlation function

$$\begin{aligned} \text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0)) &= \mathbb{E}(\mathcal{A}_2(t)\mathcal{A}_2(0)) - \mathbb{E}(\mathcal{A}_2(t))\mathbb{E}(\mathcal{A}_2(0)) \\ &= \int_{\mathbb{R}^2} s_1 s_2 \frac{\partial^2 \mathbb{P}(\mathcal{A}_2(t) \leq s_1, \mathcal{A}_2(0) \leq s_2)}{\partial s_1 \partial s_2} ds_1 ds_2 - c_1^2, \end{aligned} \quad (8.8)$$

where c_1 denotes the expectation value of the Tracy–Widom distribution (7.2). We have calculated this correlation function for $0 \leq t \leq 100$ in steps of 0.1 to an absolute error of $\pm 10^{-10}$, see Figure 8.²⁰ Here are some details about the numerical procedure:

²⁰A table can be obtained from the author upon request. Sasamoto (2005, Fig. 2) shows a plot (which differs by a scaling factor of two in both the function value and the time t) of the closely related function

$$g_2(t) = \sqrt{\text{var}(\mathcal{A}_2(t) - \mathcal{A}_2(0))/2} = \sqrt{\text{var}(\mathcal{A}_2(0)) - \text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0))}$$

—without, however, commenting on either the numerical procedure used or on the accuracy obtained.

- Infinite intervals of integration, such as in the definition (8.7) of the kernels or for the domain of the integral operators (8.6) themselves, are handled by a transformation to the finite interval $[0, 1]$ as in Section 7.2.
- The kernels (8.7) are evaluated, after transformation, by a Gauss–Legendre quadrature.
- The joint probability distribution (8.6) is then evaluated, after transformation, by the Nyström-type method of this section, based on Gauss–Legendre quadrature.
- To avoid numerical differentiation, the expectation values defining the two-point correlation (8.8) are evaluated by truncation of the integrals, partial integration, and using a Gauss–Legendre quadrature once more.

Because of analyticity, the convergence is always exponential. With parameters carefully (i.e., adaptively) adjusted to deliver an absolute error of $\pm 10^{-10}$, the evaluation of the two-point correlation takes, for a single time t and using a 2 GHz PC, about 20 minutes on average. The results were cross-checked, for small t , with the asymptotic expansion (Prähofer and Spohn 2002, Hägg 2007)

$$\begin{aligned} \text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0)) &= \text{var}(\mathcal{A}_2(0)) - \frac{1}{2}\text{var}(\mathcal{A}_2(t) - \mathcal{A}_2(0)) \\ &= \text{var}(\mathcal{A}_2(0)) - t + O(t^2) \quad (t \rightarrow 0), \end{aligned}$$

and, for large t , with the asymptotic expansion²¹ (Widom 2004, Adler and van Moerbeke 2005)

$$\text{cov}(\mathcal{A}_2(t), \mathcal{A}_2(0)) = t^{-2} + ct^{-4} + O(t^{-6}) \quad (t \rightarrow \infty),$$

where the constant $c = -3.542 \dots$ can explicitly be expressed in terms of the Hastings–McLeod solution (7.6) of Painlevé II.

Two-point correlation of the Airy₁ process. Sasamoto (2005) and Borodin et al. (2007) have introduced the Airy₁ process $\mathcal{A}_1(t)$ for which, once again, the joint probability distribution can be given in terms of a Fredholm determinant; namely

$$\mathbb{P}(\mathcal{A}_1(t) \leq s_1, \mathcal{A}_1(0) \leq s_2) = \det \left(I - \begin{pmatrix} A_0 & A_t \\ A_{-t} & A_0 \end{pmatrix} \Big|_{L^2(s_1, \infty) \oplus L^2(s_2, \infty)} \right)$$

with integral operators A_t that are now induced by the kernel functions

$$K_t(x, y) = \begin{cases} \text{Ai}(x + y + t^2) e^{t(x+y) + 2t^3/3} - \frac{\exp(-(x-y)^2/(4t))}{\sqrt{4\pi t}}, & t > 0, \\ \text{Ai}(x + y + t^2) e^{t(x+y) + 2t^3/3}, & \text{otherwise.} \end{cases}$$

²¹Adler and van Moerbeke (2005) have derived this asymptotic expansion from the masterfully obtained result that $G(t, x, y) = \log \mathbb{P}(\mathcal{A}_2(t) \leq x, \mathcal{A}_2(0) \leq y)$ satisfies the following nonlinear 3rd order PDE with certain (asymptotic) boundary conditions:

$$\begin{aligned} t \frac{\partial}{\partial t} \left(\frac{\partial^2}{\partial x^2} - \frac{\partial^2}{\partial y^2} \right) G &= \frac{\partial^3 G}{\partial x^2 \partial y} \left(2 \frac{\partial^2 G}{\partial y^2} + \frac{\partial^2 G}{\partial x \partial y} - \frac{\partial^2 G}{\partial x^2} + x - y - t^2 \right) \\ &\quad - \frac{\partial^3 G}{\partial y^2 \partial x} \left(2 \frac{\partial^2 G}{\partial x^2} + \frac{\partial^2 G}{\partial x \partial y} - \frac{\partial^2 G}{\partial y^2} - x + y - t^2 \right) + \left(\frac{\partial^3 G}{\partial x^3} \frac{\partial}{\partial y} - \frac{\partial^3 G}{\partial y^3} \frac{\partial}{\partial x} \right) \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right) G. \end{aligned}$$

The reader should contemplate a numerical calculation of the two-point correlation based on this PDE, rather than directly treating the Fredholm determinant as suggested by us.




FIG. 9. Values of the two-point correlation function $\text{cov}(\mathcal{A}_1(t), \mathcal{A}_1(0))$ of the Airy_1 process $\mathcal{A}_1(t)$.

By basically employing the same numerical procedure as for the Airy process, we have succeeded in calculating the two-point correlation function $\text{cov}(\mathcal{A}_1(t), \mathcal{A}_1(0))$ for $0 \leq t \leq 2.5$ in steps of 0.025 to an absolute error of $\pm 10^{-10}$, see Figure 9.²² For a single time t the evaluation takes about 5 minutes on average (using a 2 GHz PC). This numerical result has been used by Bornemann, Ferrari and Prähofer (2008) as a strong evidence that the Airy_1 process is, unlike previously conjectured, *not* the limit of the largest eigenvalue in GOE matrix diffusion.

A. Appendices.

A.1. Quadrature Rules. For the ease of reference, we collect in this appendix some classical facts about quadrature rules in one and more dimensions.

Quadrature rules in one dimension. We consider quadrature rules of the form

$$Q(f) = \sum_{j=1}^m w_j f(x_j) \quad (\text{A.1})$$

which are meant to approximate $\int_a^b f(x) dx$ for continuous functions f on some finite interval $[a, b] \subset \mathbb{R}$. We define the norm of a quadrature rule by

$$\|Q\| = \sum_{j=1}^m |w_j|$$

Convergence of a sequence of quadrature rules is characterized by the following theorem of Pólya (Davis and Rabinowitz 1984, p. 130).

THEOREM A.1. *A sequence Q_n of quadrature rules converges for continuous functions,*

$$\lim_{n \rightarrow \infty} Q_n(f) = \int_a^b f(x) dx \quad (f \in C[a, b]),$$

²²A table can be obtained from the author upon request. Sasamoto (2005, Fig. 2) shows a plot (which differs by a scaling factor of two in both the function value and the time t) of the closely related function

$$g_1(t) = \sqrt{\text{var}(\mathcal{A}_1(t) - \mathcal{A}_1(0))/2} = \sqrt{\text{var}(\mathcal{A}_1(0)) - \text{cov}(\mathcal{A}_1(t), \mathcal{A}_1(0))}$$

—without, however, commenting on either the numerical procedure used or on the accuracy obtained.

if and only if the sequence $\|Q_n\|$ of norms is bounded by some stability constant Λ and if

$$\lim_{n \rightarrow \infty} Q_n(x^k) = \int_a^b x^k dx \quad (k = 0, 1, 2, \dots). \quad (\text{A.2})$$

If the weights are all positive, then (A.2) already implies the boundedness of $\|Q_n\| = Q_n(1)$.

A quadrature rule Q is of order $\nu \geq 1$, if it is exact for all polynomials of degree at most $\nu - 1$. Using results from the theory polynomial best approximation one can prove quite strong error estimates (Davis and Rabinowitz 1984, §4.8).

THEOREM A.2. *If $f \in C^{k-1,1}[a, b]$, then for each quadrature rule Q of order $\nu \geq k$ with positive weights there holds the error estimate*

$$\left| Q(f) - \int_a^b f(x) dx \right| \leq c_k (b-a)^{k+1} \nu^{-k} \|f^{(k)}\|_{L^\infty(a,b)},$$

with a constant²³ c_k depending only on k .

If f is bounded analytic in the ellipse \mathcal{E}_ρ with foci at a, b and semiaxes of lengths $s > \sigma$ such that

$$\rho = \sqrt{\frac{s+\sigma}{s-\sigma}},$$

then for each quadrature rule Q of order ν with positive weights there holds the error estimate

$$\left| Q(f) - \int_a^b f(x) dx \right| \leq \frac{4(b-a)\rho^{-\nu}}{1-\rho^{-1}} \|f\|_{L^\infty(\mathcal{E}_\rho)}.$$

Quadrature rules in two and more dimensions. For the n -dimensional integral

$$\int_{[a,b]^n} f(t_1, \dots, t_n) dt_1 \cdots dt_n$$

we consider the product quadrature rule Q^n that is induced by an one dimensional quadrature rule Q of the form (A.1), namely

$$Q^n(f) = \sum_{j_1, \dots, j_n=1}^m w_{j_1} \cdots w_{j_n} f(x_{j_1}, \dots, x_{j_n}). \quad (\text{A.3})$$

We introduce some further notation for two classes of functions f . First, for $f \in C^{k-1,1}([a, b]^n)$, we define the seminorm

$$|f|_k = \sum_{i=1}^n \|\partial_i^k f\|_{L^\infty([a,b]^n)}. \quad (\text{A.4})$$

Second, if $f \in C([a, b]^n)$ is sectional analytic—that is, analytic with respect to each variable t_i while the other variables are fixed in $[a, b]$ —in the ellipse \mathcal{E}_ρ (defined in Theorem A.2), and if f is uniformly bounded there, we call f to be of class \mathcal{C}_ρ with norm

$$\|f\|_{\mathcal{C}_\rho} = \sum_{i=1}^n \max_{(t_1, \dots, t_{i-1}, t_{i+1}, \dots, t_n) \in [a,b]^{n-1}} \|f(t_1, \dots, t_{i-1}, \cdot, t_{i+1}, \dots, t_n)\|_{L^\infty(\mathcal{E}_\rho)}. \quad (\text{A.5})$$

²³Taking Jackson's inequality as given in Cheney (1998, p. 147), $c_k = 2(\pi e/4)^k / \sqrt{2\pi k}$ will do the job.

By a straightforward reduction argument (Davis and Rabinowitz 1984, p. 361) to the quadrature errors of the one-dimensional coordinate sections of f , Theorems A.1 and A.2 can now be generalized to n dimensions.

THEOREM A.3. *If a sequence of quadrature rules converges for continuous functions, then the same holds for the induced n -dimensional product rules.*

If $f \in C^{k-1,1}([a, b]^n)$, then for each one-dimensional quadrature rule Q of order $\nu \geq k$ with positive weights there holds the error estimate

$$\left| Q^n(f) - \int_{[a,b]^n} f(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \leq c_k (b-a)^{n+k} \nu^{-k} |f|_k,$$

with the same constant c_k depending only on k as in Theorem A.2.

If $f \in C([a, b]^n)$ is of class C_ρ , then for each one-dimensional quadrature rule Q of order ν with positive weights there holds the error estimate

$$\left| Q^n(f) - \int_{[a,b]^n} f(t_1, \dots, t_n) dt_1 \cdots dt_n \right| \leq \frac{4(b-a)^n \rho^{-\nu}}{1-\rho^{-1}} \|f\|_{C_\rho}.$$

Notes on Gauss–Legendre and Curtis–Clenshaw quadrature. Arguably, the most interesting families of quadrature rules, with *positive* weights, are the Clenshaw–Curtis and Gauss–Legendre rules. With m points, the first is of order $\nu = m$, the second of order $\nu = 2m$. Thus, Theorems A.1 and A.2 apply. The cost of computing the weights and points of Clenshaw–Curtis is $O(m \log m)$ using FFT, that of Gauss–Legendre is $O(m^2)$ using the Golub–Welsh algorithm; for details see (Waldvogel 2006) and (Trefethen 2008). The latter paper studies in depth the reasons why the Clenshaw–Curtis rule, despite having only half the order, performs essentially as well as Gauss–Legendre for most integrands. To facilitate reproducibility we offer the Matlab code (which is just a minor variation of the code given in the papers mentioned above) that has been used in our numerical experiments:

```
function [w,c] = ClenshawCurtis(a,b,m)
m = m-1;
c = cos((0:m)*pi/m);
M = [1:2:m-1]'; l = length(M); n = m-1;
v0 = [2./M./(M-2); 1/M(end); zeros(n,1)];
v2 = -v0(1:end-1)-v0(end:-1:2);
g0 = -ones(m,1); g0(1+1)=g0(1+1)+m; g0(1+n)=g0(1+n)+m;
g = g0/(m^2+mod(m,2));
w = ifft(v2+g); w(m+1) = w(1);
c = ((1-c)/2*a+(1+c)/2*b)';
w = ((b-a)*w/2)';
```

for Clenshaw–Curtis; and

```
function [w,c] = GaussLegendre(a,b,m)
k = 1:m-1; beta = k./sqrt((2*k-1).*(2*k+1));
T = diag(beta,-1) + diag(beta,1);
[V,L] = eig(T);
c = (diag(L)+1)/2; c = (1-c)*a+c*b;
w = (b-a)*V(1,:).^2;
```

for Gauss–Legendre, respectively. Note, however, that the code for Gauss–Legendre is, unfortunately, suboptimal in requiring $O(m^3)$ rather than $O(m^2)$ operations, since it establishes the full matrix V of eigenvectors of the Jacobi matrix T instead of directly calculating just their first components $V(1, :)$ as in the fully fledged Golub–Welsh algorithm. Even then, there may well be more accurate, and more efficient, alternatives of computing the points and weights of Gauss–Legendre quadrature, see the discussions in Laurie (2001, §2) and Swarztrauber (2002, §4) and the literature cited therein.

A.2. Determinantal bounds. In Section 6, for a continuous kernel $K \in C([a, b]^2)$ of an integral operator, we need some bounds on the derivatives of the induced n -dimensional function

$$K_n(t_1, \dots, t_n) = \det (K(t_p, t_q))_{p,q=1}^n.$$

To this end, if $K \in C^{k-1,1}([a, b]^2)$ we define the norm

$$\|K\|_k = \max_{i+j \leq k} \|\partial_1^i \partial_2^j K\|_{L^\infty}. \quad (\text{A.6})$$

LEMMA A.4. *If $K \in C([a, b]^2)$, then $K_n \in C([a, b]^n)$ with*

$$\|K_n\|_{L^\infty} \leq n^{n/2} \|K\|_{L^\infty}^n. \quad (\text{A.7})$$

If $K \in C^{k-1,1}([a, b]^2)$, then $K_n \in C^{k-1,1}([a, b]^n)$ with the seminorm (defined in (A.4))

$$|K_n|_k \leq 2^k n^{(n+2)/2} \|K\|_k^n. \quad (\text{A.8})$$

If K is bounded analytic on $\mathcal{E}_\rho \times \mathcal{E}_\rho$ (with the ellipse \mathcal{E}_ρ defined in Theorem A.2), then K_n is of class C_ρ (defined in (A.5)) and satisfies

$$\|K_n\|_{C_\rho} \leq n^{(n+2)/2} \|K\|_{L^\infty(\mathcal{E}_\rho \times \mathcal{E}_\rho)}^n. \quad (\text{A.9})$$

Proof. Using the multilinearity of the determinant we have

$$\begin{aligned} & \frac{\partial^k}{\partial t_i^k} \frac{\partial^l}{\partial s_j^l} \begin{vmatrix} K(t_1, s_1) & \cdots & K(t_1, s_j) & \cdots & K(t_1, s_n) \\ \vdots & & \vdots & & \vdots \\ K(t_i, s_1) & \cdots & K(t_i, s_j) & \cdots & K(t_i, s_n) \\ \vdots & & \vdots & & \vdots \\ K(t_n, s_1) & \cdots & K(t_n, s_j) & \cdots & K(t_n, s_n) \end{vmatrix} \\ &= \begin{vmatrix} K(t_1, s_1) & \cdots & \partial_2^l K(t_1, s_j) & \cdots & K(t_1, s_n) \\ \vdots & & \vdots & & \vdots \\ \partial_1^k K(t_i, s_1) & \cdots & \partial_1^k \partial_2^l K(t_i, s_j) & \cdots & \partial_1^k K(t_i, s_n) \\ \vdots & & \vdots & & \vdots \\ K(t_n, s_1) & \cdots & \partial_2^l K(t_n, s_j) & \cdots & K(t_n, s_n) \end{vmatrix}, \end{aligned}$$

which is, by Hadamard's inequality (Meyer 2000, p. 469),²⁴ bounded by the expression (see also Lax (2002, p. 262))

$$n^{n/2} \left(\max_{i+j \leq k+l} \|\partial_1^i \partial_2^j K\|_{L^\infty} \right)^n.$$

Now, with

$$\partial_j^k K_n(t_1, \dots, t_n) = \sum_{l=0}^k \binom{k}{l} \frac{\partial^{k-l}}{\partial t_j^{k-l}} \frac{\partial^l}{\partial s_j^l} \left| \begin{array}{ccc} K(t_1, s_1) & \cdots & K(t_1, s_n) \\ \vdots & & \vdots \\ K(t_n, s_1) & \cdots & K(t_n, s_n) \end{array} \right|_{s_1=t_1, \dots, s_n=t_n}$$

we thus get

$$\|\partial_j^k K_n\|_{L^\infty} \leq \sum_{l=0}^k \binom{k}{l} n^{n/2} \left(\max_{i+j \leq k} \|\partial_1^i \partial_2^j K\|_{L^\infty} \right)^n = 2^k n^{n/2} \left(\max_{i+j \leq k} \|\partial_1^i \partial_2^j K\|_{L^\infty} \right)^n.$$

This proves the asserted bounds (A.7) and (A.8) with $k = 0$ and $k \geq 1$, respectively. The class \mathcal{C}_ρ bound (A.9) follows analogously to the case $k = 0$. \square

A.3. Properties of a certain function used in Theorem 6.2. The power series

$$\Phi(z) = \sum_{n=1}^{\infty} \frac{n^{(n+2)/2}}{n!} z^n \quad (\text{A.10})$$

defines an entire function on \mathbb{C} (as the following lemma readily implies).

LEMMA A.5. *Let Ψ be the entire function given by the expression*

$$\Psi(z) = 1 + \frac{\sqrt{\pi}}{2} z e^{z^2/4} \left(1 + \operatorname{erf} \left(\frac{z}{2} \right) \right).$$

If $x > 0$, then the series $\Phi(x)$ is enclosed by:²⁵

$$\sqrt{\frac{e}{\pi}} x \Psi(x\sqrt{2e}) \leq \Phi(x) \leq x \Psi(x\sqrt{2e}).$$

Proof. For $x > 0$ we have

$$\Phi(x) = x \sum_{n=1}^{\infty} \frac{n^{n/2}}{\Gamma(n)} x^{n-1}.$$

By Stirling's formula and monotonicity we get for $n \geq 1$

$$\sqrt{\frac{e}{\pi}} \leq \frac{n^{n/2}}{\Gamma((n+1)/2) (\sqrt{2e})^{n-1}} \leq 1;$$

in fact, the upper bound is obtained for $n = 1$ and the lower bound for $n \rightarrow \infty$. Thus, by observing

$$\sum_{n=1}^{\infty} \frac{\Gamma((n+1)/2)}{\Gamma(n)} z^{n-1} = 1 + \frac{\sqrt{\pi}}{2} z e^{z^2/4} \left(1 + \operatorname{erf} \left(\frac{z}{2} \right) \right) = \Psi(z)$$

we get the asserted enclosure. \square

²⁴This inequality, discovered by Hadamard in 1893, was already of fundamental importance to Fredholm's original theory (Fredholm 1900, p. 41).

²⁵Note the sharpness of this enclosure: $\sqrt{e/\pi} = 0.93019\dots$

Acknowledgements. It is a pleasure to acknowledge that this work has started when I attended the programme on “Highly Oscillatory Problems” at the Isaac Newton Institute in Cambridge. It was a great opportunity meeting Percy Deift there, who introduced me to numerical problems related to random matrix theory (even though he was envisioning a general numerical treatment of Painlevé transcendents, but not of Fredholm determinants). I am grateful for his advice as well as for the communication with Herbert Spohn and Michael Prähofer who directed my search for an “open” numerical problem to the two-point correlation functions of the Airy and Airy_1 processes. I thank Patrik Ferrari who pointed me to the (2005) paper of Sasamoto. Given the discovery that Theorem 6.1, which was pivotal to my study, is essentially a long forgotten (see my discussion on p. 4) result of Hilbert’s 1904 work, I experienced much reconciliation—please allow me this very personal statement—from reading the poem “East Coker” (1940), in which T. S. Eliot, that “radical traditionalist”, described the nature of the human struggle for progress in life, art, or science:

... And so each venture
Is a new beginning ...
... And what there is to conquer
By strength and submission, has already been discovered
Once or twice, or several times, by men whom one cannot hope
To emulate—but there is no competition—
There is only the fight to recover what has been lost
And found and lost again and again ...

REFERENCES

- Ablowitz, M. J. and Fokas, A. S.: 2003, *Complex variables: introduction and applications*, 2nd edn, Cambridge University Press, Cambridge.
- Adler, M. and van Moerbeke, P.: 2005, PDEs for the joint distributions of the Dyson, Airy and sine processes, *Ann. Probab.* **33**, 1326–1361.
- Albeverio, S. and Høegh-Krohn, R.: 1977, Oscillatory integrals and the method of stationary phase in infinitely many dimensions, with applications to the classical limit of quantum mechanics. I, *Invent. Math.* **40**, 59–106.
- Axler, S.: 1995, Down with determinants!, *Amer. Math. Monthly* **102**, 139–154.
- Axler, S.: 1997, *Linear algebra done right*, 2nd edn, Springer-Verlag, New York.
- Baker, C. T. H.: 1977, *The numerical treatment of integral equations*, Clarendon Press, Oxford.
- Birkhoff, G. (ed.): 1973, *A source book in classical analysis*, Harvard University Press, Cambridge.
- Bornemann, F., Ferrari, P. L. and Prähofer, M.: 2008, The Airy_1 process is not the limit of the largest eigenvalue in GOE matrix diffusion, [arXiv:0806.3410](https://arxiv.org/abs/0806.3410).
- Borodin, A., Ferrari, P. L., Prähofer, M. and Sasamoto, T.: 2007, Fluctuation properties of the TASEP with periodic initial configuration, *J. Stat. Phys.* **129**, 1055–1080.
- Carleman, T.: 1918, Über die Fourierkoeffizienten einer stetigen Function, *Acta Math.* **41**, 377–384.
- Carleman, T.: 1921, Zur Theorie der linearen Integralgleichungen, *Math. Zeitschr.* **9**, 196–217.
- Cheney, E. W.: 1998, *Introduction to approximation theory*, reprint of the 2nd (1982) edn, AMS Chelsea Publishing, Providence.
- Courant, R. and Hilbert, D.: 1953, *Methods of mathematical physics. Vol. I*, Interscience Publishers, Inc., New York.
- Davis, P. J. and Rabinowitz, P.: 1984, *Methods of numerical integration*, 2nd edn, Academic Press, Orlando.
- Deift, P. A.: 1999, *Orthogonal polynomials and random matrices: a Riemann-Hilbert approach*, American Mathematical Society, Providence.
- Deift, P. A., Its, A. R. and Zhou, X.: 1997, A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics, *Ann. of Math.* **146**, 149–235.

- Deift, P., Its, A. and Krasovsky, I.: 2008, Asymptotics of the Airy-kernel determinant, *Comm. Math. Phys.* **278**, 643–678.
- Delves, L. M. and Mohamed, J. L.: 1985, *Computational methods for integral equations*, Cambridge University Press, Cambridge.
- DeVore, R. A. and Lorentz, G. G.: 1993, *Constructive approximation*, Springer-Verlag, Berlin.
- Dieng, M.: 2005, *Distribution Functions for Edge Eigenvalues in Orthogonal and Symplectic Ensembles: Painlevé Representations*, PhD thesis, University of Davis. [arXiv:math/0506586v2](https://arxiv.org/abs/math/0506586v2).
- Dieudonné, J.: 1981, *History of functional analysis*, North-Holland Publishing Co., Amsterdam.
- Dunford, N. and Schwartz, J. T.: 1963, *Linear operators. Part II: Spectral theory*, John Wiley & Sons.
- Dyson, F. J.: 1976, Fredholm determinants and inverse scattering problems, *Comm. Math. Phys.* **47**, 171–183.
- Eastham, M.: 1973, *The spectral theory of periodic differential equations*, Scottish Academic Press, Edinburgh.
- Falloon, P. E., Abbott, P. C. and Wang, J. B.: 2003, Theory and computation of spheroidal wavefunctions, *J. Phys. A* **36**, 5477–5495.
- Fenyő, S. and Stolle, H.-W.: 1982–1984, *Theorie und Praxis der linearen Integralgleichungen. Vol. I–IV*, Birkhäuser, Basel.
- Fredholm, I.: 1900, Sur une nouvelle méthode pour la résolution du problème de Dirichlet, *Öfversigt Kongl. Vetenskaps-Akad. Förhandlingar* **57**, 39–46.
- Fredholm, I.: 1903, Sur une classe d'équations fonctionnelles, *Acta Math.* **27**, 365–390.
- Fredholm, I.: 1909, Les équations intégrales linéaires, *C. R. Congrès des Math. tenu à Stockholm 1909*.
- Gaudin, M.: 1961, Sur la loi limite de l'espacement des valeurs propres d'une matrice aléatoire, *Nucl. Phys.* **25**, 447–458.
- Gautschi, W.: 2002, Computation of Bessel and Airy functions and of related Gaussian quadrature formulae, *BIT* **42**, 110–118.
- Gohberg, I. C. and Krein, M. G.: 1969, *Introduction to the theory of linear nonselfadjoint operators*, American Mathematical Society, Providence.
- Gohberg, I., Goldberg, S. and Kaashoek, M. A.: 1990, *Classes of linear operators. Vol. I*, Birkhäuser Verlag, Basel.
- Gohberg, I., Goldberg, S. and Krupnik, N.: 2000, *Traces and determinants of linear operators*, Birkhäuser Verlag, Basel.
- Golub, G. H. and Van Loan, C. F.: 1996, *Matrix computations*, 3rd edn, Johns Hopkins University Press, Baltimore.
- Greub, W. H.: 1967, *Multilinear algebra*, Springer-Verlag, New York.
- Grothendieck, A.: 1956, La théorie de Fredholm, *Bull. Soc. Math. France* **84**, 319–384.
- Hackbusch, W.: 1995, *Integral equations: Theory and numerical treatment*, Birkhäuser Verlag, Basel.
- Hadamard, J.: 1893, Résolution d'une question relative aux déterminants, *Bull. Sci. Math.* **17**, 240–246.
- Hastings, S. P. and McLeod, J. B.: 1980, A boundary value problem associated with the second Painlevé transcendent and the Korteweg-de Vries equation, *Arch. Rational Mech. Anal.* **73**, 31–51.
- Hägg, J.: 2007, Local Gaussian fluctuations in the Airy and discrete PNG processes, [arXiv:math/0701880](https://arxiv.org/abs/math/0701880).
- Higham, N. J.: 2002, *Accuracy and stability of numerical algorithms*, 2nd edn, Society for Industrial and Applied Mathematics, Philadelphia.
- Hilbert, D.: 1904, Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen. (Erste Mitteilung), *Nachr. Ges. Wiss. Göttingen* **1904**, 49–91.
- Hilbert, D.: 1912, *Grundzüge einer allgemeinen Theorie der linearen Integralgleichungen*, Teubner, Leipzig, Berlin.
- Hille, E. and Tamarkin, J. D.: 1931, On the characteristic values of linear integral equations, *Acta Math.* **57**, 1–76.
- Hochstadt, H.: 1973, *Integral equations*, John Wiley & Sons, New York.
- Jimbo, M., Miwa, T., Mōri, Y. and Sato, M.: 1980, Density matrix of an impenetrable Bose gas and the fifth Painlevé transcendent, *Phys. D* **1**, 80–158.
- Johansson, K.: 2000, Shape fluctuations and random matrices, *Comm. Math. Phys.* **209**, 437–476.
- Johansson, K.: 2003, Discrete polynuclear growth and determinantal processes, *Comm. Math. Phys.* **242**, 277–329.
- Jost, R. and Pais, A.: 1951, On the scattering of a particle by a static potential, *Physical Rev.* **82**, 840–851.
- Katz, N. M. and Sarnak, P.: 1999, *Random matrices, Frobenius eigenvalues, and monodromy*, American Mathematical Society, Providence.
- Kline, M.: 1972, *Mathematical thought from ancient to modern times*, Oxford University Press, New York.
- Knopp, K.: 1964, *Theorie und Anwendung der unendlichen Reihen*, 5th edn, Springer-Verlag, Berlin.
- Kress, R.: 1999, *Linear integral equations*, 2nd edn, Springer-Verlag, New York.
- Laurie, D. P.: 2001, Computation of Gauss-type quadrature formulas, *J. Comput. Appl. Math.* **127**, 201–217.
- Lax, P. D.: 2002, *Functional analysis*, John Wiley & Sons, New York.

- Mehta, M. L.: 2004, *Random matrices*, 3rd edn, Elsevier/Academic Press, Amsterdam.
- Meyer, C.: 2000, *Matrix analysis and applied linear algebra*, Society for Industrial and Applied Mathematics, Philadelphia.
- Moiseiwitsch, B.: 1977, Recent progress in atomic collisions theory, *Rep. Prog. Phys.* **40**, 843–904.
- Nyström, E.: 1930, Über die praktische Auflösung von Integralgleichungen mit Anwendungen auf Randwertaufgaben, *Acta Math.* **54**, 185–204.
- Pietsch, A.: 2007, *History of Banach spaces and linear operators*, Birkhäuser, Boston.
- Plemelj, J.: 1904, Zur Theorie der Fredholmschen Funktionalgleichung, *Monatsh. f. Math.* **15**, 93–128.
- Porter, D. and Stirling, D. S. G.: 1990, *Integral equations*, Cambridge University Press, Cambridge.
- Prähofer, M. and Spohn, H.: 2002, Scale invariance of the PNG droplet and the Airy process, *J. Statist. Phys.* **108**, 1071–1106.
- Prähofer, M. and Spohn, H.: 2004, Exact scaling functions for one-dimensional stationary KPZ growth, *J. Statist. Phys.* **115**, 255–279.
- Prössdorf, S. and Silbermann, B.: 1991, *Numerical analysis for integral and related operator equations*, Birkhäuser Verlag, Basel.
- Reinhardt, W. P. and Szabo, A.: 1970, Fredholm method. I. A numerical procedure for elastic scattering, *Phys. Rev. A* **1**, 1162–1169.
- Rezende, J.: 1994, Feynman integrals and Fredholm determinants, *J. Math. Phys.* **35**, 4357–4371.
- Riess, R. D. and Johnson, L. W.: 1972, Error estimates for Clenshaw-Curtis quadrature, *Proc.* **18**, 345–353.
- Sasamoto, T.: 2005, Spatial correlations of the 1D KPZ surface on a flat substrate, *J. Phys. A* **38**, L549–L556.
- Simon, B.: 1977, Notes on infinite determinants of Hilbert space operators, *Advances in Math.* **24**, 244–273.
- Simon, B.: 2005, *Trace ideals and their applications*, 2nd edn, American Mathematical Society, Providence.
- Smithies, F.: 1937, The eigen-values and singular values of integral equations, *Proc. London Math. Soc.* **43**, 255–279.
- Smithies, F.: 1958, *Integral equations*, Cambridge University Press, Cambridge.
- Spohn, H.: 2008, Personal communication.
- Stewart, G. W.: 1998, *Matrix algorithms. Vol. I: Basic decompositions*, Society for Industrial and Applied Mathematics, Philadelphia.
- Stratton, J. A., Morse, P. M., Chu, L. J., Little, J. D. C. and Corbató, F. J.: 1956, *Spheroidal wave functions, including tables of separation constants and coefficients*, John Wiley & Sons, New York.
- Swarztrauber, P. N.: 2002, On computing the points and weights for Gauss-Legendre quadrature, *SIAM J. Sci. Comput.* **24**, 945–954.
- Tracy, C. A. and Widom, H.: 1994, Level-spacing distributions and the Airy kernel, *Comm. Math. Phys.* **159**, 151–174.
- Tracy, C. A. and Widom, H.: 2000, Universality of the distribution functions of random matrix theory, *Integrable systems: from classical to quantum (Montréal, QC, 1999)*, Vol. 26 of CRM Proc. Lecture Notes, Amer. Math. Soc., Providence, pp. 251–264.
- Trefethen, L. N.: 2008, Is Gauss quadrature better than Clenshaw-Curtis?, *SIAM Rev.* **50**, 67–87.
- Tricomi, F. G.: 1957, *Integral equations*, Interscience Publishers, Inc., New York.
- Vallée, O. and Soares, M.: 2004, *Airy functions and applications to physics*, Imperial College Press, London.
- von Koch, H.: 1892, Sur les déterminants infinis et les équations différentielles linéaires, *Acta Math.* **16**, 217–295.
- Waldvogel, J.: 2006, Fast construction of the Fejér and Clenshaw-Curtis quadrature rules, *BIT* **46**, 195–202.
- Webster, A. G.: 1927, *Partial differential equations of mathematical physics*, G. E. Stechert & Co., New York.
- Whittaker, E. T. and Watson, G. N.: 1927, *A course of modern analysis*, 4th edn, Cambridge University Press, Cambridge.
- Widom, H.: 2004, On asymptotics for the Airy process, *J. Statist. Phys.* **115**, 1129–1134.
- Wilkinson, D.: 1978, Continuum derivation of the Ising model two-point function, *Phys. Rev. D* **17**, 1629–1636.