

# Does Church-Kleene ordinal $\omega_1^{CK}$ exist?

Hitoshi Kitada

Graduate School of Mathematical Sciences

University of Tokyo

Komaba, Meguro-ku, Tokyo 153-8914, Japan

e-mail: kitada@ms.u-tokyo.ac.jp

July 7, 2003

*Abstract.* A question is proposed if a nonrecursive ordinal, the so-called Church-Kleene ordinal  $\omega_1^{CK}$  really exists.

We consider the systems  $S^{(\alpha)}$  defined in [2].

Let  $\tilde{q}(\alpha)$  denote the Gödel number of Rosser formula or its negation  $A_{(q(\alpha))}$  ( $= A_{q(\alpha)}(\mathbf{q}^{(\alpha)})$  or  $\neg A_{q(\alpha)}(\mathbf{q}^{(\alpha)})$ ), if the Rosser formula  $A_{q(\alpha)}(\mathbf{q}^{(\alpha)})$  is well-defined.

By “recursive ordinals” we mean those defined by Rogers [4]. Then that  $\alpha$  is a recursive ordinal means that  $\alpha < \omega_1^{CK}$ , where  $\omega_1^{CK}$  is the Church-Kleene ordinal.

**Lemma.** The number  $\tilde{q}(\alpha)$  is recursively defined for countable recursive ordinals  $\alpha < \omega_1^{CK}$ . Here ‘recursively defined’ means that  $\tilde{q}(\alpha)$  is defined inductively starting from 0.

**Remark.** The original meaning of ‘recursive’ is ‘inductive.’ The meaning of the word ‘recursive’ in the following is the one that matches the spirit of Kleene [3] (especially, the spirit of the inductive construction of metamathematical predicates described in section 51 of [3]).

*Proof.* The well-definedness of  $\tilde{q}(0)$  is assured by Rosser-Gödel theorem as explained in [2].

We make an induction hypothesis that for each  $\delta < \alpha$ , the Gödel number  $\tilde{q}(\gamma)$  of the formula  $A_{(q(\gamma))}$  ( $= A_{q(\gamma)}(\mathbf{q}^{(\gamma)})$  or  $\neg A_{q(\gamma)}(\mathbf{q}^{(\gamma)})$ ) with  $\gamma \leq \delta$  is recursively defined for  $\gamma \leq \delta$ .

We want to prove that the Gödel number  $\tilde{q}(\gamma)$  is recursively well-defined for  $\gamma \leq \alpha$ .

i) When  $\alpha = \delta + 1$ , by induction hypothesis we can determine recursively whether or not a given formula  $A_r$  with Gödel number  $r$  is equal to one of the axiom formulas  $A_{(\gamma)}$  ( $\gamma \leq \delta$ ) of  $S^{(\alpha)}$ . In fact, we have only to see, for a finite number of  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma \leq \delta$ , if we have  $A_{(\gamma)} = A_r$  or not. By induction hypothesis that  $\tilde{q}(\gamma)$  is recursively well-defined for  $\gamma \leq \delta$ , this is then decided recursively.

Thus Gödel predicate  $\mathbf{A}^{(\alpha)}(a, b)$  and Rosser predicate  $\mathbf{B}^{(\alpha)}(a, c)$  with superscript  $\alpha$  are recursively defined, and hence are numeralwise expressible in  $S^{(\alpha)}$ . Then the Rosser formula  $A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  is well-defined, and the Gödel number  $\tilde{q}(\alpha)$  of Rosser formula or its negation  $A_{(\alpha)}$  ( $= A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  or  $\neg A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$ ) is defined recursively. Thus  $\tilde{q}(\gamma)$  is recursively well-defined for  $\gamma \leq \alpha$ .

ii) If  $\alpha$  is a countable *recursive* limit ordinal, then there is an increasing sequence of recursive ordinals  $\alpha_n < \alpha$  such that

$$\alpha = \bigcup_{n=0}^{\infty} \alpha_n. \quad (1)$$

In the system  $S^{(\alpha)}$ , the totality of the added axioms  $A_{(\gamma)}$  ( $\gamma < \alpha$ ) is the sum of the added axioms  $A_{(\gamma)}$  ( $\gamma < \alpha_n$ ) of  $S^{(\alpha_n)}$ . By induction hypothesis,  $\tilde{q}(\gamma)$  is recursively defined for  $\gamma < \alpha_n$ . Thus in each  $S^{(\alpha_n)}$  we can determine recursively whether or not a given formula  $A_r$  is an axiom of  $S^{(\alpha_n)}$  by seeing, for a finite number of  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha_n$ , if  $A_{(\gamma)} = A_r$  or not.

This is extended to  $S^{(\alpha)}$ . To see this, we have only to see the  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$ , and determine for those finite number of  $\gamma$ 's if  $A_{(\gamma)} = A_r$  or not. By (1),

$$\tilde{q}(\gamma) \leq r \text{ and } \gamma < \alpha \Leftrightarrow \exists n \text{ such that } \tilde{q}(\gamma) \leq r \text{ and } \gamma < \alpha_n.$$

Then by induction on  $n$  with using the result in the above paragraph for  $S^{(\alpha_n)}$  and noting that the bound  $r$  on  $\tilde{q}(\gamma)$  is uniform in  $n$ , we can show that the condition whether or not  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$  is recursively determined. Whence the question whether or not a given formula  $A_r$  is one of the axioms  $A_{(\gamma)}$  of  $S^{(\alpha)}$  with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$  is determined recursively. Thus Gödel predicate  $\mathbf{A}^{(\alpha)}(a, b)$  and Rosser predicate  $\mathbf{B}^{(\alpha)}(a, c)$  with superscript  $\alpha$  are recursively defined, and hence are numeralwise expressible

in  $S^{(\alpha)}$ . Therefore the Rosser formula  $A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  is well-defined, and the Gödel number  $\tilde{q}(\alpha)$  of Rosser formula or its negation  $A_{(\alpha)}$  ( $= A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  or  $\neg A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$ ) is defined recursively. Thus  $\tilde{q}(\gamma)$  is recursively well-defined for  $\gamma \leq \alpha$ . This completes the proof of the lemma.

Assume now that  $\alpha$  is a countable limit ordinal such that there is an increasing sequence of recursive ordinals  $\alpha_n < \alpha$  with

$$\alpha = \bigcup_{n=0}^{\infty} \alpha_n. \quad (2)$$

An actual example of such an  $\alpha$  is the Church-Kleene ordinal  $\omega_1^{CK}$ .

In the system  $S^{(\alpha)}$ , the totality of the added axioms  $A_{(\gamma)}$  ( $\gamma < \alpha$ ) is the sum of the added axioms  $A_{(\gamma)}$  ( $\gamma < \alpha_n$ ) of  $S^{(\alpha_n)}$ . By the lemma,  $\tilde{q}(\gamma)$  is recursively defined for  $\gamma < \alpha_n$ . Thus in each  $S^{(\alpha_n)}$  we can determine recursively whether or not a given formula  $A_r$  is an axiom of  $S^{(\alpha_n)}$  by seeing, for a finite number of  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha_n$ , if  $A_{(\gamma)} = A_r$  or not.

This is extended to  $S^{(\alpha)}$ . To see this, we have only to see the  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$ , and determine for those finite number of  $\gamma$ 's if  $A_{(\gamma)} = A_r$  or not. By (2),

$$\tilde{q}(\gamma) \leq r \text{ and } \gamma < \alpha \Leftrightarrow \exists n \text{ such that } \tilde{q}(\gamma) \leq r \text{ and } \gamma < \alpha_n.$$

Then by induction on  $n$  with using the above result for  $S^{(\alpha_n)}$  in the preceding paragraph and noting that the bound  $r$  on  $\tilde{q}(\gamma)$  is uniform in  $n$ , we can show that the condition whether or not  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$  is recursively determined. Then within those finite number of  $\gamma$ 's with  $\tilde{q}(\gamma) \leq r$  and  $\gamma < \alpha$ , we can decide recursively if for some  $\gamma < \alpha$  with  $\tilde{q}(\gamma) \leq r$ , we have  $A_r = A_{(\gamma)}$  or not. Therefore we can determine recursively whether or not a given formula  $A_r$  is an axiom of  $S^{(\alpha)}$ .

Therefore Gödel predicate  $\mathbf{A}^{(\alpha)}(a, b)$  and Rosser predicate  $\mathbf{B}^{(\alpha)}(a, c)$  are recursively defined, and hence are numeralwise expressible in  $S^{(\alpha)}$ . Then the Gödel number  $q^{(\alpha)}$  of the formula

$$\forall b[\neg A^{(\alpha)}(a, b) \vee \exists c(c \leq b \ \& \ B^{(\alpha)}(a, c))]$$

is well-defined, and hence Rosser formula  $A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  is well-defined and Rosser-Gödel theorem applies to the system  $S^{(\alpha)}$ . Therefore we can extend

$S^{(\alpha)}$  consistently by adding one of Rosser formula or its negation  $A_{(\alpha)}$  ( $= A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$  or  $\neg A_{q^{(\alpha)}}(\mathbf{q}^{(\alpha)})$ ) to the axioms of  $S^{(\alpha)}$  and get a consistent system  $S^{(\alpha+1)}$ .

In particular if we assume a least nonrecursive ordinal  $\omega_1^{CK}$  exists and take  $\alpha = \omega_1^{CK}$ , we get a consistent system  $S^{(\omega_1^{CK}+1)}$ . This contradicts the case ii) of the theorem in [2]. We now arrive at

**Question.** The least nonrecursive ordinal, the so-called Church-Kleene ordinal  $\omega_1^{CK}$  has been assumed to give a bound on recursive construction of formal systems (see [1], [5], [6]). However the above argument seems to question if  $\omega_1^{CK}$  really exists in usual set theoretic sense. How should we think?

## References

- [1] S. Feferman, *Transfinite recursive progressions of axiomatic theories*, Journal Symbolic Logic, **27** (1962), 259-316.
- [2] H. Kitada, *Is mathematics consistent?*, 2003 (<http://arXiv.org/abs/math.GM/0306007>).
- [3] S. C. Kleene, *Introduction to Metamathematics*, North-Holland Publishing Co. Amsterdam, P. Noordhoff N. V., Groningen, 1964.
- [4] H. Rogers Jr., *Theory of Recursive Functions and Effective computability*, McGraw-Hill, 1967.
- [5] U. R. Schmerl, *Iterated reflection principles and the  $\omega$ -rule*, Journal Symbolic Logic, **47** (1982), 721-733.
- [6] A. M. Turing, *Systems of logic based on ordinals*, Proc. London Math. Soc., ser. 2, **45** (1939), 161-228.