

**Oral creatine monohydrate supplementation improves brain performance: a double-blind, placebo-controlled, cross-over trial.**

Caroline Rae<sup>1</sup>, Alison L. Digney<sup>1</sup>, Sally R. McEwan<sup>1</sup> & Timothy C. Bates<sup>2</sup>.

1. Discipline of Biochemistry, School of Molecular and Microbial Biosciences G08, The University of Sydney, Sydney, NSW, 2006, Australia.
2. Macquarie Centre for Cognitive Science, Macquarie University, NSW, 2109, Australia

Corresponding author: Dr Caroline Rae, Dept of Biochemistry G08, The University of Sydney, NSW, 2006, Australia. Ph +61 2 93513904; Fax +61 2 93514726; Email [crae@biochem.usyd.edu.au](mailto:crae@biochem.usyd.edu.au)

**Summary**

Creatine supplementation is in widespread use to enhance sports/fitness performance, and has been trialed successfully in treatment of neurological, neuromuscular and atherosclerotic disease. Creatine plays a pivotal role in brain energy homeostasis, being a temporal and spatial buffer for cytosolic and mitochondrial pools of the cellular energy currency, adenosine triphosphate (ATP) and its regulator, adenosine diphosphate. In this work, we tested the hypothesis that oral creatine supplementation (5 g per day for 6 weeks) would enhance intelligence test scores and working memory performance in 45 young adult, vegetarian subjects in a double-blind, placebo-controlled, cross-over design. Creatine supplementation had a significant positive effect ( $P < 0.0001$ ) on both working memory (backward digit span) and intelligence (Ravens Advanced Progressive Matrices), both tasks which require speed of processing. These findings underline a dynamic and significant role of brain energy capacity in influencing brain performance.

**Keywords:** creatine, oral supplementation, intelligence, memory, brain bioenergetics

Induction of work in the brain induces a cascade of biochemical and physiological sequelae in order to provide the active brain area with the major fuels glucose and oxygen. When the induced workload is heavy, the activated brain area may be temporarily fuel-limited (Fox *et al.*, 1988; Silver & Erecinska, 1994; Dienel & Hertz, 2001). Neuronal depolarisation results in restoration of  $\text{Na}^+/\text{K}^+$  gradients via activation of the  $\text{Na}^+/\text{K}^+$  ATPase. The hydrolysis of ATP by this pump is a major driver of energy metabolism in the active brain (Mata *et al.*, 1980) and an association of creatine kinase with this and other ATP-dependent ion pumps has been demonstrated (Sappey-Marinier *et al.*, 1992; Walliman *et al.*, 1992).

ATP is strongly buffered in the brain by conversion via creatine kinase catalysed reaction to phosphocreatine, the phosphorylated analogue of the guanidino amino acid creatine. ATP can be resynthesised from phosphocreatine 12 times faster than via oxidative phosphorylation and more than 70 times faster than de novo pathways (Walliman *et al.*, 1992). Phosphocreatine levels can decrease acutely upon brain activation (Sappey-Marinier *et al.*, 1992; Rango *et al.*, 1997) while ATP levels remain relatively constant. This pattern of acute decrease in local levels is also displayed by glucose (Merboldt *et al.*, 1992; Silver & Erecinska, 1994) and oxygen (Lowry & Fillenz, 1997; Madsen *et al.*, 1999).

These data indicate that the brain may be temporarily fuel-limited upon activation and suggest that brain performance might benefit from an increased supply of fuel in the initial stages of the workload. This has been demonstrated to be so for glucose (Benton *et al.*, 1994; Parker & Benton, 1995; Kennedy & Scholey, 2000; Scholey *et al.*, 2001) and oxygen (Moss *et al.*, 1998). Due to its pivotal role in energy homeostasis, creatine may similarly boost brain performance. Indeed, creatine has

been shown to be neuroprotective in various neurological conditions (Wyss & Schulze, 2002). Recently, brain creatine levels in humans have been shown to increase in response to mental training (Valenzuela *et al.*, 2002) and acute oral creatine supplementation has been shown to reduce mental fatigue and decrease task-responsive oxygen delivery (demand) to activated areas on performance of a serial calculation task (Watanabe *et al.*, 2002).

We sought to determine whether a beneficial effect on mental performance could occur if the creatine reservoir were increased by oral supplementation. Oral creatine (5 g/day for 6 weeks) has previously been shown to increase (by ~9%) creatine levels in the brains of omnivores (Dechent *et al.*, 1999). Brain creatine is derived from both synthesis and diet (Wyss & Kaddurah-Daouk, 2000). The dietary source is restricted to animal sources, such as meat. However, supplementation can only increase levels to a saturated value, beyond which excess creatine is excreted. To obtain maximal increase by supplementation, we chose to examine the effect of oral creatine monohydrate supplementation on young adults with vegetarian diets, in whom creatine levels are lower than they are in omnivores (Delanghe *et al.*, 1989). We note here that creatine supplementation in vegetarians results in the same ergonomic increases in muscle performance as it does in omnivores, despite vegetarians having lower tissue creatine prior to supplementation (Shomrat *et al.*, 2000).

## **Materials and Methods**

**Subjects.** 45 vegan or vegetarian subjects (12 males (median age 27.5, range 19-37 y), 33 females (median age 24.9, range 18-40 y); 18 vegan (median duration 4.6 y, range 0.7 – 17 y) and 27 vegetarian (median duration 14.3, range 1 – 23 y)) were recruited with informed consent from amongst the student population of The University of Sydney. Subjects with a medical history of drug/alcohol abuse, diagnosed psychiatric disorders, diabetes or who showed evidence of renal

insufficiency were excluded from the study. This study was approved by the University of Sydney and Macquarie University Human Ethics committees.

**Study Design.** The study followed a double-blind, placebo-controlled, cross-over design. Subjects were seen on 4 separate occasions, at 6 week intervals, following an overnight fast to minimise fluctuations in blood glucose. At each visit, a blood sample was taken to measure red cell and plasma creatine levels and blood glucose concentration. Red blood cells contain no creatine transporter so their creatine content represents creatine content in the tissue where the red blood cell was made. Red cells live for an average of 120 days so red cell creatine content is a delayed, averaged measure of tissue creatine levels and represents the best sample of tissue creatine when tissue biopsy is not possible.

A cognitive test battery was also administered. At the end of the first and third test sessions, subjects were given an envelope marked with their study number and containing 5 g doses of supplement (creatine monohydrate ((2-methylguanido)acetic acid; Pan Pharmaceuticals, Australia)) or placebo (maltodextrin, (Manildra Starches, Australia)) in plastic vials. Subjects were asked to consume this supplement at the same time each day for the next 6 weeks and received advice on how best to take this supplement to ensure maximum solubility and absorption. Subjects returned the envelope with unused vials at the end of each 6 week period and the number of vials remaining was used to assess compliance, validated against increases in red cell (tissue) creatine. Between visits 2 and 3, subjects consumed no supplement. Six weeks has been shown to be an adequate “wash out” period (Harris *et al.*, 1992).

**Cognitive tests.** Subjects completed timed (10 min) parallel versions of Raven's Advanced Progressive Matrices (RAPM) constructed to have equal levels of difficulty based on the published normative performance data and verified by us on an independent sample of 20 subjects. This intelligence test is well validated as a measure of general ability with minimal dependence on cultural factors (Raven *et al.*, 1988). Versions of these tests were assigned evenly across the four visits. Subjects also completed the Wechsler Auditory Backward Digit Span (BDS) task (Wechsler, 1955). This subtest of the Wechsler Adult Intelligence Scales loads on both short term storage and verbal working memory functions supported by mid ventro-lateral and dorsolateral prefrontal cortex, respectively (Owen *et al.*, 2000).

**Creatine and glucose measurements.** Creatine concentrations in both blood plasma and in red blood cells were measured (Griffiths, 1968) on the morning of testing. Blood glucose levels were measured randomly, by finger prick using a hand-held glucometer, to monitor compliance with the request for an overnight fast prior to presenting for testing. None of the blood glucose levels recorded suggested that any of the subjects had violated this request.

## **Results**

Measured by the number of vials of creatine returned, compliance for creatine supplement was mean 91% (range 85 – 100%) and 97% (range 88- 100%) for placebo. Red blood cell creatine levels, indicative of tissue creatine levels, increased significantly with supplementation (  $P = 0.001$ ) compared to placebo indicating that tissue creatine levels had been increased by the supplement. Plasma creatine levels, more indicative of acute creatine ingestion, did not vary significantly with supplementation.

Supplementation with oral creatine monohydrate significantly increased intelligence (as measured by Ravens Advanced Progressive Matrices done under time pressure, Fig. 1A) compared to placebo ( $F(3, 33) = 32.3$ ,  $P < 0.0001$ ; repeated measures ANOVA). There was no significant effect of treatment order ( $F(1,33) = 1.62$ ,  $P = 0.21$ ) although there was a significant interaction with treatment order ( $F(3,99) = 6.7$ ,  $P = 0.0004$ ). The mean Ravens Advanced Progressive Matrices raw score under placebo was 9.7 (SD = 3.8) items correct in 10 min versus 13.7 (SD = 4.1) items correct under the experimental treatment. Supplementation with oral creatine monohydrate (Fig. 1B) significantly affected performance on backward digit span ( $F(3,34) = 29.0$ ,  $P < 0.0001$ ), with no effect of order ( $F(3,102) = 0.98$ ,  $P = 0.40$ ). Mean backward digit span under placebo was 7.05 items (SD = 1.19), compared to a mean of 8.5 items under creatine treatment (SD = 1.76).

## Discussion

This study showed that increasing creatine intake by oral supplementation resulted in improved brain function, similar to effects shown previously in muscle and heart. These results are in agreement with observations showing that brain creatine levels correlate positively with recognition memory (Ferrier *et al.*, 2001) and that creatine supplementation reduces mental fatigue on a serial calculation task (Watanabe *et al.*, 2002). This latter study also showed reduced activation-stimulated oxyhaemoglobin delivery to the activated area following creatine supplementation. This suggests that creatine supplementation is acting to smooth fluctuations in the blood oxygen level dependent response curve which results from brain activation (Gjedde *et al.*, 1999; Madsen *et al.*, 1999), possibly by altering rates of ATP synthesis in the mitochondrion through the mitochondrial creatine kinase/adenine nucleotide translocase/porin complex (Walliman *et al.*, 1992; Saks *et al.*, 2000). Most recently, deletion of cytosolic brain-type creatine kinase in mice was shown to result in slower learning of a spatial task and diminished open-field habituation as well as increased intra and infra-pyramidal hippocampal mossy fibre area suggesting that the creatine/creatine kinase

network is also involved in brain plasticity in addition to metabolism (Jost *et al.*, 2002).

It is not currently known whether brain creatine levels are lower in vegetarian subjects than omnivores although consideration of factors affecting brain creatine levels suggests that this ought certainly to be the case. It is known controlled by both the rates of synthesis and transport of creatine that plasma levels are lower in vegetarians (Delanghe *et al.*, 1989) but creatine supplementation of nto cells in the brain. This is illustrated by the fact that deficits in the creatine transporter or in the enzymes which synthesise creatine result in similar neurological outcomes (Stöckler *et al.*, 1996; Salomons *et al.*, 2001). It has also been suggested that a “creatin cycle” exists between cells in the brain so that a combination of synthesis and transport routes would be required to maintain brain creatine levels (Möller & Hamprecht, 1989; Dringen *et al.*, 1998). In vegetarians it is likely that the synthesis route is upregulated. Chronic ingestion of creatine has been shown to result in down-regulation of the vegetarians has been shown to produce similar increases in muscle performance to that seen in omnivores (Shomrat *et al.*, 2000). Creatine levels in the brain are into the brain and icreatine transporter (Guerrero-Ontiveros & Wallimann, 1998). We would therefore expect to see a beneficial effect of creatine supplementation on brain performance in the vast majority of omnivores apart from those who consume very high amounts of meat (~2 kg day).

Given that Raven’s Advanced Progressive Matrices is a rather pure measure of *g*, or general ability, the data suggest that general ability was improved by creatine supplementation. These results support efficiency models of intelligence (Bates & Stough, 1998). On this view, differences in intelligence test performance reflect individual differences in underlying biochemical and structural factors influencing the energetic and temporal resources of the central nervous system. Increasing the energy available for computation increases the speed and (in a distributed computational

system such as the brain) power of computational resource, reflected directly in improved general ability.

Long term supplementation with creatine has yet to be declared truly safe, with reported effects on glucose homeostasis (Rooney *et al.*, 2003) and other side effects (Terjung *et al.*, 2000). This trial of creatine supplementation showed beneficial effects of creatine on mental performance. These effects may add to the physical enhancement gained by athletes supplementing creatine levels and may be of use to those requiring boosted mental performance in the short term.

### **Acknowledgements**

This work was supported by the Australian National Health and Medical Research Council.

## References

- Bates, T. and Stough, C. (1998). Improved reaction time method, information processing speed and intelligence. *Intelligence* **26**.
- Benton, D., Owens, D. S. and Parker, P. Y. (1994). Blood glucose influences memory and attention in young adults. *Neuropsychologia* **32**, 595-607.
- Dechent, P., Pouwels, P. J. W., Wilken, B., Hanefeld, F. and Frahm, J. (1999). Increase of total creatine in human brain after oral supplementation of creatine-monohydrate. *Am J Physiol* **277**, R698-704.
- Delanghe, J., De Slypere, J. P., De Buyzere, M., Robbrecht, J., Wieme, R. and Vermeulen, A. (1989). Normal reference values for creatine, creatinine and carnitine are lower in vegetarians. *Clin Chem* **35**, 1802-1803.
- Dienel, G. A. and Hertz, L. (2001). Glucose and lactate metabolism during brain activation. *J Neurochem Res* **66**, 824-838.
- Dringen, R., Verleysdonk, S., Hamprecht, B., Wilker, W., Leibfritz, D. and Brand, A. (1998). Metabolism of glycine in primary astroglial cells: synthesis of creatine, serine and glutathione. *J Neurochem* **70**, 835-840.
- Ferrier, C. H., Alarcon, G., Glover, A., Koutroumanidis, M., Morris, R. G., Simmons, A., Elwes, R. D., Cox, T., Binnie, C. D. and Polkey, C. E. (2001). NAA and creatine levels measured by <sup>1</sup>H MRS relate to recognition memory. *Neurology* **55**, 1874-1883.
- Fox, P. T., Raichle, M. E., Mintun, M. A. and Dence, C. (1988). Non-oxidative glucose consumption during focal physiological neuronal activity. *Science* **241**, 462-464.
- Gjedde, A., Poulsen, P. H. and Østergaard, L. (1999). On the oxygenation of hemoglobin in the human brain. *Adv Exp Med Biol* **471**, 97-81.

- Griffiths, W. J. (1968). Estimation of creatine in red cells. *J Clin Path* **21**, 412-413.
- Guerrero-Ontiveros, M. L. and Wallimann, T. (1998). Creatine supplementation in health and disease. Effects of chronic creatine ingestion in vivo: Down-regulation of the expression of creatine transporter isoforms in skeletal muscle. *Mol Cell Biochem* **184**, 427-437.
- Harris, R. C., Söderlund, K. and Hultman, E. (1992). Elevation of creatine in resting and exercised muscle of normal subjects by creatine supplementation. *Clin Sci* **83**, 367-374.
- Jost, C. R., Van der Zee, C. E. E. M., in 'tZandt, H. J. A., Oerlemans, F., Verheij, M., Streijger, F., Fransen, J., Heerschap, A., Cools, A. R. and Weiringa, B. (2002). Creatine kinase B-driven energy transfer in the brain is important for habituation and spatial learning behaviour, mossy fibre field size and determination of seizure susceptibility. *Eur J Neurosci* **15**, 1692-1706.
- Kennedy, D. O. and Scholey, A. B. (2000). Glucose administration, heart rate and cognitive performance: effects of increasing mental effort. *Psychopharmacol* **149**, 63-71.
- Lowry, J. P. and Fillenz, M. (1997). Evidence for uncoupling of oxygen and glucose utilisation during neuronal activation in rat striatum. *J Physiol* **498.2**, 497-501.
- Madsen, P. L., Cruz, N. F., Sokoloff, L. and Dienel, G. A. (1999). Cerebral oxygen/glucose ratio is low during sensory stimulation and rises about normal during recovery: excess glucose consumption during stimulation is not accounted for by lactate efflux from or accumulation in brain tissue. *J Cereb Blood Flow Metab* **19**, 393-400.
- Mata, M., Fink, D. J., Gainer, H., Smith, C. B., Davidsen, L., Savaki, H., Schwartz, W. J. and Sokoloff, L. (1980). Activity-dependent energy metabolism in rat

- posterior pituitary primarily reflects sodium pump activity. *J Neurochem* **34**, 213-215.
- Merboldt, K.-D., Bruhn, H., Hänicke, W., Michaelis, T. and Frahm, J. (1992).  
Decrease of glucose in the human visual cortex during photic stimulation.  
*Magn Reson Med* **25**, 187-194.
- Möller, A. and Hamprecht, B. (1989). Creatine transport in cultured cells of rat and mouse brain. *J Neurochem* **52**, 544-550.
- Moss, M. C., Scholey, A. B. and Wesnes, K. (1998). Oxygen administration selectively enhances cognitive performance in healthy young adults; a placebo controlled double blind crossover study. *Psychopharmacol* **138**, 27-33.
- Owen, A. M., Lee, A. C. H. and Williams, E. J. (2000). Dissociating aspects of verbal working memory within the human frontal lobe: Further evidence for a "process specific" model of lateral frontal organization. *Psychobiol* **28**, 146-155.
- Parker, P. Y. and Benton, D. (1995). Blood glucose levels selectively influence memory for word lists dichotically presented to the right ear.  
*Neuropsychologia* **33**, 843-854.
- Rango, M., Castelli, A. and Scarlato, G. (1997). Energetics of 3.5 s neural activation in humans: a <sup>31</sup>P spectroscopy study. *Magn Reson Med* **38**, 878-883.
- Raven, J. C., Court, J. H. and Raven, J. (1988). *Manual for Raven's Progressive Matrices and Vocabulary Scales*. London, H.K. Lewis and Co, Ltd.
- Rooney, K., Bryson, J., Digney, A. L., Rae, C. D. and Thompson, C. H. (2003). Creatine supplementation affects glucose homeostasis but not insulin secretion in vegetarians. *Annal Nutr Metab* **47**

- Saks, V. A., Kongas, O., Vendelin, M. and Kay, L. (2000). Role of the creatine/phosphocreatine system in the regulation of mitochondrial respiration. *Acta Physiol Scand* **168**, 635-641.
- Salomons, G. S., van Dooren, S. J. M., Verhoeven, N. M., Cecil, K. M., Ball, W. S., Degrauw, T. J. and Jakobs, C. (2001). X-linked creatine-transporter gene (SLC6A8) defect: A new creatine-deficiency syndrome. *Am J Hum Genet* **68**, 1497-1500.
- Sappey-Marinier, D., Calabrese, G., Fein, G., Hugg, J. W., Biggins, C. and Weiner, M. W. (1992). Effect of photic stimulation on human visual cortex lactate and phosphates using <sup>1</sup>H and <sup>31</sup>P magnetic resonance spectroscopy. *J Cereb Blood Flow Metab* **12**, 584-592.
- Scholey, A. B., Harper, S. and Kennedy, D. O. (2001). Cognitive demand and blood glucose. *Physiol Behav* **73**, 585-592.
- Shomrat, A., Weinstein, Y. and Katz, A. (2000). Effect of creatine feeding on maximal exercise performance in vegetarians. *Eur J Appl Physiol* **82**, 321-325.
- Silver, I. A. and Erecinska, M. (1994). Extracellular glucose concentration in mammalian brain: continuous monitoring of changes during increased neuronal activity and upon limitation in oxygen supply in normo-, hypo- and hyperglycaemic animals. *J Neurosci* **14**, 5068-5076.
- Stöckler, S., Hanefeld, F. and Frahm, J. (1996). Creatine replacement therapy in guanidinoacetate methyltransferase deficiency, a novel inborn error of metabolism. *Lancet* **348**, 789-790.
- Terjung, R. L., Clarkson, P., Eichner, E. R., Greenhaff, P. L., Hespel, P. J., Israel, R. G., Kraemer, W. J., Meyer, R. A., Spriet, L. L., Tarnopolsky, M. A.,

- Wagenmakers, A. J. and Williams, M. H. (2000). The physiological and health effects of oral creatine supplementation. *Med Sci Sport Exercise* **32**, 706-717.
- Valenzuela, M. J., Jones, M., Sachdev, P., Wen, W., Rae, C. and Scott, G. (2003). Memory exercise alters creatine and choline in hippocampus of healthy elderly. *Neuroreport* In press.
- Walliman, T., Wyss, M., Brdiczka, D., Nicolay, K. and Eppenberger, H. M. (1992). Intracellular compartmentation, structure and function of creatine kinase isoenzyme in tissues with high and fluctuating energy demands: the phosphocreatine circuit for cellular energy homeostasis. *Biochem J* **281**, 21-40.
- Watanabe, A., Kato, N. and Kato, T. (2002). Effects of creatine on mental fatigue and cerebral hemoglobin oxygenation. *Neurosci Res* **42**, 279-285.
- Wechsler, D. (1955). *Adult Intelligence Scale Manual*. New York, Psychological Corporation.
- Wyss, M. and Kaddurah-Daouk, R. (2000). Creatine and creatinine metabolism. *Physiol Rev* **80**, 1107-1212.
- Wyss, M. and Schulze, A. (2002). Health implications of creatine: can oral creatine supplementation protect against neurological and atherosclerotic disease? *Neurosci* **112**, 243-260.

### Figure captions

**Fig. 1. Variation in mean test scores with supplement and placebo** A. Raven's Progressive Matrices under time pressure and B, Backward Digit Span. Square points represent group 1 and triangle points represent group 2 subjects. Bars represent standard errors.

**Short title:** Oral creatine improves brain performance

