

BERNOULLI NUMBERS, HOMOTOPY GROUPS, AND A THEOREM OF ROHLIN

By JOHN W. MILNOR AND MICHEL A. KERVAIRE

A homomorphism $J: \pi_{k-1}(\mathbf{SO}_m) \rightarrow \pi_{m+k-1}(S^m)$ from the homotopy groups of rotation groups to the homotopy groups of spheres has been defined by H. Hopf and G. W. Whitehead^[16]. This homomorphism plays an important role in the study of differentiable manifolds. We will study its relation to one particular problem: the question of possible Pontrjagin numbers of an 'almost parallelizable' manifold.

Definition. A connected differentiable manifold M^k with base point x_0 is *almost parallelizable* if $M^k - x_0$ is parallelizable. If M^k is imbedded in a high-dimensional Euclidean space R^{m+k} ($m \geq k+1$) then this is equivalent to the condition that the normal bundle ν , restricted to $M^k - x_0$, be trivial (compare the argument given by Whitehead^[17], or Kervaire^[5], § 8).

The following theorem was proved by Rohlin in 1952 (see Rohlin^[11, 12], Kervaire^[6]).

Theorem (Rohlin). Let M^4 be a compact oriented differentiable 4-manifold with Stiefel-Whitney class w_2 equal to zero. Then the Pontrjagin number $p_1[M^4]$ is divisible by 48.

Rohlin's proof may be sketched as follows. It may be assumed that M^4 is a connected manifold imbedded in R^{m+4} , $m \geq 5$.

Step 1. It is shown that M^4 is almost parallelizable.

Let f be a cross-section of the normal \mathbf{SO}_m -bundle ν restricted to $M^4 - x_0$. The obstruction to extending f is an element

$$\vartheta(\nu, f) \in H^4(M^4; \pi_3(\mathbf{SO}_m)) \approx \pi_3(\mathbf{SO}_m).$$

Step 2. It is shown that $J\vartheta(\nu, f) = 0$.

Since J carries the infinite cyclic group $\pi_3(\mathbf{SO}_m)$ onto the cyclic group $\pi_{m+3}(S^m)$ of order 24, this implies that $\vartheta(\nu, f)$ is divisible by 24. Now identify the group $\pi_3(\mathbf{SO}_m)$ with the integers.

Step 3. It is shown that the Pontrjagin class $p_1(\nu)$ is equal to $\pm 2\vartheta(\nu, f)$.

Since by Whitney duality $p_1(\nu) = -p_1$ (tangent bundle), it follows that $p_1[M^4]$ is divisible by 48.

The first step in this argument does not generalize to higher dimensions. However Step 2, the assertion that $J\vartheta(\nu, f) = 0$, generalizes immediately. In fact we have:

Lemma 1. Let $\alpha \in \pi_{k-1}(\mathbf{SO}_m)$; then $J\alpha = 0$ if and only if there exists an almost parallelizable manifold $M^k \subset R^{m+k}$ and a cross-section f of the induced normal \mathbf{SO}_m -bundle ν over $M^k - x_0$ such that $\alpha = \nu(\nu, f)$.

Step 3 can be replaced by the following. Identify the group $\pi_{4n-1}(\mathbf{SO}_m)$, $m > 4n$, with the integers (compare Bott^[2]). Define a_n to be equal to 2 for n odd and 1 for n even.

Lemma 2. Let ξ be a stable \mathbf{SO}_m -bundle over a complex K ($\dim K < m$), and let f be a cross-section of ξ restricted to the skeleton $K^{(4n-1)}$. Then the obstruction class $\nu(\xi, f) \in H^{4n}(K; \pi_{4n-1}(\mathbf{SO}_m))$ is related to the Pontrjagin class $p_n(\xi)$ by the identity $p_n(\xi) = \pm a_n \cdot (2n - 1)! \nu(\xi, f)$.

Combining Lemmas 1 and 2, we obtain the following theorems.

Define j_n as the order of the finite cyclic group $J\pi_{4n-1}(\mathbf{SO}_m)$ in the stable range $m > 4n$.

Theorem 1. The Pontrjagin number $p_n[M^{4n}]$ of an almost parallelizable $4n$ -manifold is divisible by $j_n a_n (2n - 1)!$.

(For $n = 1$, this gives Rohlin's assertion, since $j_1 = 24$, $a_1 = 2$.)

Proof. This follows since $\nu(\nu, f)$ must be divisible by j_n .

Conversely:

Theorem 2. There exists an almost parallelizable manifold M_0^{4n} with

$$p_n[M_0^{4n}] = j_n a_n \cdot (2n - 1)!.$$

The proof is clear.

Proof of Lemma 1. Given an imbedding $i: V^{k-1} \rightarrow R^{m+k-1}$ of a compact differentiable manifold V^{k-1} into Euclidean space, and given a cross-section f of the normal \mathbf{SO}_m -bundle over V^{k-1} , a well-known procedure due to Thom associates with i and f a sphere mapping $\phi: S^{m+k-1} \rightarrow S^m$ (compare Kervaire^[5], p. 223).

The map ϕ is homotopic to zero if and only if there exists a bounded manifold Q^k with boundary V^{k-1} imbedded in R^{m+k} on one side of R^{m+k-1} such that:

(i) the restriction to V^{k-1} of the imbedding of Q^k is the given imbedding of V^{k-1} in R^{m+k-1} ;

(ii) Q^k meets R^{m+k-1} orthogonally so that the restriction to V^{k-1} of the normal bundle of Q^k is just the normal bundle of V^{k-1} in R^{m+k-1} ; and

(iii) the cross-section f can be extended throughout Q^k as a cross-section f' of the normal \mathbf{SO}_m -bundle.

These facts follow from Thom^[15], ch. I, § 2 and Lemmas IV, 5, IV.5'.

To obtain Lemma 1 above, take $V^{k-1} = S^{k-1}$ and take $i(S^{k-1})$ to be the unit sphere in $R^k \subset R^{m+k-1}$. Since the normal m -plane at each point of $i(S^{k-1})$ in R^{m+k-1} admits a natural basis (consisting of the radius vector followed by the vectors of a basis for R^{m+k-1}/R^k), the cross-section f

provides a mapping $a: S^{k-1} \rightarrow \mathbf{SO}_m$. Let $\alpha \in \pi_{k-1}(\mathbf{SO}_m)$ be its homotopy class. It is easily seen (compare Kervaire^[7], §1.8) that the map $\phi: S^{m+k-1} \rightarrow S^m$ associated with i and f represents $J\alpha$ up to sign.

If $J\alpha = 0$ then there exists a bounded manifold $Q^k \subset R^{m+k}$ satisfying conditions (i), (ii) and (iii). Let $M^k \subset R^{m+k}$ denote the unbounded manifold obtained from Q^k by adjoining a k -dimensional hemisphere, which lies on the other side of R^{m+k-1} and has the same boundary $i(S^{k-1})$. Since the normal bundle ν restricted to Q^k has a cross-section f' , it follows that M^k is almost parallelizable. Clearly the obstruction class $\nu(\nu, f')$ is equal to α .

Conversely, let M^k be a manifold imbedded in R^{m+k} and let f be a cross-section of the normal bundle ν restricted to $M^k - x_0$. After modifying this imbedding by a diffeomorphism of R^{m+k} we may assume that some neighborhood of x_0 in M^k is a hemisphere lying on one side of the hyperplane R^{m+k-1} , and that the rest of M^k lies on the other side. Removing this neighborhood we obtain a bounded manifold $Q^k \subset R^{m+k}$ just as above, having the unit sphere $S^{k-1} \subset R^k \subset R^{m+k-1}$ as boundary. The cross-section f restricted to S^{k-1} gives rise to a map $a: S^{k-1} \rightarrow \mathbf{SO}_m$ which represents the homotopy class $\nu(\nu, f)$. The argument above shows that $J\nu(\nu, f) = 0$; which completes the proof of Lemma 1.

Remark. Lemma 1 could also be proved using the interpretation of J given in Milnor^[9].

Proof of Lemma 2. (Compare Kervaire^[8].) The \mathbf{SO}_m -bundle ξ induces a \mathbf{U}_m -bundle ξ' and hence a $\mathbf{U}_m/\mathbf{U}_{2n-1}$ -bundle ξ'' . Similarly, the partial cross-section f induces partial cross-sections f' and f'' . By definition the obstruction class $\nu(\xi'', f'')$ is equal to the Chern class $c_{2n}(\xi')$ and hence to the Pontrjagin class $\pm p_n(\xi)$. Therefore $p_n(\xi)$ equals $\pm q_* h_* \nu(\xi, f)$, where

$$h: \pi_{4n-1}(\mathbf{SO}_m) \rightarrow \pi_{4n-1}(\mathbf{U}_m) \quad \text{and} \quad q: \pi_{4n-1}(\mathbf{U}_m) \rightarrow \pi_{4n-1}(\mathbf{U}_m/\mathbf{U}_{2n-1})$$

are the natural homomorphisms and h_*, q_* are the homomorphisms in the cohomology of K induced by the coefficient homomorphisms h, q .

Using the following computations of Bott^[2]:

$$\pi_{4n-1}(\mathbf{U}_m) \approx \mathbf{Z}, \quad \pi_{4n-1}(\mathbf{U}_m/\mathbf{SO}_m) \approx \mathbf{Z}_{a_n}, \quad \pi_{4n-2}(\mathbf{SO}_m) = 0,$$

it follows that h carries a generator into a_n times a generator. Similarly, using the fact that

$$\pi_{4n-2}(\mathbf{U}_{2n-1}) \approx \mathbf{Z}_{(2n-1)!} \quad (\text{see } [3]) \quad \text{and} \quad \pi_{4n-2}(\mathbf{U}_m) = 0,$$

it follows that q carries a generator into $(2n-1)!$ times a generator. Therefore $p_n(\xi) = \pm a_n(2n-1)! \nu(\xi, f)$. This completes the proof of Lemma 2.

Hirzebruch's index theorem^[4] states that the index $I(M^{4n})$ of any $4n$ -manifold is equal to

$$2^{2n}(2^{2n-1} - 1) B_n p_n[M^{4n}]/(2n)! + (\text{terms involving lower Pontrjagin classes}).$$

Here B_n denotes the n th Bernoulli number. For an almost parallelizable manifold the lower Pontrjagin classes are zero. Therefore

Corollary. The index $I(M_0^{4n})$ is equal to $2^{2n-1}(2^{2n-1} - 1) B_n j_n a_n/n$; and the index of any almost parallelizable $4n$ -manifold is a multiple of this number.

The fact that $I(M_0^{4n})$ is an integer can be used to estimate the number j_n (compare Milnor^[9]). However, a sharper estimate, which includes the prime 2, can be obtained as follows, using a new generalization of Rohlin's theorem.

Borel and Hirzebruch (^[1], §§ 23.1 and 25.4) define a rational number

$$\hat{A}[M^{4n}] = -B_n p_n[M^{4n}]/2(2n)! + (\text{terms involving } p_1, \dots, p_{n-1});$$

and prove that the denominator of $\hat{A}[M^{4n}]$ is a power of 2.

Theorem 3. If the Stiefel-Whitney class w_2 of M^{4n} is zero then $\hat{A}[M^{4n}]$ is actually an integer.†

The proof will be given in a subsequent paper by Borel and Hirzebruch. It is based on the methods of ^[1], together with the assertion that the Todd genus of a generalized almost complex manifold is an integer (Milnor^[10]).

Applying this theorem to the manifold M_0^{4n} of Theorem 2 it follows that $B_n j_n a_n/4n$ is an integer. Therefore:

Theorem 4. The order j_n of the stable group $J\pi_{4n-1}(\mathbf{SO}_m)$ is a multiple of the denominator of the rational number $B_n a_n/4n$.

As examples, for $n = 1, 2, 3$, the number $B_n a_n/4n$ is equal to $1/12$, $1/240$, and $1/252$ respectively. Since $\pi_{m+7}(S^m)$ is cyclic of order 240, it follows that $j_2 = 240$. Since $\pi_{m+11}(S^m)$ is cyclic of order 504, it follows that j_3 is either 252 or 504. It may be conjectured that j_n is always equal to the denominator of $B_n/4n$.

The theorems of von Staudt^[13,14] can be used to compute such denominators (compare Milnor^[9]).

Lemma 3. The denominator of $B_n/2n$ can be described as follows. A prime power p^{i+1} divides this denominator if and only if

$$2n \equiv 0 \pmod{p^i(p-1)}.$$

† See note at the end of the paper.

Combining Lemma 3 with Theorem 4, we see that the stable homotopy groups of spheres contain elements of arbitrary finite order. In fact:

Corollary. *If $2n$ is a multiple of the Euler Φ function $\Phi(r)$, then the stable group $\pi_{m+4n-1}(S^m)$ contains an element of order r .*

REFERENCES

- [1] Borel, A. and Hirzebruch, F. Characteristic classes and homogeneous spaces. *Amer. J. Math.* 80, 458–538 (1958).
- [2] Bott, R. The stable homotopy of the classical groups. *Proc. Nat. Acad. Sci., Wash.*, 43, 933–935 (1957).
- [3] Bott, R. and Milnor, J. W. On the parallelizability of the spheres. *Bull. Amer. Math. Soc.* 64, 87–89 (1958).
- [4] Hirzebruch, F. *Neue topologische Methoden in der algebraischen Geometrie*. Springer, 1956.
- [5] Kervaire, M. A. Courbure intégrale généralisée et homotopie. *Math. Ann.* 131, 219–252 (1956).
- [6] Kervaire, M. A. On the Pontryagin classes of certain $SO(n)$ -bundles over manifolds. *Amer. J. Math.* 80, 632–638 (1958).
- [7] Kervaire, M. A. An interpretation of G. Whitehead's generalization of Hopf's invariant. *Ann. Math. (2)*, 69, 345–365 (1959).
- [8] Kervaire, M. A. A note on obstructions and characteristic classes. (In preparation.)
- [9] Milnor, J. W. On the Whitehead homomorphism J . *Bull. Amer. Math. Soc.* 64, 79–82 (1958).
- [10] Milnor, J. W. On the cobordism ring Ω^* , and a complex analogue. (In preparation.)
- [11] Rohlin, V. A. Classification of mappings of S^{n+3} onto S^n . (Russian.) *Dokl. Akad. Nauk, SSSR*, 81, 19–22 (1951).
- [12] Rohlin, V. A. New results in the theory of 4-dimensional manifolds. (Russian.) *Dokl. Akad. Nauk, SSSR*, 84, 221–224 (1952).
- [13] von Staudt. *J. Math.* 21, 372–374 (1840).
- [14] von Staudt. *De numeris Bernoullianis commentatio*. Erlangen, 1845.
- [15] Thom, R. Quelques propriétés globales des variétés différentiables. *Comment. math. Helvet.* 28, 17–86 (1954).
- [16] Whitehead, G. W. On the homotopy groups of spheres and rotation groups. *Ann. Math. (2)*, 43, 634–640 (1942).
- [17] Whitehead, J. H. C. On the homotopy type of manifolds. *Ann. Math. (2)*, 41, 825–832 (1940).

[*Added in proof.*] For the case n odd, Hirzebruch has since sharpened Theorem 3, showing that $\hat{A}[M^n]$ is an even integer. Thus the factor a_n in Theorem 4 can be cancelled.