## ROBUST SOLUTIONS TO UNCERTAIN SEMIDEFINITE PROGRAMS\*

FRANCOIS OUSTRY, LAURENT EL GHAOUI<sup>†</sup> AND HERVÉ LEBRET ECOLE NATIONALE SUPÉRIEURE DE TECHNIQUES AVANCÉES, 32, BD. VICTOR, 75739 PARIS, FRANCE.

INTERNET: (OUSTRY, ELGHAOUI, LEBRET)@ENSTA.FR

Abstract. In this paper we consider semidefinite programs (SDPs) whose data depends on some unknown-but-bounded perturbation parameters. We seek "robust" solutions to such programs, that is, solutions which minimize the (worst-case) objective while satisfying the constraints for every possible values of parameters within the given bounds. Assuming the data matrices are rational functions of the perturbation parameters, we show how to formulate sufficient conditions for a robust solution to exist, as SDPs. When the perturbation is "full", our conditions are necessary and sufficient. In this case, we provide sufficient conditions which guarantee that the robust solution is unique, and continuous (Hölder-stable) with respect to the unperturbed problems' data. The approach can thus be used to regularize ill-conditioned SDPs. We illustrate our results with examples taken from linear programming, maximum norm minimization, polynomial interpolation and integer programming.

**Key Words.** convex optimisation, semidefinite programming, uncertainty, robustness, regularization.

**Notation.** For a matrix X, ||X|| denotes the largest singular value. If X is square,  $X \geq 0$  (resp. X > 0) means X is symmetric, and positive semidefinite (resp. definite). For  $X \geq 0$ ,  $X^{1/2}$  denotes the symmetric square root of X. The notation  $I_p$  denotes the  $p \times p$  identity matrix; the subscript is omitted when it can be inferred from context.

1. Introduction. A semidefinite program (SDP) consists of minimizing a linear objective under a linear matrix inequality (LMI) constraint, precisely

(1) 
$$\mathcal{P}_0$$
: minimize  $c^T x$  subject to  $F(x) = F_0 + \sum_{i=1}^m x_i F_i \ge 0$ 

where  $c \in \mathbf{R}^m - \{0\}$ , and the symmetric matrices  $F_i = F_i^T \in \mathbf{R}^{n \times n}$ , i = 0, ..., m are given. SDPs are convex optimization problems and can be solved in polynomial-time with e.g. primal-dual interior-point methods [22, 33, 24, 17, 2]. SDPs include linear programs and convex quadratically constrained quadratic programs, and arise in a wide range of engineering applications, see e.g. [10, 1, 33, 20].

In the SDP (1), the "data" consists of the objective vector c and the matrices  $F_0, \ldots, F_m$ . In practice, this data is subject to uncertainty. An extensive body of work has concentrated on the sensitivity issue, in which the perturbations are assumed infinitesimal, and regularity of optimal values and solution(s), as functions of the data matrices, is studied. Recent references on sensitivity analysis include [28, 29, 8] for general nonlinear programs, [31] for semiinfinite programs and [30] for semidefinite programs.

<sup>\*</sup> Accepted for publication in SIAM J. Optimization, September 1997.

<sup>†</sup> Author to whom correspondence should be sent.

When the perturbation affecting the data of the problem is not necessarily small, a sensitivity analysis is not sufficient. For general optimisation problems, a whole field of study (stochastic programming) concentrates on the case when the perturbation is stochastic, with known statistics. One object of this field is to study the impact of say, a random objective on the distribution of optimal values (this problem is called the "distribution problem"). References relevant to this approach to the perturbation problem include [13, 7, 27]. We are not aware of special references for general SDPs with randomly perturbed data, except for the last section of [28], some exercices in the course notes [11], and a section in the paper [21].

The main objective of this paper is to quantify the effect on solutions of unknown-but-bounded, deterministic perturbation on problem data. In our framework, the perturbation is not necessarily small, and we seek a solution that is "robust", that is, remains feasible despite the allowable, not necessarily small, perturbation. Our aim is to obtain (approximate) robust solutions via SDP. Links between regularity of solutions and robustness are of course expected. One of our side objectives is to clarify these links to some extent. This paper extends results given in [14] for the least-squares problem.

The approach developed here can be viewed as a special case of stochastic programming, in which the distribution of the perturbation is uniform.

The ideas developed in this paper draw mainly from two sources: control theory, in which we have found the tools for robustness analysis [34, 15, 10], and some recent work on sensitivity analysis of optimization problems due to Shapiro, Bonnans and Cominetti [29, 8].

Shortly after completion of our manuscript, we became aware of the ongoing work of Ben-Tal and Nemirovskii on the same subject. In [6], the authors apply similar ideas to a truss topology design problem, and derive very efficient algorithms for solving the corresponding robustness problem. In [4], the general problem of tractability of obtaining a robust solution is studied, and "tractable counterparts" of a large class of uncertain SDPs is given. The case of robust linear programming, under quite general assumptions on the perturbation bounds, is studied in detail in [5]. Our paper can be seen as a complement of [4], giving ways to cope with (non necessarily) tractable robust SDPs by means of upper bounds.

The paper is divided as follows. Our problem is defined in §2. In §3, we show how to compute upper bounds on our problem via SDP. We give special attention to the so-called full perturbations case, for which our results are non conservative. In §4, we examine sensitivity of the robust solutions in the full perturbations case. We provide conditions which guarantee that the robust solution is unique and a regular function of the data matrices. We then consider several interesting examples in §5, such as robust linear programming, robust norm minimization and error-in-variables SDPs.

### 2. Problem Definition.

**2.1. Robust SDPs.** Let  $\mathbf{F}(x,\Delta)$  be a symmetric matrix-valued function of two variables  $x \in \mathbf{R}^m$ ,  $\Delta \in \mathbf{R}^{p \times q}$ . In the sequel, we consider x as the decision variable, and we think of  $\Delta$  as a perturbation. We assume that  $\Delta$  is unknown-but-bounded.

Precisely, we assume that  $\Delta$  is known to belong to a given linear subspace  $\mathcal{D}$  of  $\mathbf{R}^{p \times q}$ , and in addition,  $\|\Delta\| \leq \rho$ , where  $\rho \geq 0$  is given.

In section 2.2, we will be more precise about the dependence of  $\mathbf{F}$  on  $\Delta$ . We define the robust feasible set by

(2) 
$$\mathcal{X}_{\rho} = \left\{ x \in \mathbf{R}^{m} \mid \text{for every } \Delta \in \mathcal{D}, \|\Delta\| \leq \rho, \\ \mathbf{F}(x, \Delta) \text{ is well-defined and } \mathbf{F}(x, \Delta) \geq 0 \right\}.$$

Now let  $\mathbf{c}(\Delta)$  be a vector-valued rational function of the perturbation  $\Delta$ , such that  $\mathbf{c}(0) = c$ . We consider the following min-max problem

(3) minimize 
$$\max_{\Delta \in \mathcal{D}, \|\Delta\| \le \rho} \mathbf{c}(\Delta)^T x$$
 subject to  $x \in \mathcal{X}_{\rho}$ .

From now on, we assume that the function  $\mathbf{c}(\Delta)$  is independent of  $\Delta$  (in other words, the objective vector c not subject to perturbation). This is done with no loss of generality: introduce a slack variable  $\lambda$  and define

$$ilde{x} = \left[ egin{array}{c} x \ \lambda \end{array} 
ight], \ \ ilde{c} = \left[ egin{array}{c} 0 \ 1 \end{array} 
ight], \ \ ilde{\mathbf{F}}( ilde{x}, \Delta) = \mathbf{diag}(\mathbf{F}(x, \Delta), \lambda - \mathbf{c}(\Delta)^T x).$$

Problem (3) can be formulated as

minimize 
$$\tilde{c}^T \tilde{x}$$
 subject to  $\tilde{x} \in \tilde{\mathcal{X}}_a$ 

where  $\tilde{\mathcal{X}}_{\rho}$  is the robust feasible set corresponding to the function  $\tilde{\mathbf{F}}$ . In the sequel, we thus consider a problem of the form

(4) 
$$\mathcal{P}_{\rho}$$
: minimize  $c^T x$  subject to  $x \in \mathcal{X}_{\rho}$ 

and refer to it as a Robust Semidefinite Problem (RSDP). In general,  $\mathcal{X}_{\rho}$  is not convex, and  $\mathcal{P}_{\rho}$  is not an SDP. Our aim is to find a convex, inner approximation of  $\mathcal{X}_{\rho}$  that is described by a linear matrix inequality constraint. This inner approximation is then used to find an upper bound on the optimal value of  $\mathcal{P}_{\rho}$ , by solving an SDP. In some cases, we can prove our results are non conservative, that is, as in the so-called "full perturbation" case.

We refer to the set  $\mathcal{X}_0$  (resp. problem  $\mathcal{P}_0$ , i.e. (1)) as the nominal feasible set (resp. nominal SDP). We shall assume that the nominal SDP is feasible, that is,  $\mathcal{X}_0 \neq \emptyset$ . Of course, the robust feasible set  $\mathcal{X}_{\rho}$  may become empty for some  $\rho > 0$ ; we return to this question later.

2.2. Linear-fractional representation. In this paper, we restrict our attention to functions **F** that are given by a "linear-fractional representation" (LFR):

(5) 
$$\mathbf{F}(x,\Delta) = F(x) + L\Delta(I - D\Delta)^{-1}R(x) + R(x)^{T}(I - \Delta^{T}D^{T})^{-1}\Delta^{T}L^{T},$$

where F(x) is defined in (1),  $R(\cdot)$  is an affine mapping taking values in  $\mathbf{R}^{q \times n}$ ,  $L \in \mathbf{R}^{n \times p}$  and  $D \in \mathbf{R}^{q \times p}$  are given matrices. Together, the mappings  $F(\cdot)$ ,  $R(\cdot)$ , the matrices

L, D, the subspace  $\mathcal{D}$  and the scalar  $\rho$  consitute our perturbation model for the nominal SDP (1).

The above class of models seem quite specialized. In fact, these models can be used in a wide variety of situations, for example in the case when the (matrix) coefficients  $F_i$  in  $\mathcal{P}_0$  are rational functions of the perturbation. The representation Lemma, given below, and also the examples of section 5, illustrate this point.

A constructive proof of the following result can be found in [35].

LEMMA 2.1. For any rational matrix function  $\mathbf{M}: \mathbf{R}^k \to \mathbf{R}^{n \times c}$ , with no singularities at the origin, there exist nonnegative integers  $r_1, \ldots, r_k$ , and matrices  $M \in \mathbf{R}^{n \times c}$ ,  $L \in \mathbf{R}^{n \times N}$ ,  $R \in \mathbf{R}^{N \times c}$ ,  $D \in \mathbf{R}^{N \times N}$ , with  $N = r_1 + \ldots + r_k$ , such that  $\mathbf{M}$  has the following Linear-Fractional Representation (LFR): For all  $\delta$  where  $\mathbf{M}$  is defined,

(6) 
$$\mathbf{M}(\delta) = M + L\Delta (I - D\Delta)^{-1} R, \text{ where } \Delta = \mathbf{diag}(\delta_1 I_{r_1}, \dots, \delta_k I_{r_k}).$$

Using the LFR lemma, we may devise LFR models for SDPs where a perturbation vector  $\delta \in \mathbf{R}^k$  enters rationally in the coefficient matrices. The resulting set  $\mathcal{D}$  of perturbation matrices  $\Delta$  is then a set of diagonal matrices of repeated elements, as in (6). Componentwise bounds on the vector  $\delta$ , such as  $|\delta|_i \leq \rho$ ,  $i = 1, \ldots, k$ , translate into a norm-bound  $||\Delta|| \leq \rho$  on the corresponding matrix  $\Delta$ .

**2.3.** A special case. We distinguish a special case for which exact (non conservative) results can be obtained via SDP. This is when  $\mathbf{F}(x, \Delta)$  is block-diagonal, each block being independently perturbed, precisely, when

(7) 
$$\mathbf{F}(x,\Delta) = \mathbf{diag}(\mathbf{F}_1(x,\Delta_1), \dots, \mathbf{F}_L(x,\Delta_L)),$$

where each  $\mathbf{F}_i(x, \Delta_i)$  assumes the form (5) for appropriate  $L_i, R_i, D_i$ , with  $\Delta_i \in \mathbf{R}^{p_i \times p_i}$ ,  $i = 1, \ldots, L$ , and  $\mathcal{D}$  consists of block-diagonal matrices of the form

$$\mathcal{D} = \left\{ \Delta = \mathbf{diag}(\Delta_1, \dots, \Delta_L), \; \; \middle| \; \Delta_i \in \mathbf{R}^{p_i imes q_i} 
ight\}.$$

We refer to this situation as the block-full perturbation case. When L=1, we speak of the full perturbation case. As seen later, all results given for L=1 can be generalized to the case L>1.

- 3. Robust solutions for SDPs. Unless otherwise specified, we fix  $\rho > 0$ .
- **3.1. Full perturbations case.** In this section, we consider the full perturbations case, that is,  $\mathcal{D} = \mathbf{R}^{p \times q}$ . We assume  $||D|| < \rho^{-1}$ , which is a necessary and sufficient condition for  $\mathbf{F}(x, \Delta)$  to be well-defined for every  $x \in \mathbf{R}^m$  and  $\Delta \in \mathbf{R}^{p \times q}$ ,  $||\Delta|| \leq \rho$ .

The following lemma is a simple corollary of a classic result on quadratic inequalities referred to as the S-procedure [10]. Its proof is detailed in e.g. [14].

LEMMA 3.1. Let  $F = F^T$ , L, R, D be real matrices of appropriate size. We have  $det(I - D\Delta) \neq 0$  and

(8) 
$$F + L\Delta(I - D\Delta)^{-1}R + R^{T}(I - D\Delta)^{-T}\Delta^{T}L^{T} \ge 0$$

for every  $\Delta$ ,  $\|\Delta\| \leq 1$ , if and only if  $\|D\| < 1$  and there exist a scalar  $\tau$  such that

(9) 
$$\left[ \begin{array}{cc} F - \tau L L^T & R^T - \tau L D^T \\ R - \tau D L^T & \tau (I - D D^T) \end{array} \right] \ge 0.$$

A direct application of the above lemma shows that, in the full perturbations case, the RSDP (4) is an SDP.

THEOREM 3.1. When  $\mathcal{D} = \mathbf{R}^{p \times q}$ , the RSDP (4) and a corresponding solution x can be computed by solving the SDP in variables  $x, \tau$ 

(10) minimize 
$$c^T x$$
 subject to 
$$\begin{bmatrix} F(x) - \tau L L^T & R(x)^T - \tau L D^T \\ R(x) - \tau D L^T & \tau(\rho^{-2} I - D D^T) \end{bmatrix} \ge 0.$$

Special barrier functions adapted to a conic formulation of the problem can be devised, and yield an interior-point algorithm that has the same complexity as the nominal problem, see [22].

We may define the maximum allowable perturbation level, which is the largest numer  $\rho_{\text{max}}$  such that  $\mathcal{X}_{\rho} \neq \emptyset$  for every  $\rho$ ,  $0 \leq \rho \leq \rho_{\text{max}}$  (note  $\rho_{\text{max}} > 0$  since  $\mathcal{X}_{0} \neq \emptyset$ ). Computing  $\rho_{\text{max}}$  is a (quasi-convex) generalized eigenvalue minimization problem [22, 9]:

minimize 
$$\lambda$$
 subject to  $\begin{bmatrix} F(x) - \tau L L^T & R(x)^T - \tau L D^T \\ R(x) - \tau D L^T & \tau (\lambda I - D D^T) \end{bmatrix} \ge 0.$ 

**Remark:** The above exact results are readily generalized to the block-full perturbation case (L > 1), as defined in §2.2.

**3.2. Structured case.** We now turn to the general case ( $\mathcal{D}$  is now an arbitrary linear subspace). In this section, we associate to  $\mathcal{D}$  the following linear subspace

(11) 
$$\mathcal{B} \stackrel{\Delta}{=} \left\{ (S, T, G) \in \mathbf{R}^{p \times p} \times \mathbf{R}^{q \times q} \times \mathbf{R}^{q \times p} \mid S\Delta = \Delta T, \ G\Delta = -\Delta^T G^T \text{ for every } \Delta \in \mathcal{D} \right\}.$$

As shown in [14], a general instance of problem (4) is NP-hard. Therefore, we look for upper bounds on its optimal value. The following lemma is a generalization of lemma 3.1, that traces back to [15]. Its proof is detailed in e.g. [14].

LEMMA 3.2. Let  $F = F^T$ , L, R, D be real matrices of appropriate size. Let  $\mathcal{D}$  a subspace of  $\mathbf{R}^{p \times q}$ , and denote by  $\mathcal{B}$  the set of matrices associated to  $\mathcal{D}$  as in (11). We have  $\det(I - D\Delta) \neq 0$  and

(12) 
$$F + L\Delta(I - D\Delta)^{-1}R + R^{T}(I - D\Delta)^{-T}\Delta^{T}L^{T} > 0$$

for every  $\Delta \in \mathcal{D}$ ,  $\|\Delta\| \le 1$ , if there exist a triple  $(S, T, G) \in \mathcal{B}$  such that S > 0, T > 0, and

(13) 
$$\left[ \begin{array}{ccc} F - LSL^T & R^T - LSD^T + LG \\ R - DSL - GL^T & T - GD^T + DG - DSD^T \end{array} \right] > 0.$$

Using lemma 3.2, we obtain the following result.

THEOREM 3.2. An upper bound on the RSDP (4) and a corresponding solution x can be computed by solving the SDP in variables x, S, T, G

$$\begin{split} &\inf c^T x \text{ subject to } (S,T,G) \in \mathcal{B}, \ S>0, \ T>0, \\ &\left[ \begin{array}{cc} F(x) - LSL^T & R(x)^T - LSD^T + LG \\ R(x) - DSL - GL^T & \rho^{-2}T - GD^T + DG - DSD^T \end{array} \right] > 0. \end{split}$$

Note that when the perturbation is full, the variable G is zero and S, T are of the form  $\tau I_p$ ,  $\tau I_q$  respectively, for some  $\tau \geq 0$ . We then recover the exact results of §3.1.

As before, we may define the maximum allowable perturbation level, which is the largest numer  $\rho_{\text{max}}$  such that  $\mathcal{X}_{\rho} \neq \emptyset$  for every  $\rho$ ,  $0 \leq \rho \leq \rho_{\text{max}}$ . Computing a lower bound on this number is a (quasi-convex) generalized eigenvalue minimization problem:

(14) 
$$\begin{aligned} & \inf \lambda \text{ subject to } (S,T,G) \in \mathcal{B}, \ S>0, \ T>0, \\ & F(x)-LSL^T \quad R(x)^T-LSD^T+LG \\ & R(x)-DSL-GL^T \quad \lambda T-GD^T+DG-DSD^T \end{aligned} \right]>0.$$

4. Unicity and Regularity of Robust Solutions. In this section, we derive unicity and regularity results for the RSDP in the case of full perturbations. As before, we first take L = 1 (one block), that is,  $\mathcal{D} = \mathbf{R}^{p \times q}$ . The results of this section remain valid in the general case L > 1 (several blocks).

We fix  $\rho$ ,  $0 < \rho < \rho_{\text{max}}$ . For simplicity of notation (and without loss of generality) we take  $\rho = 1$  (and thus,  $\rho_{\text{max}} > 1$ ). For well-posedness reasons, we must assume ||D|| < 1. We make the further assumption that D = 0 (in other words,  $\mathbf{F}(x, \Delta)$  is affine in  $\Delta$ ). In §4.5, we show how the case  $D \neq 0$  can be treated.

For full perturbations and D = 0, the RSDP is the SDP

(15) minimize 
$$c^T x$$
 subject to  $\begin{bmatrix} F(x) - \tau L L^T & R(x)^T \\ R(x) & \tau I \end{bmatrix} \ge 0$ .

- **4.1. Hypotheses.** We assume that the SDP (15) (with D = 0) satisfies the following hypotheses:
  - H1: The Slater condition holds, that is, the problem is strictly feasible.
  - **H2:** The problem is *inf-compact*, meaning that any unbounded sequence  $(x_k)$  of feasible points (if any) produces an unbounded sequence of objectives. An equivalent condition is that the Slater condition holds for the dual problem [26, p.317,Thm.30.4].
  - H3: (a) The nullspace of the matrix

$$\lambda R_0 + \sum_{i=1}^m x_i R_i$$

is independent of  $(\lambda, x) \neq (0, 0)$ , and not equal to the whole space.

(c) For every x,

$$\begin{bmatrix} L^T \\ R(x) \end{bmatrix}$$
 is full column-rank.

Hypotheses **H1-2** ensure in particular the existence of optimal points for problem (15) and its dual. Hypothesis **H3(a)-(b)** are difficult to check in general, but can be sometimes easily tested in practical examples, as seen in §5. We note that **H3(a)** implies that  $R(x) \neq 0$  for every x.

Hypothesis **H1** is equivalent to Robinson's condition [25], which can be expressed in terms of

$$\mathcal{F}(x,\tau) = \begin{bmatrix} F(x) - \tau L L^T & R(x)^T \\ R(x) & \tau I \end{bmatrix}.$$

Robinson's condition is stated in [25] as the existence of  $x_0 \in \mathbf{R}^m$ ,  $\tau_0 \in \mathbf{R}$  such that

$$0 \in \operatorname{int} \left( \mathcal{F}(x_0, \tau_0) + d\mathcal{F}(x_0, \tau_0) \mathbf{R}^{m+1} - \mathcal{S}_{n+q}^+ \right),$$

where  $d\mathcal{F}$  is the differential of  $\mathcal{F}$ , and  $\mathcal{S}_{n+q}^+$  is the set of positive semidefinite matrices of order n+q. The equivalence between  $\mathbf{H}\mathbf{1}$  and Robinson's assumption is not true in general. Here, this equivalence stems from the fact that the problem is convex and that the cone  $\mathcal{S}_{n+q}^+$  has non empty interior.

**Remark**: Hypothesis **H1** holds if and only if it holds for the nominal problem (1) (recall our assumption  $\rho_{\text{max}} > 1$ ). Also, hypothesis **H2** implies  $L \neq 0$  (otherwise, we can let  $\tau \to \infty$  without affecting the objective value). If **H2** holds for the nominal problem and  $L \neq 0$ , then **H2** holds for the RSDP (15).

**4.2.** An equivalent nonlinear program. Let  $x_{\text{opt}}$ ,  $\tau_{\text{opt}}$  be optimal for (15). Hypothesis  $\mathbf{H3}(\mathbf{a})$  ensures that any  $\tau$  that is feasible for (15) is nonzero (otherwise, R(x) would be zero for some x). We thus have  $\tau_{\text{opt}} > 0$ .

We introduce some notation: for  $x \in \mathbf{R}^m$ ,  $Z \in \mathbf{R}^{n \times n}$ ,  $\tau > 0$  and  $\mu \in \mathbf{R}$ , define

$$\begin{split} d &= \left[ \begin{array}{c} c \\ 0 \end{array} \right], \ \ y = \left[ \begin{array}{c} x \\ \tau \end{array} \right], \ \ Y = \mathbf{diag}(Z, \mu), \\ G(y) &= F(x) - \tau L L^T - \frac{1}{\tau} R(x)^T R(x), \ \ \mathcal{G}(y) = \mathbf{diag}(G(y), \tau - .99 \tau_{\mathrm{opt}}), \\ \mathcal{L}(y, Y) &= d^T y - \mathbf{Tr} Y G(y). \end{split}$$

Using Schur complements, and using  $\tau_{\text{opt}} > 0$ , we obtain that problem (15) can be rewritten as

(16) minimize 
$$d^T y$$
 subject to  $G(y) \ge 0$ ,

and that  $y_{\text{opt}} = [x_{\text{opt}}^T \tau_{\text{opt}}]^T$  is optimal for (16). Our aim is first to prove that the so-called quadratic growth condition [8] holds at  $y_{\text{opt}}$  for problem (16). Then, we will apply the results of [8] to obtain unicity and regularity theorems.

4.3. Checking the quadratic growth condition. Following [8], we say that the quadratic growth condition (QGC) holds at  $y_{\text{opt}}$  if there exists a scalar  $\alpha > 0$  such that, for every feasible y,

$$d^T y \ge d^T y_{\text{opt}} + \alpha ||y - y_{\text{opt}}||^2 + o(||y - y_{\text{opt}}||^2)$$

Roughly speaking, this condition guarantees that  $y_{\text{opt}}$  is not on a facet on the boundary of the feasible set.

Define the set of dual variables associated with  $y_{\text{opt}}$  by

$$\mathcal{Y}(y_{\mathrm{opt}}) = \left\{ Y = \mathbf{diag}(Z, \mu) \mid Y \geq 0, \ \mathbf{Tr}Y \frac{\partial G}{\partial y_i}(y_{\mathrm{opt}}) = d_i, \ i = 1, \dots, m+1 \right\}.$$

The following result is a direct consequence of a general result due to Bonnans, Cominetti and Shapiro [8]. Roughly speaking, this result states that, if an optimization problem satisfies Robinson's condition and has an optimal point, and if a certain "curvature" condition is satisfied, then the QGC holds at that point.

THEOREM 4.1. With the notations above, if H1-2 hold, and if

(17) 
$$\exists Y \in \mathcal{Y}(y_{\text{opt}}) \text{ such that } \nabla^2_{yy} \mathcal{L}(y_{\text{opt}}, Y) > 0$$

then problem (16) satisfies the QGC.

The following theorem is proven in appendix A.

THEOREM 4.2. If **H1-3** hold, problem (15) satisfies the quadratic growth condition at every optimal point  $y_{\text{opt}}$ . Consequently, there exist a unique solution to the SDP (15).

**Remark:** Note that the QGC is satisfied independent of the objective vector. This means that the boundary of the feasible set is strictly convex (it contains no facets).

**4.4. Regularity results.** In problem (15), the data consists of the matrices L, and  $F_i$ ,  $R_i$ , i = 0, ..., m. We seek to examine the sensitivity of the problem with respect to small variations in  $F_i$ ,  $L_i$  and  $R_i$ .

In this section, we consider matrices L(u),  $R_i(u)$  and  $F_i(u)$ , i = 0, ..., m that are functions of class  $C^1$  of a (small) parameter vector u. Define

$$F(x,u) = F_0(u) + \sum_{i=1}^m x_i F_i(u), \quad R(x,u) = R_0(u) + \sum_{i=1}^m x_i R_i(u).$$

We denote by  $\mathcal{P}(u)$  the corresponding problem (15), where  $F(\cdot)$ ,  $R(\cdot)$  and L are replaced by  $F(\cdot,u)$ ,  $R(\cdot,u)$  and L(u). We assume that  $F(\cdot,0) = F(\cdot)$ ,  $R(\cdot,0) = R(\cdot)$  and L(0) = L, so that  $\mathcal{P}(0)$  is (15).

We first note that, in the vicinity of u = 0, problem  $\mathcal{P}(u)$  satisfies the hypotheses **H1-2** if  $\mathcal{P}(0)$  does. In this case, for every  $\epsilon > 0$  we may define the set  $\mathcal{S}_{\epsilon}(u)$  of  $\epsilon$ -suboptimal points of  $\mathcal{P}(u)$ :

$$S_{\epsilon}(u) = \left\{ x \mid x \text{ is feasible for } \mathcal{P}(u) \text{ and } c^T x \leq v(u) + \epsilon \right\},$$

where v(u) is the optimal value of  $\mathcal{P}(u)$ .

Recall that, if  $\mathcal{P}_0$  satisfies hypotheses **H1-H2**, the optimal value v(u) is continuous, and even directionnally differentiable, at u = 0 [30, Thm. 5.1]. With the QGC in force, and using [29, Thm.4.1], we can give quite complete regularity results for the robust solutions.

THEOREM 4.3. If hypotheses **H1-3** hold for  $\mathcal{P}(0)$ , then for every  $\epsilon = O(u)$ , there exists a  $\gamma > 0$  and a neighborhood V of u = 0 such that for every  $u \in V$  and  $x \in \mathcal{S}_{\epsilon}(u)$ , we have

(18) 
$$||x - x(0)|| \le \gamma ||u||^{1/2}.$$

When **H1-3** hold for  $\mathcal{P}(0)$ , the above theorem states that every (sufficiently) suboptimal solution to  $\mathcal{P}(0)$  is Hölder-stable (with coefficient 1/2). This is true in particular for any optimal solution of  $\mathcal{P}(u)$  (that is, for  $\epsilon = 0$ ). The fact that the theorem remains true for  $\epsilon > 0$  guarantees regularity of numerical solutions to the RSDP. The main consequence is that even if the nominal SDP is ill-conditioned (with respect to variations in the  $F_i$ 's), the RSDP becomes well-conditioned for every  $\rho > 0$ .

Now assume  $\rho \neq 1$ . We seek to examine the behavior of problem (10) (with D=0) when the uncertainty level  $\rho$  for  $0 < \rho < \rho_{\text{max}}$  varies. This is a special case of the problem examined above, with  $u = \rho$ ,  $F(\cdot, u) = F(\cdot)$ ,  $R(\cdot, u) = R(\cdot)$ ,  $L(u) = \rho L$ .

COROLLARY 4.1. For every  $\rho$ ,  $0 < \rho < \rho_{\text{max}}$ , the solution to (10) (with D = 0) is unique and satisfies the regularity results (written with  $u = \rho$ ) of theorem 4.3.

**Remark**: The results of this section are all valid in the block-full perturbation case (L > 1), as defined in §2.2. Of course, the conditions given in **H3** should be understood block-wise.

**4.5.** Case  $D \neq 0$ . When  $D \neq 0$ , we can get back to the case D = 0 as follows. Recall we have ||D|| < 1 in order to ensure that  $\mathbf{F}(x, \Delta)$  is everywhere defined on  $\mathcal{D}$ . With this assumption, we can define for  $x \in \mathbf{R}^m$  and  $\tau > 0$ 

$$\begin{split} \tilde{L} &= L(I - D^T D)^{-1/2}, \\ \tilde{R}(x) &= (I - DD^T)^{-1/2} R(x), \\ \tilde{F}(x) &= F(x) - LD^T (I - DD^T)^{-1} R(x) - R(x)^T (I - DD^T)^{-1} L^T, \\ \tilde{G}(y) &= \tilde{F}(x) - \tau \tilde{L} \tilde{L}^T - \frac{1}{\tau} \tilde{R}(x)^T \tilde{R}(x) \end{split}$$

Using Schur complements, we have for every x and  $\tau > 0$ ,

$$\tilde{G}(y) \ge 0$$
 if and only if  $\begin{bmatrix} F(x) - \tau LL^T & R(x)^T - \tau LD^T \\ R(x) - \tau DL^T & \tau (I - DD^T) \end{bmatrix} \ge 0$ .

Hypothesis **H3** holds for  $\tilde{L}$ ,  $\tilde{R}(\cdot)$  if and only if it holds for L,  $R(\cdot)$ . We can then follow the steps detailed previously.

COROLLARY 4.2. If the SDP (10) (with  $\rho = 1$ ) satisfies **H1-3** and if ||D|| < 1, the results of theorem 4.3 hold.

### 5. Examples.

# 5.1. Unstructured perturbations. Assume

(19) 
$$\mathbf{F}(x,\Delta) = F(x) + \Delta_0 + \Delta_0^T + \sum_{i=1}^m x_i (\Delta_i + \Delta_i^T),$$

where  $\Delta = [\Delta_0 \dots \Delta_m]$ . This case corresponds to (5), with

(20) 
$$L = I, \quad R(x) = \begin{bmatrix} 1 \\ x \end{bmatrix} \otimes I, \quad D = 0, \quad \mathcal{D} = \mathbf{R}^{n \times nm}.$$

Using lemma 3.2, we obtain that problem (4) is equivalent to the SDP

(21) minimize 
$$c^T x$$
 subject to 
$$\begin{bmatrix} F(x) - \tau I & \begin{bmatrix} 1 & x^T \end{bmatrix} \otimes \rho I \\ \begin{bmatrix} 1 & x^T \end{bmatrix}^T \otimes \rho I & \tau I \end{bmatrix} \geq 0.$$

It turns out that we may get rid of the variable  $\tau$  and get back to a convex problem of the same size as that of the unperturbed problem (1). To see this, first note that every feasible variable  $\tau$  in problem (21) is strictly positive. Use Schur complements to rewrite the matrix inequality in (21) as

$$F(x) \ge \left(\tau + \rho^2 \frac{1 + ||x||^2}{\tau}\right) I, \quad \tau > 0.$$

Minimizing (over variable  $\tau$ ) the scalar in the left-hand side of the above inequality shows that the RSDP (1) is equivalent to

(22) minimize 
$$c^T x$$
 subject to  $F(x) \ge 2\rho \sqrt{\|x\|^2 + 1} \cdot I$ .

Formulation (22) is more advantageous than (21), since (22) involves a (convex) matrix inequality constraint of same size as the original problem. As noted before, special barrier functions can be devised for this problem, and yield an interior-point algorithm that has the same complexity as the original problem, see [22].

We note that, with the above choice for L, R, hypothesis **H3** holds, which yields the following result.

THEOREM 5.1. The optimal value of the RSDP (21) can be computed by solving the convex problem (22). If (22) satisfies hypotheses **H1-2**, then for every  $\rho > 0$ , the solution is unique and satisfies the regularity conditions of theorem 4.3.

**Remark**: A sufficient condition for hypotheses **H1-2** to hold for (22) is that they hold for the nominal problem. A more restrictive sufficient condition is that the nominal feasible set  $\mathcal{X}_0$  is non empty and bounded, and  $\rho < \rho_{\text{max}}$ .

**5.2. Robust center of a linear matrix inequality.** In this section, for  $\rho > 0$ , we consider the SDP (22) and corresponding feasible (convex) set  $\mathcal{X}_{\rho}$ . We assume that  $\mathcal{X}_0$  is non empty and bounded, and that  $\mathcal{P}_0$  is strictly feasible. Then, for every  $\rho$ ,  $0 < \rho < \rho_{\text{max}}$ ,  $\mathcal{X}_{\rho}$  is non empty and bounded, and we can define a (unique) solution  $x(\rho)$  to the strictly convex problem (22).

In view of corollary 4.1,  $x(\rho)$  is a continuous function of  $\rho$  in  $]0 \rho_{\text{max}}[$ . Since  $(\mathcal{X}_{\rho})$  is a decreasing family of bounded sets, we may define

$$(23) x^* = \lim_{\rho \to \rho_{\text{max}}} x(\rho)$$

Note that  $x^*$  is independent on the objective vector c.

Thus, to the matrix inequality  $F(x) \geq 0$ , we may associate the robust center, defined by (23). The robust center has the property to be the most tolerant (with respect to unstructured perturbation) among the feasible points.

An example is depicted in Fig. 1. The nominal feasible set  $\mathcal{X}_0$  is described by a linear matrix inequality  $F(x) \geq 0$ , where F is a  $5 \times 5$  matrix. For various values of  $\rho$ , we seek to minimize  $x_2$ . The dashed lines correspond to the optimal objectives. As  $\rho$  increases, we observe the robust feasible sets shrink. A crucial property of these robust sets is that they do not posess any straight faces, as observed in the figure. For  $\rho = \rho_{\text{max}} \simeq 5$ , the robust feasible set is a singleton (in this example,  $x^* = 0$ ). When  $\rho = 0$ , the optimal solution is not unique, and not continuous with respect to changes in the coefficient matrices  $F_i$ , i = 0, 1, 2 (although the optimal value is continuous). Since the sets  $\mathcal{X}_{\rho}$  become strictly convex as soon as  $\rho > 0$ , the resulting robust solutions are continuous.




Fig. 1. Nominal and robust solutions of an SDP, with a  $5 \times 5$  matrix F(x). Here  $\rho_{\text{max}} = 5$ .

**5.3. Robust linear programs.** An interesting special case arises with Linear Programming (LP). Consider the LP

minimize  $c^T x$  subject to  $a_i^T x \ge b_i$ , i = 1, ..., L.

Assume that the  $a_i$ 's and  $b_i$ 's are subject to unstructured perturbations. The perturbed value of  $[a_i^T b_i]^T$  is  $[a_i^T b_i]^T + \delta_i$ , where  $\|\delta_i\|_2 \leq \rho$ , i = 1, ..., L. We seek a robust solution to our problem, wich is a special case of the block-full perturbation case referred to in §2.2, with **F** given by (7), and

$$\mathbf{F}_{i}(x, \Delta_{i}) = a_{i}^{T} x - b_{i} + 2[x^{T} - 1]\Delta_{i}, \quad i = 1, \dots, L,$$

where  $\Delta_i = \delta_i/2$ , and  $\mathcal{D}$  the set of diagonal,  $L \times L$  matrices. The robust LP is

(24) minimize 
$$c^T x$$
 subject to  $a_i^T x - \rho \sqrt{\|x\|_2^2 + 1} \ge b_i, \ i = 1, \dots, L.$ 

The above program is readily written as an SDP, by introducing slack variables. In fact, the robust LP is second-order cone program (SOCP), for which efficient special-purpose interior-point methods are available [22, 18, 21].

We note that hypothesis **H3** holds block-wise. This yields the following result.

THEOREM 5.2. The optimal value of the robust LP can be computed by solving the convex problem (24). If the latter satisfies hypotheses **H1-2**, then for every  $\rho$ ,  $0 < \rho < \rho_{\text{max}}$ , the solution is unique and satisfies the regularity conditions of theorem 4.3.

In [5], robust linear programming is studied in detail. For a wide class of perturbation models, where the data of every linear constraint vary in an ellipsoid, explicit robust solutions are constructed using convex (second-order cone) programs (SOCP). Reference [21] also provides examples of robust linear programs solved via SOCP.

5.4. Robust eigenvalue minimization. Consider the case when the nominal problem consists of minimizing the largest eigenvalue of a matrix-valued function F(x):

(25) minimize 
$$\lambda_{\max}(F(x))$$
.

When  $F(\cdot)$  is subject to unstructured perturbations (as defined in §5.1), the robust version of the problem is

minimize 
$$\lambda + 2\rho \sqrt{\|x\|^2 + 1}$$
 subject to  $\lambda I \ge F(x)$ ,

or equivalently

(26) minimize 
$$\lambda_{\max}(F(x)) + 2\rho\sqrt{\|x\|^2 + 1}$$
.

Let  $\rho > 0$ . When written in an SDP form, the above problem satisfies the hypotheses **H1-3**. From theorem 4.3 we obtain that the solution is unique. If we consider that the data of the above problem consists of the matrices  $F_i$ , i = 0, ..., m, then we know that the corresponding solution is Hölder stable (with coefficient 1/2). Since the problem is unconstrained, we can use a result of Shapiro [29, Thm.3.1], by which we conclude that the solution is actually Lipschitz stable (inequality (18) holds with the exponent 1/2 replaced by 1). Finally, using the results from Attouch [3], we can show that computing the solution for  $\rho \to 0$  amounts to select the minimum norm solution among the solutions of the nominal problem.

THEOREM 5.3. The optimal value of the min-max problem (25) can be computed by solving the convex problem (26). For every  $\rho > 0$ , the solution is unique and is Lipschitz stable with respect to perturbations in  $F_i$ , i = 0, ..., m. When  $\rho \to 0$ , the solution converges to the minimum norm solution of the nominal problem (25).

Remark: In this case, the RSDP is a regularized version of the nominal SDP, which belongs to the class of Tikhonov regularizations [32]. The regularization parameter is  $2\rho$ , and is chosen according to some a priori information on uncertainty associated with the nominal problem's data. Taking  $\rho$  close to zero can be used as a selection procedure, for choosing a particular (minimum norm, regular) solution among the (non necessarily unique and/or regular) solutions of the nominal problem.

Problem (26) is particularly suitable to the recent so-called  $\mathcal{U}$ -Newton algorithms for solving problem (25). These methods, described in [19, 23], require that the Hessian of the "smooth part" (the so-called  $\mathcal{U}$ -Hessian) of the objective of (25) to be positive-definite. For general mappings  $F(\cdot)$ , this property is not guaranteed. However, when looking at the robust problem (26), we see that the modified  $\mathcal{U}$ -Hessian is guaranteed to be positive-definite for every x and  $\rho > 0$ . This indicates that the RSDP approach may be used to devise robust algorithms for solving SDPs.

**5.5.** Robust second-order cone programs. A second-order cone program (SOCP) is a problem of the form

(27) minimize 
$$c^T x$$
  
subject to  $||C_i x + d_i|| \le e_i^T x + f_i, \quad i = 1, \dots, L,$ 

where  $C_i \in \mathbf{R}^{n_i \times m}$ ,  $d_i \in \mathbf{R}^{n_i}$ ,  $e_i \in \mathbf{R}^m$ ,  $f_i \in \mathbf{R}$ , i = 1, ..., L. SOCP can be formulated as SDPs, but special-purpose, more efficient algorithms can be devised for them, see [22, 4, 21].

Assuming that  $C_i$ ,  $d_i$ ,  $e_i$ ,  $f_i$  are subject to linear—or even rational—uncertainty, we may formulate corresponding RSDP, as an SDP. This SDP can be written as an SOCP if the uncertainty is unstructured, and affects each constraint independently.

The subject of robust SOCP is explored in [4] in detail. Explicit SDPs that yield robust counterpart to SOCPs non conservatively are given, for a wide class of uncertainty structures. In some cases, albeit not all, the robust counterpart is itself an SOCP. In [14, 12], the special case of least squares problems with uncertainty in the data is studied at length.

5.6. Robust maximum norm minimization. Several engineering problems take the form

(28) minimize 
$$||H(x)||$$

where

$$H(x) = H_0 + \sum_{i=1}^{m} x_i H_i,$$

and  $H_i$ ,  $i=1,\ldots,m$  are given  $p\times q$  matrices. A well-known instance of this problem is the linear least-squares problem, with H(x)=Ax-b. Another example is a minimal norm extension problem for an Hankel operator studied in [16], in which  $H_0$  is a given (arbitrary)  $n\times n$  Hankel matrix, and  $H_i$ ,  $i=1,\ldots,m$ , is the  $n\times n$  Hankel matrix associated with the polynomial  $1/z^i$ . In practice, the matrices  $H_i$ ,  $i=0,\ldots,m$  are subject to perturbation, which motivates a study of the robust version of problem (28). Note that the least-squares case is extensively studied in [14].

Consider the full perturbation case, which occurs when each  $H_i$  is perturbed independently in a linear manner. Precisely, consider the matrix-valued function

$$\mathbf{H}(x,\Delta) = H_0 + \Delta_0 + \sum_{i=1}^{m} x_i (H_i + \Delta_i),$$

where  $\Delta = [\Delta_0 \dots \Delta_m]$ . For a given  $\rho > 0$ , we address the min-max problem

(29) 
$$\min_{x} \max_{\|\Delta\| \le \rho} \|\mathbf{H}(x, \Delta)\|$$

This problem is a RSDP for which we can get exact results using SDP. Indeed, for every  $x \in \mathbf{R}^m$  and  $\lambda \geq 0$ , the property

$$\max_{\|\Delta\| \le \rho} \|\mathbf{H}(x, \Delta)\| \le \lambda$$

is equivalent to  $\mathbf{F}(x,\lambda,\Delta) \geq 0$  for every  $\Delta$ ,  $\|\Delta\| \leq \rho$ , where

$$\mathbf{F}(x,\lambda,\Delta) = F(x,\lambda) + L\Delta R(x) + R(x)^T \Delta^T L^T,$$

where

$$F(x,\lambda) = \left[ \begin{array}{cc} \lambda I & H(x) \\ H(x)^T & \lambda I \end{array} \right], \quad L = \left[ \begin{array}{c} I \\ 0 \end{array} \right], \quad R(x) = \left[ \begin{array}{cc} 0 & \left[ \begin{array}{c} 1 \\ x \end{array} \right] \otimes I \end{array} \right].$$

We thus write problem (29) as (4), where the perturbation set  $\mathcal{D}$  is  $\mathbf{R}^{p \times q}$ .

Applying theorem 3.2, we obtain that the RSDP above is equivalent to the SDP (15) (with D = 0). As in §5.1, we may get rid of the variable  $\tau$ , and obtain the equivalent formulation

(30) minimize 
$$||H(x)|| + \rho \sqrt{||x||^2 + 1}$$
.

This RSDP satisfies hypotheses **H1-3**, so we conclude that the results of theorem 4.3 hold. As in robust eigenvalue minimization, we can get improved results using [29, 3, Thm.3.1].

THEOREM 5.4. The optimal value of the min-max problem (29) can be computed by solving the convex problem (30). For every  $\rho > 0$ , the solution is unique and is Lipschitz stable with respect to perturbations in  $H_i$ , i = 0, ..., m. When  $\rho \to 0$ , the solution converges to the minimum norm solution of the nominal problem (28). **Remark**: As for the RSDP arising in robust eigenvalue minimisation, the robust minimum norm minimization probem is a regularized version of the nominal problem, which belongs to the class of Tikhonov regularizations.

We now consider the general case when each matrix  $H_i$  in (28) is perturbed in a structured manner. To be specific, we concentrate on the minimal norm extension problem mentioned above.

In practice, the matrix  $H_0$  is obtained from measurement, and is thus subject to error. We may assume that this matrix is constructed from a  $n \times 1$  vector  $h_0(\delta) = h_0 + \delta$ , where  $\delta$  is unknown-but-bounded. The perturbed matrix  $H_0$  is of the form

$$H_0(\Delta) = H_0 + L\Delta R,$$

where L, R are given matrices (the exact form of which we do not detail), and

$$\Delta \in \mathcal{D} = \{ \mathbf{diag}(\delta_1 I_1, \dots, \delta_n I_n) \mid \delta_i \in \mathbf{R}, \ i = 1, \dots, n \}.$$

(In the above, each  $\delta_i$  corresponds to the uncertainty associated with the *i*-th antidiagonal of  $H_0$ .) We address the min-max problem

(31) minimize 
$$\max_{\Delta \in \mathcal{D}, \|\Delta\| \le \rho} \|H(x) + L\Delta R\|$$

where  $\rho \geq 0$  is given.

This problem is amenable to the robustness analysis technique. Defining

$$\mathcal{S} \stackrel{\Delta}{=} \left\{ \operatorname{diag}(S_1, \dots, S_n) \mid S_i \in \mathbf{R}^{i \times i}, \quad i = 1, \dots, n \right\},\,$$

we obtain the following result.

Theorem 5.5. An upper bound on the objective value of the min-max problem (31) can be computed by solving the SDP in variables x, S, G:

inf 
$$\lambda$$
 subject to  $S = S^T$ ,  $G = -G^T \in \mathcal{S}$ , 
$$\begin{bmatrix} \lambda I - LSL^T & H(x) & LG \\ H(x)^T & \lambda I & \rho R \\ G^T L^T & \rho R^T & S \end{bmatrix} > 0.$$

**5.7. Polynomial interpolation.** This example is one taken from [14], that can be formulated as RSDP with rational dependence. For given integers  $n \geq 1$ , k, we seek a polynomial of degree n-1  $p(t) = x_1 + \ldots + x_n t^{n-1}$  that interpolates given points  $(a_i, b_i), i = 1, \ldots, k$ , that is

$$p(a_i) = b_i, i = 1, ..., k.$$

If we assume that  $(a_i, b_i)$  are known exactly, we obtain a linear equation in the unknown x, with a Vandermonde structure:

$$\begin{bmatrix} 1 & a_1 & \dots & a_1^{n-1} \\ \vdots & \vdots & & \vdots \\ 1 & a_k & \dots & a_k^{n-1} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} b_1 \\ \vdots \\ b_n \end{bmatrix},$$

which can be solved via standard least-squares techniques.

Now assume that the interpolation points are not known exactly. For instance, we may assume that the  $b_i$ 's are known, while the  $a_i$ 's are parameter-dependent:

$$a_i(\delta) = a_i + \delta_i, \quad i = 1, \dots, k,$$

where the  $\delta_i$ 's are unknown-but-bounded:  $|\delta_i| \leq \rho$ , i = 1, ..., k, where  $\rho \geq 0$  is given. We seek a robust interpolant, that is, solution x that minimizes

(32) 
$$\max_{\|\delta\|_{\infty} \le \rho} \|\mathbf{A}(\delta)x - b\|,$$

where

$$\mathbf{A}(\delta) = \begin{bmatrix} 1 & a_1(\delta) & \dots & a_1(\delta)^{n-1} \\ \vdots & \vdots & & \vdots \\ 1 & a_k(\delta) & \dots & a_k(\delta)^{n-1} \end{bmatrix}.$$

The above problem is a RSDP. Indeed, it can be shown that

$$\begin{bmatrix} \mathbf{A}(\delta) & b \end{bmatrix} = \begin{bmatrix} \mathbf{A}(0) & b \end{bmatrix} + L\Delta(I - D\Delta)^{-1}R,$$

where

$$L = \bigoplus_{i=1}^{k} \begin{bmatrix} 1 & a_i & \dots & a_i^{n-2} \end{bmatrix}, R = \begin{bmatrix} R_1 \\ \vdots \\ R_k \end{bmatrix}, D = \bigoplus_{i=1}^{k} D_i, \Delta = \bigoplus_{i=1}^{k} \delta_i I_{n-1}.$$

and, for each  $i, i = 1, \ldots, k$ ,

$$R_{i} = \begin{bmatrix} 0 & 1 & a_{i} & \dots & a_{i}^{n-2} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & a_{i} \\ 0 & \dots & \dots & 0 & 1 \end{bmatrix} \in \mathbf{R}^{(n-1)\times n},$$

$$D_{i} = \begin{bmatrix} 0 & 1 & a_{i} & \dots & a_{i}^{n-3} \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ \vdots & & \ddots & \ddots & a_{i} \\ 0 & \dots & \dots & 0 & 1 \end{bmatrix} \in \mathbf{R}^{(n-1)\times (n-1)}.$$

(Note that  $det(I - D\Delta) \neq 0$ , since D is strictly upper triangular.) With the above notation, if we define  $\mathbf{F}(x,\Delta)$  as in (5), then problem (32) can be formulated as the RSDP (4).

With the approach described in this paper, one can compute an upper bound for the minimizing value of (32), and a corresponding suboptimal x. We do not know if the problem can be solved exactly in polynomial-time, e.g. using SDP. We conjecture (as the reviewers of this paper did ) that the answer is no. To motivate this claim, note that the solution to the problem of computing (32) for arbitrary affine functions  $\mathbf{A}$  is already NP-hard [14].

5.8. Error-in-Variables RSDPs. In many SDPs that arise in engineering, the variable x represents physical parameters that can be implemented with finite absolute precision only. A typical example is integer programming, where integer solutions to (linear) programs are sought. These problems (which are equivalent to integer programming) are NP-hard. We now show that we may find upper bounds on these problems using robustness analysis.

Consider for instance the problem of finding a solution x to the feasibility SDP

(33) find 
$$x$$
 integer such that  $F(x) \ge 0$ .

Now, consider the robust SDP

maximize  $\lambda$  subject to

(34) 
$$\lambda I \leq F_0 + \sum_{i=1}^m (x_i + \Delta x_i) F_i \text{ for every } \Delta x, \|\Delta x\|_{\infty} \leq 1/2.$$

Assume there exist a feasible pair  $(x_{\text{feas}}, \lambda)$  to the above problem, with  $\lambda \geq 0$ . By construction,  $x_{\text{feas}}$  satisfies  $F(x_{\text{feas}}) \geq 0$ . Furthermore, any vector x chosen such that  $||x - x_{\text{feas}}||_{\infty} \leq 1/2$  is guaranteed to satisfy  $F(x) \geq 0$ . This is true in particular for  $x_{\text{int}}$ , the integer closest to  $x_{\text{feas}}$ . Thus, if we know a positive lower bound  $\lambda$ , and corresponding feasible point for problem (34), then we can compute an integer solution to our original problem.

Finding a lower bound for (34) and associated feasible point can be done as follows. For  $i, 1 \leq i \leq m$ , define  $F_i = 2L_iR_i$ , where  $L_i, R_i^T \in \mathbf{R}^{n \times r_i}$ ,  $r_i = \mathbf{Rank}F_i$ . Let

$$L = \begin{bmatrix} L_1 & \dots & L_m \end{bmatrix}, \quad R = \begin{bmatrix} R_1 \\ \vdots \\ R_m \end{bmatrix} \text{ and } \mathcal{D} = \left\{ \Delta = \bigoplus_{i=1}^m \Delta x_i I_{r_i}, \quad \Delta x_i \in \mathbf{R} \right\}.$$

Problem (34) can be formulated as

(35) maximize 
$$\lambda$$
 subject to  $\lambda I \leq F(x) + L\Delta R + R^T \Delta^T L^T$  for every  $\Delta \in \mathcal{D}$ ,  $\|\Delta\| \leq 1/2$ .

The above is a special instance of the structured problem examined in §3.2. Define

$$\mathcal{S} \stackrel{\Delta}{=} \left\{ \mathbf{diag}(S_1, \dots, S_m) \mid S_i \in \mathbf{R}^{r_i \times r_i}, \ i = 1, \dots, n \right\},\,$$

Theorem 5.6. A sufficent condition for an integer solution to the feasibility SDP (33) is that the constraints

$$\lambda \geq 0, \quad S = S^T \in \mathcal{S}, \quad G = -G^T \in \mathcal{S}, \quad \left[ \begin{array}{cc} F(x) - \lambda I - LSL^T & \frac{1}{2}R^T + LG \\ \frac{1}{2}R - GL^T & S \end{array} \right] > 0$$

are feasible. If  $x_{\text{feas}}$  is feasible for the above constraints, then any integer vector closest to  $x_{\text{feas}}$  (in the maximum norm sense) is feasible for (33).

6. Conclusions. In this paper, we considered semidefinite programs subject to uncertainty. Assuming the latter is unknown-but-bounded, we have provided sufficient conditions that guarantee "robust" solutions to exist, via SDPs. Under some conditions (detailed in §4), the robust solution is unique, and not surprisingly, stable. The method can then be used to regularize possibly ill-conditioned problems. For some perturbation structures (as for unstructured perturbations), the conditions are also necessary. That is, there is no conservatism induced by the method.

The paper raises several open questions.

In our description, we have considered the problem of robustifying the primal SDP, thereby obtaining upper bounds on a SDP subject to uncertainty. The dual point of view should be very interesting. One might be interested in applying the approach to the dual problem instead. Does this lead lower bounds on the perturbed problem? Also, in some cases, the RSDP approach leads to a unique (and stable) primal solution. May we obtain a unique solution to the dual problem, by robustifying the latter? (This would lead to analycity of the primal solution, see [30]).

As seen in §5.2 the notion of robust center has certainly connections with the well-known analytic center; is the latter related to some robustness characterization?

It seems that the RSDP method could be useful for deriving fast and robust (stable) algorithms for solving SDPs (see §5.4), especially in connection with maximum eigenvalue minimization.

Finally, as said in section 2.2 (lemma 2.1), an SDP with coefficient matrices depending rationnaly on a perturbation vector can always be represented by an LFR model. Now, this LFR model is not unique. However, the results given here (for example, theorem 3.2) hinge on a particular linear-fractional representation for a perturbed SDP. Hence the question: are our results independent of the chosen representation? We partially answer this difficult question in appendix B.

Acknowledgements. This paper has benefited from many stimulating discussions with several colleagues, in particular, Aharon Ben-Tal, Stephen Boyd, Arkadii Nemirovskii, Michael Overton and Lieven Vandenberghe. Last but not least, the authors would like to thank the Editor and Reviewers for their very helpful comments for revising the paper.

#### REFERENCES

- [1] F. ALIZADEH, Interior point methods in semidefinite programming with applications to combinatorial optimization, SIAM J. Optimization, 5 (1995), pp. 13-51.
- [2] F. ALIZADEH, J.-P. A. HAEBERLY, AND M. L. OVERTON, Primal-dual interior-point methods for semidefinite programming: Convergence rates, stability and numerical results, Tech. Rep. 721, NYU Computer Science Dept, May 1996.
- [3] H. Attouch, Viscosity solutions of optimization problems, Tech. Rep. 07, Dept des Sciences Mathématiques, Université Montpellier 2, France, 1994.
- [4] A. BEN-TAL AND A. NEMIROVSKI, Robust convex programming, Tech. Rep. 1/95, Optimization Laboratory, Faculty of Industrial Engineering and Management, Technion, Israel Institute of Technology, Technion City, Haifa 32000, Israel, 1995.

- [5] ——, Robust solutions to uncertain linear programs, Tech. Rep. 6/95, Optimization Laboratory, Faculty of Industrial Engineering and Management, Technion, Israel Institute of Technology, Technion City, Haifa 32000, Israel, 1995.
- [6] ——, Robust truss topology design, SIAM J. on Optimization, (1997). To appear.
- [7] B. Bereanu, Some numerical methods in stochastic programming under risk and uncertainty, Academic Press, 1980, ch. 11, pp. 169-205.
- [8] J. F. Bonnans, R. Cominetti, and A. Shapiro, Sensivity analysis of optimization problems under second order regular constraints, Tech. Rep. 2989, INRIA, 1996.
- [9] S. Boyd and L. El Ghaoui, Method of centers for minimizing generalized eigenvalues, Linear Algebra and Applications, special issue on Numerical Linear Algebra Methods in Control, Signals and Systems, 188 (1993).
- [10] S. BOYD, L. EL GHAOUI, E. FERON, AND V. BALAKRISHNAN, Linear Matrix Inequalities in System and Control Theory, Studies in Applied Mathematics, SIAM, Philadelphia, PA, June 1994.
- [11] S. P. BOYD AND L. VANDENBERGHE, Introduction to convex optimization with engineering applications, lecture notes for ee392x, Stanford University, 1996. Available via anonymous ftp into isl.stanford.edu, under /pub/boyd.
- [12] S. CHANDRASEKARAN, G. H. GOLUB, M. Gu, and A. H. Sayed, A new linear least-squares type model for parameter estimation in the presence of data uncertainties. Preliminary draft., 1996.
- [13] M. Dempster, Stochastic programming, Academic Press, 1980.
- [14] L. El Ghaoui and H. Lebret, Robust solutions to least-squares problems with uncertain data matrices, SIAM J. Matrix Anal. Appl., (1997).
- [15] M. K. H. Fan, A. L. Tits, and J. C. Doyle, Robustness in the presence of mixed parametric uncertainty and unmodeled dynamics, IEEE Trans. Aut. Control, 36 (1991), pp. 25-38.
- [16] J. W. HELTON AND H. J. WOERDEMAN, Symmetric Hankel operators: Minimal norm extensions and eigenstructures, Linear Algebra and Appl., 185 (1993), pp. 1-19.
- [17] M. KOJIMA, M. SHIDA, AND S. SHINDOH, Global and local convergence of predictor-correctorinterior-point agorithm for semidefinite programming, Tech. Rep. B-308, Dept. of Information Sciences, Tokyo Institute of Technology, 2-12-1 Oh-Okayama, Meguru-ku, Tokyo 152, Japan, 1995.
- [18] H. Lebret, Synthèse de diagrammes de réseaux d'antennes par optimisation convexe, PhD thesis, Université de Rennes I, Nov. 1994.
- [19] C. LEMARÉCHAL, F. OUSTRY, AND C. SAGASTIZÁBAL, The *U-Lagrangian of a convex function*, Transactions of the American Mathematical Society, (1996). Submitted.
- [20] A. S. LEWIS AND M. L. OVERTON, Eigenvalue optimization, Acta Numerica, (1996), pp. 149–190
- [21] M. S. Lobo, L. Vandenberghe, S. Boyd, and H. Lebret, Second-order cone programming: interior-point methods and engineering applications, Linear Algebra Appl., (1997). Submitted.
- [22] Y. Nesterov and A. Nemirovsky, Interior point polynomial methods in convex programming: Theory and applications, SIAM, Philadelphia, PA, 1994.
- [23] F. Oustry, The *U-Lagrangian of the maximum eigenvalue function*, SIAM J. Optimization, (1996). Submitted.
- [24] F. A. Potra and R. Sheng, Homogeneous interior-point agorithms for semi-definite programming, Tech. Rep. 82, Department of Mathematics, The University of Iowa, Iowa City, Iowa 52242, USA, November 1995.
- [25] S. Robinson, Stability theorems for systems of inequalities, part II: differentiable nonlinear systems, SIAM J. Numer. Analysis, 13 (1976), pp. 497-513.
- [26] R. T. ROCKAFELLAR, Convex Analysis, Princeton Univ. Press, Princeton, second ed., 1970.
- [27] R. Rubinstein and A. Shapiro, Discrete event systems, Wiley & Sons, 1993.
- [28] A. Shapiro, Perturbation theory of nonlinear programs when the set of optimal solutions is not a singleton, Applied Mathematics and Optimization, 18 (1988), pp. 215-229.
- [29] ——, Perturbation analysis of optimization problems in Banach spaces, Numer. Funct. Anal. and Optimiz., 13 (1992), pp. 97-116.

- [30] —, First and second order analysis of nonlinear semidefinite programs, Mathematical Programming (Series B), (1996). to appear.
- [31] A. Shapiro and M. K. H. Fan, On eigenvalue optimization, SIAM J. Optimization, 5 (1995), pp. 552-568.
- [32] A. TIKHONOV AND V. ARSENIN, Solutions of Ill-Posed Problems, Wiley, New York, 1977.
- [33] L. VANDENBERGHE AND S. BOYD, Semidefinite programming, SIAM Review, 38 (1996), pp. 49–95.
- [34] V. A. Yakubovich, The solution of certain matrix inequalities in automatic control theory, Soviet Math. Dokl., 3 (1962), pp. 620-623. In Russian, 1961.
- [35] K. Zhou, J. Doyle, and K. Glover, Robust and Optimal Control, Prentice Hall, New Jersey, 1995.

**A. Proof of theorem 4.2.** We take the notations of §4. Let  $Y = \operatorname{diag}(Z, \mu)$  be dual variables associated to  $(x_{\text{opt}}, \tau_{\text{opt}})$  that are optimal (their existence is guaranteed by **H1-2**). Then,  $Y \in \mathcal{Y}(y_{\text{opt}})$ . Let us show that condition (17) holds for this choice of Y.

Since the problem satisfies **H1-2**, the complementarity conditions hold, therefore the (optimal) dual variable  $\mu$  associated with the constraint  $\tau = \tau_{\rm opt}$  is zero. Consequently the variable Z is nonzero (recall  $c \neq 0$ ). Using

$$\operatorname{Tr} Y \frac{\partial G}{\partial y_{m+1}}(y_{\text{opt}}) = d_{m+1} = 0,$$

we obtain

$$\tau_{\text{opt}}^2 \mathbf{Tr} L L^T Z = \mathbf{Tr} R(x_{\text{opt}})^T R(x_{\text{opt}}) Z.$$

From  $\tau_{\text{opt}} \neq 0$  (implied by  $\mathbf{H3(a)}$ ), and using hypothesis  $\mathbf{H3(b)}$ , we can show that

$$\mathbf{Tr}LL^TZ = 0$$
 and  $\mathbf{Tr}R(x_{\text{opt}})^TR(x_{\text{opt}})Z = 0$ 

is impossible for  $Z \geq 0$ ,  $Z \neq 0$ . This yields  $\mathbf{Tr} R(x_{\text{opt}})^T R(x_{\text{opt}}) Z > 0$ . Now let  $\xi \in \mathbf{R}^m$  and  $\lambda \in \mathbf{R}$ , and define

(36) 
$$\Phi(\xi,\lambda) = d^{2}\mathcal{G}(x_{\text{opt}}, \tau_{\text{opt}})[(\xi,\lambda), (\xi,\lambda)],$$
$$\phi(\xi,\lambda) = \begin{bmatrix} \xi \\ \lambda \end{bmatrix}^{T} \nabla_{yy}^{2} \mathcal{L}(y_{\text{opt}}, Y) \begin{bmatrix} \xi \\ \lambda \end{bmatrix} = -\text{Tr} Z \Phi(\xi,\lambda).$$

We have, for every  $i, j, 1 \le i, j \le m$ ,

$$\frac{\partial G}{\partial x_i} = F_i - \frac{1}{\tau} (R(x)^T R_i + R_i^T R(x)), \quad \frac{\partial G}{\partial \tau} = -LL^T + \frac{1}{\tau^2} R(x)^T R(x),$$

$$\frac{\partial^2 G}{\partial x_i \partial x_j} = -\frac{1}{\tau} (R_j^T R_i + R_i^T R_j), \quad \frac{\partial^2 G}{\partial x_i \partial \tau} = \frac{1}{\tau^2} (R(x)^T R_i + R_i^T R(x)),$$

$$\frac{\partial^2 G}{\partial \tau^2} = -\frac{2}{\tau^3} R(x)^T R(x).$$

By summation, we have

$$\begin{split} -\Phi(\xi,\lambda) &= \frac{2}{\tau_{\rm opt}} (R(\xi) - R(0))^T (R(\xi) - R(0)) + 2 \frac{\lambda^2}{\tau_{\rm opt}^3} R(x_{\rm opt})^T R(x_{\rm opt}) \\ &- \frac{\lambda}{\tau_{\rm opt}^2} \left( R(x_{\rm opt})^T (R(\xi) - R(0)) + (R(\xi) - R(0))^T R(x_{\rm opt}) \right) \\ &= \frac{\lambda^2}{\tau_{\rm opt}^3} R(x_{\rm opt})^T R(x_{\rm opt}) + \frac{1}{\tau_{\rm opt}} \mathcal{R}^T \mathcal{R} + \frac{1}{\tau_{\rm opt}} (R(\xi) - R(0))^T (R(\xi) - R(0)), \end{split}$$

where  $\mathcal{R} = R(\xi) - R(0) - \frac{\lambda}{\tau_{\text{opt}}} R(x_{\text{opt}})$ . We obtain finally

$$\begin{split} \phi(\xi,\lambda) &= \frac{1}{\tau_{\text{opt}}} \mathbf{Tr} Z \mathcal{R}^T \mathcal{R} + \frac{\lambda^2}{\tau_{\text{opt}}^3} \mathbf{Tr} Z R(x_{\text{opt}})^T R(x_{\text{opt}}) \\ &+ \frac{1}{\tau_{\text{opt}}} \mathbf{Tr} Z (R(\xi) - R(0))^T (R(\xi) - R(0)). \end{split}$$

If  $\phi(\xi, \lambda) = 0$ , then  $\lambda = 0$  (from  $\operatorname{Tr} R(x_{\operatorname{opt}})^T R(x_{\operatorname{opt}}) Z > 0$ ), and thus  $\operatorname{Tr} Z \mathcal{R}^T \mathcal{R} = 0$  with  $\mathcal{R} = R(\xi) - R(0)$ . Since  $Z \geq 0$ , this means that every column of  $Z^{1/2}$  belongs to the nullspace of  $R(\xi) - R(0)$ . Assume now  $\xi \neq 0$ . By hypothesis  $\operatorname{H3}(\mathbf{a})$ , we obtain that every column of  $Z^{1/2}$  also belongs to the nullspace of  $R(x_{\operatorname{opt}})$ , which contradicts  $\operatorname{Tr} R(x_{\operatorname{opt}})^T R(x_{\operatorname{opt}}) Z > 0$ . We conclude that  $\nabla^2_{yy} \mathcal{L}$  is positive-definite at  $(y_{\operatorname{opt}}, Y)$ . Thus, problem (15) satisfies the QCG.

**B.** Invariance with respect to LFR model. In this section, we show that the sufficient conditions obtained in this paper are, in some sense, independent of the LFR model used to describe the perturbation structure.

Consider a function **F** taking values in the set of symmetric matrices having an LFR such as (5). This function can be written in a more symmetric form

(37) 
$$\mathbf{F}(\Delta) = F + \tilde{L}\tilde{\Delta}(I - D\tilde{\Delta})^{-1}\tilde{L}^{T}$$

where we have dropped the dependence on x for convenience, and

$$\tilde{L} = \begin{bmatrix} L & R^T \end{bmatrix}, \quad \tilde{D} = \begin{bmatrix} 0 & D^T \\ D & 0 \end{bmatrix}, \quad \tilde{\Delta} = \begin{bmatrix} 0 & \Delta \\ \Delta^T & 0 \end{bmatrix}.$$

It is easy to check that, if an invertible matrix Z satisfies the relation  $Z\tilde{\Delta}Z^T = \tilde{\Delta}$  for every  $\Delta \in \mathcal{D}$ , then

$$\mathbf{F}(\Delta) = F + (\tilde{L}Z)\tilde{\Delta}(I - (Z^T\tilde{D}Z)\tilde{\Delta})^{-1}(\tilde{L}Z)^T.$$

In other words, the "scaled" triple  $(F, (\tilde{L}Z), (Z^T\tilde{D}Z))$  can be used to represent  $\mathbf{F}$  instead of  $F, \tilde{L}, \tilde{D}$  in (37). By spanning valid scaling matrices Z, we span a subset of all LFR models that describe  $\mathbf{F}$ .

A valid scaling matrix Z can be constructed as follows. Let  $(S,T,G)\in\mathcal{B},$  and define

$$Z = \begin{bmatrix} T^{-1/2} & 0 \\ 0 & S^{1/2} \end{bmatrix} \begin{bmatrix} I & G \\ 0 & I \end{bmatrix}.$$

It turns out that such a Z satisfies the relation  $Z\tilde{\Delta}Z^T = \tilde{\Delta}$  for every  $\Delta \in \mathcal{D}$ .

Using the above facts, we can show that if condition (13) is true for the original LFR model F, L, R, D with appropriate S, T, G, then it is also true for the scaled LFR obtained using any scaling matrix Z such as above, for appropriate matrices  $\tilde{S}$ ,  $\tilde{G}$ ,  $\tilde{T}$ . That is, the condition is independent of the scaling Z.

In this sense, the conditions we obtained are independent of the LFR used to represent the perturbation structure.