

ORIGINAL ARTICLE

**STRESS AND ITS RELATIONSHIP WITH BODY MASS INDEX
AMONG BIOMEDICAL SCIENCE STUDENTS IN
KUALA LUMPUR, MALAYSIA**

*Nur Zakiah Mohd Saat, Ismarulyusda Ishak, Syarif Husin Lubis,
See Huong Wen, Siti Nor Lily Suriyanee Mohd, Norul Sahira Zakaria,
Tan Xin Ee, Marlina Othman, Tang Hui Shein, Zahirah Zulkifli, Tan Liy Si,
Law Ngik Ling, Surayya Razali, Umi Romaizatul Amiera Zainudin,
Yeoh Thong Wei, Luqman Osman, Kholilul Syafiq Baharuddin*

Department of Biomedical Science, Faculty of Allied Health Science,
Universiti Kebangsaan Malaysia, Kuala Lumpur Campus,
Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia

Abstract

Objective: The objective of this study was to assess the relationship between stress score and body mass index (BMI) of students of Science in Kuala Lumpur. **Methods:** A cross-sectional study was carried out among 126 students. Data about stress score were collected using a questionnaire, the Student-Life Stress Inventory (SLSI) Data about BMI were calculated from height and weight of the respondents. **Results:** A total of 117 (92.85%) out of 126 students had responded. The mean difference of stress score among first, second and third year students were 87.6 ± 17.0 , 83.4 ± 16.9 and 86.7 ± 15.4 respectively. The mean difference of stress scores between the Bumiputra and non-Bumiputra students were 87.9 ± 16.5 and 83.2 ± 15.9 . The mean of body mass Index(BMI) between first, second and third year students were 21.0 ± 3.5 , 20.3 ± 2.7 and 21.8 ± 4.3 respectively. **Conclusions:** This study showed that the stress level is higher in first year, female and bumiputra student. There was no significant relationship between stress score and BMI. *ASEAN Journal of Psychiatry Vol. 11 (2): July-Dec 2010: XX XX.*

Keywords: Stress, Body Mass Index, Student-Life Stress Inventory

Introduction

Stress is one of a factor which may influence behaviors and health especially when an individual faces challenges that surpass his or her coping skills [1]. Another research by

Torres et al.(2007) indicated that stress may change overall food intake such as, under- or overeating according to stress severity. Stress-induced eating may be one of the factors that lead to obesity[2].

Obesity, as indicated by a high body mass index (BMI, kg/m²), is a well-established

risk factor for major causes of morbidity and mortality such as cardiovascular disease, hypertension, stroke, type 2 diabetes, and certain forms of cancer [3-6]. The cause of obesity is not well understood [7, 8], but it is related to an interaction between biological and environmental factors [9]. For the past few decades, obesity has grown to epidemic proportions, and its prevalence is still increasing rapidly [3, 4]. Globally, already more than 1 billion adults are overweight and at least 300 million of them are clinically obese [4].

Chronic psychosocial stress may lead to obesity through physiological effects, such as excess cortisol secretion, or could contribute to the development of obesity through an association with poorer behavioral risk factors, increased caloric intake and sedentary lifestyle [10]. Dietary habits are major aspects of people's lifestyles that influence health, morbidity, and mortality for a range of conditions. Carbohydrate consumption has been hypothesized to relieve depressive moods [11], and this has been considered as part of the causal link for developing obesity [12]. However, the association has also been seen in the opposite direction, with stress resulting in poorer food choices [13]. Stress may increase food consumption in certain individuals but also may change their food choices from lower fat to higher fat foods [14].

Although there are many studies that have been done on obesity and stress, however very few studies relate stress and obesity especially among university students. Therefore, the objective of this study is to

determine the relationship between stress score and BMI changes across from first to third year of study.

Methods

A cross-sectional study was conducted among 126 Science students, in Kuala Lumpur, Malaysia. The participants were selected by using stratified random sampling followed by systematic random sampling method. The questionnaire consists of two parts: (i) sociodemographic questions and (ii) questions designed to gain information of stress score. Sociodemographic information includes years of study, gender and race. The Student-Life Stress Inventory (SLSI) [15] comprises of 33 questions that were already modified. SLSI is a questionnaire with a Likert type response format (1 = never, 2 = seldom, 3 = occasionally, 4 = often, 5 = most of the time). Examples of questions: As a student, I fail to achieve the self-set goals. As a student, I am not socially accepted. As an individual, I always worry about all the things and people around me. The SLSI score was calculated by adding up all the marks according to participants' rating.

Body mass index (BMI) were measured using height and weight of the respondents. Data for both stress score and BMI were collected one week before mid-semester examination. Questionnaires were administered in the same week to minimize the effect of varying stress levels that may occur during the mid-semester examination.

Pilot study was carried out on 30 students. With data from the pilot study, Cronbach's alpha value for modified SLSI was calculated, and the value was 0.96. This shows that the modified SLSI is reliable

since acceptable level of reliability is 0.80 or higher. [16]. Data were analyzed using Statistical Package for Social Sciences (SPSS) version 16.

Results

A total of 117 (92.85%) out of 126 students had responded in which 95 (81.2%) of them were female students. All years of study were approximately equally represented: 37 students (31.7%) from first year, 39 students (33.3%) from second year and 41 students (35.0%) from third year. According to race, 67 (57.3%) students were Bumiputras while 50 (42.7%) students were non-Bumiputras.

In this study we found that the stress score among study year, gender and race were showed in Table 2. First year students

showed highest mean stress score (87.6 ± 17.0) among the three study years, followed by third year (86.7 ± 15.4) and second year (83.4 ± 16.9) students. There was no significant mean difference of stress score among study years ($p=0.509$, $p>0.05$). Male students showed lower stress score (83.4 ± 17.9) compared to female students (86.5 ± 16.0).

However there was no significant difference of stress score between male and female students ($p=0.366$, $p>0.05$). Besides, bumiputra students (87.9 ± 16.5) showed higher stress score compared to non-Bumiputra students (83.2 ± 15.9). There was no significant mean difference of stress score between Bumiputra and non-Bumiputra ($p=0.123$, $p>0.05$).

Table 1: Description of demographic factors

Demographic factors	Variables	Frequency (n)	Percentage (%)
Study year	First year	37	31.7
	Second year	39	33.3
	Third year	41	35.0
Gender	Male	22	18.8
	Female	95	81.2
Race	Bumiputra	67	57.3
	Non-Bumiputra	50	42.7

Table 2: Comparison of stress score among study year, gender and race

Variables	Mean ± Standard Deviation	test statistic	p value
First year	87.6 ± 17.0	0.680	0.509
Second year	83.4 ± 16.9		
Third year	86.7 ± 15.4		
Male	83.4 ± 17.9	915.500	0.366
Female	86.474± 16.044		
Bumiputra	87.910 ± 16.520	1.555	0.123
Non-bumiputra	83.180± 15.940		

Table 3: Comparison of BMI among study year, gender and race

Variables	Mean ± Standard Deviation	test statistic	p value
First year	21.0 ± 3.5	1.268	0.530
Second year	20.3 ± 2.7		
Third year	21.8 ± 4.3		
Male	21.9 ± 5.3	983.500	0.668
Female	20.8 ± 3.1		
Bumiputra	21.5 ± 3.7	1321.000	0.051
Non-bumiputra	20.3 ± 3.3		

Table 3 shows a comparison of BMI among study year, gender and race. Third year students showed highest BMI (21.8 ± 4.3) among the three study years, followed by first year (21.0 ± 3.5) and second year (20.3 ± 2.7) students. However, there was no significant difference of BMI among study years ($p=0.530$, $p>0.05$). Meanwhile, male students (21.9 ± 5.3) showed slightly higher BMI compared to female students (20.8 ± 3.1) but there was no significant difference of BMI between them ($p=0.668$, $p>0.05$). Moreover, Bumiputra (21.5 ± 3.7) showed slightly higher BMI compared to non-Bumiputra (20.3 ± 3.3). There was no significant difference of BMI between Bumiputra and non-Bumiputra ($p=0.051$, $p>0.05$).

In order to find out the relationship between overall stress score and BMI, Spearman's correlation was employed. The result showed that there was no relationship between the stress score and the BMI ($r=0.056$, $p>0.05$). Furthermore, according to year of study the results from Spearman's correlation shows that there was no significant relationship between stress score and BMI of first ($r=0.125$, $p>0.05$), second ($r=0.125$, $p>0.05$) and third year students ($r=-0.185$, $p>0.05$) respectively. Thus, there was also no significant relationship between the stress score and BMI according to first year, second year and third year students respectively. Furthermore, according to gender there was no association between stress score and BMI for both male ($r=0.26$, $p>0.05$) and female student ($r=0.002$, $p>0.05$). Moreover according to race, there was no significant relationship between stress score and BMI for Bumiputra ($r=-0.058$, $p>0.05$) and non-Bumiputra respectively ($r=0.198$, $p>0.05$).

Discussion

Previous studies in Mexico using SLSI found that academic stress among college students varies across the years academic. Specifically, freshman and sophomores had higher academic stress level than juniors and seniors, among pharmacy students [17]. On the other hand, our study had shown that first year students had the highest stress score. This suggests that different courses and places will have different stress score among the study years. Besides, their mean stress scores were higher compared to our finding. This study had shown that there was no significant difference of BMI among study years. This is because narrow age ranges does not affect the BMI. This is supported by other studies which found that width of the age ranges was often affected by the sample size. Hence, combining subjects of different ages is needed to have a significant difference of BMI [18]

The comparison of mean stress score between races was not significant in our study. This result is consistent with previous studies claiming that other factors such as race did not contribute to stress level in students [19]. Besides that, other studies find that there was significant difference between races with Malays having the highest mean BMI, followed by the Indians and Chinese [20]. The differences may be due to our research was too centralized at one place. However, the Bumiputra with majority being Malays, still showed higher mean BMI compared to non-Bumiputra which were Chinese and Indians.

Our finding had also shown that there was no significant difference for mean stress score between male and female students and this is different compared to other studies.

They had found that females had significantly higher stress scores compared to males [17]. However, females' stress score was still slightly higher than the males. This could be due to females being more emotional and sensitive than males. There was no significant difference in the BMI between male and female students and this should not be comparable to other studies because the subjects university student. Other study found that the mean BMI of adult women was significantly higher than the mean BMI of adult men [20]. Possible reason to explain this is that our sample was too localized at one place

Our study had shown that there was no significant relationship between the overall stress score and BMI. Besides, the relationship of stress score and BMI of first year, second year, third year, male, female, Bumiputra and non-Bumiputra students respectively had also shown no significant relationships. This is different from other studies [1] where there was a weak relationship between work stress and BMI. Besides, the previous study had measured the stress score of people in working sector while this study measured stress among students in education sector. This is because the stress at workplace differed with academic stress. In the workplace, stress is influenced by skill, team work and work load especially in the professional areas such as medical doctors, engineer, teachers and nurse [21-23]. Mean while for stress among students maybe due to concern with academic performance, relationship with colleagues, friends and family [24]

Conclusion

In conclusion, this study showed that the level of stress score varies according to

years of study, gender and race among students of Science Kuala Lumpur Student in first year, female and bumiputra showed a higher level of stress. Mean while the BMI is higher among third year, male and bumiputras. However in this study, there was no significant relationship between stress and BMI. Other factors may also contribute to the stress score such as time allocation, social support and individual characteristic.

Acknowledgements

The authors wished to thank the Department of Biomedical Science, Faculty and Health Sciences, Universiti Kebangsaan Malaysia for the assistance and support for this study.

References

1. Kouvonen A, Kivimäki M, Cox SJ, Cox T, Vahtera J. Relationship Between Work Stress and Body Mass Index Among 45,810 Female and Male Employees. *Psychosom Med.* 2005;67:577-583.
2. Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Int J Appl Basic Nutr Sci.* 2007;23(1112):887-894.
3. Visscher TLS, Seidell JC. The public health impact of obesity. *Ann Rev Public Health.* 2001;22:355-75.
4. Kannel WB. Effect of weight on cardiovascular disease. *Nutr.* 1997; 13:157-8.
5. Jousilahti P, Tuomilehto J, Vartiainen E, Pekkanen J, Puska P. Body weight, cardiovascular risk factors, and coronary mortality: 15-year follow-up of

middle-aged men and women in Eastern Finland. *Circulation*. 1996;93:1372–9.

6. Ogden J. *Health Psychology. A Textbook*, 3rd ed. Maidenhead: Open University Press; 2004.

7. Landsbergis PA, Schnall PL, Deitz DK, Warren K, Pickering TG, Schwartz JE. Job strain and health behaviors: results of a prospective study. *Am J Health Promot*. 1998;12:237-45.

8. Goodrick GK, Poston WSC II, Foreyt JP. Methods for voluntary weight loss and control: update 1996. *Nutr* 1996;12:672– 6.

9. Schulz AJ, House JS, Israel BA et al. Relational pathways between socioeconomic position and cardiovascular risk in a multiethnic urban sample: complexities and their implications for improving health in economically disadvantaged populations. *J Epidemiol Community Health*. 2008;62:638–646.

10. Wurtman RJ, Wurtman JJ. Carbohydrates and depression. *Sci Am*. 1989;260:68-75.

11. Arnow B, Kenardy J, Agras WS. Binge eating among the obese: a descriptive study. *J Behav Med*. 1992;15:155-170.

12. Liu C, Xie B, Chou CP, Koprowski C, Zhou D, Palmer P et al. Perceived stress, depression and food consumption frequency in the college students of China Seven Cities. *Physiol Behav*. 2007;92:748-754.

13. Zellner DA, Loaiza S, Gonzalez Z, Pita J, Morales J, Pecora D et al. Food

selection changes under stress. *Physiol Behav*. 2006;87:789-793.

14. Oliver G, Wardle J, Gibson L. Stress and Food Choice: A Laboratory Study. *Psychosom Med* 2000; 62:853-865.

15. Gadzella, B. M. 1991. *Student-Life stress inventory*. Commerce, TX: Gadzella.

16. Dawson B, Trapp RG. *Basic & Clinical Biostatistics*. 4rd ed. United States of America, McDraw-Hill; 2004.

17. Gupchup GH, Borrego ME, Konduri N. The impact of student life stress on health related quality of life among doctor of pharmacy students. *Coll Student J*. 2004;38(2): 292-302.

18. Stevens J. Studies of the impact of age on optimal body weight. *J. Nutr. Biochem* 1998; 9(9):501-510.

19. Muhamad SBY, Ahmad FAR, Mohd JY. Prevalence and sources of stress among Universiti Sains Malaysia Medical Students. *Malays J. Med Sci*. 2009;17(1):30-37.

20. Azmi MY, Junidah R, Siti MA, Safiah MY, Fatimah S, Norimah AK et al. Body Mass Index (BMI) of Adults: Findings of the Malaysian Adult Nutrition Survey (MANS). *Malaysian J. Nutr.* 1998;15(2):97 – 119.

21. McVicar, A., *Integrative Literature Reviews and Meta-Analyses Workplace Stress in Nursing: A Literature Review*. *J Adv Nurs* 2003; 44(6):633-642.

22. Kyriacou C. Teacher stress: directions for future research. *Edu Rev* 2001; 53,(1); 27–35.

23. McManus IC, Keeling A, Paice, E. Stress, burnout and doctors' attitudes to work are determined by personality and learning style: A twelve year longitudinal study of UK medical graduates. *BMC Med* 2004; 2-29.

24. Westerman GH, Grandy TG, Ocanto RA, Erskine CG. Perceived sources of stress in the dental school environment. *J Dent Educ* 1993;57:225-31.

Corresponding Author: *Nur Zakiah Mohd Saat, Department of Biomedical Science, Faculty of Allied Health Science, Universiti Kebangsaan Malaysia, Jalan Raja Muda Abdul Aziz, 50300 Kuala Lumpur, Malaysia.*

Email: nur14504@gmail.com

Received: 27 July 2010

Accepted: 13 August 2010