LAYOUT OF GRAPHS WITH BOUNDED TREE-WIDTH *

VIDA DUJMOVIĆ^{‡§†}, PAT MORIN^{§†}, AND DAVID R. WOOD^{§¶†}

Abstract. A queue layout of a graph consists of a total order of the vertices, and a partition of the edges into queues, such that no two edges in the same queue are nested. The minimum number of queues in a queue layout of a graph is its *queue-number*. A three-dimensional (straight-line grid) drawing of a graph represents the vertices by points in \mathbb{Z}^3 and the edges by non-crossing line-segments. This paper contributes three main results:

(1) It is proved that the minimum volume of a certain type of three-dimensional drawing of a graph G is closely related to the queue-number of G. In particular, if G is an n-vertex member of a proper minor-closed family of graphs (such as a planar graph), then G has a $\mathcal{O}(1) \times \mathcal{O}(1) \times \mathcal{O}(n)$ drawing if and only if G has $\mathcal{O}(1)$ queue-number.

(2) It is proved that queue-number is bounded by tree-width, thus resolving an open problem due to Ganley and Heath (2001), and disproving a conjecture of Pemmaraju (1992). This result provides renewed hope for the positive resolution of a number of open problems in the theory of queue lavouts.

(3) It is proved that graphs of bounded tree-width have three-dimensional drawings with $\mathcal{O}(n)$ volume. This is the most general family of graphs known to admit three-dimensional drawings with $\mathcal{O}(n)$ volume.

The proofs depend upon our results regarding track layouts and tree-partitions of graphs, which may be of independent interest.

Key words. queue layout, queue-number, three-dimensional graph drawing, tree-partition, treepartition-width, tree-width, k-tree, track layout, track-number, acyclic colouring, acyclic chromatic number.

AMS subject classifications. 05C62 (graph representations)

1. Introduction. A queue layout of a graph consists of a total order of the vertices, and a partition of the edges into queues, such that no two edges in the same queue are nested. The dual concept of a *stack layout*, introduced by Ollmann [73] and commonly called a *book embedding*, is defined similarly, except that no two edges in the same stack may cross. The minimum number of queues (respectively, stacks) in a queue layout (stack layout) of a graph is its queue-number (stack-number). Queue layouts have been extensively studied [41, 53, 54, 58, 76, 80, 86, 88] with applications in parallel process scheduling, fault-tolerant processing, matrix computations, and sorting networks (see [76] for a survey). Queue layouts of directed acyclic graphs [9, 56, 57, 76] and posets [55, 76] have also been investigated. Our motivation for studying queue layouts is a connection with three-dimensional graph drawing.

Graph drawing is concerned with the automatic generation of aesthetically pleasing geometric representations of graphs. Graph drawing in the plane is well-studied (see [24, 64]). Motivated by experimental evidence suggesting that displaying a graph in three dimensions is better than in two [90, 91], and applications including information visualisation [90], VLSI circuit design [66], and software engineering [92], there is a growing body of research in three-dimensional graph drawing. In this paper

[‡]School of Computer Science, McGill University, Montréal, Canada. E-mail: vida@cs.mcgill.ca [§]School of Computer Science, Carleton University, Ottawa, Canada.

E-mail: {morin,davidw}@scs.carleton.ca

[¶]Department of Applied Mathematics, Charles University, Prague, Czech Republic. Research supported by COMBSTRU.

[†]Research supported by NSERC.

^{*}Submitted October 14, 2002. Revised June 16, 2004. Results in this paper were presented at the GD '02 [34], FST TCS '02 [93], and WG '03 [36] conferences. 1

we study three-dimensional straight-line grid drawings, or three-dimensional drawings for short. In this model, vertices are positioned at grid-points in \mathbb{Z}^3 , and edges are drawn as straight line-segments with no crossings [17, 21, 25, 27, 28, 42, 53, 78, 75]. We focus on the problem of producing three-dimensional drawings with small volume. Three-dimensional drawings with the vertices in \mathbb{R}^3 have also been studied [39, 47, 19, 16, 18, 61, 22, 63, 60, 62, 69, 74]. Aesthetic criteria besides volume that have been considered include symmetry [60, 61, 62, 63], aspect ratio [19, 47], angular resolution [47, 19], edge-separation [19, 47], and convexity [18, 19, 39, 87].

The first main result of this paper reduces the question of whether a graph has a three-dimensional drawing with small volume to a question regarding queue layouts (Theorem 2.10). In particular, we prove that every *n*-vertex graph from a proper minor-closed graph family \mathcal{G} has a $\mathcal{O}(1) \times \mathcal{O}(1) \times \mathcal{O}(n)$ drawing if and only if \mathcal{G} has a $\mathcal{O}(1)$ queue-number, and this result holds true when replacing $\mathcal{O}(1)$ by $\mathcal{O}(\text{polylog}n)$. Consider the family of planar graphs, which are minor-closed. (In the conference version of their paper) Felsner *et al.* [42] asked whether every planar graph has a three-dimensional drawing with $\mathcal{O}(n)$ volume? Heath *et al.* [58, 54] asked whether every planar graph has $\mathcal{O}(1)$ queue-number? By our result, these two open problems are almost equivalent in the following sense. If every planar graph has $\mathcal{O}(1)$ queue-number, then every planar graph has a $\mathcal{O}(1) \times \mathcal{O}(1) \times \mathcal{O}(1) \times \mathcal{O}(n)$ drawing, then every planar graph has $\mathcal{O}(1)$ queue-number. It is possible, however, that planar graphs have unbounded queue-number, yet have say $\mathcal{O}(n^{1/3}) \times \mathcal{O}(n^{1/3})$ drawings.

Our other main results regard three-dimensional drawings and queue layouts of graphs with bounded tree-width. Tree-width, first defined by Halin [50], although largely unnoticed until independently rediscovered by Robertson and Seymour [81] and Arnborg and Proskurowski [7], is a measure of the similarity of a graph to a tree (see §2.1 for the definition). Tree-width (or its special case, path-width) has been previously used in the context of graph drawing by Dujmović *et al.* [33], Hliněný [59], and Peng [77], for example.

The second main result is that the queue-number of a graph is bounded by its tree-width (Corollary 2.8). This solves an open problem due to Ganley and Heath [45]. who proved that stack-number is bounded by tree-width, and asked whether a similar relationship holds for queue-number. This result has significant implications for the above open problem (does every planar graph have $\mathcal{O}(1)$ queue-number), and the more general question (since planar graphs have stack-number at most four [94]) of whether queue-number is bounded by stack-number. Heath et al. [58, 54] originally conjectured that both of these questions have an affirmative answer. More recently however, Pemmaraju [76] conjectured that the 'stellated K_3 ', a planar 3-tree, has $\Theta(\log n)$ queue-number, and provided evidence to support this conjecture (also see [45]). This suggested that the answer to both of the above questions was negative. In particular, Pemmaraju [76] and Heath [private communication, 2002] conjectured that planar graphs have $\mathcal{O}(\log n)$ queue-number. However, our result provides a queuelayout of any 3-tree, and thus the stellated K_3 , with $\mathcal{O}(1)$ queues. Hence our result disproves the first conjecture of Pemmaraju [76] mentioned above, and renews hope in an affirmative answer to the above open problems.

The third main result is that every graph of bounded tree-width has a threedimensional drawing with $\mathcal{O}(n)$ volume. The family of graphs of bounded tree-width includes most of the graphs previously known to admit three-dimensional drawings with $\mathcal{O}(n)$ volume (for example, outerplanar graphs), and also includes many graph families for which the previous best volume bound was $\mathcal{O}(n^2)$ (for example, seriesparallel graphs). Many graphs arising in applications of graph drawing do have small tree-width. Outerplanar and series-parallel graphs are the obvious examples. Another example arises in software engineering applications. Thorup [89] proved that the control-flow graphs of go-to free programs in many programming languages have treewidth bounded by a small constant; in particular, 3 for Pascal and 6 for C. Other families of graphs having bounded tree-width (for constant k) include: almost trees with parameter k, graphs with a feedback vertex set of size k, band-width k graphs, cut-width k graphs, planar graphs of radius k, and k-outerplanar graphs. If the size of a maximum clique is a constant k then chordal, interval and circular arc graphs also have bounded tree-width. Thus, by our result, all of these graphs have threedimensional drawings with $\mathcal{O}(n)$ volume, and $\mathcal{O}(1)$ queue-number.

To prove our results for graphs of bounded tree-width, we employ a related structure called a tree-partition, introduced independently by Seese [85] and Halin [51]. A *tree-partition* of a graph is a partition of its vertices into 'bags' such that contracting each bag to a single vertex gives a forest (after deleting loops and replacing parallel edges by a single edge). In a result of independent interest, we prove that every k-tree has a tree-partition such that each bag induces a connected (k - 1)-tree, amongst other properties. The second tool that we use is a *track layout*, which consists of a vertex-colouring and a total order of each colour class, such that between any two colour classes no two edges cross.

The remainder of the paper is organised as follows. In §2 we introduce the required background material, and state our results regarding three-dimensional drawings and queue layouts, and compare these with results in the literature. In §3 we establish a number of results concerning track layouts. That three-dimensional drawings and queue-layouts are closely related stems from the fact that three-dimensional drawings and queue layouts are both closely related to track layouts, as proved in §4 and §5, respectively. In §6 we prove the above-mentioned theorem for tree-partitions of k-trees, which is used in §7 to construct track layouts of graphs with bounded tree-width. We conclude in §8 with a number of open problems.

2. Background and Results. Throughout this paper all graphs G are undirected, simple, and finite with vertex set V(G) and edge set E(G). The number of vertices and the maximum degree of G are respectively denoted by n = |V(G)| and $\Delta(G)$. The subgraph induced by a set of vertices $A \subseteq V(G)$ is denoted by G[A]. For all disjoint subsets $A, B \subseteq V(G)$, the bipartite subgraph of G with vertex set $A \cup B$ and edge set $\{vw \in E(G) : v \in A, w \in B\}$ is denoted by G[A, B].

A graph H is a *minor* of a graph G if H is isomorphic to a graph obtained from a subgraph of G by contracting edges. A family of graphs closed under taking minors is *proper* if it is not the class of all graphs.

A graph parameter is a function α that assigns to every graph G a non-negative integer $\alpha(G)$. Let \mathcal{G} be a family of graphs. By $\alpha(\mathcal{G})$ we denote the function $f : \mathbb{N} \to \mathbb{N}$, where f(n) is the maximum of $\alpha(G)$, taken over all *n*-vertex graphs $G \in \mathcal{G}$. We say \mathcal{G} has bounded α if $\alpha(\mathcal{G}) \in \mathcal{O}(1)$. A graph parameter α is bounded by a graph parameter β (for some graph family \mathcal{G}), if there exists a function g such that $\alpha(G) \leq g(\beta(G))$ for every graph G (in \mathcal{G}).

2.1. Tree-Width. Let G be a graph and let T be a tree. An element of V(T) is called a *node*. Let $\{T_x \subseteq V(G) : x \in V(T)\}$ be a set of subsets of V(G) indexed by the nodes of T. Each T_x is called a *bag*. The pair $(T, \{T_x : x \in V(T)\})$ is a

tree decomposition of G if:

- 1. $\bigcup_{x \in V(T)} T_x = V(G)$ (that is, every vertex of G is in at least one bag),
- 2. \forall edge vw of G, \exists node x of T such that $v \in T_x$ and $w \in T_x$, and

3. \forall nodes x, y, z of T, if y is on the path from x to z in T, then $T_x \cap T_z \subseteq T_y$. The width of a tree-decomposition is one less than the maximum cardinality of a bag. A path-decomposition is a tree-decomposition where the tree T is a path $T = (x_1, x_2, \ldots, x_m)$, which is simply identified by the sequence of bags T_1, T_2, \ldots, T_m where each $T_i = T_{x_i}$. The path-width (respectively, tree-width) of a graph G, denoted by pw(G) (tw(G)), is the minimum width of a path- (tree-) decomposition of G. Graphs with tree-width at most one are precisely the forests. Graphs with tree-width at most two are called series-parallel¹, and are characterised as those graphs with no K_4 minor (see [10]).

A k-tree for some $k \in \mathbb{N}$ is defined recursively as follows. The empty graph is a k-tree, and the graph obtained from a k-tree by adding a new vertex adjacent to each vertex of a clique with at most k vertices is also a k-tree. This definition of a k-tree is by Reed [79]. The following more restrictive definition of a k-tree, which we call 'strict', was introduced by Arnborg and Proskurowski [7], and is more often used in the literature. A k-clique is a strict k-tree, and the graph obtained from a strict k-tree by adding a new vertex adjacent to each vertex of a k-clique is also a strict k-tree. Obviously the strict k-trees are a proper sub-class of the k-trees. A subgraph of a k-tree is called a partial k-tree, and a subgraph of a strict k-tree is called a partial strict k-tree. The following result is well known (see for example [10, 79]). A chord of a cycle C is an edge not in C whose end-vertices are both in C. A graph is chordal if every cycle on at least four vertices has a chord.

LEMMA 2.1. Let G be a graph. The following are equivalent:

- 1. G has tree-width $\mathsf{tw}(G) \leq k$,
- 2. G is a partial k-tree,
- 3. G is a partial strict k-tree,
- 4. G is a subgraph of a chordal graph that has no clique on k+2 vertices.

Proof. Scheffler [83] proved that (1) and (3) are equivalent. That (1) and (4) are equivalent is due to Robertson and Seymour [81]. That (2) and (4) are equivalent is the characterisation of chordal graphs in terms of 'perfect elimination' vertex-orderings due to Fulkerson and Gross [44].

2.2. Tree-Partitions. As in the definition of a tree-decomposition, let G be graph and let $\{T_x \subseteq V(G) : x \in V(T)\}$ be a set of subsets of V(G) (called *bags*) indexed by the nodes of a tree T. The pair $(T, \{T_x : x \in V(T)\})$ is a *tree-partition* of G if

- 1. \forall distinct nodes x and y of T, $T_x \cap T_y = \emptyset$, and
- 2. \forall edge vw of G, either
- (i) \exists node x of T with $v \in T_x$ and $w \in T_x$ (vw is called an *intra-bag* edge), or
- (ii) \exists edge xy of T with $v \in T_x$ and $w \in T_y$ (vw is called an inter-bag edge).

The main property of tree-partitions that has been studied in the literature is the maximum cardinality of a bag, called the *width* of the tree-partition [11, 51, 85, 31, 32]. The minimum width over all tree-partitions of a graph G is the *tree-partition-width*²

¹'Series-parallel digraphs' are often defined in terms of certain 'series' and 'parallel' composition operations. The underlying undirected graph of such a digraph has tree-width at most two (see [10]).

²Tree-partition-width has also been called *strong tree-width* [85, 11].

of G, denoted by tpw(G). A graph with bounded degree has bounded tree-partitionwidth if and only if it has bounded tree-width [32]. In particular, for every graph G, Ding and Oporowski [31] proved that $tpw(G) \leq 24 tw(G)\Delta(G)$, and Seese [85] proved that $tw(G) \leq 2 tpw(G) - 1$.

Theorem 6.1 provides a tree-partition of a k-tree G with additional features besides small width. First, the subgraph induced by each bag is a connected (k-1)-tree. This allows us to perform induction on k. Second, in each non-root bag T_x the set of vertices in the parent bag of x with a neighbour in T_x form a clique. This feature is crucial in the intended application (Theorem 7.3). Finally the tree-partition has width at most max $\{1, k(\Delta(G) - 1)\}$, which represents a constant-factor improvement over the above result by Ding and Oporowski [31] in the case of k-trees.

2.3. Track Layouts. Let G be a graph. A colouring of G is a partition $\{V_i : i \in I\}$ of V(G), where I is a set of colours, such that for every edge vw of G, if $v \in V_i$ and $w \in V_j$ then $i \neq j$. Each set V_i is called a colour class. A colouring of G with c colours is a c-colouring, and we say that G is c-colourable. The chromatic number of G, denoted by $\chi(G)$, is the minimum c such that G is c-colourable.

If $\langle i$ is a total order of a colour class V_i , then we call the pair $(V_i, \langle i)$ a *track*. If $\{V_i : i \in I\}$ is a colouring of G, and $(V_i, \langle i)$ is a track, for each colour $i \in I$, then we say $\{(V_i, \langle i) : i \in I\}$ is a *track assignment* of G indexed by I. Note that at times it will be convenient to also refer to a colour $i \in I$ and the colour class V_i as a *track*. The precise meaning will always be clear from the context. A *t*-track assignment is a track assignment with t tracks.

As illustrated in Fig. 2.1, an X-crossing in a track assignment consists of two edges vw and xy such that $v <_i x$ and $y <_j w$, for distinct tracks V_i and V_j . A t-track assignment with no X-crossing is called a t-track layout. The track-number of a graph G, denoted by $\operatorname{tn}(G)$, is the minimum t such that G has a t-track layout.



FIG. 2.1. An example of an X-crossing in a track assignment.

Let $\{(V_i, <_i) : i \in I\}$ be a *t*-track layout of a graph *G*. The *span* of an edge *vw* of *G*, with respect to a numbering of the tracks $I = \{1, 2, ..., t\}$, is defined to be |i - j| where $v \in V_i$ and $w \in V_j$.

Track layouts will be central in most of our proofs. To enable comparison of our results to those in the literature we now introduce the notion of an 'improper' track layout. A *improper colouring* of a graph G is simply a partition $\{V_i : i \in I\}$ of V(G). Here adjacent vertices may be in the same colour class. A track of an improper colouring is defined as above. Suppose $\{V_i : i \in I\}$ is an improper colouring of G, and $(V_i, <_i)$ is a track, for each colour $i \in I$. An edge with both end-vertices in the same track is called an *intra-track* edge; otherwise it is called an *inter-track* edge. We say $\{(V_i, <_i) : i \in I\}$ is an *improper track assignment* of G if, for all intra-track edges $vw \in E(G)$ with $v \in V_i$ and $w \in V_i$ for some $i \in I$, there is no vertex x with $v <_i x <_i w$. That is, adjacent vertices in the same track are consecutive in that track. An improper t-track assignment with no X-crossing is called an *improper*

t-track layout³

LEMMA 2.2. If a graph G has an improper t-track layout, then G has a 2t-track layout.

Proof. For every track V_i of an improper t-track layout of G, let V'_i be a new track. Move every second vertex from V_i to V'_i , such that V'_i inherits its total order from the original V_i . Clearly there is no intra-track edge and no X-crossing. Thus we obtain a 2t-track layout of G

Hence the track-number of a graph is at most twice its 'improper track-number'. The following lemma, which was jointly discovered with Giuseppe Liotta, gives a compelling reason to only consider proper track layouts. Similar ideas can be found in [42, 27]. Let vw be an edge of a graph G. Let G' be the graph obtained from G by adding a new vertex x only adjacent to v and w. We say x is an *ear*, and G' is obtained from G by adding an ear to vw.

LEMMA 2.3. Let \mathcal{G} be a class of graphs closed under the addition of ears (for example, series-parallel graphs or planar graphs). If every graph in \mathcal{G} has an improper t-track layout for some constant t, then every graph in \mathcal{G} has a (proper) t-track layout.

Proof. For any graph $G \in \mathcal{G}$, let G' be the graph obtained from G by adding t ears to every edge of G. By assumption, G' has an improper t-track layout. Suppose that there is an edge vw of G such that v and w are in the same track. None of the ears added to vw are on the same track, as otherwise adjacent vertices would not be consecutive in that track. Thus there is a track containing at least two of the ears added to vw. However, this implies that there is an X-crossing, which is a contradiction. Thus the end-vertices of every edge of G are in distinct tracks. Hence the improper t-track layout of G' contains a t-track layout of G. п

Lemmata 2.2 and 2.3 imply that only for relatively small classes of graphs will the distinction between track layouts and improper track layouts be significant. We therefore chose to work with the less cumbersome notion of a track layout. The following theorem summarises our bounds on the track-number of a graph.

THEOREM 2.4. Let G be a graph with maximum degree $\Delta(G)$, path-width pw(G), tree-partition-width tpw(G), and tree-width tw(G). The track-number of G satisfies:

- (a) $tn(G) < pw(G) + 1 < 1 + (tw(G) + 1) \log n$,
- (b) $\operatorname{tn}(G) \leq 3 \operatorname{tpw}(G) \leq 72 \Delta(G) \operatorname{tw}(G),$ (c) $\operatorname{tn}(G) \leq 3^{\operatorname{tw}(G)} \cdot 6^{(4^{\operatorname{tw}(G)} 3 \operatorname{tw}(G) 1)/9}.$

Proof. Part (a) follows from Lemma 3.2, and the fact that $pw(G) \leq (tw(G) + C)$ 1) $\log n$ (see [10]). Note that $\operatorname{tn}(G) \leq 1 + (\operatorname{tw}(G) + 1) \log n$ can be proved directly using a separator-based approach similar to that used to prove $pw(G) \leq (tw(G) + 1) \log n$. Part (b) follows from Lemma 3.3 in §3, and the result of Ding and Oporowski [31] discussed in §2.2. Part (c) is Theorem 7.3. П

2.4. Vertex-Orderings. Let G be a graph. A total order $\sigma = (v_1, v_2, \ldots, v_n)$ of V(G) is called a *vertex-ordering* of G. Suppose G is connected. The *depth* of a vertex v_i in σ is the graph-theoretic distance between v_1 and v_i in G. We say σ is a breadth-first vertex-ordering if for all vertices v and w with $v <_{\sigma} w$, the depth of v in σ is no more than the depth of w in σ . Vertex-orderings, and in particular, vertex-orderings of trees will be used extensively in this paper. Consider a breadthfirst vertex-ordering σ of a tree T such that vertices at depth $d \geq 1$ are ordered with respect to the ordering of vertices at depth d-1. In particular, if v and x are vertices

³In [34, 35, 93] we called a track layout an ordered layering with no X-crossing and no intra-layer edges, and an improper track layout was called an ordered layering with no X-crossing.



FIG. 2.2. A lexicographical breadth-first vertex-ordering of a tree.

2.5. Queue Layouts. A queue layout of a graph G consists of a vertex-ordering σ of G, and a partition of E(G) into queues, such that no two edges in the same queue are nested with respect to σ . That is, there are no edges vw and xy in a single queue with $v <_{\sigma} x <_{\sigma} y <_{\sigma} w$. The minimum number of queues in a queue layout of G is called the queue-number of G, and is denoted by qn(G). A similar concept is that of a stack layout (or book embedding), which consists of a vertex-ordering σ of G, and a partition of E(G) into stacks (or pages) such that there are no edges vw and xy in a single stack with $v <_{\sigma} x <_{\sigma} w <_{\sigma} y$. The minimum number of stacks in a stack layout of G is called the stack-number (or page-number or book-thickness) of G, and is denoted by sn(G). A queue (respectively, stack) layout with k queues (stacks) is called a k-queue (k-stack) layout, and a graph that admits a k-queue (k-stack) layout is called a k-queue (k-stack) graph.

Heath and Rosenberg [58] characterised 1-queue graphs as the 'arched levelled planar' graphs, and proved that it is \mathcal{NP} -complete to recognise such graphs. This result is in contrast to the situation for stack layouts — 1-stack graphs are precisely the outerplanar graphs [8], which can be recognised in polynomial time. Heath *et al.* [54] proved that 1-stack graphs are 2-queue graphs (rediscovered by Rengarajan and Veni Madhavan [80]), and that 1-queue graphs are 2-stack graphs.

While it is \mathcal{NP} -hard to minimise the number of stacks in a stack layout given a fixed vertex-ordering [46], the analogous problem for queue layouts can be solved as follows. A *k*-rainbow in a vertex-ordering σ consists of a matching $\{v_iw_i : 1 \leq i \leq k\}$ such that $v_1 <_{\sigma} v_2 <_{\sigma} \cdots <_{\sigma} v_k <_{\sigma} w_k <_{\sigma} w_{k-1} <_{\sigma} \cdots <_{\sigma} w_1$, as illustrated in Fig. 2.3.



FIG. 2.3. A rainbow of five edges in a vertex-ordering.

A vertex-ordering containing a k-rainbow needs at least k queues. A straightforward application of Dilworth's Theorem [30] proves the converse. That is, a fixed vertex-ordering admits a k-queue layout where k is the size of the largest rainbow. (Heath and Rosenberg [58] describe a $\mathcal{O}(m \log \log n)$ time algorithm to compute the queue assignment.) Thus determining qn(G) can be viewed as the following vertexordering problem.

LEMMA 2.5 ([58]). The queue-number qn(G) of a graph G is the minimum, taken over all vertex-orderings σ of G, of the maximum size of a rainbow in σ .

Stack and/or queue layouts of k-trees have previously been investigated in [20,80, 45]. A 1-tree is a 1-queue graph, since in a lexicographical breadth-first vertexordering of a tree no two edges are nested (see Fig. 2.2). Chung et al. [20] proved that in a depth-first vertex-ordering of a tree no two edges cross. Thus 1-trees are 1-stack graphs. Rengarajan and Veni Madhavan [80] proved that graphs with tree-width at most two (the series parallel graphs) are 2-stack and 3-queue graphs⁴. Improper track layouts are implicit in the work of Heath et al. [54] and Rengarajan and Veni Madhavan [80]. In §5 we prove the following fundamental relationship between queue and track layouts.

THEOREM 2.6. For every graph G, $qn(G) \leq tn(G) - 1$. Moreover, if G is any proper minor-closed graph family, then \mathcal{G} has queue-number $qn(\mathcal{G}) \in \mathcal{F}(n)$ if and only if \mathcal{G} has track-number $tn(\mathcal{G}) \in \mathcal{F}(n)$, where $\mathcal{F}(n)$ is any family of functions closed under multiplication (such as $\mathcal{O}(1)$ or $\mathcal{O}(\operatorname{polylog} n)$).

Ganley and Heath [45] proved that every graph G has stack-number sn(G) < $\mathsf{tw}(G) + 1$ (using a depth-first traversal of a tree-decomposition), and asked whether queue-number is bounded by tree-width? One of the principal results of this paper is to solve this question in the affirmative. Applying Theorems 2.4 and 2.6 we have the following.

THEOREM 2.7. Let G be a graph with maximum degree $\Delta(G)$, path-width pw(G). tree-partition-width tpw(G), and tree-width tw(G). The queue-number qn(G) satis $fies^5$:

(a) $\operatorname{qn}(G) \leq \operatorname{pw}(G) \leq (\operatorname{tw}(G) + 1) \log n$,

(b) $qn(G) \leq 3 tpw(G) - 1 \leq 72\Delta(G) tw(G) - 1,$ (c) $qn(G) \leq 3 tw(G) \cdot 6^{(4^{tw(G)} - 3 tw(G) - 1)/9} - 1.$

A similar upper bound to Theorem 2.7(a) is obtained by Heath and Rosenberg [58], who proved that every graph G has $qn(G) \leq \lfloor \frac{1}{2}bw(G) \rfloor$, where bw(G)is the band-width of G. In many cases this result is weaker than Theorem 2.7(a) since pw(G) < bw(G) (see [29]). More importantly, we have the following corollary of Theorem 2.7(c).

COROLLARY 2.8. Queue-number is bounded by tree-width, and hence graphs with bounded tree-width have bounded queue-number.

2.6. Three-Dimensional Drawings. A three-dimensional straight-line grid drawing of a graph, henceforth called a three-dimensional drawing, represents the vertices by distinct points in \mathbb{Z}^3 (called *grid-points*), and represents each edge as a line-segment between its end-vertices, such that edges only intersect at common end-vertices, and an edge only intersects a vertex that is an end-vertex of that edge.

In contrast to the case in the plane, a folklore result states that every graph has a three-dimensional drawing. Such a drawing can be constructed using the 'moment curve' algorithm in which vertex v_i , $1 \leq i \leq n$, is represented by the grid-point (i, i^2, i^3) . It is easily seen — compare with Lemma 4.2 — that no two edges cross. (Two edges *cross* if they intersect at some point other than a common end-vertex.)

 $^{^{4}}$ In [35] we give a simple proof based on Theorem 6.1 for the result by Rengarajan and Veni Madhavan [80] that every series-parallel graph has a 3-queue layout.

⁵In [93] we obtained an alternative proof that $qn(G) \leq pw(G)$ using the 'vertex separation number' of a graph (which equals its path-width), and applying Lemma 2.5 directly we proved that $qn(G) \leq \frac{3}{2} tpw(G)$, and thus $qn(G) \leq 36 \Delta(G) tw(G)$.

Since every graph has a three-dimensional drawing, we are interested in optimising certain measures of the aesthetic quality of a drawing. If a three-dimensional drawing is contained in an axis-aligned box with side lengths X - 1, Y - 1 and Z - 1, then we speak of an $X \times Y \times Z$ drawing with volume $X \cdot Y \cdot Z$ and aspect ratio $\max\{X, Y, Z\}/\min\{X, Y, Z\}$. This paper considers the problem of producing a threedimensional drawing of a given graph with small volume, and with small aspect ratio as a secondary criterion.

Observe that the drawings produced by the moment curve algorithm have $\mathcal{O}(n^6)$ volume. Cohen *et al.* [21] improved this bound, by proving that if p is a prime with $n , and each vertex <math>v_i$ is represented by the grid-point $(i, i^2 \mod p, i^3 \mod p)$, then there is still no crossing. This construction is a generalisation of an analogous two-dimensional technique due to Erdős [40]. Furthermore, Cohen *et al.* [21] proved that the resulting $\mathcal{O}(n^3)$ volume bound is asymptotically optimal in the case of the complete graph K_n . It is therefore of interest to identify fixed graph parameters that allow for three-dimensional drawings with small volume.

The first such parameter to be studied was the chromatic number [17, 75]. Calamoneri and Sterbini [17] proved that every 4-colourable graph has a three-dimensional drawing with $\mathcal{O}(n^2)$ volume. Generalising this result, Pach *et al.* [75] proved that graphs of bounded chromatic number have three-dimensional drawings with $\mathcal{O}(n^2)$ volume, and that this bound is asymptotically optimal for the complete bipartite graph with equal sized bipartitions. If p is a suitably chosen prime, the main step of this algorithm represents the vertices in the *i*th colour class by grid-points in the set $\{(i, t, it) : t \equiv i^2 \pmod{p}\}$. It follows that the volume bound is $\mathcal{O}(k^2n^2)$ for k-colourable graphs.

The lower bound of Pach *et al.* [75] for the complete bipartite graph was generalised by Bose *et al.* [14] for all graphs. They proved that every three-dimensional drawing with *n* vertices and *m* edges has volume at least $\frac{1}{8}(n+m)$. In particular, the maximum number of edges in an $X \times Y \times Z$ drawing is exactly (2X - 1)(2Y - 1)(2Z - 1) - XYZ. For example, graphs admitting three-dimensional drawings with $\mathcal{O}(n)$ volume have $\mathcal{O}(n)$ edges.

The first non-trivial $\mathcal{O}(n)$ volume bound was established by Felsner *et al.* [42] for outerplanar graphs. Their elegant algorithm 'wraps' a two-dimensional drawing around a triangular prism to obtain an improper 3-track layout (see Lemmata 3.1 and 3.4 for more on this method). Poranen [78] proved that series-parallel digraphs have upward three-dimensional drawings with $\mathcal{O}(n^3)$ volume, and that this bound can be improved to $\mathcal{O}(n^2)$ and $\mathcal{O}(n)$ in certain special cases. Di Giacomo [27] proved that series-parallel graphs with maximum degree three have three-dimensional drawings with $\mathcal{O}(n)$ volume.

In §4 we prove the following intrinsic relationship between three-dimensional drawings and track layouts.

THEOREM 2.9. Every graph G has a $\mathcal{O}(\operatorname{tn}(G)) \times \mathcal{O}(\operatorname{tn}(G)) \times \mathcal{O}(n)$ drawing. Moreover, G has a $\mathcal{F}(n) \times \mathcal{F}(n) \times \mathcal{O}(n)$ drawing if and only if G has track-number $\operatorname{tn}(G) \in \mathcal{F}(n)$, where $\mathcal{F}(n)$ is a family of functions closed under multiplication.

Of course, every graph has an *n*-track layout — simply place a single vertex on each track. Thus Theorem 2.9 matches the $\mathcal{O}(n^3)$ volume bound discussed in §2.6. In fact, the drawings of K_n produced by our algorithm, with each vertex in a distinct track, are identical to those produced by the algorithm of Cohen *et al.* [21].

Theorems 2.6 and 2.9 immediately imply the following result, which reduces the problem of producing a three-dimensional drawing with small volume to that of pro-

ducing a queue layout of the same graph with few queues.

THEOREM 2.10. Let \mathcal{G} be a proper minor-closed family of graphs, and let $\mathcal{F}(n)$ be a family of functions closed under multiplication. The following are equivalent:

(a) every n-vertex graph in \mathcal{G} has a $\mathcal{F}(n) \times \mathcal{F}(n) \times \mathcal{O}(n)$ drawing,

(b) \mathcal{G} has track-number $\operatorname{tn}(\mathcal{G}) \in \mathcal{F}(n)$, and

(c) \mathcal{G} has queue-number $qn(\mathcal{G}) \in \mathcal{F}(n)$.

Graphs with constant queue-number include de Bruijn graphs, FFT and Beneš network graphs [58]. By Theorem 2.10, these graphs have three-dimensional drawings with $\mathcal{O}(n)$ volume. Applying Theorems 2.4 and 2.9 we have the following result.

THEOREM 2.11. Let G be a graph with maximum degree $\Delta(G)$, path-width pw(G), tree-partition-width tpw(G), and tree-width tw(G). Then G has a three-dimensional drawing with the following dimensions:

(a) $\mathcal{O}(\mathsf{pw}(G)) \times \mathcal{O}(\mathsf{pw}(G)) \times \mathcal{O}(n)$, which is $\mathcal{O}(\mathsf{tw}(G) \log n) \times \mathcal{O}(\mathsf{tw}(G) \log n) \times \mathcal{O}(n)$,

(b) $\mathcal{O}(\mathsf{tpw}(G)) \times \mathcal{O}(\mathsf{tpw}(G)) \times \mathcal{O}(n)$, which is $\mathcal{O}(\Delta(G) \mathsf{tw}(G)) \times \mathcal{O}(\Delta(G) \mathsf{tw}(G)) \times \mathcal{O}(n)$,

(c) $\mathcal{O}(3^{\mathsf{tw}(G)} \cdot 6^{(4^{\mathsf{tw}(G)} - 3 \mathsf{tw}(G) - 1)/9}) \times \mathcal{O}(3^{\mathsf{tw}(G)} \cdot 6^{(4^{\mathsf{tw}(G)} - 3 \mathsf{tw}(G) - 1)/9}) \times \mathcal{O}(n)$. Most importantly, we have the following corollary of Theorem 2.11(c).

COROLLARY 2.12. Every graph with bounded tree-width has a three-dimensional drawing with $\mathcal{O}(n)$ volume.

Note that bounded tree-width is not necessary for a graph to have a threedimensional drawing with $\mathcal{O}(n)$ volume. The $\sqrt{n} \times \sqrt{n}$ plane grid graph has $\Theta(\sqrt{n})$ tree-width, and has a $\sqrt{n} \times \sqrt{n} \times 1$ drawing with *n* volume. It also has a 3-track layout, and thus, by Lemma 4.2, has a $\mathcal{O}(1) \times \mathcal{O}(1) \times \mathcal{O}(n)$ drawing.

Since a planar graph is 4-colourable, by the results of Calamoneri and Sterbini [17] and Pach [75] discussed above, every planar graph has a three-dimensional drawing with $\mathcal{O}(n^2)$ volume. This result also follows from the classical algorithms of de Fraysseix *et al.* [23] and Schnyder [84] for producing $\mathcal{O}(n) \times \mathcal{O}(n)$ plane grid drawings. All of these methods produce $\mathcal{O}(n) \times \mathcal{O}(n) \times \mathcal{O}(1)$ drawings, which have $\Theta(n)$ aspect ratio. Since every planar graph G has $pw(G) \in \mathcal{O}(\sqrt{n})$ [10], we have the following corollary of Theorem 2.11(a).

COROLLARY 2.13. Every planar graph has a three-dimensional drawing with $\mathcal{O}(n^2)$ volume and $\Theta(\sqrt{n})$ aspect ratio.

This result matches the above $\mathcal{O}(n^2)$ volume bounds with an improvement in the aspect ratio by a factor of $\Theta(\sqrt{n})$. As discussed in §1, it is an open problem whether every planar graph has a three-dimensional drawing with $\mathcal{O}(n)$ volume. Subsequent to this research, Dujmović and Wood [37] proved that graphs excluding a clique minor on a fixed number of vertices, such as planar graphs, have three-dimensional drawings with $\mathcal{O}(n^{3/2})$ volume, as do graphs with bounded degree.

Our final result regarding three-dimensional drawings, which is proved in §4, examines the apparent trade-off between aspect ratio and volume.

THEOREM 2.14. For every graph G and for every r, $1 \le r \le n/\operatorname{tn}(G)$, G has a three-dimensional drawing with $\mathcal{O}(n^3/r^2)$ volume and aspect ratio 2r.

3. Track Layouts. In this section we describe a number of methods for producing and manipulating track layouts. The following result is implicit in the proof by Felsner *et al.* [42] that every outerplanar graph has an improper 3-track layout.

LEMMA 3.1 ([42]). Every tree T has a 3-track layout.

Proof. Root T at an arbitrary node r. Let σ be a lexicographical breadth-first vertex-ordering of T starting at r, as described in §2.4. For $i \in \{0, 1, 2\}$, let V_i be the

set of nodes of T with depth $d \equiv i \pmod{3}$ in σ . With each V_i ordered by σ , we have a 3-track assignment of T. Clearly adjacent vertices are on distinct tracks. Since no two edges are nested in σ , there is no X-crossing (see Fig. 3.1).



FIG. 3.1. A 3-track layout of a tree.

LEMMA 3.2. Every graph G with path-width pw(G) has track-number $tn(G) \le pw(G) + 1$.

Proof. Let k = pw(G) + 1. It is well known that a G is the subgraph of a kcolourable interval graph [10, 48]. That is, there is a set of intervals $\{[\ell(v), r(v)] \subseteq \mathbb{R} : v \in V(G)\}$ such that $[\ell(v), r(v)] \cap [\ell(w), r(w)] \neq \emptyset$ for every edge vw of G. Let $\{V_i : 1 \leq i \leq k\}$ be a k-colouring of G. Consider each colour class V_i to be an ordered track (v_1, v_2, \ldots, v_p) , where $\ell(v_1) < r(v_1) < \ell(v_2) < r(v_2) < \cdots < \ell(v_p) < r(v_p)$. Suppose there is an X-crossing between edges vw and xy with $v, x \in V_i$ and $w, y \in V_j$ for some pair of tracks V_i and V_j . Without loss of generality, $r(v) < \ell(x)$ and $r(y) < \ell(w)$. Since vw is an edge, $\ell(w) \leq r(v)$. Thus $r(y) < \ell(w) \leq r(v) < \ell(x)$, which implies that xy is not an edge of G. This contradiction proves that there is no X-crossing, and G has a k-track layout.



FIG. 3.2. A 4-track layout of a 4-colourable interval graph.

The next lemma uses a tree-partition to construct a track layout.

LEMMA 3.3. Every graph G with maximum degree $\Delta(G)$, tree-width tw(G), and tree-partition-width tpw(G), has track-number tn(G) ≤ 3 tpw(G) $\leq 72 \Delta(G)$ tw(G).

Proof. Let $(T, \{T_x : x \in V(T)\})$ be a tree-partition of G with width tpw(G). By Lemma 3.1, T has a 3-track layout. Replace each track by tpw(G) 'sub-tracks', and for each node x in T, place the vertices in bag T_x on the sub-tracks replacing the track containing x, with at most one vertex in T_x in a single track. For all nodes x and yof T, if x < y in a single track of the 3-track layout of T, then for all vertices $v \in T_x$ and $w \in T_y$, v < w whenever v and w are assigned to the same track. There is no X-crossing, since in the track layout of T, adjacent nodes are on distinct tracks and there is no X-crossing. Thus we have a track layout of G. The number of tracks is 3 tpw(G), which is at most $72 \Delta(G) tw(G)$ by the theorem of Ding and Oporowski [31] discussed in §2.2.

In the remainder of this section, we prove two results that show how track layouts can be manipulated without introducing an X-crossing. The first is a generalisation of the 'wrapping' algorithm of Felsner *et al.* [42], who implicitly proved the case s = 1. LEMMA 3.4. If a graph G has an (improper) track layout $\{(V_i, <_i) : 1 \le i \le t\}$

with maximum edge span s, then G has an (improper) (2s + 1)-track layout. Proof. Let $\ell = 2s + 1$. Construct an ℓ -track assignment of G by merging the tracks $\{V_i : i \equiv j \pmod{t}\}$ for each $j, 0 \leq j \leq t - 1$, with vertices in V_{α} appearing

tracks $\{v_i : i \equiv j \pmod{t}\}$ for each $j, 0 \leq j \leq t-1$, with vertices in v_{α} appearing before vertices in V_{β} in the new track j, for all $\alpha, \beta \equiv j \pmod{t}$ with $\alpha < \beta$. The given order of each V_i is preserved in the new tracks. It remains to prove that there is no X-crossing. Consider two edges vw and xy. Let i_1 and $i_2, 1 \leq i_1 < i_2 \leq t$, be the minimum and maximum tracks containing v, w, x or y in the given t-track layout of G.

First consider the case that $i_2 - i_1 > 2s$. Then without loss of generality v is in track i_2 and y is in track i_1 . Thus w is in a greater track than x, and even if x (or y) appear on the same track as v (or w) in the new ℓ -track assignment, x (or y) will be to the left of v (or w). Thus these edges do not form an X-crossing in the ℓ -track assignment. Otherwise $i_2 - i_1 \leq 2s$. Thus any two of v, w, x or y will appear on the same track in the ℓ -track assignment if and only if they are on the same track in the given t-track layout (since $\ell > 2s$). Hence the only way for these four vertices to appear on exactly two tracks in the ℓ -track assignment is if they were on exactly two layers in the given t-track layout, in which case, by assumption vw and xy do not form an X-crossing. Therefore there is no X-crossing, and we have an ℓ -track layout of G.

The next result shows that the number of vertices in different tracks of a track layout can be balanced without introducing an X-crossing. The proof is based on an idea due to Pach *et al.* [75] for balancing the size of the colour classes in a colouring.

LEMMA 3.5. If a graph G has an (improper) t-track layout, then for every t' > 0, G has an (improper) $\lfloor t + t' \rfloor$ -track layout with at most $\lceil \frac{n}{t'} \rceil$ vertices in each track.

Proof. For each track with $q > \lfloor \frac{n}{t'} \rfloor$ vertices, replace it by $\lfloor q/\lfloor \frac{n}{t'} \rfloor$ 'sub-tracks' each with exactly $\lfloor \frac{n}{t'} \rfloor$ vertices except for at most one sub-track with $q \mod \lfloor \frac{n}{t'} \rfloor$ vertices, such that the vertices in each sub-track are consecutive in the original track, and the original order is maintained. There is no X-crossing between sub-tracks from the same original track as there is at most one edge between such sub-tracks. There is no X-crossing between sub-tracks from different original tracks as otherwise there would be an X-crossing in the original. There are at most $\lfloor t' \rfloor$ tracks with $\lfloor \frac{n}{t'} \rfloor$ vertices. Since there are at most t tracks with less than $\lfloor \frac{n}{t'} \rfloor$ vertices, one for each of the original tracks, there is a total of at most $\lfloor t + t' \rfloor$ tracks.

4. Three-Dimensional Drawings and Track Layouts. In this section we prove Theorem 2.9, which states that three-dimensional drawings with small volume are closely related to track layouts with few tracks.

LEMMA 4.1. If a graph G has an $A \times B \times C$ drawing, then G has an improper AB-track layout, and G has a 2AB-track layout.

Proof. Let $V_{x,y}$ be the set of vertices of G with an X-coordinate of x and a Y-coordinate of y, where without loss of generality $1 \le x \le A$ and $1 \le y \le Y$. With each set $V_{x,y}$ ordered by the Z-coordinates of its elements, $\{V_{x,y} : 1 \le x \le A, 1 \le y \le Y\}$ is an improper AB-track assignment. There is no X-crossing, as otherwise there would be a crossing in the original drawing, and hence we have an improper AB-track layout. By Lemma 2.2, G has a 2AB-track layout.

We now prove the converse of Lemma 4.1. The proof is inspired by the generalisations of the moment curve algorithm by Cohen *et al.* [21] and Pach *et al.* [75], described in §2.6. Loosely speaking, Cohen *et al.* [21] allow three 'free' dimensions, whereas Pach *et al.* [75] use the assignment of vertices to colour classes to 'fix' one dimension with two dimensions free. We use an assignment of vertices to tracks to fix two dimensions with one dimension free. The style of three-dimensional drawing produced by our algorithm, where tracks are drawn vertically, is illustrated in Fig. 4.1.



FIG. 4.1. A three-dimensional drawing produced from a track layout.

LEMMA 4.2. If a graph G has a (possibly) improper k-track layout, then G has a $k \times 2k \times 2k \cdot n'$ three-dimensional drawing, where n' is the maximum number of vertices in a track.

Proof. Suppose $\{(V_i, <_i) : 1 \le i \le k\}$ is the given improper k-track layout. Let p be the smallest prime such that p > k. Then $p \le 2k$ by Bertrand's postulate. For each $i, 1 \le i \le k$, represent the vertices in V_i by the grid-points

$$\{(i, i^2 \mod p, t) : 1 \le t \le p \cdot |V_i|, t \equiv i^3 \pmod{p}\}$$

such that the Z-coordinates respect the given total order $\langle i \rangle$. Draw each edge as a line-segment between its end-vertices. Suppose two edges e and e' cross such that their end-vertices are at distinct points $(i_{\alpha}, i_{\alpha}^2 \mod p, t_{\alpha}), 1 \leq \alpha \leq 4$. Then these points are coplanar, and if M is the matrix

$$M = \begin{pmatrix} 1 & i_1 & i_1^2 \mod p & t_1 \\ 1 & i_2 & i_2^2 \mod p & t_2 \\ 1 & i_3 & i_3^2 \mod p & t_3 \\ 1 & i_4 & i_4^2 \mod p & t_4 \end{pmatrix}$$

then the determinant det(M) = 0. We proceed by considering the number of distinct tracks $N = |\{i_1, i_2, i_3, i_4\}|$.

• N = 1: By the definition of an improper track layout, e and e' do not cross.

• N = 2: If either edge is intra-track then e and e' do not cross. Otherwise neither edge is intra-track, and since there is no X-crossing, e and e' do not cross.

• N = 3: Without loss of generality $i_1 = i_2$. It follows that $det(M) = (t_2 - t_1) \cdot det(M')$, where

$$M' = \begin{pmatrix} 1 & i_2 & i_2^2 \mod p \\ 1 & i_3 & i_3^2 \mod p \\ 1 & i_4 & i_4^2 \mod p \end{pmatrix}$$

Since $t_1 \neq t_2$, det(M') = 0. However, M' is a Vandermonde matrix modulo p, and thus

$$\det(M') \equiv (i_2 - i_3)(i_2 - i_4)(i_3 - i_4) \pmod{p},$$

which is non-zero since i_2 , i_3 and i_4 are distinct and p is a prime, a contradiction.

• N = 4: Let M' be the matrix obtained from M by taking each entry modulo p. Then det(M') = 0. Since $t_{\alpha} \equiv i_{\alpha}^3 \pmod{p}$, $1 \leq \alpha \leq 4$,

$$M' \equiv \begin{pmatrix} 1 & i_1 & i_1^2 & i_1^3 \\ 1 & i_2 & i_2^2 & i_2^3 \\ 1 & i_3 & i_3^2 & i_3^3 \\ 1 & i_4 & i_4^2 & i_4^3 \end{pmatrix} \pmod{p}$$

Since each $i_{\alpha} < p, M'$ is a Vandermonde matrix modulo p, and thus

$$\det(M') \equiv (i_1 - i_2)(i_1 - i_3)(i_1 - i_4)(i_2 - i_3)(i_2 - i_4)(i_3 - i_4) \pmod{p}$$

which is non-zero since $i_{\alpha} \neq i_{\beta}$ and p is a prime. This contradiction proves there are no edge crossings. The produced drawing is at most $k \times 2k \times 2k \cdot n'$.

Proof of Theorem 2.9. Let $\mathcal{F}(n)$ be a family of functions closed under multiplication. Let G be an n-vertex graph with a t-track layout, where $t \in \mathcal{F}(n)$. By Lemma 3.5 with t' = t, G has a 2t-track layout with at most $\lceil \frac{n}{t} \rceil$ vertices in each track. By Lemma 4.2, G has a $2t \times 4t \times 4t \cdot \lceil \frac{n}{t} \rceil$ drawing, which is $\mathcal{O}(t) \times \mathcal{O}(t) \times \mathcal{O}(n)$. Conversely, suppose an n-vertex graph G has a $A \times B \times \mathcal{O}(n)$ drawing, where $A, B \in \mathcal{F}(n)$. By Lemma 4.1, G has a track layout with $2AB \in \mathcal{F}(n)$ tracks.

Proof of Theorem 2.14. Let $t = \operatorname{tn}(G)$, and suppose $1 \le r \le n/t$. By Lemma 3.5 with $t' = \frac{n}{r}$, G has a $\lfloor \frac{n}{r} + t \rfloor$ -track layout with at most r vertices in each track. By assumption $t \le \frac{n}{r}$, and the number of tracks is at most $\frac{2n}{r}$. By Lemma 4.2, G has a $\frac{2n}{r} \times \frac{4n}{r} \times 4n$ three-dimensional drawing, which has volume $32n^3/r^2$ and aspect ratio 2r.

5. Queue Layouts and Track Layouts. In this section we prove Theorem 2.6, which states that track and queue layouts are closely related. Our first lemma highlights this fact — its proof follows immediately from the definitions (see Fig. 5.1).

LEMMA 5.1. A bipartite graph G = (A, B; E) has a 2-track layout with tracks A and B if and only if G has a 1-queue layout such that in the corresponding vertex-ordering, the vertices in A appear before the vertices in B.



FIG. 5.1. A 2-track layout and a 1-queue layout of a bipartite graph.

We now show that a queue layout can be obtained from a track layout. This result can be viewed as a generalisation of the construction of a 2-queue layout of an outerplanar graph by Heath *et al.* [54] and Rengarajan and Veni Madhavan [80] (with s = 1).

LEMMA 5.2. If a graph G has a (possibly) improper t-track layout $\{(V_i, <_i) : 1 \le i \le t\}$ with maximum edge span s $(\le t-1)$, then $qn(G) \le s+1$, and if the given track layout is not improper, then $qn(G) \le s$.

Proof. First suppose that there are no intra-track edges. Let σ be the vertex ordering (V_1, V_2, \ldots, V_t) of G. Let E_{α} be the set of edges with span α in the given track layout. As in Lemma 5.1, two edges from the same pair of tracks are nested in σ if and only if they form an X-crossing in the track layout. Since no two edges form an X-crossing in the track layout, no two edges that are between the same pair of tracks have the same pair of tracks are nested in σ . If two edges not from the same pair of tracks have the same span then they are not nested in σ . (This idea is due to Heath and Rosenberg [58].) Thus no two edges are nested in each E_{α} , and we have an s-queue layout of G. If there are intra-track edges, then they all form one additional queue in σ .

We now set out to prove the converse of Lemma 5.2. It is well known that the subgraph induced by any two tracks of a track layout is a forest of caterpillars [52]. A colouring of a graph is *acyclic* if every bichromatic subgraph is a forest; that is, every cycle receives at least three distinct colours. Thus a *t*-track layout of a graph G defines an acyclic *t*-colouring of G. The minimum number of colours in an acyclic colouring of G is the *acyclic chromatic number* of G, denoted by $\chi_{\mathbf{a}}(G)$. Thus,

$$\chi_{a}(G) \leq \operatorname{tn}(G)$$

Acyclic colourings were introduced by Grünbaum [49], who proved that every planar graph is acyclically 9-colourable. This result was steadily improved [1, 65, 68] until Borodin [12] proved that every planar graph is acyclically 5-colourable, which is the best possible bound. Many other graph families have bounded acyclic chromatic number, including graphs embeddable on a fixed surface [2, 3, 6], 1-planar graphs [13], graphs with bounded maximum degree [5], and graphs with bounded tree-width. A folklore result states that $\chi_a(G) \leq \operatorname{tw}(G) + 1$ (see [43]). More generally, Nešetřil and Ossona de Mendez [71] proved that every proper minor-closed graph family has bounded acyclic chromatic number. In fact, Nešetřil and Ossona de Mendez [71] proved that every graph G has a star k-colouring (every bichromatic subgraph is a forest of stars), where k is a (small) quadratic function of the maximum chromatic number of a minor of G.

LEMMA 5.3. Every graph G with acyclic chromatic number $\chi_a(G) \leq c$ and queuenumber $qn(G) \leq q$ has track-number $tn(G) \leq c (2q)^{c-1}$.

Proof. Let $\{V_i : 1 \leq i \leq c\}$ be an acyclic colouring of G. Let σ be the vertexordering in a q-queue layout of G. Consider an edge vw with $v \in V_i$, $w \in V_j$, and i < j. If $v <_{\sigma} w$ then vw is forward, and if $w <_{\sigma} v$ then vw is backward. Consider the edges to be coloured with 2q colours, where each colour class consists of the forward edges in a single queue, or the backward edges in a single queue.

Alon and Marshall [4] proved that given a (not necessarily proper) edge k-colouring of a graph G, any acyclic c-colouring of G can be refined to a ck^{c-1} -colouring so that the edges between any pair of (vertex) colour classes are monochromatic, and each (vertex) colour class is contained in some original colour class. (Nešetřil and Raspaud [72] generalised this result for coloured mixed graphs.) Apply this result with the given acyclic c-colouring of G and the edge 2q-colouring discussed above. Consider the resulting $c(2q)^{c-1}$ colour classes to be tracks ordered by σ . The edges between any two tracks are from a single queue, and are all forward or all backward.

Suppose that there are edges vw and xy that form an X-crossing. Since each track is a subset of some V_i , we can assume that $v, x \in V_i$, $w, y \in V_j$ and i < j. Suppose that vw and xy are both forward. The case in which vw and xy are both backward is symmetric. Thus $v <_{\sigma} w$ and $x <_{\sigma} y$. Since vw and xy form an X-crossing, and the tracks are ordered by σ , we have $v <_{\sigma} x$ and $y <_{\sigma} w$. Hence $v <_{\sigma} x <_{\sigma} y <_{\sigma} w$. That is, vw and xy are nested. This is the desired contradiction, since edges between any pair of tracks are from a single queue. Thus we have a $c(2q)^{c-1}$ -track layout of G.

Proof of Theorem 2.6. Let $\mathcal{F}(n)$ be a family of functions closed under multiplication. Let G be an n-vertex graph from a proper minor-closed graph family \mathcal{G} . First, suppose that G has a t-track layout, where $t \in \mathcal{F}(n)$. By Lemma 5.2, G has queue-number $qn(G) \leq t-1 \in \mathcal{F}(n)$. Conversely, suppose G has queuenumber $qn(G) = q \in \mathcal{F}(n)$. By the above-mentioned result of Nešetřil and Ossona de Mendez [71], G has bounded acyclic chromatic number $\chi_a(G) \leq c \in \mathcal{O}(1)$. By Lemma 5.3, G has a t-track layout, where $t \leq c(2q)^{c-1} \in \mathcal{F}(n)$.

6. Tree-Partitions of k-Trees. In this section we prove our theorem regarding tree-partitions of k-trees mentioned in §2.2. This result forms the cornerstone of the proof of Theorem 7.3.

THEOREM 6.1. Let G be a k-tree with maximum degree Δ . Then G has a rooted tree-partition $(T, \{T_x : x \in V(T)\})$ such that for all nodes x of T,

(a) if x is a non-root node of T and y is the parent node of x, then the set of vertices in T_y with a neighbour in T_x form a clique C_x of G, and

(b) the induced subgraph $G[T_x]$ is a connected (k-1)-tree.

Furthermore the width of $(T, \{T_x : x \in V(T)\})$ is at most $\max\{1, k(\Delta - 1)\}$.

Proof. We assume G is connected, since if G is not connected then a tree-partition of G that satisfies the theorem can be determined by adding a new root node with an empty bag, adjacent to the root node of a tree-partition of each connected component of G.

It is well-known that G is a connected k-tree if and only if G has a vertex-ordering $\sigma = (v_1, v_2, \ldots, v_n)$, such that for all $i \in \{1, 2, \ldots, n\}$,

(i) if G^i is the induced subgraph $G[\{v_1, v_2, \ldots, v_i\}]$, then G^i is connected and the vertex-ordering of G^i induced by σ is a breadth-first vertex-ordering of G^i , and

(ii) the neighbours of v_i in G^i form a clique $C_i = \{v_j : v_i v_j \in E(G), j < i\}$ with $1 \le |C_i| \le k$ (unless i = 1 in which case $C_i = \emptyset$).

In the language of chordal graphs, σ is a (reverse) 'perfect elimination' vertexordering and can be determined, for example, by the Lex-BFS algorithm by Rose *et al.* [82] (also see [48]). Moreover, we can choose v_1 to be any vertex in G.

Let r be a vertex of minimum degree⁶ in G. Then deg(r) $\leq k$. Let $\sigma = (v_1, v_2, \ldots, v_n)$ be a vertex-ordering of G with $v_1 = r$, and satisfying (i) and (ii). By (i), the depth of each vertex v_i in σ is the same as the depth of v_i in the vertex-ordering of G^j induced by σ , for all $j \geq i$. We therefore simply speak of the depth of v_i . Let V_d be the set of vertices of G at depth d.

CLAIM 1. For all $d \ge 1$, and for every connected component Z of $G[V_d]$, the set of vertices at depth d-1 with a neighbour in Z form a clique of G.

Proof. The claim in trivial for d = 1 or d = 2. Now suppose that $d \ge 3$. Assume for the sake of contradiction that there are two non-adjacent vertices x and y at depth d-1, such that x has a neighbour in Z and y has a neighbour in Z. Let P_1 be a shortest path between x and y with its interior vertices in Z. Let P_2 be a shortest path between x and y with its interior vertices at depth at most d-2. Since the interior vertices of P_1 are at depth d, there is no edge between an interior vertex of

⁶We choose r to have minimum degree to obtain a slightly improved bound on the width of the tree-partition. If we choose r to be an arbitrary vertex then the width is at most $\max\{1, \Delta, k(\Delta - 1)\}$, and the remainder of Theorem 6.1 holds.

 P_1 and an interior vertex of P_2 . Thus $P_1 \cup P_2$ is a chordless cycle of length at least four, contradicting the fact that G is chordal (by Lemma 2.1).

Define a graph T and a partition $\{T_x : x \in V(T)\}$ of V(G) indexed by the nodes of T as follows. There is one node x in T for every connected component of each $G[V_d]$, whose bag T_x is the vertex-set of the corresponding connected component. We say x and T_x are at depth d. Clearly a vertex in a depth-d bag is also at depth d. The (unique) node of T at depth zero is called the *root* node. Let two nodes x and y of Tbe connected by an edge if there is an edge vw of G with $v \in T_x$ and $w \in T_y$. Thus $\{T_x : x \in V(T)\}$ is a 'graph-partition'.

We now prove that in fact T is a tree. First observe that T is connected since G is connected. By definition, nodes of T at the same depth d are not adjacent. Moreover nodes of T can be adjacent only if their depths differ by one. Thus T has a cycle only if there is a node x in T at some depth d, such that x has at least two distinct neighbours in T at depth d-1. However this is impossible since by Claim 1, the set of vertices at depth d-1 with a neighbour in T_x form a clique (which we call C_x), and are hence in a single bag at depth d-1. Thus T is a tree and $(T, \{T_x : x \in V(T)\})$ is a tree-partition of G (see Fig. 6.1).



FIG. 6.1. Illustration for Theorem 6.1 in the case of k = 3.

We now prove that each bag T_x induces a connected (k-1)-tree. This is true for the root node which only has one vertex. Suppose x is a non-root node of T at depth d. Each vertex in T_x has at least one neighbour at depth d-1. Thus in the vertex-ordering of T_x induced by σ , each vertex $v_i \in T_x$ has at most k-1 neighbours $v_j \in T_x$ with j < i. Thus the vertex-ordering of T_x induced by σ satisfies (i) and (ii) for k-1, and $G[T_x]$ is (k-1)-tree. By definition each $G[T_x]$ is connected.

Finally, consider the cardinality of a bag in T. We claim that each bag contains at most $\max\{1, k(\Delta - 1)\}$ vertices. The root bag has one vertex. Let x be a non-root

node of T with parent node y. Suppose y is the root node. Then $T_y = \{r\}$, and thus $|T_x| \leq \deg(r) \leq k \leq k(\Delta - 1)$ assuming $\Delta \geq 2$. If $\Delta \leq 1$ then all bags have one vertex. Now assume y is a non-root node. The set of vertices in T_y with a neighbour in T_x forms the clique C_x . Let $k' = |C_x|$. Thus $k' \geq 1$, and since $C_x \subseteq T_y$ and $G[T_y]$ is a (k-1)-tree, $k' \leq k$. A vertex $v \in C_x$ has k'-1 neighbours in C_x and at least one neighbour in the parent bag of y. Thus v has at most $\Delta - k'$ neighbours in T_x . Hence the number of edges between C_x and T_x is at most $k'(\Delta - k')$. Every vertex in T_x is adjacent to a vertex in C_x . Thus $|T_x| \leq k'(\Delta - k') \leq k(\Delta - 1)$. This completes the proof.

7. Tree-Width and Track Layouts. In this section we prove that tracknumber is bounded by tree-width. Let $\{(V_i, <_i) : i \in I\}$ be a track layout of a graph G. We say a clique C of G covers the set of tracks $\{i \in I : C \cap V_i \neq \emptyset\}$. Let S be a set of cliques of G. Suppose there exists a total order \preceq on S such that for all cliques $C_1, C_2 \in S$, if there exists a track $i \in I$, and vertices $v \in V_i \cap C_1$ and $w \in V_i \cap C_2$ with $v <_i w$, then $C_1 \prec C_2$. In this case, we say \preceq is nice, and S is nicely ordered by the track layout.

LEMMA 7.1. Let $L \subseteq I$ be a set of tracks in a track layout $\{(V_i, <_i) : i \in I\}$ of a graph G. If S is a set of cliques, each of which covers L, then S is nicely ordered by the given track layout.

Proof. Define a relation \leq on S as follows. For every pair of cliques $C_1, C_2 \in S$, define $C_1 \leq C_2$ if $C_1 = C_2$ or there exists a track $i \in L$ and vertices $v \in C_1$ and $w \in C_2$ with $v <_i w$. Clearly all cliques in S are comparable.

Suppose that \leq is not antisymmetric; that is, there exists distinct cliques $C_1, C_2 \in S$, distinct tracks $i, j \in L$, and distinct vertices $v_1, w_1 \in C_1$ and $v_2, w_2 \in C_2$, such that $v_1 <_i v_2$ and $w_2 <_j w_1$. Since C_1 and C_2 are cliques, the edges v_1w_1 and v_2w_2 form an X-crossing, which is a contradiction. Thus \leq is antisymmetric.

We claim that \leq is transitive. Suppose there exist cliques $C_1, C_2, C_3 \in S$ such that $C_1 \leq C_2$ and $C_2 \leq C_3$. We can assume that C_1, C_2 and C_3 are pairwise distinct. Thus there are vertices $u_1 \in C_1$, $u_2 \in C_2$, $v_2 \in C_2$ and $v_3 \in C_3$, such that $u_1 <_i u_2$ and $v_2 <_j v_3$ for some pair of (not necessarily distinct) tracks $i, j \in L$. Since C_3 has a vertex in V_i and since $C_3 \leq C_2$, there is a vertex $u_3 \in C_3$ with $u_2 \leq_i u_3$. Thus $u_1 <_i u_3$, which implies that $C_1 \leq C_3$. Thus \leq is transitive.

Hence \leq is a total order on S, which by definition is nice.

Consider the problem of partitioning the cliques of a graph into sets such that each set is nicely ordered by a given track layout. The following immediate corollary of Lemma 7.1 says that there exists such a partition where the number of sets does not depend upon the size of the graph.

COROLLARY 7.2. Let G be a graph with maximum clique size k. Given a t-track layout of G, there is a partition of the cliques of G into $\sum_{i=1}^{k} {t \choose i}$ sets, each of which is nicely ordered by the given track layout.

We do not actually use Corollary 7.2 in the following result, but the idea of partitioning the cliques into nicely ordered sets is central to its proof.

THEOREM 7.3. For every integer $k \ge 0$, there is a constant $t_k = 3^k \cdot 6^{(4^k - 3k - 1)/9}$ such that every graph G with tree-width $\mathsf{tw}(G) \le k$ has a t_k -track layout.

Proof. If the input graph G is not a k-tree then add edges to G to obtain a k-tree containing G as a subgraph. It is well-known that a graph with tree-width at most k is a *spanning* subgraph of a k-tree. These extra edges can be deleted once we are done. We proceed by induction on k with the following hypothesis:

For all $k \in \mathbb{N}$, there exists a constant s_k , and sets \mathcal{I}_k and \mathcal{S}_k such that

1. $|\mathcal{I}_k| = t_k \text{ and } |\mathcal{S}_k| = s_k$,

2. each element of S_k is a subset of I_k , and

3. every k-tree G has a t_k -track layout indexed by \mathcal{I}_k , such that for every clique C of G, the set of tracks that C covers is in \mathcal{S}_k .

Consider the base case with k = 0. A 0-tree G has no edges and thus has a 1-track layout. Let $\mathcal{I}_0 = \{1\}$ and order $V_1 = V(G)$ arbitrarily. Thus $t_0 = 1$, $s_0 = 1$, and $\mathcal{S}_0 = \{\{1\}\}$ satisfy the hypothesis for every 0-tree. Now suppose the result holds for k - 1, and G is a k-tree.

Let $(T, \{T_x : x \in V(T)\})$ be a tree-partition of G described in Theorem 6.1, where T is rooted at r. Each induced subgraph $G[T_x]$ is a (k-1)-tree. Thus, by induction, there are sets \mathcal{I}_{k-1} and \mathcal{S}_{k-1} with $|\mathcal{I}_{k-1}| = t_{k-1}$ and $|\mathcal{S}_{k-1}| = s_{k-1}$, such that for every node x of T, the induced subgraph $G[T_x]$ has a t_{k-1} -track layout indexed by \mathcal{I}_{k-1} . For every clique C of $G[T_x]$, if C covers $L \subseteq \mathcal{I}_{k-1}$ then $L \in \mathcal{S}_{k-1}$. Assume $\mathcal{I}_{k-1} = \{1, 2, \ldots, t_{k-1}\}$ and $\mathcal{S}_{k-1} = \{X_1, X_2, \ldots, X_{s_{k-1}}\}$. By Theorem 6.1, for each non-root node x of T, if p is the parent node of x, then the set of vertices in T_p with a neighbour in T_x form a clique C_x . Let $\alpha(x) = i$ where C_x covers X_i . For the root node r of T, let $\alpha(r) = 1$.

Track layout of T**.** To construct a track layout of G we first construct a track layout of the tree T indexed by the set $\{(d, i) : d \ge 0, 1 \le i \le s_{k-1}\}$, where the track $L_{d,i}$ consists of nodes x of T at depth d with $\alpha(x) = i$. Here the *depth* of a node x is the distance in T from the root node r to x. We order the nodes of T within the tracks by increasing depth. There is only one node at depth d = 0. Suppose we have determined the orders of the nodes up to depth d - 1 for some $d \ge 1$.

Let $i \in \{1, 2, \ldots, s_{k-1}\}$. The nodes in $L_{d,i}$ are ordered primarily with respect to the relative positions of their parent nodes (at depth d-1). More precisely, let $\rho(x)$ denote the parent node of each node $x \in L_{d,i}$. For all nodes x and y in $L_{d,i}$, if $\rho(x)$ and $\rho(y)$ are in the same track and $\rho(x) < \rho(y)$ in that track, then x < y in $L_{d,i}$. For x and y with $\rho(x)$ and $\rho(y)$ on distinct tracks, the relative order of x and y is not important. It remains to specify the order of nodes in $L_{d,i}$ with a common parent.

Suppose P is a set of nodes in $L_{d,i}$ with a common parent node p. By construction, for every node $x \in P$, the parent clique C_x covers X_i in the track layout of $G[T_p]$. By Lemma 7.1 the cliques $\{C_x : x \in P\}$ are nicely ordered by the track layout of $G[T_p]$. Let the order of P in track $L_{d,i}$ be specified by a nice ordering of $\{C_x : x \in P\}$, as illustrated in Fig. 7.1.

This construction defines a partial order on the nodes in track $L_{d,i}$, which can be arbitrarily extended to a total order. Hence we have a track assignment of T. Since the nodes in each track are ordered primarily with respect to the relative positions of their parent nodes in the previous tracks, there is no X-crossing, and hence we have a track layout of T.

Track layout of G. To construct a track assignment of G from the track layout of T, replace each track $L_{d,i}$ by t_{k-1} 'sub-tracks', and for each node x of T, insert the track layout of $G[T_x]$ in place of x on the sub-tracks corresponding to the track containing x in the track layout of T. More formally, the track layout of G is indexed by the set

$$\{(d, i, j) : d \ge 0, 1 \le i \le s_{k-1}, 1 \le j \le t_{k-1}\}$$

Each track $V_{d,i,j}$ consists of those vertices v of G such that, if T_x is the bag containing v, then x is at depth d in T, $\alpha(x) = i$, and v is in track j in the track layout of $G[T_x]$.



FIG. 7.1. Track layout of nodes with a common parent p.

If x and y are distinct nodes of T with x < y in $L_{d,i}$, then v < w in $V_{d,i,j}$, for all vertices $v \in T_x$ and $w \in T_y$ in track j. If v and w are vertices of G in track j in bag T_x at depth d, then the relative order of v and w in $V_{d,\alpha(x),j}$ is the same as in the track layout of $G[T_x]$.

Clearly adjacent vertices of G are in distinct tracks. Thus we have defined a track assignment of G. We claim there is no X-crossing. Clearly an intra-bag edge of Gis not in an X-crossing with an edge not in the same bag. By induction, there is no X-crossing between intra-bag edges in a common bag. Since there is no X-crossing in the track layout of T, inter-bag edges of G which are mapped to edges of T without a common parent node, are not involved in an X-crossing.

Consider a parent node p in T. For each child node x of p, the set of vertices in T_p adjacent to a vertex in T_x forms the clique C_x . Thus there is no X-crossing between a pair of edges both from C_x to T_x , since the vertices of C_x are on distinct tracks. Consider two child nodes x and y of p. For there to be an X-crossing between an edge from T_p to T_x and an edge from T_p to T_y , the nodes x and y must be on the same track in the track layout of T. Suppose x < y in this track. By construction, C_x and C_y cover the same set of tracks, and $C_x \leq C_y$ in the corresponding nice ordering. Thus for any track containing vertices $v \in C_x$ and $w \in C_y$, $v \leq w$ in that track. Since all the vertices in T_x are to the left of the vertices in T_y (in a common track), there is no X-crossing between an edge from T_p to T_x and an edge from T_p to T_y . Therefore there is no X-crossing, and hence we have a track layout of G.

Wrapped track layout of G. As illustrated in Fig. 7.2, we now 'wrap' the track layout of G in the spirit of Lemma 3.1. In particular, define a track assignment of G indexed by

 $\{(d', i, j) : d' \in \{0, 1, 2\}, 1 \le i \le s_{k-1}, 1 \le j \le t_{k-1}\}$,

where each track

$$W_{d',i,j} = \bigcup \{ V_{d,i,j} : d \equiv d' \pmod{3} \}$$

If $v \in V_{d,i,j}$ and $w \in V_{d+3,i,j}$ then v < w in the order of $W_{d',i,j}$ (where $d' = d \mod 3$). The order of each $V_{d,i,j}$ is preserved in $W_{d',i,j}$. The set of tracks $\{W_{d',i,j} : d' \in \{0,1,2\}, 1 \le i \le s_{k-1}, 1 \le j \le t_{k-1}\}$ forms a track assignment of G. For every edge vw of G, the depths of the bags in T containing v and w differ by at most one. Thus in the wrapped track assignment of G, adjacent vertices remain on distinct tracks, and there is no X-crossing. The number of tracks is $3 \cdot s_{k-1} \cdot t_{k-1}$.

Every clique C of G is either contained in a single bag of the tree-partition or is contained in two adjacent bags. Let

$$\mathcal{S}' = \left\{ \left\{ (d', i, h) : h \in X_j \right\} : d' \in \{0, 1, 2\}, 1 \le i, j \le s_{k-1} \right\}$$

For every clique C of G contained in a single bag, the set of tracks containing C is in \mathcal{S}' . Let

$$\mathcal{S}'' = \left\{ \{ (d', i, \ell) : \ell \in X_j \} \cup \{ ((d'+1) \mod 3, p, h) : h \in X_q \} : d' \in \{0, 1, 2\}, 1 \le i, j, p, q \le s_{k-1} \right\}.$$

For every clique C of G contained in two bags, the set of tracks containing C is in \mathcal{S}'' . Observe that $\mathcal{S}' \cup \mathcal{S}''$ is independent of G. Hence $\mathcal{S}_k = \mathcal{S}' \cup \mathcal{S}''$ satisfies the hypothesis for k.

Now $|\mathcal{S}'| = 3s_{k-1}^2$ and $|\mathcal{S}''| = 3s_{k-1}^4$, and thus $|\mathcal{S}' \cup \mathcal{S}''| = 3s_{k-1}^2(s_{k-1}^2 + 1)$. Therefore any solution to the following set of recurrences satisfies the theorem:

$$s_0 \ge 1, \quad t_0 \ge 1, \quad s_k \ge 3s_{k-1}^2(s_{k-1}^2+1), \quad t_k \ge 3s_{k-1} \cdot t_{k-1}$$
 (7.1)

We claim that

$$s_k = 6^{(4^k - 1)/3}$$
 and $t_k = 3^k \cdot 6^{(4^k - 3k - 1)/9}$

is a solution to (7.1). Observe that $s_0 = 1$ and $t_0 = 1$. Now

$$3s_{k-1}^2(s_{k-1}^2+1) \leq 6s_{k-1}^4$$

and

$$6(6^{(4^{k-1}-1)/3})^4 = 6^{1+4(4^{k-1}-1)/3} = 6^{(4^k-1)/3} = s_k \ .$$

Thus the recurrence for s_k is satisfied. Now

$$3 \cdot s_{k-1} \cdot t_{k-1} = 3 \cdot 6^{(4^{k-1}-1)/3} \cdot 3^{k-1} \cdot 6^{(4^{k-1}-3(k-1)-1)/9}$$

= $3^k \cdot 6^{(3 \cdot 4^{k-1}-3+4^{k-1}-3k+3-1)/9}$
= $3^k \cdot 6^{(4^k-3k-1)/9}$
= t_k .

Thus the recurrence for t_k is satisfied. This completes the proof.

In the proof of Theorem 7.3 we have made little effort to reduce the bound on t_k , beyond that it is a doubly exponential function of k. In [35] we describe a number of refinements that result in improved bounds on t_k . One such refinement uses strict k-trees. From an algorithmic point of view, the disadvantage of using strict k-trees is that at each recursive step, extra edges must be added to enlarge the graph from a partial strict k-tree into a strict k-tree, whereas when using (non-strict) k-trees, extra edges need only be added at the beginning of the algorithm.

For small values of k, much-improved results can be obtained. For example, we prove that every series-parallel graph (that is, with tree-width at most two) has an



FIG. 7.2. Wrapped track layout in Theorem 7.3.

18-track layout [35], whereas $t_2 = 54$. This bound has recently been improved to 15 by Di Giacomo *et al.* [26]. Their method is based on Theorems 6.1 and 7.3, and in the general case, still gives a doubly exponential upper bound on the track-number of graphs with tree-width k. For other particular classes of graphs, Di Giacomo and Meijer [25, 28] recently improved the constants in our results.

Our doubly exponential upper bound is probably not best possible. Di Giacomo *et al.* [26] constructed graphs with tree-width k and track-number at least 2k + 1. The following construction establishes a quadratic lower bound. It is similar to a graph due to Albertson [3], which gives a tight lower bound on the star chromatic number of graphs with tree-width k.

THEOREM 7.4. For all $k \ge 0$, there is a graph G_k with tree-width at most k and track-number $\operatorname{tn}(G_k) = \frac{1}{2}(k+1)(k+2)$.

Proof. Let $G_0 = K_1$. Obviously G_0 has tree-width 0. Construct G_k from G_{k-1} as follows. Start with a k-clique $\{v_1, v_2, \ldots, v_k\}$. Let $n = 2(\frac{1}{2}(k+1)(k+2)-1-k)+1$. Add n vertices $\{w_1, w_2, \ldots, w_n\}$ each adjacent to every v_i . Let H_1, H_2, \ldots, H_n be copies of G_{k-1} . For all $1 \leq j \leq n$, add an edge between w_j and each vertex of H_j . It is easily seen that from a tree decomposition of G_{k-1} of width k-1, we can construct a tree decomposition of G_k has tree-width at most k.

To prove that $\operatorname{tn}(G_k) \geq \frac{1}{2}(k+1)(k+2)$, we proceed by induction on $k \geq 0$. Obviously $\operatorname{tn}(G_0) = 1$. Suppose that $\operatorname{tn}(G_{k-1}) \geq \frac{1}{2}k(k+1)$, but $\operatorname{tn}(G_k) \leq \frac{1}{2}(k+1)(k+2) - 1$. Since $\{v_1, v_2, \ldots, v_k\}$ is a clique, we can assume that v_i is in track i. Since each vertex w_j is adjacent to each v_i , no w_j is in tracks $\{1, 2, \ldots, k\}$. There are $\frac{1}{2}(k+1)(k+2) - 1 - k$ remaining tracks. Since n is more than twice this number, there are at least three w_j vertices in a single track. Without loss of generality, $w_1 < w_2 < w_3$ in track k+1. No vertex x of H_2 is in track $i \in \{1, 2, \ldots, k\}$, as otherwise xw_2 would form an X-crossing with v_iw_1 or v_iw_3 . No vertex x of H_2 is in track k+1. Thus all



FIG. 7.3. The graph G_k .

the vertices of H_2 are in tracks $\{k+2, k+3, \ldots, \frac{1}{2}(k+1)(k+2) - 1\}$. There are $\frac{1}{2}(k+1)(k+2) - 1 - (k+1) = \frac{1}{2}k(k+1) - 1$ such tracks. This contradicts the assumption that $\operatorname{tn}(G_{k-1}) \geq \frac{1}{2}k(k+1)$. Therefore $\operatorname{tn}(G_k) \geq \frac{1}{2}(k+1)(k+2)$.

It remains to prove that $\operatorname{tn}(G_k) \leq \frac{1}{2}(k+1)(k+2)$. Suppose we have a $\frac{1}{2}k(k+1)$ -track layout of G_{k-1} . Thus each H_j has a $\frac{1}{2}k(k+1)$ -track layout. Put each vertex v_i of G_k in track *i*. Put the vertices $\{w_1, w_2, \ldots, w_n\}$ in track k+1 in this order. Put the track layout of each H_j in tracks $k+2, k+3, \ldots, \frac{1}{2}(k+1)(k+2)$, such that the vertices of H_j precede the vertices of H_{j+1} . Clearly there are no X-crossings.

Also note that Theorem 7.4 (for $k \ge 2$) can be extended using the proof technique of Lemma 2.3 to give the same lower bound for improper track layouts.

8. Open Problems.

1. (In the conference version of their paper) Felsner [42] asked whether every planar graph has a three-dimensional drawing with $\mathcal{O}(n)$ volume? By Theorem 2.9, this question has an affirmative answer if every planar graph has $\mathcal{O}(1)$ track-number. Whether every planar graph has $\mathcal{O}(1)$ track-number is an open problem due to H. de Fraysseix [private communication, 2000], and by Theorem 2.6, is equivalent to the following question.

2. Heath *et al.* [58, 54] asked whether every planar graph has $\mathcal{O}(1)$ queuenumber? The best known upper bound on the queue-number of a planar graph is $\mathcal{O}(\sqrt{n})$. In general, Dujmović and Wood [38] proved that every *m*-edge graph has queue-number at most $e\sqrt{m}$, where *e* is the base of the natural logarithm.

3. Heath *et al.* [58, 54] asked whether stack-number is bounded by queue-number (and vice-versa)? Note that there is a family of graphs \mathcal{G} with $\mathsf{sn}(G) \in \Omega(3^{\Omega(\mathsf{qn}(G))-\epsilon})$, for all $G \in \mathcal{G}$ [54].

4. Is the queue-number of a graph bounded by a polynomial (or even singly exponential) function of its tree-width?

Acknowledgements. The authors are grateful for stimulating discussions with Prosenjit Bose, Jurek Czyzowicz, Hubert de Fraysseix, Stefan Langerman, Giuseppe Liotta, Patrice Ossona de Mendez, and Matthew Suderman. Thanks to an anonymous referee for many helpful comments.

REFERENCES

- M. O. ALBERTSON AND D. M. BERMAN, Every planar graph has an acyclic 7-coloring, Israel J. Math., 28 (1977), pp. 169-174.
- [2] , An acyclic analogue to Heawood's theorem, Glasgow Math. J., 19 (1978), pp. 163-166.

- [3] M. O. ALBERTSON, G. G. CHAPPELL, H. A. KIERSTEAD, A. KÜNDGEN, AND R. RAMAMURTHI, Coloring with no 2-colored P₄'s, Electron. J. Combin., 11 #R26 (2004).
- [4] N. ALON AND T. H. MARSHALL, Homomorphisms of edge-colored graphs and Coxeter groups, J. Algebraic Combin., 8 (1998), pp. 5-13.
- [5] N. ALON, C. MCDIARMID, AND B. REED, Acyclic coloring of graphs, Random Structures Algorithms, 2 (1991), pp. 277-288.
- [6] N. ALON, B. MOHAR, AND D. P. SANDERS, On acyclic colorings of graphs on surfaces, Israel J. Math., 94 (1996), pp. 273-283.
- [7] S. ARNBORG AND A. PROSKUROWSKI, Linear time algorithms for NP-hard problems restricted to partial k-trees, Discrete Appl. Math., 23 (1989), pp. 11-24.
- [8] F. BERNHART AND P. C. KAINEN, The book thickness of a graph, J. Combin. Theory Ser. B, 27 (1979), pp. 320-331.
- S. N. BHATT, F. R. K. CHUNG, F. T. LEIGHTON, AND A. L. ROSENBERG, Scheduling tree-dags using FIFO queues: A control-memory trade-off, J. Parallel Distrib. Comput., 33 (1996), pp. 55-68.
- [10] H. L. BODLAENDER, A partial k-arboretum of graphs with bounded treewidth, Theoret. Comput. Sci., 209 (1998), pp. 1-45.
- [11] H. L. BODLAENDER AND J. ENGELFRIET, Domino treewidth, J. Algorithms, 24 (1997), pp. 94– 123.
- [12] O. V. BORODIN, On acyclic colorings of planar graphs, Discrete Math., 25 (1979), pp. 211-236.
- [13] O. V. BORODIN, A. V. KOSTOCHKA, A. RASPAUD, AND É. SOPENA, Acyclic colouring of 1-planar graphs, Discrete Appl. Math., 114 (2001), pp. 29-41.
- [14] P. BOSE, J. CZYZOWICZ, P. MORIN, AND D. R. WOOD, The maximum number of edges in a three-dimensional grid-drawing, J. Graph Algorithms Appl., (to appear).
- [15] F. J. BRANDENBURG, ed., Proc. International Symp. on Graph Drawing (GD '95), vol. 1027 of Lecture Notes in Comput. Sci., Springer, 1996.
- [16] I. BRUß AND A. FRICK, Fast interactive 3-D graph visualization, in Brandenburg [15], pp. 99-110.
- [17] T. CALAMONERI AND A. STERBINI, 3D straight-line grid drawing of 4-colorable graphs, Inform. Process. Lett., 63 (1997), pp. 97-102.
- [18] K. CHILAKAMARRI, N. DEAN, AND M. LITTMAN, Three-dimensional Tutte embedding, in Proc. 26th Southeastern International Conf. on Combinatorics, Graph Theory and Computing, vol. 107 of Cong. Numer., 1995, pp. 129–140.
- [19] M. CHROBAK, M. GOODRICH, AND R. TAMASSIA, Convex drawings of graphs in two and three dimensions, in Proc. 12th Annual ACM Symp. on Comput. Geom., 1996, pp. 319-328.
- [20] F. R. K. CHUNG, F. T. LEIGHTON, AND A. L. ROSENBERG, Embedding graphs in books: a layout problem with applications to VLSI design, SIAM J. Algebraic Discrete Methods, 8 (1987), pp. 33-58.
- [21] R. F. COHEN, P. EADES, T. LIN, AND F. RUSKEY, Three-dimensional graph drawing, Algorithmica, 17 (1996), pp. 199-208.
- [22] I. F. CRUZ AND J. P. TWAROG, 3D graph drawing with simulated annealing, in Brandenburg [15], pp. 162-165.
- [23] H. DE FRAYSSEIX, J. PACH, AND R. POLLACK, How to draw a planar graph on a grid, Combinatorica, 10 (1990), pp. 41-51.
- [24] G. DI BATTISTA, P. EADES, R. TAMASSIA, AND I. G. TOLLIS, Graph Drawing: Algorithms for the Visualization of Graphs, Prentice-Hall, 1999.
- [25] E. DI GIACOMO, Drawing series-parallel graphs on restricted integer 3D grids, in Liotta [67], pp. 238-246.
- [26] E. DI GIACOMO, G. LIOTTA, AND H. MEIJER, 3D straight-line drawings of k-tree, Tech. Report 2003-473, School of Computing, Queens's University, Kingston, Canada, 2003.
- [27] E. DI GIACOMO, G. LIOTTA, AND S. WISMATH, Drawing series-parallel graphs on a box, in Proc. 14th Canadian Conf. on Computational Geometry (CCCG '02), The University of Lethbridge, Canada, 2002, pp. 149–153.
- [28] E. DI GIACOMO AND H. MEIJER, Track drawings of graphs with constant queue number, in Liotta [67], pp. 214-225.
- [29] J. DÍAZ, J. PETIT, AND M. SERNA, A survey of graph layout problems, ACM Comput. Surveys, 34 (2002), pp. 313-356.
- [30] R. P. DILWORTH, A decomposition theorem for partially ordered sets, Ann. of Math. (2), 51 (1950), pp. 161-166.
- [31] G. DING AND B. OPOROWSKI, Some results on tree decomposition of graphs, J. Graph Theory, 20 (1995), pp. 481-499.
- [32] —, On tree-partitions of graphs, Discrete Math., 149 (1996), pp. 45-58.

- [33] V. DUJMOVIĆ, M. FELLOWS, M. HALLETT, M. KITCHING, G. LIOTTA, C. MCCARTIN, N. NISHIMURA, P. RAGDE, F. ROSEMAND, M. SUDERMAN, S. WHITESIDES, AND D. R. WOOD, On the parameterized complexity of layered graph drawing, in Proc. 5th Annual European Symp. on Algorithms (ESA '01), F. Meyer auf der Heide, ed., vol. 2161 of Lecture Notes in Comput. Sci., Springer, 2001, pp. 488-499.
- [34] V. DUJMOVIĆ, P. MORIN, AND D. R. WOOD, Path-width and three-dimensional straight-line grid drawings of graphs, in Proc. 10th International Symp. on Graph Drawing (GD '02), M. T. Goodrich and S. G. Kobourov, eds., vol. 2528 of Lecture Notes in Comput. Sci., Springer, 2002, pp. 42-53.
- [35] V. DUJMOVIĆ AND D. R. WOOD, Tree-partitions of k-trees with applications in graph layout, Tech. Report TR-2002-03, School of Computer Science, Carleton University, Ottawa, Canada, 2002.
- [36] _____, Tree-partitions of k-trees with applications in graph layout, in Proc. 29th Workshop on Graph Theoretic Concepts in Computer Science (WG'03), H. Bodlaender, ed., vol. 2880 of Lecture Notes in Comput. Sci., Springer, 2003, pp. 205-217.
- [37] —, Three-dimensional grid drawings with sub-quadratic volume, in Towards a Theory of Geometric Graphs, J. Pach, ed., vol. 342 of Contemporary Mathematics, Amer. Math. Soc., 2004, pp. 55-66.
- [38] _____, On linear layouts of graphs, Discrete Math. Theor. Comput. Sci., (to appear).
- [39] P. EADES AND P. GARVAN, Drawing stressed planar graphs in three dimensions, in Brandenburg [15], pp. 212-223.
- [40] P. ERDŐS, Appendix, in K. F. ROTH, On a problem of Heilbronn, J. London Math. Soc., 26 (1951), pp. 198-204.
- [41] S. EVEN AND A. ITAI, Queues, stacks, and graphs, in Proc. International Symp. on Theory of Machines and Computations, Z. Kohavi and A. Paz, eds., Academic Press, 1971, pp. 71–86.
- [42] S. FELSNER, G. LIOTTA, AND S. WISMATH, Straight-line drawings on restricted integer grids in two and three dimensions, J. Graph Algorithms Appl., 7 (2003), pp. 363-398.
- [43] G. FERTIN, A. RASPAUD, AND B. REED, On star coloring of graphs, in Proc. 27th International Workshop on Graph-Theoretic Concepts in Computer Science (WG '01), A. Branstädt and V. B. Le, eds., vol. 2204 of Lecture Notes in Comput. Sci., Springer, 2001, pp. 140-153.
- [44] D. R. FULKERSON AND O. A. GROSS, Incidence matrices and interval graphs, Pacific J. Math., 15 (1965), pp. 835-855.
- [45] J. L. GANLEY AND L. S. HEATH, The pagenumber of k-trees is O(k), Discrete Appl. Math., 109 (2001), pp. 215-221.
- [46] M. R. GAREY, D. S. JOHNSON, G. L. MILLER, AND C. H. PAPADIMITRIOU, The complexity of coloring circular arcs and chords, SIAM J. Algebraic Discrete Methods, 1 (1980), pp. 216– 227.
- [47] A. GARG, R. TAMASSIA, AND P. VOCCA, Drawing with colors, in Proc. 4th Annual European Symp. on Algorithms (ESA '96), J. Diaz and M. Serna, eds., vol. 1136 of Lecture Notes in Comput. Sci., Springer, 1996, pp. 12–26.
- [48] M. C. GOLUMBIC, Algorithmic graph theory and perfect graphs, Academic Press, 1980.
- [49] B. GRÜNBAUM, Acyclic colorings of planar graphs, Israel J. Math., 14 (1973), pp. 390-408.
- [50] R. HALIN, S-functions for graphs, J. Geometry, 8 (1976), pp. 171-186.
- [51] R. HALIN, Tree-partitions of infinite graphs, Discrete Math., 97 (1991), pp. 203-217.
- [52] F. HARARY AND A. SCHWENK, A new crossing number for bipartite graphs, Utilitas Math., 1 (1972), pp. 203-209.
- [53] T. HASUNUMA, Laying out iterated line digraphs using queues, in Liotta [67], pp. 202-213.
- [54] L. S. HEATH, F. T. LEIGHTON, AND A. L. ROSENBERG, Comparing queues and stacks as mechanisms for laying out graphs, SIAM J. Discrete Math., 5 (1992), pp. 398-412.
- [55] L. S. HEATH AND S. V. PEMMARAJU, Stack and queue layouts of posets, SIAM J. Discrete Math., 10 (1997), pp. 599-625.
- [56] —, Stack and queue layouts of directed acyclic graphs. II, SIAM J. Comput., 28 (1999), pp. 1588-1626.
- [57] L. S. HEATH, S. V. PEMMARAJU, AND A. N. TRENK, Stack and queue layouts of directed acyclic graphs. I, SIAM J. Comput., 28 (1999), pp. 1510-1539.
- [58] L. S. HEATH AND A. L. ROSENBERG, Laying out graphs using queues, SIAM J. Comput., 21 (1992), pp. 927-958.
- [59] P. HLINĚNÝ, Crossing-number critical graphs have bounded path-width, J. Combin. Theory Ser. B, 88 (2003), pp. 347-367.
- [60] S.-H. HONG, Drawing graphs symmetrically in three dimensions, in Mutzel et al. [70], pp. 189– 204.
- [61] S.-H. HONG AND P. EADES, An algorithm for finding three dimensional symmetry in series par-

allel digraphs, in Proc. 11th International Conf. on Algorithms and Computation (ISAAC '00), D. Lee and S.-H. Teng, eds., vol. 1969 of Lecture Notes in Comput. Sci., Springer, 2000, pp. 266-277.

[62] , Drawing trees symmetrically in three dimensions, Algorithmica, 36 (2003), pp. 153-178.

- [63] S.-H. HONG, P. EADES, A. QUIGLEY, AND S.-H. LEE, Drawing algorithms for series-parallel digraphs in two and three dimensions, in Proc. 6th International Symp. on Graph Drawing (GD '98), S. Whitesides, ed., vol. 1547 of Lecture Notes in Comput. Sci., Springer, 1998, pp. 198-209.
- [64] M. KAUFMANN AND D. WAGNER, eds., Drawing Graphs: Methods and Models, vol. 2025 of Lecture Notes in Comput. Sci., Springer, 2001.
- [65] A. V. KOSTOČKA, Acyclic 6-coloring of planar graphs, Diskret. Analiz, 28 (1976), pp. 40-54.
- [66] F. T. LEIGHTON AND A. L. ROSENBERG, Three-dimensional circuit layouts, SIAM J. Comput., 15 (1986), pp. 793-813.
- [67] G. LIOTTA, ed., Proc. 11th International Symp. on Graph Drawing (GD '03), vol. 2912 of Lecture Notes in Comput. Sci., Springer, 2004.
- [68] J. MITCHEM, Every planar graph has an acyclic 8-coloring, Duke Math. J., 41 (1974), pp. 177-181.
- [69] B. MONIEN, F. RAMME, AND H. SALMEN, A parallel simulated annealing algorithm for generating 3D layouts of undirected graphs, in Brandenburg [15], pp. 396-408.
- [70] P. MUTZEL, M. JÜNGER, AND S. LEIPERT, eds., Proc. 9th International Symp. on Graph Drawing (GD '01), vol. 2265 of Lecture Notes in Comput. Sci., Springer, 2002.
- [71] J. NEŠETŘIL AND P. OSSONA DE MENDEZ, Colorings and homomorphisms of minor closed classes, in Discrete and Computational Geometry, The Goodman-Pollack Festschrift, B. Aronov, S. Basu, J. Pach, and M. Sharir, eds., vol. 25 of Algorithms and Combinatorics, Springer, 2003, pp. 651-664.
- [72] J. NEŠETŘIL AND A. RASPAUD, Colored homomorphisms of colored mixed graphs, J. Combin. Theory Ser. B, 80 (2000), pp. 147-155.
- [73] L. T. OLLMANN, On the book thicknesses of various graphs, in Proc. 4th Southeastern Conference on Combinatorics, Graph Theory and Computing, F. Hoffman, R. B. Levow, and R. S. D. Thomas, eds., vol. VIII of Congressus Numerantium, 1973, p. 459.
- [74] D. I. OSTRY, Some three-dimensional graph drawing algorithms, master's thesis, Department of Computer Science and Software Engineering, The University of Newcastle, Australia, 1996.
- [75] J. PACH, T. THIELE, AND G. TÓTH, Three-dimensional grid drawings of graphs, in Advances in discrete and computational geometry, B. Chazelle, J. E. Goodman, and R. Pollack, eds., vol. 223 of Contemporary Mathematics, Amer. Math. Soc., 1999, pp. 251-255.
- [76] S. V. PEMMARAJU, Exploring the Powers of Stacks and Queues via Graph Layouts, PhD thesis, Virginia Polytechnic Institute and State University, U.S.A., 1992.
- [77] Z. PENG, Drawing graphs of bounded treewidth/pathwidth, master's thesis, Department of Computer Science, University of Auckland, New Zealand, 2001.
- [78] T. PORANEN, A new algorithm for drawing series-parallel digraphs in 3D, Tech. Report A-2000-16, Dept. of Computer and Information Sciences, University of Tampere, Finland, 2000.
- [79] B. A. REED, Algorithmic aspects of tree width, in Recent Advances in Algorithms and Combinatorics, B. A. Reed and C. L. Sales, eds., Springer, 2003, pp. 85-107.
- [80] S. RENGARAJAN AND C. E. VENI MADHAVAN, Stack and queue number of 2-trees, in Proc. 1st Annual International Conf. on Computing and Combinatorics (COCOON '95), D. Ding-Zhu and L. Ming, eds., vol. 959 of Lecture Notes in Comput. Sci., Springer, 1995, pp. 203-212.
- [81] N. ROBERTSON AND P. D. SEYMOUR, Graph minors. II. Algorithmic aspects of tree-width, J. Algorithms, 7 (1986), pp. 309-322.
- [82] D. J. ROSE, R. E. TARJAN, AND G. S. LEUKER, Algorithmic aspects of vertex elimination on graphs, SIAM J. Comput., 5 (1976), pp. 266-283.
- [83] P. SCHEFFLER, Die Baumweite von Graphen als ein Maß für die Kompliziertheit algorithmischer Probleme, PhD thesis, Akademie der Wissenschaften der DDR, Berlin, Germany, 1989.
- [84] W. SCHNYDER, Planar graphs and poset dimension, Order, 5 (1989), pp. 323-343.
- [85] D. SEESE, Tree-partite graphs and the complexity of algorithms, in Proc. International Conf. on Fundamentals of Computation Theory, L. Budach, ed., vol. 199 of Lecture Notes in Comput. Sci., Springer, 1985, pp. 412-421.
- [86] F. SHAHROKHI AND W. SHI, On crossing sets, disjoint sets, and pagenumber, J. Algorithms, 34 (2000), pp. 40-53.

- [87] E. STEINITZ, Polyeder und Raumeinteilungen, Encyclopädie der Mathematischen Wissenschaften, 3AB12 (1922), pp. 1-139.
- [88] R. TARJAN, Sorting using networks of queues and stacks, J. Assoc. Comput. Mach., 19 (1972), pp. 341-346.
- [89] M. THORUP, All structured programs have small tree-width and good register allocation, Information and Computation, 142 (1998), pp. 159-181.
- [90] C. WARE AND G. FRANCK, Viewing a graph in a virtual reality display is three times as good as a 2D diagram, in Proc. IEEE Symp. Visual Languages (VL '94), A. L. Ambler and T. D. Kimura, eds., IEEE, 1994, pp. 182-183.
- [91] —____, Evaluating stereo and motion cues for visualizing information nets in three dimensions, ACM Trans. Graphics, 15 (1996), pp. 121–140.
- [92] C. WARE, D. HUI, AND G. FRANCK, Visualizing object oriented software in three dimensions, in Proc. IBM Centre for Advanced Studies Conf. (CASCON '93), 1993, pp. 1-11.
- [93] D. R. WOOD, Queue layouts, tree-width, and three-dimensional graph drawing, in Proc. 22nd Foundations of Software Technology and Theoretical Computer Science (FST TCS '02), M. Agrawal and A. Seth, eds., vol. 2556 of Lecture Notes in Comput. Sci., Springer, 2002, pp. 348-359.
- [94] M. YANNAKAKIS, Embedding planar graphs in four pages, J. Comput. System Sci., 38 (1986), pp. 36-67.