

A "Schrödinger Cat" Superposition State of an Atom

C. Monroe, D.M. Meekhof, B.E. King, and D.J. Wineland

Time and Frequency Division, MS 847, National Institute of Standards and Technology

Boulder, CO 80303

A "Schrödinger cat"-like state of matter has been generated at the single atom level. A trapped ${}^9\text{Be}^+$ ion was laser-cooled to the zero-point energy and then prepared in a superposition of spatially separated coherent harmonic oscillator states. We create this state by applying a sequence of laser pulses, which entangle internal (electronic) and external (motional) states of the ion. We verify the Schrödinger cat superposition by detecting the quantum mechanical interference between the localized wavepackets. This mesoscopic system may provide insight into the fuzzy boundary between classical and quantum worlds by allowing controlled studies of quantum measurement and quantum decoherence.

Work of U.S. Government; not subject to U.S. copyright.

Quantum mechanics allows the possibility of preparing physical systems in superposition states, or states which are "smeared" between two or more distinct values. This curious principle of quantum mechanics [1] has been extremely successful at describing physical behavior in the microscopic world - from interactions of atoms with photons to interactions at the sub-nuclear level. But what happens when we extend the quantum superposition principle to macroscopic systems conventionally described by classical physics? Here, superpositions introduce a great amount of conceptual difficulty, as pointed out in 1935 by the celebrated Einstein-Podolsky-Rosen [2] and Schrödinger cat [3] paradoxes. For example, in Schrödinger's thought experiment [3], an unfortunate cat is placed in a quantum superposition of being dead and alive (correlated with a single radioactive atom which has and has not decayed). The state of the system can be represented by the entangled quantum mechanical wave-function,

$$\Psi = \frac{|\odot\rangle|\uparrow\rangle + |\otimes\rangle|\downarrow\rangle}{\sqrt{2}} \quad (1)$$

where $|\odot\rangle$ and $|\otimes\rangle$ refer to the states of a live and dead cat, and $|\downarrow\rangle$ and $|\uparrow\rangle$ refer to the internal states of an atom which has and has not radioactively decayed. This situation defies our sense of reality, because we only observe live or dead cats, and we expect that cats are either alive or dead independent of our observation [4]. Schrödinger's cat paradox is a classic illustration of the conflict between the existence of quantum superpositions and our real world experience of observation and measurement.

Although superposition states such as Schrödinger's cat do not appear in the macroscopic world, there is great interest in the realization of "Schrödinger cat-like" states in mesoscopic systems, or systems that have both macroscopic and microscopic features. In this context, the "cat" is generalized to represent a physical system whose attributes are normally associated with classical concepts, such as the distinguishable position of a particle (instead of the state of livelihood of a real cat). In this article, we report the creation of a mesoscopic "Schrödinger cat" state at the single atom level - an atom is prepared in a quantum superposition of two spatially separated but localized positions. In analogy to Schrödinger's original proposition given by Eq. 1, we create the state:

$$\Psi = \frac{|x_1\rangle|\uparrow\rangle + |x_2\rangle|\downarrow\rangle}{\sqrt{2}} \quad (2)$$

where $|x_1\rangle$ and $|x_2\rangle$ denote classical-like wavepacket states corresponding to separated spatial positions of the atom, and $|\downarrow\rangle$ and $|\uparrow\rangle$ refer to distinct internal electronic quantum states of the atom [5]. The wavepackets are separated by a mesoscopic distance of more than 80 nm, which is large compared to the size of the individual wavepackets (≈ 7 nm) as well as the atomic dimension (≈ 0.1 nm).

Mesoscopic Schrödinger cats may provide an interesting testing ground for the controversial theory of quantum measurement [6]. At the core of this historical issue is the question of the universality of quantum mechanics. The "Copenhagen interpretation" of Bohr [7] and Heisenberg [8] holds that the measuring apparatus always involves classical concepts, thus forcing a seemingly arbitrary division between the quantum and classical worlds. Einstein [2] on the other hand argued that for quantum mechanics to be complete, it should describe physical behavior at all scales. One practical approach toward resolving this controversy is the introduction of quantum decoherence, or the environmentally-induced reduction of quantum superpositions into statistical mixtures and classical behavior [9]. Decoherence is commonly interpreted as a way of quantifying the elusive boundary between classical and quantum worlds, and almost always precludes the existence of macroscopic Schrödinger cat states, except at extremely short time scales [9,10]. The creation of mesoscopic Schrödinger cat states may allow controlled studies of quantum decoherence and the quantum/classical boundary. We note that quantum decoherence has received much interest lately due to its importance in proposals for quantum computation [11] and quantum cryptography [12].

Macroscopic superposition states of matter have been realized for electron [13], neutron [14], and atom [15] beamsplitters, where these particles are split into superpositions of separated paths. The matter wavepackets in these experiments spread in time, since the particles are unbound. Spatially separated superpositions of electrons within atoms have been demonstrated by exciting electrons to Rydberg states with pulsed lasers [16]. Here, the electron wavepacket is also dispersive due to its anharmonic binding potential. There have been related proposals for the creation of

macroscopic superposition states of vibration in molecules or crystals [17] and of electrical currents flowing in superconducting rings [18].

The appeal of creating a Schrödinger cat state in a harmonic oscillator is that wavepacket dispersion can be negligible. The simple time evolution of a coherent harmonic oscillator wavepacket preserves the separation of the superposition, and aids visualization and interpretation of experiments. There have been several proposals to create mesoscopic Schrödinger cat states in a single mode of the electromagnetic field, which is formally equivalent to a harmonic oscillator. For instance, these states are expected to evolve from the amplitude dispersion of a laser beam propagating in an anharmonic Kerr medium [19]. In cavity-quantum-electrodynamics, these states are predicted to emerge by driving a coherent state with a Jaynes-Cummings interaction to the point of collapse [20], by continuously pumping a single cavity mode with polarized two-level atoms [21], or by realizing a dispersive interaction between a single atom and a single cavity mode [22,23]. The creation of Schrödinger cat states of a single harmonically bound atom has been proposed by driving the atom with a strong laser field and relying on a measurement to project the desired superposition state [24], or by optically pumping the atom to a "dark" state with multiple laser beams [25].

EXPERIMENTAL APPROACH

In the present work, we create a Schrödinger cat state of the harmonic oscillator by forming a superposition of two coherent state wavepackets of a single trapped atom with a sequence of laser pulses. Each wavepacket is correlated with a particular internal state of the atom. To analyze this state we apply an additional laser pulse to couple the internal states and we measure the resulting interference of the distinct wavepackets [26,27]. The key features of our approach are that (i) we control the harmonic motion of the trapped atom to a high degree by exciting the motion from initial zero-point wavepackets to coherent state wavepackets of well-defined amplitude and phase; (ii) we do not rely on a conditional measurement to project out the desired Schrödinger cat state; and (iii) wavepacket dispersion of the atomic motion is negligible.

The experimental apparatus is described elsewhere [28,29]. A single ${}^9\text{Be}^+$ ion is confined in a coaxial-resonator rf-ion trap [28] that provides harmonic oscillation frequencies of $(\omega_x, \omega_y, \omega_z)/2\pi \approx (11.2, 18.2, 29.8)$ MHz along the principal axes of the trap. We laser-cool the ion to the quantum

ground state of motion [29], and then coherently manipulate its internal (electronic) and external (motional) state by applying pairs of off-resonant laser beams, which drive two-photon stimulated Raman transitions [29,30]. As shown in Fig. 1a, the two internal states of interest are the stable $^2S_{1/2}(F = 2, m_F = -2)$ and $^2S_{1/2}(F = 1, m_F = -1)$ hyperfine ground states (denoted by $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ respectively), separated in frequency by $\omega_{\text{HF}}/2\pi \approx 1.250$ GHz. Here, F and m_F are quantum numbers representing the total internal angular momentum of the atom and its projection along a quantization axis. The Raman beams are detuned by $\Delta \approx -12$ GHz from the $^2P_{1/2}(F = 2, m_F = -2)$ excited state, which acts as the virtual level providing the Raman coupling. The external motional states are characterized by the quantized vibrational harmonic oscillator states $|n\rangle_e$ in the x -dimension, separated in frequency by $\omega_x/2\pi \approx 11.2$ MHz.

When we tune the Raman beam difference frequency near ω_{HF} and apply the “carrier beams” A and B of Fig. 1, the ion experiences a coherent Rabi oscillation between the internal states $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$. By adjusting the exposure time of the carrier beams, we can for example “flip” the internal state (a π -pulse, or $1/2$ of a Rabi cycle), or “split/recombine” the internal state (a $\pi/2$ -pulse, or $1/4$ of a Rabi cycle). Transitions on the carrier do not significantly affect the state of motion, because beams A and B are copropagating. When we tune the Raman beam difference frequency near ω_x and apply the “displacement” beams B and C of Fig. 1, the effect is formally equivalent to applying the displacement operator to the state of motion [30]. Alternatively, the displacement beams can be thought of as producing a “walking wave” pattern whose time-dependent dipole force resonantly excites the harmonic motion [31]. This force promotes an initial zero-point state of motion $|0\rangle_e$ to a coherent state $|\beta\rangle_e = e^{-|\beta|^2/2} \sum_n \beta^n / (n!)^{1/2} |n\rangle_e$ [32], where $\beta = \alpha e^{i\theta}$ is a dimensionless complex number which represents the amplitude and phase of the motion in the harmonic potential [33]. The probability distribution of vibrational levels in a coherent state is Poissonian with mean number of vibrational quanta $\langle n \rangle = \alpha^2$. The coherent state of motion is much like classical motion in a harmonic potential with amplitude $2\alpha x_0$, where $x_0 = (\hbar/2m\omega_x)^{1/2} = 7.1(1)$ nm is the root-mean-square Gaussian size of the oscillating wavepacket, m is the mass of the ion, and \hbar is Planck's constant divided by 2π .

The polarizations of the three Raman beams A, B, and C produce π , σ^+/σ^- , and σ^- couplings, respectively with respect to a quantization axis defined by an applied 0.20 mT magnetic field, as indicated in Fig. 1b. As a result, the displacement beams (B and C) affect only the motional state

correlated with the $|\uparrow\rangle_i$ internal state, since the σ^- -polarized beam C cannot couple the internal state $|\downarrow\rangle_i$ to any virtual ${}^2P_{1/2}$ states [34]. This selectivity of the displacement force provides quantum entanglement of the internal state with the external motional state. Although the motional state can be thought of as nearly classical, its entanglement with the internal atomic quantum levels precludes any type of semi-classical analysis.

Each Raman beam contains ≈ 1 mW of power at ≈ 313 nm. This results in a two-photon Rabi frequency of $\Omega/2\pi \approx 250$ kHz for the copropagating Raman carrier beams A and B, or a π -pulse exposure time of about 1 μ s. We apply the displacement Raman beams (B and C) to the ion in directions such that their wavevector difference $\delta\mathbf{k}$ points nearly along the x-axis of the trap. Motion in the y or z dimensions is therefore highly insensitive to the displacement beams. When we apply the displacement beams to a zero-point wavepacket (correlated with the $|\uparrow\rangle_i$ state) for time τ , we expect to create a coherent state of amplitude $\alpha = \eta\Omega_d\tau$. Here, $\eta = 0.205(5)$ is the Lamb-Dicke parameter [30] and $\Omega_d/2\pi \approx 300$ kHz is the coupling strength of the displacement beams (numbers in parentheses are the standard errors in the last digit). After each preparation cycle (described below), we detect which internal state ($|\downarrow\rangle_i$ or $|\uparrow\rangle_i$) the atom occupies independent of its state of motion. This is accomplished by applying a few microwatts of σ^- -polarized light ("detection" beam D of Fig. 1a) resonant with the cycling $|\downarrow\rangle_i \rightarrow {}^2P_{3/2}(F=3, m_F=-3)$ transition (radiative linewidth $\gamma/2\pi \approx 19.4$ MHz at $\lambda \approx 313$ nm) and observing the resulting ion fluorescence. Because this radiation does not appreciably couple to the $|\uparrow\rangle_i$ state, the fluorescence reading is proportional to the probability P_\downarrow the ion is in state $|\downarrow\rangle_i$. We collect on average ≈ 1 photon per measurement cycle when the ion is in the $|\downarrow\rangle_i$ state [29].

CREATION AND DETECTION OF THE SCHRÖDINGER CAT STATE

The ion is first laser-cooled so that the $|\downarrow\rangle_i|n_x=0\rangle_e$ state is occupied $\approx 95\%$ of the time as described in [29]. We then apply five sequential pulses of the Raman beams (the evolving state of the system is summarized in Table I and Fig. 2): (1) A $\pi/2$ -pulse on the carrier splits the wavefunction into an equal superposition of states $|\downarrow\rangle_i|0\rangle_e$ and $|\uparrow\rangle_i|0\rangle_e$. (2) The displacement beams excite the motion correlated with the $|\uparrow\rangle_i$ component to a coherent state $|\alpha e^{-i\phi/2}\rangle_e$. (3) A π -pulse on the carrier swaps the internal states of the superposition. (4) The displacement beams excite the motion

correlated with the $|\uparrow\rangle_i$ component to a second coherent state $|\alpha e^{i\phi/2}\rangle_e$. (5) A final $\pi/2$ -pulse on the carrier combines the two coherent states. The relative phases of the above steps are determined by the phases of the rf difference frequencies of the Raman beams [29,30], which are easily controlled by phase-locking the rf sources.

The state created after step 4 is a superposition of two independent coherent states each correlated with an internal state of the ion, in the spirit of Schrödinger's original thought experiment (Eqs. 1 and 2). We verify this superposition by recombining the coherent wavepackets in the final step 5. This creates the entangled state:

$$|\Psi\rangle = |\downarrow\rangle_i |S_-\rangle_e - i |\uparrow\rangle_i |S_+\rangle_e, \quad \text{with}$$

$$|S_{\pm}\rangle_e \equiv \frac{|\alpha e^{-i\phi/2}\rangle_e \pm e^{i\delta} |\alpha e^{i\phi/2}\rangle_e}{2}. \quad (3)$$

For $\phi=\pi$ and $\delta=0$, the states $|S_{\pm}\rangle_e$ (when properly normalized) are known as "even" and "odd" Schrödinger cats [35].

The relative populations of $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ depend on the motional phase difference ϕ between the two coherent wavepackets, because of the quantum interference between the two coherent states contained in $|S_{\pm}\rangle_e$. We directly measure this interference by detecting the probability $P_i(\phi)$ that the ion is in the $|\downarrow\rangle_i$ internal state for a given value of ϕ . We continuously repeat the experiment - cooling, state preparation, detection - while slowly sweeping the relative coherent state motional phase ϕ . Figure 3 depicts the expected position basis wavepacket $|\langle x|S_-\rangle_e|^2$ correlated with the $|\downarrow\rangle_i$ internal state as a function of ϕ for $\delta=0$ and $\alpha=3$. The calculated wavepackets in the figure are snapshots in time, as each part of the superposition oscillates in the harmonic trap. The measured signal $P_i(\phi)$ is just the integral of the complete $|\downarrow\rangle_i$ wavepacket over space and is time-independent:

$$P_i(\phi) = \int_{-\infty}^{+\infty} |\langle x|S_-\rangle_e|^2 dx = \frac{1 - e^{-\alpha^2(1-\cos\phi)} \cos(\delta + \alpha^2 \sin\phi)}{2} \quad (4)$$

The wavepackets of the superposition are maximally separated in phase space for $\phi \approx \pm\pi$, where the

signal is approximately $1/2$ (for large α). However, as ϕ approaches zero, the wavepackets of the superposition begin to overlap, finally interfering completely at $\phi=0$. For large α , the signal $P_1(\phi)$ acquires oscillations near $\phi=0$, with the width of the central interference fringe (in ϕ -space) proportional to $1/\alpha^2$. If the two pieces of the wavepacket are not phase-coherent or if the state is a statistical mixture (δ random between preparations) instead of a coherent superposition of wavepackets, the signal would remain constant, $P_1(\phi) = 1/2$. We experimentally set the phase δ associated with the internal state superposition by blocking the displacement beams ($\alpha=0$) and measuring $P_1 = \sin^2(\delta/2)$.

SUPERPOSITIONS VERSUS MIXTURES

In Fig. 4, we display the measured $P_1(\phi)$ for $\delta=0$ and a few different values of the coherent state amplitude α , which is set by changing the duration τ of application of the displacement beams (steps 2 and 4 of Table I). The presence of the interference feature near $\phi=0$ verifies that we are producing superposition states instead of statistical mixtures, and the feature clearly narrows as α is increased. We have verified that the interference feature vanishes [$P_1(\phi) = 1/2$] when δ is randomized between preparations. In Fig. 5, we present $P_1(\phi)$ for three different values of the phase δ while fixing τ . The shape of the interference at $\phi \approx 0$ indicates the parity of the cat state at $\phi = \pm\pi$. Here, we see the transition from an even cat ($\delta \approx \pi$) to the "Yurke-Stoler" [19] cat ($\delta \approx \pi/2$) to an odd cat state ($\delta \approx 0$) correlated with the $|\downarrow\rangle_i$ state.

We extract the amplitude of the Schrödinger cat state by fitting the interference data to the parameter α appearing in Eq. 4. The extracted values of α agree with the independently measured value $\eta\Omega_d\tau$ for short displacement beam durations ($\tau \lesssim 10 \mu\text{s}$) [36]. We measure coherent state amplitudes as high as $\alpha \approx 2.97(6)$, corresponding to an average of $\langle n \rangle \approx 9$ vibrational quanta in the state of motion. This indicates a maximum spatial separation of $4\alpha x_0 = 83(3)$ nm, which is significantly larger than the single wavepacket size of $x_0 = 7.1(1)$ nm. The individual wavepackets are thus clearly separated in phase space.

For longer displacement beam durations ($\tau \gtrsim 10 \mu\text{s}$), the interference signal loses contrast, as evident in Fig. 4d. We believe this is partly due to fluctuations of the ion oscillation frequency ω_x , which causes the motional phase difference ϕ to fluctuate from measurement to measurement and

wash out narrow interference features. The measured interference signal is sensitive to fluctuations of ω_x at a time scale which is longer than the time to create the cat ($\tau_c \approx \tau \approx 10 \mu\text{s}$) but shorter than the integrated measurement time ($\approx 1 \text{ s}$ per data point in Figs. 4 and 5). The observed loss of contrast indicates a phase fluctuation of $\delta\phi \approx 0.1 \text{ rad}$, which would be consistent with a fractional ion oscillation frequency fluctuation of $\delta\omega_x/\omega_x \approx 10^{-4}$ in a $\approx 100 \text{ kHz}$ bandwidth. Anharmonicities of the trap [28] are expected to contribute to a phase dispersion of only $\approx 10^{-6} \text{ rad}$ during the creation of the cat.

DECOHERENCE

When a Schrödinger cat consisting of two separated coherent states is coupled to a thermal reservoir, the superposition decays exponentially to a statistical mixture with a rate initially proportional to α^2 , or the square of the separation of the wavepackets [9,10,27]. As the separation is made larger (more classical), the lifetime of the superposition shortens. This decoherence process underlies the reason quantum superpositions are not generally seen in the macroscopic world, and also illustrates the experimental difficulty in preparing and maintaining even mesoscopic superpositions.

In the experiment, the quantum interference signal is only sensitive to decoherence during the period of time τ_c between the generation of the two coherent states (steps 2 and 4 of Table I). This is because only the internal atomic state is detected, and once the second coherent state is produced (step 4), the internal and motional states do not interact - even if the motion equilibrates with an external reservoir. We therefore expect the interference signal (Eq. 4) to exhibit a contrast of $\exp(-\alpha^2\lambda\tau_c)$, where λ is the temperature-dependent relaxation rate to the thermal reservoir [10]. The loss of contrast we observe may involve the onset of decoherence, although it is difficult to make a quantitative comparison because we do not know the spectrum and effective temperature of the supposed reservoir. We note that we have previously measured a heating rate of $\partial\langle n \rangle/\partial t \approx 10^3/\text{s}$ [29], but because the source of this heating is not understood at the present time, it is difficult to characterize its effect on decoherence.

We are currently devoting effort to deliberately induce decoherence of the Schrödinger cat by coupling the system to “engineered” reservoirs during the interval τ_c . For instance, we can apply

a uniform stochastic electric field, whose coupling to the ion could simulate a thermal reservoir at a controllable temperature. Alternatively, we can pulse the Raman beams and controllably allow spontaneous emission to occur during the interval τ_c (similar to stimulated Raman cooling [29]). With this coupling, we can simulate thermal, zero-temperature, squeezed, and other reservoirs [37]. By monitoring the contrast of the interference signal, we should then be able to study the decoherence of the Schrödinger cat to these known reservoirs. We might also measure the effects of decoherence by mapping the complete density matrix of the Schrödinger cat state, as proposed in recent tomographic schemes [38].

We finally note that our technique for preparing Schrödinger cat superpositions of two coherent states in one dimension can easily be extended to create superpositions of more than two coherent states, and superpositions in two and three dimensions. This technique may also be useful for the creation of superposition states of the collective motion of many trapped atoms.

This work is supported by the US Office of Naval Research and the US Army Research Office. We acknowledge key assistance from W.M. Itano, D. Leibfried, J.C. Bergquist, and J. Erickson, and useful discussions with J.I. Cirac, P. Zoller, and D. Walls. We thank M. Young, S. Mechels, and D. Lee for critical comments on the manuscript

Table I. Raman beam pulse sequence for the generation of a Schrödinger cat state. The magnitude (phase) of the coherent state is controlled by the duration (phase) of the applied displacement beams in steps 2 and 4. The phases of internal state carrier operations in steps 1, 3, 5 are relative to step 5. The states created after each step do not include overall phase factors, and the phase appearing in the final state is $\delta \equiv \mu - 2\nu + \pi$.

step	function	approximate duration (μs)	phase	state created (see Fig. 2) (initial state: $ \downarrow\rangle_i 0\rangle_e$)
1.	carrier $\pi/2$ -pulse	0.5	μ	$[\downarrow\rangle_i 0\rangle_e - ie^{-i\mu} \uparrow\rangle_i 0\rangle_e]/\sqrt{2}$
2.	displacement	$\tau \approx 10.0$	$-\phi/2$	$[\downarrow\rangle_i 0\rangle_e - ie^{-i\mu} \uparrow\rangle_i \alpha e^{-i\phi/2}\rangle_e]/\sqrt{2}$
3.	carrier π -pulse	1.0	ν	$[e^{i(\nu-\mu)} \downarrow\rangle_i \alpha e^{-i\phi/2}\rangle_e + ie^{-i\nu} \uparrow\rangle_i 0\rangle_e]/\sqrt{2}$
4.	displacement	$\tau \approx 10.0$	$\phi/2$	$[e^{i(\nu-\mu)} \downarrow\rangle_i \alpha e^{-i\phi/2}\rangle_e + ie^{-i\nu} \uparrow\rangle_i \alpha e^{i\phi/2}\rangle_e]/\sqrt{2}$
5.	carrier $\pi/2$ -pulse	0.5	0	$1/2 \downarrow\rangle_i[\alpha e^{-i\phi/2}\rangle_e - e^{i\delta} \alpha e^{i\phi/2}\rangle_e]$ $-i/2 \uparrow\rangle_i[\alpha e^{-i\phi/2}\rangle_e + e^{i\delta} \alpha e^{i\phi/2}\rangle_e]$

FIGURE CAPTIONS

Fig 1. (a) Electronic (internal) and motional (external) energy levels (not to scale) of the trapped ${}^9\text{Be}^+$ ion, coupled by indicated laser beams A through D. The difference frequency of the "carrier" Raman beams A and B is set near $\omega_{\text{HF}}/2\pi \approx 1.250$ GHz, providing a two-photon Raman coupling between the ${}^2\text{S}_{1/2}(F=2, m_F=-2)$ and ${}^2\text{S}_{1/2}(F=1, m_F=-1)$ hyperfine ground states (denoted by $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ respectively). The difference frequency of the "displacement" Raman beams B and C is set to $\omega_x/2\pi \approx 11.2$ MHz. This excites the motion of the ion to a coherent state $|\alpha e^{i\theta}\rangle_e$ from an initial zero-point state of motion $|0\rangle_e$ in the harmonic potential. Because of the polarization of beams B and C, they do not affect motion correlated with the $|\downarrow\rangle_i$ internal state. The three Raman beams (A, B, and C) are detuned $\Delta \approx -12$ GHz from the ${}^2\text{P}_{1/2}(F=2, m_F=-2)$ excited state (radiative linewidth $\gamma/2\pi \approx 19.4$ MHz). Detection of the internal state is accomplished by illuminating the ion with σ^- -polarized "detection" beam D, which drives the cycling ${}^2\text{S}_{1/2}(F=2, m_F=-2) \rightarrow {}^2\text{P}_{3/2}(F=3, m_F=-3)$ transition, and observing the scattered fluorescence. (b) Geometry of the three Raman laser beams A, B, and C, with polarizations indicated. The quantization axis defined by the applied magnetic field **B** is 45° from the x-axis of the harmonic trap potential.

Fig 2. Evolution of the position-space atomic wavepacket entangled with the internal states $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ during creation of a Schrödinger cat state with $\alpha=3$ and $\phi=\pi$ (see Table I). The wavepackets are snapshots in time, taken when the atom is at the extrema of motion in the harmonic trap (represented by the parabolas). The area of the wavepackets corresponds to the probability of finding the atom in the given internal state. (a) The initial wavepacket corresponds to the quantum ground state of motion following laser-cooling. (b) The wavepacket is split following a $\pi/2$ -pulse on the carrier. (c) The $|\uparrow\rangle_i$ wavepacket is excited to a coherent state by the force **F** of the displacement beams. Note the force **F** acts only on the $|\uparrow\rangle_i$ wavepacket, thereby entangling the internal and motional systems. (d) The $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ wavepackets are exchanged following a π -pulse on the carrier. (e) The $|\uparrow\rangle_i$ wavepacket is excited to a coherent state by the displacement beam force **-F**, which is out of phase with respect to the force in (c). The state shown in (e) corresponds most closely to Schrödinger's cat (Eqs. 1 and 2). (f) The $|\downarrow\rangle_i$ and $|\uparrow\rangle_i$ wavepackets are finally combined following a $\pi/2$ -pulse on the carrier.

Fig 3. Evolution of the position-space wavepacket superposition correlated with the $|\downarrow\rangle_i$ internal state as the phase separation ϕ of the two coherent states is varied, for $\alpha=3$ and $\delta=0$. The expected signal $P_i(\phi)$ is the integrated area under these wavepackets. Each trace is a snapshot in time, taken when one of the wavepackets is at the right-most turning point in the harmonic trap. The wavepackets are maximally separated at $\phi=\pi$ [$P_i(\phi) \approx 1/2$], but begin to overlap as ϕ gets smaller [$P_i(\phi)$ oscillates]. Finally, the wavepackets destructively interfere at $\phi=0$ [$P_i(\phi) = 0$]. This vanishing interference signal is a signature of an odd Schrödinger cat state associated with the $|\downarrow\rangle_i$ state, since $\delta=0$. Probability conservation is ensured by a similar but constructive interference in the $|\uparrow\rangle_i$ state.

Fig 4. Measured and fit interference signal $P_i(\phi)$ versus the phase difference ϕ of two coherent states for $\delta=0$. Curves (a) to (d) represent measurements for various values of τ (2, 3, 5, and 15 μs , respectively). As τ grows, the feature near $\phi=0$ narrows. The lines are fits of the measurements to

the parameter α (Eq. (4)), yielding $\alpha = 0.84, 1.20, 1.92,$ and 2.97 . The fit in curve (d) includes a loss of contrast and represents a superposition of two $x_0 \approx 7$ nm wavepackets with a maximum separation of $4\alpha x_0 \approx 80$ nm. Curve (e) is a theoretical plot for a pair of coherent states with $\alpha=6$. Each data point in (a) to (d) represents an average of ≈ 4000 measurements, or 1 s of integration.

Fig 5. Measured interference signal $P_1(\phi)$ for three values of δ ($\alpha \approx 1.5$). The solid curve corresponds to $\delta = 1.03\pi$ (approximate even cat state correlated with $|\downarrow\rangle_i$ exhibiting constructive interference), the dashed curve to $\delta = 0.48\pi$ (approximate "Yurke-Stoler" cat state [19]), and the dotted curve to $\delta = 0.06\pi$ (approximate odd cat state exhibiting destructive interference). Each data point represents an average of ≈ 4000 measurements, or 1 s of integration.

REFERENCES

1. P.A.M. Dirac, *The Principles of Quantum Mechanics* (Clarendon, Oxford, ed. 4, 1984), p. 12.
2. A. Einstein, B. Podolsky, N. Rosen, *Phys. Rev.* **47**, 777 (1935).
3. E. Schrödinger, *Naturwissenschaften* **23**, 807, 823, 844 (1935); translation in [6].
4. Issues involving observation by the cat itself (by virtue of its consciousness) are beyond the scope of this paper. For our purposes, the cat can be replaced for example by an object placed in a superposition of macroscopically separated positions.
5. In the literature, there is not universal agreement about the definition of the “cat” state. As opposed to the definition used in Eqs. 1 and 2 of this article, some authors prefer to define a Schrödinger cat state as $(|x_1\rangle + |x_2\rangle)/\sqrt{2}$, or a superposition of two classical-like states that are not correlated with the state of a second system. In the analysis of the state shown in Eq. 2, we in fact create the state corresponding to this alternate definition.
6. *Quantum Theory and Measurement*, J.A. Wheeler and W.H. Zurek, Eds. (Princeton Univ. Press, Princeton, NJ, 1983).
7. N. Bohr, *Nature* **121**, 580 (1928); *Phys. Rev.* **48**, 696 (1935).
8. W. Heisenberg, *Z. Physik* **43**, 172 (1927); translation in [6].
9. W.H. Zurek, *Physics Today* **44**, 36 (October, 1991).
10. D.F. Walls and G.J. Milburn, *Phys. Rev.* **A31**, 2403 (1985); C.M. Savage and D.F. Walls, *Phys. Rev.* **A32**, 2316 (1985); M.J. Collett, *Phys. Rev.* **A38**, 2233 (1988).
11. D.P. DiVincenzo, *Science* **270**, 255 (1995); I.L. Chuang, R. Laflamme, P.W. Shor, W.H. Zurek, *Science* **270**, 1633 (1995).
12. C.H. Bennett, *Physics Today* **48**, 24 (October, 1995).
13. L. Marton, J. Arol Simson, J.A. Suddeth, *Phys. Rev.* **90**, 490 (1954).
14. H. Maier-Leibnitz and T. Springer, *Z. Physik* **167**, 368 (1962).
15. O. Carnal and J. Mlynek, *Phys. Rev. Lett.* **66**, 2689 (1991); D.W. Keith, C.R. Ekstrom, Q.A. Turchette, D.E. Pritchard, *Phys. Rev. Lett.* **66**, 2693 (1991); M. Kasevich and S. Chu, *Phys. Rev. Lett.*, **67** 181 (1991); J. Lawall, S. Kulin, B. Saubamea, N. Bigelow, M. Leduc, and C. Cohen-Tannoudji, *Phys. Rev. Lett.* **75**, 4194 (1995).
16. L.D. Noordam, D.I. Duncan, T.F. Gallagher, *Phys. Rev.* **A45**, 4734 (1992); R.R. Jones, C.S. Raman, D.W. Schumacher, P.H. Bucksbaum, *Phys. Rev. Lett.* **71**, 2575 (1993); M.W. Noel and C.R. Stroud, Jr., *Phys. Rev. Lett.* **75**, 1252 (1995).
17. J. Janszky, A.V. Vinogradov, T. Kobayashi, and Z. Kis, *Phys. Rev.* **A50**, 1777 (1994); I.A. Walmsley and M.G. Raymer, *Phys. Rev.* **A52**, 681 (1995).
18. A.J. Leggett and A. Garg, *Phys. Rev. Lett.* **54**, 857 (1985).
19. B. Yurke and D. Stoler, *Phys. Rev. Lett.* **57**, 13 (1986); *Phys. Rev.* **A35**, 4846 (1987); *Phys. Rev.* **A36**, 1955 (1987).
20. S.J.D. Phoenix and P.L. Knight, *Ann. Phys.* **186**, 381 (1988); J. Gea-Banacloche, *Phys. Rev. Lett.* **65**, 3385 (1990); V. Bužek, H. Moya-Cessa, P.L. Knight, S.J.D. Phoenix, *Phys. Rev.* **A45**, 8190 (1992).
21. J.J. Slosser, P. Meystre, and E.M. Wright, *Opt. Lett.*, **15**, 233 (1990).
22. C.M. Savage, S.L. Braunstein, D.F. Walls, *Opt. Lett.* **15**, 628 (1990).

23. M. Brune, S. Haroche, J.M. Raimond, L. Davidovich, N. Zagury, *Phys. Rev.* **A45**, 5193 (1992); S. Haroche and J.M. Raimond, in *Cavity Quantum Electrodynamics*, P. Berman, Ed. (Academic Press, NY, 1994), p. 123.
24. J.F. Poyatos, J.I. Cirac, R. Blatt, P. Zoller (to be published).
25. R.L. de Matos Filho and W. Vogel, *Phys. Rev. Lett.* **76**, 608 (1996).
26. L. Davidovich, A. Maali, M. Brune, J.M. Raimond, S. Haroche, *Phys. Rev. Lett.*, **71**, 2360 (1993).
27. P. Goetsch, R. Graham, F. Haake, *Phys. Rev.* **A51**, 136 (1995).
28. S.R. Jefferts, C. Monroe, E.W. Bell, D.J. Wineland, *Phys. Rev.* **A51**, 3112 (1995).
29. C. Monroe, D.M. Meekhof, B.E. King, S.R. Jefferts, W.M. Itano, D.J. Wineland, P. Gould, *Phys. Rev. Lett.* **75**, 4011 (1995).
30. D.M. Meekhof, C. Monroe, B.E. King, W.M. Itano, D.J. Wineland, *Phys. Rev. Lett.* **76**, 1796 (1996).
31. D.J. Wineland, J.C. Bergquist, J.J. Bollinger, W.M. Itano, F.L. Moore, J.M. Gilligan, M.G. Raizen, D.J. Heinzen, C.S. Weimer, C.H. Manney, in *Laser manipulation of Atoms and Ions*, E. Arimondo, W.D. Phillips, F. Strumia, Eds. (North-Holland, Amsterdam, 1992), p. 553.
32. R.J. Glauber, *Phys. Rev.* **131**, 2766; **131**, 2529 (1963).
33. In the laboratory frame, the coherent state parameter β has the periodic time dependence $\beta_L = \alpha e^{i(\theta - \omega t)}$, where ω is the frequency of the harmonic oscillator. Throughout the article, this time dependence is suppressed by working in an interaction frame where $\beta = \beta_L e^{i\omega t}$.
34. The effects of coupling to the $^2P_{3/2}$ excited state are small (detuning ≈ -210 GHz) and can be neglected here.
35. J. Peřina, *Quantum Statistics of Linear and Nonlinear Optical Phenomena* (Reidel, Dordrecht, 1984).
36. For large values of $\eta\Omega_d\tau$, we expect $\alpha(\tau)$ to saturate due to nonlinear correction factors which limit the amplitude of the coherent state and introduce amplitude squeezing (see [30] and W. Vogel and R.L. de Matos Filho, *Phys. Rev.* **A52**, 4214 (1995)). For the experimental Lamb-Dicke parameter of ~ 0.2 , these corrections are not significant for $\alpha \lesssim 5$.
37. J.F. Poyatos, J.I. Cirac, P. Zoller (private communication).
38. D.T. Smithey, M. Beck, M.G. Raymer, A. Faridani, *Phys. Rev. Lett.* **70**, 1244 (1993); S. Wallentowitz and W. Vogel, *Phys. Rev. Lett.* **75**, 2932 (1995); J.I. Cirac and P. Zoller (to be published).