

Presented at the European Bioelectromagnetics Association annual meeting, on November 12, 2003, Budapest Hungary

DOES GROUNDING THE HUMAN BODY TO EARTH REDUCE CHRONIC INFLAMMATION AND RELATED CHRONIC PAIN?

Clinton OBER, Earth Tether International, Ventura CA and Roger W COGHILL, Coghill Research Laboratories, Pontypool Wales NP4 5UH

INTRODUCTION

In modern societies our bodies are not normally grounded to earth because synthetic soled shoes, elevated sleeping places, motorcars, etc insulate us from the earth's surface. Accordingly, the human body can become charged with static electricity exacerbated by exposure to extraneous stray fields and currents from artificial sources such as cell phone masts, cell phones, power lines, and domestic appliances. On the surface of the earth a flow of electrons encompasses the entire planet. This natural flow of electrons and its related electric field also exists on the surface of all conductive objects (including people, plants and animals) in physical contact with the earth. The earth's electron current flow and its associated electric field pulsate at approximately 10 Hz (similar to alpha brain waves) and follow a rhythmic 24-hour circadian rise and fall in amplitude (Natural Electric Currents of the Earth, 43(1):47-57).

It is known that exposure to environmental 50-60 Hz electromagnetic fields increases concentrations of free radicals, lengthens their lifespan, and enhances the probability that they can do damage to the body (Bonnafous 1999, Brezitskaia 2000, Cannistraro 1980, Eveson 2000, Fernie 2001, Fiorani 1997, Hanel, 2000, Jajte 2000, Koana 1997, Roy 1995, Simko 2001(A), Simko 2001(B), Scaiano 1995, Scaiano 1994(A), Scaiano 1994(B), Supino 2001, Varani 2000, Yoshikawa 2000, Zmyslony 1998).

We propose that physical contact with the earth maintains the human body at the natural electrical potential (voltage) of the earth. When the body is electrically coupled with the earth, the DC electrons of the earth and/or its related electric field, then residing on the surface of the body, absorb the excitation effects of environmental electric fields. By transferring (offsetting) the attraction of an electric field from the body (which has a limited supply of electrons) to the earth (which has an infinite supply of electrons), perturbing excitation of the body's endogenous electrons is significantly reduced. When the body is shielded with the DC of the earth, the electrons of the body are then protected from perturbation.

GROUNDING AND FREE RADICALS

Free radicals are unpaired electrons which can have damaging effects on organic cells, hence the ubiquitous existence in all organisms of free radical scavengers such as catalase, superoxydismutase, glutathione and hormones like melatonin. Free radicals can arise from faulty metabolism (e.g. when insufficient molecular oxygen is available to act as final electron acceptor in the ox-phos pathway). The effects of grounding on free radical formation appear to assist in the antioxidant scavenging process by supplying additional electrons from the unlimited reservoir on

the earth's surface (we dub this "the Mother Earth effect"). All electrons carry a negative charge, but friction (e.g. on water travelling through a subterranean aquifer, or on a warm wind traversing a desert), can remove some electrons from the passing molecules, leaving positively charged ions which interact with organisms with biological sequelae. (*Webster's Dictionary, se-que-la: an abnormal condition resulting from a previous disease.*) This generally adverse effect can be mitigated by contact with the earth. Similar adverse effects are claimed from exposure to ambient, electromagnetic fields (EMFs).


One indicator of such sequelae is the release of Cortisol, a natural stress hormone signalling oxidative stress. Cortisol is a steroid hormone made in the adrenal glands, which are small glands adjacent to the kidneys. It can easily be measured by a standard saliva test. Among its important functions in the body include roles in the regulation of blood pressure and cardiovascular function as well as regulation of the body's use of proteins, carbohydrates, and fats. Cortisol secretion increases in response to any stress in the body, whether physical (such as illness, trauma, surgery, or temperature extremes) or psychological.

When Cortisol is secreted, it causes a breakdown of muscle protein, leading to release of amino acids (the "building blocks" of protein) into the bloodstream. These amino acids are then used by the liver to synthesize glucose for energy, in a process called *gluconeogenesis*. This process raises the blood sugar level so the brain will have more glucose for energy. At the same time the other tissues of the body decrease their use of glucose as fuel. Cortisol also leads to the release of so-called fatty acids, an energy source from fat cells, for use by the muscles. Taken together, these energy-directing processes prepare the individual to deal with stressors and insure that the brain receives adequate energy sources. Excess Cortisol is associated with sleep disorders, inflammation, pain and other adverse symptoms.

EXPERIMENTAL STUDIES

A new study (Ghaly, Teplitz, 2004) reports that electrical grounding during the night significantly reduced Cortisol levels in eight female subjects.

Cortisol levels in normal and grounded subjects*



*Cortisol measurements by Sabre Sciences Laboratory of San Diego, CA using a standard radioimmunoassay

In this study, conducted to identify the biological effects of grounding the human body during sleep, a grounding method of sleeping on a grounded bed pad was used.

Table 1: AC electric field-induced voltage measured on subjects' bodies while lying in their own beds

Subject	Before Grounding	After Grounding
1	3.940 V	0.003 V
2	1.470 V	0.001 V
3	2.700 V	0.004 V
4	1.200 V	0.002 V
5	2.700 V	0.005 V
6	1.670 V	0.005 V
7	5.950 V	0.008 V
8	3.940 V	0.008 V

During a six-week period, eight female subjects (age range 24-52) were grounded to the earth during sleep. Conductive bed pads were placed on their own beds (under the fitted sheet) and were connected to a fuse-protected ground wire attached to a ground rod that was placed directly in the earth outdoors (grounding via water pipes or internal ground wiring was not considered adequate in view of stray currents present in such systems). The electric field-induced body voltage (from exposure to common electrical wiring and cords near the bed), created on subjects' bodies while in bed averaged 2.9 volts, pre-grounding. Levels were significantly reduced, averaging 0.004 volts, when subjects slept on the earthed bed pads.

In addition to the objective measured reduction in body voltage potential, subjective responses were also obtained from the eight subjects. Though these clearly indicated an improvement in sleeping pattern and depth, cramps, numbness, PMS, aches and pain, and flatus the study was not controlled or blinded, hence not amenable to statistical treatment, and does not measure any placebo effect. A follow up three months later however still reported the same improvements.

Other case histories indicate that electrical grounding can alleviate ailments such as diabetes and inflammation:



Diabetic symptoms before (left) and after grounding for 7 nights (right).



Inflammation before (left) and after grounding for 7 nights (right).

FURTHER EXPERIMENTAL SUPPORT

In another study (Ober, 2000) the changes following electrical grounding in 60 sleep-disturbed and/or chronic muscle and joint pained subjects were investigated. Participants slept on grounded mattress pads for 30 consecutive nights. Subjective results of that study are reported below.

Categories	Test Subjects*		Control Subjects**	
	Same	Improved	Same	Improved
Time to fall asleep	4 - 15%	23 - 85%	20 - 87%	3 - 13%
Quality of sleep	2 - 7%	25 - 93%	20 - 87%	3 - 13%
Wake feeling rested	0 - 0%	27 - 100%	20 - 87%	3 - 13%
Muscles stiffness & pain	5 - 18%	22 - 82%	23 - 100%	0 - 0%
Chronic back and/or joint pain	7 - 26%	20 - 74%	23 - 100%	0 - 0%
General well-being	6 - 22%	21 - 78%	20 - 87%	3 - 13%

**Reports not received from three participants. **Reports not received from seven participants.*

This study provides evidence that when the human body is electrically grounded to the earth it is naturally protected from static electricity and ambient electric fields. Meter readings of the grounded subjects support the finding. The benefit of grounding the body appears to relax muscles and improve sleep.

Several of the 60 subjects in this study also experienced significant relief from asthmatic and respiratory conditions, rheumatoid arthritis, PMS, sleep apnoea and hypertension, while sleeping grounded. These unexpected results indicate that loss of grounding contact plays a much larger role in overall health than was anticipated.

DISSCUSSION

In any home the ambient, *extremely low frequency* (ELF) electric fields can vary surprisingly, being especially high near to appliances such as water heaters, fish tanks, storage heaters and other domestic electric wiring and/or appliances. A typical range is 1-10 Volts/metre, though near to such appliances the ELF fields can exceed 100V/m. Epidemiological studies of exposure to electric fields are few, since most have concentrated on the magnetic component, without any clearly accepted outcome except a persistent elevation of health disorders being reported (e.g. in childhood leukaemia where the focus of studies has been directed).

During one 1996 study of 56 participants (Coghill, Stewart et al., 1996), ALL cases and matched controls reported a nigh fivefold elevation of childhood leukaemia where the mean 12 hour bedplace exposure to ELF electric fields exceeded 20V/m. A later and much larger study (Skinner et al., 2000) however found no effects, though this relied only on spot (3 mins.) daytime measurements and 48 hour continuous measurements, both of which could arguably be unrepresentative of bedplace exposures.

The results of grounding experiments may implicate an improvement of circulation. Blood cells are normally net negatively charged on their surface, as are the surfaces of the blood vessels through which they travel. In consequence, the mutually repulsive action of the two tissues will ensure an easier flow – hence better tissue oxygenation. If the level of inhaled positive ions is high however this will cause a more attractive cohesion or “stickiness” in the bloodstream and inhibit bio-availability of molecular oxygen, leading to increased levels of free radicals, consequent oxidative stress, and symptoms such as arthritic inflammation, increased Cortisol, and other indicators of stress.

Grounding the body directly to earth permits the introduction of negatively charged electrons from the earth’s unlimited reservoir, and thus may restore the equilibrium, reducing free radical action. It is noteworthy that people suffering with arthritic conditions appear particularly sensitive to incoming positive air ionisation occurring with the build up of storms, and at times of the full moon (we dub this the Lunar effect).

According to the American Institute of Stress, over 75% of all visits to primary care physicians are now for stress-related health conditions. The description of stress is: a state of continuous anxiety and nervousness in which muscles become and remain tensed. Stress is now confirmed to be a primary contributor to cardiovascular disease, cancer, gastrointestinal, skin, neurological and emotional disorders, and a host of disorders linked to immune system disturbances ranging from the common cold and herpes, to arthritis and AIDS (see American Institute of Stress website, www.stress.org/problems).

According to the National Sleep Foundation's “Sleep-2000” report (see website: www.sleepfoundation.org/pressarchives), nearly two thirds of American adults [62%] now suffer from sleep problems. Americans have the most comfortable beds and the most protected sleep environments in the world. Yet, in traditional societies where most humans sleep on animal skins, grass mats or directly on the ground, sleep problems do not exist (see *Slumbers Unexplored Landscape* [1999] Carol M. Worthman, Anthropologist, Emory University Atlanta, GA).

As for Americans and others in industrialised societies, most now sleep within 12 inches of electrical wires hidden in the wall at the head of their bed and with electric cords around or near the bed, which emanate ELF electric fields throughout the night and create weak electric currents in the body (see National Institute of Environmental Health Sciences and the U.S. Department of Energy, *Questions and Answers about EMF, electric and magnetic fields associated with use of electric power*, 1995).

It is noteworthy that the electric component of the ELF EM field is present *all the time the wiring is switched on* at the mains, whether the appliance or light bulb is in use or not, whereas the magnetic component is only present when an appliance or other load is energised. This fact may well have confounded epidemiological studies of domestic magnetic field exposure. Moreover, a recent study

reported that over half UK children sleep with a light switched on in their bedroom, possibly disrupting synthesis of melatonin, an important oncostatic hormone produced almost entirely in the brain's light-sensitive pineal gland.

REFERENCES and BIBLIOGRAPHY

1. Bonnafous P, Vernhes M, Teissie J, Gabriel B. The generation of reactive-oxygen species associated with long-lasting pulse-induced electropermeabilisation of mammalian cells is based on a non-destructive alteration of the plasma membrane. *Biochim Biophys Acta*. 1999 Nov 9;1461(1):123-34.
2. Brezitskaia HV, Timchenko OI. On the mechanism of cytogenetic effect of electromagnetic radiation: a role of oxidation homeostasis. *Radiats Biol Radioecol*. 2000 Mar-Apr;40(2):149-53.
3. Cannistraro S, Martino G, Sportelli L. Effects of pulsed electric fields on rat liver homogenate paramagnetic species. *Radiat Environ Biophys*. 1980;18(2):123-8.
4. Coghill & Stewart. Measured ELF electric and magnetic fields in the bedplace of children diagnosed with leukemia: a case-controlled study, *Europ. J. Cancer Prev*, 5: 153-158 June 1996.
5. Eveson RW, Timmel CR, Brocklehurst B, Hore PJ, McLauchlan KA. The effects of weak magnetic fields on radical recombination reactions in micelles. *Int. J. Radiat. Biol*. 2000 Nov;76(11):1509-22.
6. Fernie KJ, Bird DM. Evidence of oxidative stress in American kestrels exposed to electromagnetic fields. *Environ Res*. 2001 Jun;86(2):198-207.
7. Fiorani M, Biagiarelli B, Vetrano F, Guidi G, Dacha M, Stocchi V. In vitro effects of 50 Hz magnetic fields on oxidatively damaged rabbit red blood cells. *Bioelectromagnetics*. 1997;18(2):125-31.
8. Ghaly, M. and Teplitz, D. The biologic effects of grounding the human body during sleep as measured by cortisol levels and subjective reporting of sleep, pain and stress. *J. Alt. Comp. Med*. 2004 Vol. 10, No.5 767-776.
9. Hanel G, Gstir B, Denifl S, Scheier P, Probst M, Farizon B, Farizon M, Illenberger E, Mark TD. Electron attachment to uracil: effective destruction at subexcitation energies. *Pathophysiology*. 2000 Jul;7(2);131-135.
10. Jajte J.M. Programmed cell death as a biological function of electromagnetic fields at a frequency of (50/60 Hz)-review. *Med Pr*. 2000;51(4):383-9.
Koana T, Okada MO, Ikehata M, Nakagawa M. Increase in the mitotic recombination frequency in *Drosophila melanogaster* by magnetic field exposure and its suppression by vitamin E supplement. *Mutat Res*. 1997 Jan 3;373(1):55-60.
11. Ober, A.C. Grounding the human body to neutralize bio-electrical stress 2000 *Electrostatics Journal* <http://www.esdjournal.com/articles/cober/ground.htm>
12. Roy S, Noda Y, Eckert V, Traber MG, Mori A, Liburdy R, Packer L. The phorbol 12-myristate 13-acetate (PMA)-induced oxidative burst in rat peritoneal neutrophils is increased by a 0.1 mT (60 Hz) magnetic field. *FEBS Lett*. 1995 Dec 4;376(3):164-6.
13. Scaiano JC, Cozens FL, McLean J, Model for the rationalization of magnetic field effects in vivo. Application of the radical-pair mechanism to biological systems. *Photochem Photobiol* 1994 Jun;59(6):585-89.
14. Scaiano JC, Cozens FL, Mohtat N. Influence of combined AC-DC magnetic fields on free radicals in organized and biological systems. Development of a model and application of the radical pair mechanism to radicals in micelles. *Photochem Photobiol*. 1995 Nov;62(5):818-29.

15. Scaiano JC, Mohtat N, Cozens FL, McLean J, Thansandote A. Application of the radical pair mechanism to free radicals in organized systems: can the effects of 60 Hz be predicted from studies under static fields? *Bioelectromagnetics*. 1994;15(6):549-54.
16. *Scientific Monthly*, The Natural Electric Currents of the Earth. Vol. 43, issue 1:47-57.
17. Simko M, Droste S, Kriehuber R, Weiss DG. Stimulation of phagocytosis and free radical production in murine macrophages by 50 Hz electromagnetic fields. *Eur J Cell Biol*. 2001 Aug;80(8):562-6.
18. Simko M, Richard D, Kriehuber R, Weiss DG. Micronucleus induction in Syrian hamster embryo cells following exposure to 50 Hz magnetic fields, benzo(a)pyrene, and TPA in vitro. *Mutat Res*. 2001 Aug 22;495(1-2):43-50.
19. Skinner, J et. al. Childhood cancer and residential proximity to power lines, (UKCCS Investigators), *British Journal of Cancer* (2000) 83 (11) 1573-1580 November 2000.
20. Supino R, Bottone MG, Pellicciari C, Caserini C, Bottiroli G, Belleri M, Veicsteinas. Sinusoidal 50 Hz magnetic fields do not affect structural morphology and proliferation of human cells in vitro. *Histol Histopathol*. 2001 Jul;16(3):719-26.
21. Varani K, Gessi S, Merighi S, Iannotta V, Cattabriga E, Spisani. Effect of low frequency electromagnetic fields on A2A adenosine receptors in human neutrophils. *Int. J. Radiat. Biol*. 2000 Nov;76(11):1509-22.
22. Yoshikawa T, Tanigawa M, Tanigawa T, Imai A, Hongo H, Kondo M. Enhancement of nitric oxide generation by low frequency electromagnetic field. *Pathophysiology*. 2000 Jul;7(2):131-135.
23. Zmyslony M, Jajte JM. The role of free radicals in mechanisms of biological function exposed to weak, constant and net magnetic fields. *Med Pr*. 1998;49(2):177-86.