

## Carotenoids Contents from Various Sources and Their Potential Health Applications

Alam Zeb and Sultan Mehmood

Department of Biotechnology, University of Malakand, Chakdara, N.W.F.P, Pakistan

E-mail: Alamzeb01@yahoo.com

**Abstract:** Carotenoids are the class of hydrocarbons, widespread in nature and important pigments in living organisms. They are present in foods such as carrots, pumpkins, sweet potatoes, tomatoes, and other deep green, yellow, orange, red fruits, sea buckthorn berries, common vegetables and some numerous vegetable oils, of which Palm oil and palm oil products contains the highest known concentration of carotenoids. Palm oil products are the good sources of carotenoids in nature. The role of carotenoids in health application has been explained. The present paper describes the chemistry of important carotenoids, their various sources and uses in health applications.

**Key words:** Carotenoids, chemistry, health applications, sources, vegetables

### Introduction

The name carotenoids, is derived from the fact that they constitute the major pigment in the carrot root, *Daucus carota*, are undoubtedly among the most widespread and important pigments in living organisms. Carotenoids are the pigments responsible for the colors of many plants, fruits and flowers. Carotenoids are fat-soluble nutrients and categorized as either xanthophylls or carotenes according to their chemical composition. Carotenoids are found in common foods and vegetables. Most xanthophylls are found in green leafy vegetables and nearly all carotenes are found in yellow vegetables. The most important carotenoids are alpha-carotene, beta-carotene, and beta-cryptoxanthin, Lutein, violaxanthin, neoxanthin, and Lycopene; their structures are given in Fig. 1.

Beta-carotene, alpha-carotene, beta-cryptoxanthin are carotenes that are converted into vitamin A or retinol in the body. They are found in many yellow fruits and vegetables. Beta-carotene is the most widely studied carotenoid. Lutein and zeaxanthin are both stored in the retina of the eye. Neither converts to vitamin A. Both are powerful antioxidants and may be very important for healthy eyes. They are found in many yellow fruits and vegetables. Lutein is also present in green vegetables, such as broccoli, cabbage, and kale. Lycopene is responsible for the red color in fruits and vegetables, including tomatoes, red grapes, watermelon, and pink grapefruit. It is also found in papayas and apricots. It does not convert to vitamin A but may have important cancer fighting properties and other health benefits. Currently, interest is focusing on the nutritional and medicinal aspects of different individual carotenoid. Since they are likely to grow in importance and value, their sources is very important. Therefore the present study aimed to collect recent information on the important carotenoids contents from various sources




Fig. 1: Structure of selected carotenoids

and their relation with health aspects, so that nutritional importance of these sources is clear for our common day life.

**Chemistry of Carotenoids:** Carotenoids are a class of

## Zeb and Mehmood: Carotenoids Contents from Various Sources and Their Potential Health Applications

Table 1: Carotenoids content of Sea buckthorn oil from different origins and variety

Sources	Carotenoids (mg/100g)		Reference	Variety
	Ranges	Average		
Carotenoids content	314 - 2139	1167	Zhang <i>et al.</i> , 1989	Caucas
Seed oil	50 - 85	67.5	Mironov, 1989	Caucas
Pulp oil	330 - 370	350	Mironov, 1989	Caucas
Seed coat oil	180 - 220	200	Mironov, 1989	Pamirs
Seed oil	Trace	--	Mironov, 1989	Pamirs
Pulp oil	900 - 1000	950	Mironov, 1989	

Table 2: Retinol equivalents of red palm oil compared with other foods products

Fruits and vegetables	Carotene derived (ppm)
Red palm oil	30,000
Carrots	2,000
Leafy vegetables	685
Apricots	250
Tomatoes	100
Bananas	30
Orange juice	8

hydrocarbons consisting of eight isoprenoid units joined in such a manner that the arrangement of isoprenoid units is reversed at the center of the molecule so that the two central methyl groups are in a 1,6-positional relationship and the remaining non-terminal methyl groups are in a 1,5-positional relationship (Fig. 1). Carotenoids are defined by their chemical structure. The majority carotenoids are derived from a 40-carbon polyene chain, which could be considered the backbone of the molecule. This chain may be terminated by cyclic end-groups (rings) and may be complemented with oxygen-containing functional groups (McCollum and Kennedy, 1916). These hydrocarbons are commonly known as carotenes, while oxygenated derivatives of these hydrocarbons are known as xanthophylls. Beta-carotene, the principal carotenoid in carrots, is a familiar carotene, while lutein, the major yellow pigment of marigold petals, is a common xanthophyll. The structure of a carotenoid ultimately determines what potential biological function that pigment may have. The characteristic pattern of alternating single and double bonds in the polyene backbone of carotenoids allows them to absorb excess energy from other molecules, while the nature of the specific end groups on carotenoids may influence their polarity. The former may account for the antioxidant properties of biological carotenoids, while the latter may explain the differences in the ways that individual carotenoids interact with biological membranes (Britton, 1995).

**Sources of Carotenoids:** About 600 carotenoids have been isolated from natural sources (Ong and Tee, 1992). The most important sources for carotenoids are plants, where often the brilliant colors of the carotenoids

are covered by the green chlorophyllic pigments (i.e. in green vegetables and leaves). In a number of cases, as plants mature, the chlorophyll content decreases leaving the carotenoids responsible for the beautiful colors of most fruits (pineapple, oranges, lemons, grapefruit, strawberry, tomatoes, paprika, rose hips) and many flowers (Eschscholtzia, Narcissus). Carotenoids are also responsible for the colors of some birds (flamingo, canary), certain insects, and marine animals (shrimp, lobster and salmon) (Mangels *et al.*, 1993).

Sea buckthorn fruits are rich in carotenoids pigments located in membranes and the fleshy mesocarp. Caroteno-lipoprotein complexes are located particularly in fruit membranes where polar lipids may function as bridge compounds between the polar (protein) and non-polar (carotenoids) moieties (Pintea *et al.*, 2001). It was found that the carotenoids to consist of nearly 20%  $\beta$ -carotene, 30%  $\alpha$ -carotene, and 30% lycopene as well as 15% oxygen containing carotenoids (Zeb, 2004). Pulp oil shows higher levels of  $\beta$ -carotene than seed oil, and *H. salicifolia* appears to have the highest level of  $\beta$ -carotene in both pulp and seed oil among species (Table 1). It was established that more the  $\beta$ -carotene greater will be stability of oils (Lutfullah *et al.*, 2003; Zeb and Ahmad 2004; Zeb *et al.*, 2004). Within *H. rhamnoides*, subsp. *mongolica* shows the lowest  $\beta$ -carotene level in seed and pulp oil (Lian *et al.*, 2000). Concentration of  $\beta$ -carotene constitute 15–55% of total carotenoids, depending on the origin (Lian *et al.*, 2000; Yang and Kallio, 2001). However the concentration of  $\beta$ -carotene and of the total carotenoids are affected substantially by berry maturity, and practices of fertilization (Zhang *et al.*, 1989; Yang, 2001).

Carotenoids are present in numerous vegetable oils, including corn, groundnut, soybean, rapeseed, linseed, olive, barley, sunflower, and cottonseed oils (NRC, 1989; Olson, 1990; Goodman, 1984; UNICEF, 1992; Int Vit A Con Group, 1989; Zeitlin *et al.*, 1992; Gibson, 1990). The concentration of carotenoids in these vegetable oils is generally low, less than 100 ppm. Of the vegetable oils that are widely consumed, Palm oil contains the highest known concentration of agriculturally derived carotenoids (Jain *et al.*, 1990). In fact, crude palm oil is the world's richest natural plant source of carotenes in terms of retinol. It contains about 15 to 300 times as many retinol

## Zeb and Mehmood: Carotenoids Contents from Various Sources and Their Potential Health Applications

Table 3: carotenoids contents (percentages) of palm oil from different species and sources

Type	<i>E. guineensis</i>			<i>E. oleifera</i>	Hybrids		Back cross	Pressed fibre oil	Second pressed oil	Carotenoid concentration	Red palm oil
	tenera	pisifera (P)	Dura (D)	(O)	O x P	O x D	ODxP				
Phytoene	1.27	1.68	2.49	1.12	1.83	2.45	1.30	11.87	6.50	1.5	2.0
Phytofluene	0.06	0.90	1.24	trace	trace	0.15	trace	0.40	1.63	0.3	1.2
Cis-\$-carotene	0.68	0.10	0.15	0.48	0.38	0.55	0.42	0.49	0.28	0.9	0.8
\$-carotene	56.02	54.39	56.02	54.08	60.53	56.42	51.64	30.95	31.10	49.9	47.4
"-carotenes	35.16	33.11	24.35	40.38	32.78	36.40	36.50	19.45	20.68	33.3	37.0
Cis-"-carotene	2.49	1.64	0.86	2.30	1.37	1.38	2.29	1.77	1.70	5.5	6.9
.-carotene <sup>a</sup>	0.69	1.12	2.31	0.36	1.13	0.70	0.36	7.56	4.62	1.7	1.3
*-carotene	0.83	0.27	2.00	0.09	0.24	0.22	0.14	6.94	2.13	0.6	0.6
(-carotene <sup>b</sup>	0.33	0.48	1.16	0.08	0.23	0.26	0.19	2.70	2.48	1.3	0.5
Neurosporene <sup>b</sup>	0.29	0.63	0.77	0.04	0.23	0.08	0.08	3.38	1.88	0.1	trace
\$-zeacarotene	0.74	0.97	0.56	0.57	1.03	0.96	1.53	0.37	0.58	1.3	0.5
"-zeacarotene	0.23	0.21	0.30	0.43	0.35	0.40	0.52	trace	0.15	0.4	0.3
Lycopene <sup>c</sup>	1.30	4.50	7.81	0.07	0.05	0.04	0.02	14.13	26.45	3.4	1.5
Total (ppm)	673	428	997	4,592	1,430	2,324	896	5,162	2,510	80,560	545

a. With two cis isomers.      b. With one as isomer.      c. With three cis isomers.

Table 4: Carotenoid content of various palm oil fractions [9]

Palm oil fractions	Carotene derived (ppm)
Crude palm oil	630-700
Crude palm olein	680-760
Crude palm stearin	380-540
Residual oil from fibre	4,000-6,000
Second-pressed oil	1,800-2,400
Red Palm oil	500-700

Total carotenoids estimated at 446 nm

equivalents as carrots, leafy green vegetables, and tomatoes, which are considered to have significant quantities of provitamin A activity (Zachman, 1988) (Table 2).

Palm forests are found in West Africa, oil palms are actually cultivated in East Africa, South America, Malaysia, and Java. Palm oil from the Far East and from Zaire contains 500-800 ppm of carotenes, whereas that from Côte d'Ivoire and especially Benin contains 1,000-1,600 ppm, but the oil yield is less (Ostrea *et al.*, 1986; Ala-Houhala *et al.*, 1988). Oil from the tenera variety that is widely planted in Malaysia has a carotenoids content of about 500-700 ppm (Zeitlin *et al.*, 1992).

Other oil palm species, such as *Elaeis oleifera*, a South American palm, have been found to contain a higher concentration of carotenes (Nichols and Nichols, 1981). Hybrids of *E. oleifera* and *E. guineensis* also produce oils with high concentrations of carotenoids. The following values have been found for the total carotenoid content of oil from different species, varieties, and hybrids: *E. guineensis* var. *pisifera* (P), 428 ppm; var. *dura* (D), 997 ppm; *E. oleifera* (O), 4,592 ppm; the hybrid O x P, 1,430 ppm; O x D, 2,324 ppm; and the back-cross OD x P, 896 ppm (Table 3). *E. oleifera* oil has the highest carotenoid

content and *E. guineensis* oil the lowest, with the hybrids and the backcross having intermediate concentrations. It can be seen from the table that the species and hybrids are comparable with regard to the major components, "- and \$-carotene. The Table 4 shows the carotenoids contents of various palm products.

The primary odor constituents derived from carotenoids are C<sub>13</sub>, C<sub>11</sub>, C<sub>10</sub> and C<sub>9</sub> derivatives formed via enzymatic oxidation and photo-oxidation of the various carotenoids found in plants, flowers and fruits. While other aroma constituents such as esters, terpenes, pyrazines, etc. are usually also present, these C<sub>9</sub> to C<sub>13</sub> compounds often are essential to the odor profile. Table 5 shows common oxidative/ degradation products of Carotenoids, responsible for the different odor of the material they contains.

**Health Applications of Carotenoids:** In human beings, carotenoids can serve several important functions. The most widely studied and well-understood nutritional role for carotenoids is their provitamin A activity. Deficiency of vitamin A is a major cause of premature death in developing nations, particularly among children. Vitamin A, which has many vital systemic functions in humans, can be produced within the body from certain carotenoids, notably beta-carotene (Britton *et al.*, 1995; Patton *et al.*, 1990; Ibrahim *et al.*, 1991). Dietary beta-carotene is obtained from a number of fruits and vegetables, such as carrots, spinach, peaches, apricots, and sweet potatoes (Mangels *et al.*, 1993). Other provitamin A carotenoids include alpha-carotene (found in carrots, pumpkin, and red and yellow peppers) and cryptoxanthin (from oranges, tangerines, peaches, nectarines, and papayas). Carotenoids also play an important potential role in

## Zeb and Mehmood: Carotenoids Contents from Various Sources and Their Potential Health Applications

Table 5: Common Carotenoids Degradation Products Found in Plants

Name	Examples of Occurrence
Beta-Ionone	Osmanthus, Rose, Black Tea, Tomato, Blackberries, Raspberries, Passion fruit, Carrots, Tobacco, Apricot, Carambola, Cherries, Mango, Bell Pepper, Plum
Alpha-Ionone	Black currant, Osmanthus, Black Tea, Blackberries, Raspberries, Carrots, Tobacco, Banana, Cherries, Plum, Celery, Peach, Popcorn, Tomato
Beta-Damascone	Rose, Osmanthus, Black tea, Mountain papaya, Rum, Tobacco
Beta-Damascenone	Apricot, Rose, Beer, Carambola, Grape, Kiwi, Mango, Tomato, Wine, Rum, Raspberries, Passion fruit, Blackberries
Oxo-Edulan I	Purple Passion fruit, Osmanthus, Burley tobacco, Virginia tobacco
Oxo-Edulan II	Purple Passion fruit, Osmanthus, Burley tobacco, Virginia tobacco
Theaspirone	Yellow Passion fruit, Black Tea, Burley tobacco
4-Oxo-beta-ionone	Red Fox, Black Tea, Osmanthus, Burley tobacco, Freesia flower, Boronia
3-Oxo-alpha-ionone	Osmanthus, Virginia tobacco,
Dihydroactinodioidide	Osmanthus, Black Tea, Tomato, Cassia, Cassie, Ambergris, Tobacco
4-Oxoisophorone	Osmanthus, Black tea, Saffron
Safranal	Saffron, Osmanthus, Black tea, Grapefruit juice, Mate, Paprika
Beta-Cyclocitral	Roasted Mate, Rum, Tea, Tomato, Cantaloupe, Paprika, Peas, Apricot, Broccoli, Melon

human health by acting as biological antioxidants, protecting cells and tissues from the damaging effects of free radicals and singlet oxygen. Lycopene, the hydrocarbon carotenoid that gives tomatoes their red color, is particularly effective at quenching the destructive potential of singlet oxygen (Di Mascio *et al.*, 1989). Lutein and zeaxanthin, xanthophylls found in corn and in leafy greens such as kale and spinach, are believed to function as protective antioxidants in the macular region of the human retina (Snodderly, 1995; Handelman *et al.* 1988; Landrum *et al.*, 1997). Astaxanthin found in salmon, shrimp, and other seafoods, is another naturally occurring xanthophyll with potent antioxidant properties (Di Mascio *et al.*, 1991). Other health benefits of carotenoids that may be related to their antioxidative potential include enhancement of immune system function (Bendich, 1989), protection from sunburn (Matthews-Roth, 1990), and inhibition of the development of certain types of cancers (Nishino, 1998). Lycopene prevents oxidation of low-density lipoprotein (LDL) cholesterol and reduces the risk of developing atherosclerosis and coronary heart disease (Agarwal and Rao, 1998). This study showed that daily consumption of tomato products providing at least 40 mg of lycopene was enough to substantially reduce low-density lipoprotein (LDL) oxidation. High LDL oxidation is associated with increased risk of atherosclerosis and coronary heart disease. This lycopene level can be achieved by drinking just two glasses of tomato juice a day. Research shows that lycopene in tomatoes can be absorbed more efficiently by the body if processed into tomato juice, sauce, paste and ketchup. The bound chemical form of lycopene found in tomatoes is converted by the temperature changes involved in processing to make it more easily absorbed by the body. A considerable wealth of research data has been

accumulated regarding the efficacy of various carotenoids in prevention, treatment and adjuvant treatment of cancers; combined with chemotherapy and/or radiotherapy (Lu *et al.*, 2003; Mannisto *et al.*, 2004; Karp and Tsao, 2003; Van Zandwijk and Hirsch, 2003; Russell, 2004; Kristal, 2004; Murtaugh *et al.*, 2004). Despite of the toxic behavior of carotenoids toward some tumor patient especially the smokers; carotenoids are good source of anti-tumor agent.

**Conclusion:** Carotenoids are present in different edible and non-edible materials; of which the highest amount are present in palm oil and their various products; secondly, the sea buckthorn berries and yellow vegetables placed on the third position. So from the easiest availability of carotenoids from the various sources, describe above and their importance in various health applications, it is therefore easy to suggest that palm oil products, sea buckthorn berries and yellow vegetables; can be used as alternative low outlay medicine in combating serious health conditions.

### Acknowledgement

The authors are thankful to Mr. Zahir Ali and Mr. Shawkat Ali, Lecturers, Department of Biotechnology, UOM, for their help during the critical evaluation of this manuscript and computer writing.

### References

- Agarwal, S. and A.V. Rao, 1998. Tomato lycopene and low-density lipoprotein oxidation: a human dietary intervention study. *Lipids*, 33: 981-984.
- Ala-Houhala, M., T. Koskinen, R. Maki and S. Rinkari, 1988. Serum vitamin A levels in mothers and their breastfed term infants with or without supplemental vitamin A. *Acta Paediatr Scand.*, 77: 198-201.

## Zeb and Mehmood: Carotenoids Contents from Various Sources and Their Potential Health Applications

- Bendich, A., 1989. Carotenoids and the immune response. *J. Nutr.*, 119: 112-115.
- Britton, G., 1995. Structure and properties of carotenoids in relation to function. *FASEB J.*, 9: 1551-1558.
- Britton, G., S. Liaaen-Jensen and H. Pfander, 1995. Carotenoids today and challenges for the future. In: Britton, G., S. Liaaen-Jensen and H. Pfander [eds], *Carotenoids vol. 1A: Isolation and Analysis*. Basel: Birkhäuser.
- Di Mascio, P., S. Kaiser and H. Sies, 1989. Lycopene as the most efficient biological carotenoid singlet oxygen quencher. *Arch. Biochem. Biophys.*, 274: 532-538.
- Di Mascio, P., M.E. Murphy and H. Sies, 1991. Antioxidant defense systems: the role of carotenoids, tocopherols, and thiols. *Am. J. Clin. Nutr.*, 53: 194S-200S.
- Gibson, R.S., 1990. *Principles of nutritional assessment*. New York: Oxford University Press, pp: 318-320.
- Goodman, D.S., 1984. Vitamin A and retinoids in health and disease. *N. Engl. J. Med.*, 310: 1023-1031.
- Handelman, G.J., E.A. Dratz., C.C. Reay and J.G. Van Kuijk, 1988. Carotenoids in the human macula and whole retina. *Invest Ophthalmol Vis Sci.*, 29: 850-5.
- Ibrahim, K., T.J. Hassan and S.N. Jafarey, 1991. Plasma vitamin A and carotene in maternal and cord blood. *Asia Oceania J. Obstet. Gynaecol.*, 17: 159-64.
- International Vitamin A Consultative Group, 1989. Guidelines for the development of a simplified dietary assessment to identify groups at risk for inadequate intake of vitamin A. Nutrition Foundation, Washington, DC.
- Jain, M.K., N.J. Mehta., M. Fonseca and N.V. Pai, 1990. Correlation of serum vitamin A and its transport protein (RBP) in malnourished and vitamin A deficient children. *J. Postgrad. Med.*, 36: 119-123.
- Karp, D.D. and A.S. Tsao, 2003. Kim Nonsmall-cell lung cancer: chemoprevention studies. *Semin Thorac Cardiovasc Surg.*, 15: 405-420.
- Kristal, A.R., 2004. Vitamin A, retinoids and carotenoids as chemo preventive agents for prostate cancer. *J. Urol.* 2004; 171(2 Pt 2):S54-8; discussion S58.
- Landrum, J.T., R.A. Bone and M.D. Kilburn, 1997. The macular pigment: a possible role in protection from age-related macular degeneration. *Adv Pharm.*, 38:537-556.
- Lian, Y.S., S.G. Lu, S.K. Xue and X.L. Chen, 2000. *Biology and Chemistry of the Genus Hippophae*. Dansu Scientific and Technological Publishing House, Lanzhou, China, pp: 88-91.
- Lu, Q.J., C.Y. Huang, S.X. Yao, R.S. Wang and X.N. Wu, 2003. Effects of fat-soluble extracts from vegetable powder and beta-carotene on proliferation and apoptosis of lung cancer cell YTMLC-90. *Biomed Environ. Sci.*, 16: 237-45.
- Lutfullah, G., A. Zeb, T. Ahmad, S. Atta and F. Khan, 2003. Changes in the Quality of Sunflower and Soybean oils Induced by high Doses of Gamma Radiation. *J. Chem. Soc. Pak.*, 25: 269-275.
- Mangels, A.R., J.M. Holden, G.R. Beecher, M.R. Forman, and E. Lanza, 1993. Carotenoid content of fruits and vegetables: an evaluation of analytic data. *J. Am. Diet. Assoc.*, 93: 284-296.
- Mannisto, S., S.A. Smith-Warner, D. Spiegelman, D. Albanes, K. Anderson, P.A. van den Brandt, J.R. Cerhan, G. Colditz, D. Feskanich, J.L. Freudenheim, E. Giovannucci, R.A. Goldbohm, S. Graham, A.B. Miller, T.E. Rohan, J. Virtamo, W.C. Willett and D.J. Hunter, 2004. Dietary carotenoids and risk of lung cancer in a pooled analysis of seven cohort studies. *Cancer Epidemiol. Biomarkers Prev.*, 13: 40-48.
- Mathews-Roth, M.M., 1990. Plasma concentration of carotenoids after large doses of beta-carotene. *Am. J. Clin. Nutr.*, 52: 500-501.
- McCollum, E.V. and C. Kennedy, 1916. *J. Biol. Chem.* 24, 491-502.
- Mironov, V.A., 1989. Chemical composition of *Hippophae rhamnoides* of different populations of the USSR. Proceedings of International Symposium on sea buckthorn (*H. rhamnoides* L.), Xian, China.
- Murtaugh, M.A., K.N. Ma, J. Benson, K. Curtin, B. Caan and M.L. Slattery, 2004. Antioxidants, carotenoids and risk of rectal cancer. *Am. J. Epidemiol.*, 159: 32-41.
- Nichols, B.L. and V.N. Nichols, 1981. Human milk: nutritional resource. In: Tsang RC, ed. *Nutrition and child health: perspectives for the 1980s*. New York: Alan R. Liss, pp: 109-146.
- Nishino, H., 1998. Cancer prevention by carotenoids. *Mutat. Res.*, 402: 159-163.
- NRC, 1989. Commission on Life Sciences, Food and Nutrition Board. *Recommended dietary allowances*. 10th ed. Washington, DC: National Academy Press.
- Olson, J.A., 1990. Vitamin A. In: Brown ML, ed. *Present knowledge in nutrition*. Washington, DC: International Life Sciences Institute, Nutrition Foundation, pp: 96-107.
- Ong, A.S.H. and E.S. Tee. (1992) Natural sources of carotenoids from plants and oils. *Meth. Enzymol.*, 213: 142-167.
- Ostrea, EM., JE. Balun., R. Winkler and T. Porter. 1986. Influence of breast-feeding on the restoration of the low serum concentration of vitamin E and  $\beta$ -carotene in the newborn infant. *Am J Obstet Gynecol.*, 154: 101-117.
- Patton, S., LM. Canfield., GE. Huston., AM. Ferris and R.G. Jensen, 1990. Carotenoids of human colostrum. *Lipids.*, 25: 159-165.
- Pintea, A., A. Marpeau, M. Faye, C. Socaciu and M. Gleizes, 2001. Polar lipid and fatty acid distribution in carotenolipoprotein complexes extracted from sea buckthorn fruits. *Phytochemical Analysis.*, 5: 293-298.

### **Zeb and Mehmood: Carotenoids Contents from Various Sources and Their Potential Health Applications**

- Russell, R.M., 2004. The enigma of beta-carotene in carcinogenesis: what can be learned from animal studies. *J. Nutr.*, 134: 262S-268S.
- Snodderly, D.M., 1995. Evidence for protection against age-related macular degeneration by carotenoids and antioxidant vitamins. *Am. J. Clin. Nutr.*, 62: 1448S-1461S.
- UNICEF., 1992. The state of the world's children 1992. New York: Oxford University Press.
- Van Zandwijk, N., and F.R. Hirsch, 2003. Chemo prevention of lung cancer: current status and future prospects. *Lung Cancer.*, 42 (Suppl) 1: S71-79.
- Yang, B.R., 2001. Lipophilic components of sea buckthorn (*Hippophae. Rhamnoids*) seeds and berries and physiological effects of sea buckthorn oils. Ph.D dissertation, Turku University, Finland.
- Yang, B.R. and H.P. Kallio., 2001. Fatty acid composition of lipids in sea buckthorn (*Hippophae rhamnoides* L.) berries of different origins. *J. Agri. Food Chem.*, 49: 1939-1947.
- Zachman, R.D., 1988. Vitamin A. In: Tsang RC, Nichols BL, eds. Nutrition during infancy. Philadelphia, Pa, USA: Hanley & Belfus; St. Louis, Mo, USA: C. V. Mosby, pp: 253-263.
- Zeb, A., 2004. Chemical and Nutritional Constituents of Sea Buckthorn Juice. *Pak. J. Nutr.*, 3: 99-106.
- Zeb, A. and T. Ahmad, 2004. The high dose irradiations affect the quality parameters of edible oils. *Pak. J. Biol. Sci.*, 7: 943-946.
- Zeb, A., T. Ahmad and G. Lutfullah, 2004. Radiolytic and Storage stability Study of Soybean and Red palm oils. *Journal of Chem., Soc. of Pak.*, (Submitted).
- Zeitlin, M.F., R. Megawangi, E.M. Kramer and H.C. Armstrong, 1992. Mothers and children's intakes of vitamin A in rural Bangladesh. *Am. J. Clin. Nutr.*, 56: 136-137.
- Zhang, W., J. Yan, J. Duo, B. Ren and J. Guo, 1989. Preliminary study of biochemical constituents of berry of sea buckthorn growing in Shanxi Province and their changing trend. *Proceeding of International Symposium on Sea Buckthorn (H. rhamnoides. L.)*, Xian, China.