



## Nutritional and functional properties of oats: An update

Seema Sangwan<sup>1</sup>, Rameshwar Singh<sup>2\*</sup>, Sudhir Kumar Tomar<sup>1</sup>

<sup>1</sup>Dairy Microbiology Division, National Dairy Research Institute, Karnal, India

<sup>2</sup>Directorate of knowledge management in Agriculture, Indian Council of Agriculture research, New Delhi, India

**Abstract:** With snowballing interest in eating for health in the developed world coupled with the endemic obesity problem, much attention is being directed towards delivering soluble fibers to the consumer through food. Cereal foods are obvious and major sources of healthy soluble and insoluble fiber grown over 73% of the total global harvested area. These contribute to over 60% of the world food production providing about 50 percent of protein and energy necessary for the human diet. Oats provide more protein, fiber, iron and zinc than other whole grains. They have high nutritive value both for people and animals because of good taste and an activity of stimulating metabolic changes in the body. Oats are said to be unique among cereals as they are therapeutically active against diabetes, dyslipidemia, hypertension, inflammatory state and vascular injury than other grains which are predominantly insoluble, such as wheat or rice. This review highlights the nutritional value of oats,  $\beta$ -glucan in them as biologically defense modifier, their mode of action against various diseases and enlists various fermented and non-fermented products of oats available in market.

**Keywords:** Hypertension;  $\beta$ -glucan; Diabetes; Dietary fibers; Cereals; health

Received: 06 January 2014 / Accepted: 27 January 2014 / Published Online: 30 March 2014

© 2014 jibresearch.com

Oats are unique among the cereals; one of the rich sources of dietary fibers among cereals belongs to the Poaceae family like all other grain varieties (Butt et al, 2008). The history of oats can be traced back to about 2000 B.C. As a cultivated crop, oats appear to be substantially later in origin than wheat and barley. The first evidence of oats was found in Egypt and among the lake dwellers of ancient Switzerland (Coffman 1961). Thirteen species and subspecies, all of the family Gramineae, genus *Avena*, are recognized as given in U.S. Department of Agriculture Technical Bulletin 1100. The most commonly cultivated and most popular species in the world today is *Avina sativa* making up more than 75% of the world's cultivars (Coffman 1961).

Oats are able to grow in acidic soil (pH up to 4.5) but perform better on soil having pH 5.3 to 5.7 (Alam and Adams 1979). They are mostly grown in cool moist climates and they are sensitive to hot, dry weather. For these reasons, world oat production is concentrated from the 40<sup>th</sup> latitude in the southern hemisphere to the 60<sup>th</sup> latitude in the northern hemisphere. Thus includes mainly Australia, China, North America,


Scandinavian, Russia and related countries (Schrickel 1986). European Union is the country that had by far the highest production of oats (8.7 million metric ton) followed by Russia, Canada, U.S.A., Poland and Finland (Agrostats 2009).

### Nutritional value of oats

Oats are generally considered "healthy", being touted commercially as nutritious which has led to wider appreciation of oats as human food. Oat grout or whole grains (after removal of hull) contain all three parts of the grain – the germ, endosperm and bran, rich in all valuable nutrients (Fig. 1.). In comparison to other cereals, these are characterized constitute large amount of total protein, carbohydrate (primary starch content), crude fat, dietary fibre (non-starch), unique antioxidants and considerable vitamins and mineral content (Table 1). A good taste and an activity of stimulating metabolic changes in the body make nutritive value of oats high for both people and animals (Brand and Merwe 1996; Lia et al. 1997; Peltonen-Sainio et al. 2004; Peterson 2004). Total carbohydrate content (including cellulose and non-starch polysaccharides)

### Corresponding Author:

Singh R. (✉) Directorate of knowledge management in Agriculture; Indian Council of Agriculture research; New Delhi, India  
Email: rsndri@gmail.com



**Figure 1** Structure of oat grain (quakeroats.com)

may reach 75-80% of the dry matter. The main component of oat is starch and its content depends on the variety and growing conditions. The amylase content of oat starch ranges from 16-18% to 28.5–28.7% (Mirmoghtadaie et al. 2009).

**Table 1** Nutritional composition of whole grain oat and oat bran (Usman et al, 2010)

Nutrients	Whole grain oat	Oat bran
Protein	15 - 17 %	15 - 18 %
Starch and sugars	59 - 70 %	10 - 50 %
Fat	~ 4.5 %	~ 6.5 %
Total dietary fibre	~ 12 %	~ 14-15 %
Ash	~ 3.5 %	~ 2.4 %
β-Glucan	2 - 6 %	5 - 20 %
Cellulose	~ 14 %	~ 2.5%
Lignin	~ 2.4 %	~ 4.5 %

Oats contain a high percentage of oat protein and balanced composition of amino acids (Table 2) which have proved them highly nutritive in comparison to other cereals (Brand and Merwe 1996; Petkov et al. 2001). Highest protein content occurs in oat grout (12.4 to 24.5%) among cereals (Miller et al. 1993; Robbins et al. 1971; Robert et al. 1983). Relatively high protein levels occur in the bran (about 20%) and lower in hull (less than 2%) (Youngs 1982). Oat is the only cereal in which the major portion of the grain protein is soluble in salt and thus classified as globulins and only a small proportion of water soluble albumins and alcohol soluble prolamin. Phenol compounds present in oats and its by-products have a considerable antioxidant potential (Sobotka et al. 2012). Biologically the most important metabolically active proteins of oat are the enzymes. The oat grout, similar to other cereal grains, contains numerous enzymes.

**Table 2** Amino acid composition of oats (USDA, 2005)

Amino acids in 1 cup oat (156 gm)	
Tryptophan	0.365 g
Threonine	0.897 g
Isoleucine	1.083 g
Leucine	2.003 g
Lysine	1.094 g
Methionine	0.487 g
Cystine	0.636 g
Phenylalanine	1.396 g
Tyrosine	0.894 g
Valine	1.462 g
Arginine	1.860 g
Histidine	0.632 g
Alanine	1.374 g
Aspartic acid	2.259 g
Glutamic acid	5.791 g
Glycine	1.312 g
Proline	1.457 g
Serine	1.170 g

Early investigations revealed the presence of proteases, maltase, α-amylase, lichenase, phenoxyacetylase and hydroxylase, phosphatase, tyrosinase and lipase as summarized by Caldwell and Pomeranz (1974). A high lipid content of oats makes them different from other cereals and is a good source of essential unsaturated fatty acids (Table 3). The percentage varies from 3.1 to 10.9% but in some high oil varieties it may be as high as 15% (Peterson and Wood, 1997; Schipper and Frey 1991). Triglycerides constitute the main component of lipids and phospholipids, glycolipids, sterols are also present in considerable quantities. The oat starch contains a high lipid content which plays an important role in decreasing the rate and amount of starch retro gradation in oats (Gudmunsson and Eliasson 1989; Paton 1986).

**Table 3** Lipid composition of oats (USDA, 2005)

Lipids in 1 cup oat (156 gm)	
Fatty acids, total saturated	1.899 g
12:0	0.037 g
14:0	0.023 g
16:0	1.613 g
18:0	0.101 g
Fatty acids, total monounsaturated	3.398 g
16:1 undifferentiated	0.020 g
18:1 undifferentiated	3.377 g
Fatty acids, total polyunsaturated	3.955 g
18:2 undifferentiated	3.781 g
18:3 undifferentiated	0.173 g
Cholesterol	0 mg

Dietary fibres (non-starch polysaccharides) also known as roughage or bulk are the edible parts of the plants and essential constituents of human nutrition. Although they are not a nutrient, they are nevertheless

an important component of our diets. Ingested dietary fibre moves along into the large intestine where it is partially or completely fermented by gut bacteria. During the fermentation process several by-products e.g. short chain fatty acids and gases are formed. It is the combined action of the fermentation process and by-products formed which contribute to the beneficial effects of dietary fibre on health. Oats comprises a very balanced profile of both soluble and insoluble dietary fibers. A high intake of dietary fibre is positively related to several preventive medical and nutritional effects (Spiller 2001) e.g. Dietary fibre complex with its antioxidants and other phytochemicals is most effective against cardiovascular disease and some types of cancer, lowering lipid levels (Jacobs et al. 1998a; Jacobs et al. 1998b; Jacobs et al. 1998; Slavin et al. 2000; Thompson 1994). Mineral content which is 2-3% in oat include phosphorus, potassium, magnesium, and calcium as main components as in other cereal (Table 4).

**Table 4** Mineral composition of oats (USDA, 2005)

Mineral content in 1 cup oat (156 gm)	
Calcium, Ca	84 mg
Iron, Fe	7.36 mg
Magnesium, Mg	276 mg
Phosphorus, P	816 mg
Potassium, K	669 mg
Sodium, Na	3 mg
Zinc, Zn	6.19 mg
Copper, Cu	0.977 mg
Manganese, Mn	7.669 mg


Water-soluble vitamin content of oat is summarized in Table V. The folate content of oat is about 20-30 µg/100 gm grain and the biotin concentration 10-15 µg/100 gm grain.

**Table 5** Vitamin composition of oats (USDA, 2005)

Vitamins content in 1 cup oat (156 gm)	
Vitamin C, total ascorbic acid	0.0 mg
Thiamin	1.190 mg
Riboflavin	0.217 mg
Niacin	1.499 mg
Pantothenic acid	2.104 mg
Vitamin B-6	0.186 mg
Folate, total	87 µg
Folic acid	0 µg
Folate, food	87 µg
Folate, DFE	87 µg
Vitamin B-12	0.00 µg
Vitamin B-12, added	0.00 µg
Vitamin A, IU	0 IU
Vitamin A, RAE	0 µg
Retinol	0 µg

### Other minor components

Oats contain various antioxidants that protect the lipids from oxidation and are important for the storage stability of various oat products. Oats are rich in tocopherols (Morrison et al. 1978; Youngs et al. 1986; Peterson 1995) which have antioxidant activities and contain about 2.3 mg tocopherols /100 gm grain. In addition to this, many low molecular weight phenolic compounds are also present in oat grains. Recent medical and nutritional investigations revealed the important role of natural plant antioxidants in the prevention of some diseases. Recent research has increased our knowledge about the phenol compounds in oats (Collins 1986; Dimberg et al. 2005). Ferulic acid, p-coumaric acid, vanillin, p-hydroxybenzoic acid are some of the compounds detected in oats. Some of these compounds (e.g., ferulic acid) are partly bound to cell wall polysaccharides or proteins. A new group of phenolic acids was obtained from oat grouts and hulls by aqueous alcoholic extraction (Collins et al. 1989). These acids occurred as conjugates linked covalently to the amine function of several different orthoamino-benzoic acids. The structure of one identified compound, designated avenalamic acid, is shown in Fig.2.




**Figure 2** Structure of Avenalamic acid

Another group of phenolic antioxidants present in oats are avenanthramides. These compounds are cinnamoyl-anthranilic acid derivatives.

The structure of two avenanthramides as elucidated by Dimberg et al (1993) is shown in Fig. 3. The antioxidative capacity of avenanthramide-2 is about 60% of that of α-tocopherol and seems to be quite heat stable. Caffeic and ferulic acid, known antioxidants, had lower activities. A comparison of 10 oat cultivars revealed that the total amount of two known avenanthramides varied between 40 and 132 µg/g grain. Oat grain also contains small quantities of saponins, designated as avenacoside. These are secondary metabolites which are derived from triterpenoid or steroid cyclisation products of 2, 3-oxidosqualene (Hostettman and Marston 1995). Since many saponins have potent antimicrobial activity the natural role of these molecules in plants is likely to be in conferring protection against attack by potential pathogens (Morrissey and Osbourn 1999). Phytic acid is the main phosphorus reservoir of oats. Lolas et al (1976) found a range of 0.82-1.01% phytic acid in 19 oat


cultivars. Miller et al (1980) reported a range of 0.95-1.41% phytic acid in oat grouts from seven cultivars grown for two years in soils with different phosphorus level. Low activity of phytase was found in oats using  $^{31}\text{P}$ -NMR spectroscopy because the phytase enzyme gets inactive by the heat treatment normally used during processing of oats for food products (Frolych et al. 1988). That is why; they are officially recognized as safe in contrast to other cereals (Lockhart and Hurt 1986).



**Figure 3** Structure of Aventhamides

#### Chemistry of $\beta$ -glucan from oats (a biological defense modifier)

$\beta$ -glucan, a type of soluble dietary fibre and a mixed-linkage polysaccharide of D-glucose units nutritionally potentiates and modulates the immune response. The bonds between D-glucopyranosyl units (Fig. 4) in  $\beta$ -glucans are primarily composed of either  $\beta$ -1,3 (cellotriosyl units) and  $\beta$ -1,4 linkages (cellotetraosyl units) but there are also regions that are more cellulose-like in character, with 4 or more consecutive  $\beta$ -1,4 linked glucose units (Wood et al. 1993). The molar ratio of celotriose to cellotetraose units (DP3:DP4) is typically 1.5 - 2.3 as far as oat  $\beta$ -glucan is considered (Miller and Fulch 1995).



**Figure 4** Linkage in Beta-glucan


The  $\beta$ -1, 3 linkages break up the uniform structure of the  $\beta$ -D-glucan molecule and make it soluble and flexible that is why it is a major component of the water extractable fibre fraction (Manthey et al. 1999). The

soluble  $\beta$ -glucans make viscous, shear thinning solutions even at low concentrations.

The viscosity is related to the molecular weight and is strongly dependent on the concentration (Åman et al. 2004; Anttila et al. 2004). Lyly et al (2004) observed that with an oat  $\beta$ -glucan preparation of high-molecular weight (average MW = 2,000,000), the maximum practicable concentration of  $\beta$ -glucan in a soup was 0.5 w-% by weight. With lower molecular weight preparations (MW  $\leq$  200,000) of oat, it was possible to add up to 2.0 w-%  $\beta$ -glucan to the soup.

#### Location of $\beta$ -glucan in oat grain

$\beta$ -glucan usually comprises 3.6–5.1 % of dry weight of the oat whole grain (Hampshire 2004). The several factors that affect the concentration of  $\beta$ -glucan include cultivar variation and the growing conditions e.g., nitrogen level, temperature and rainfall (Asp et al. 1992, Brunner and Freed 1994).  $\beta$ -glucans are located throughout the starchy endosperm. They are concentrated in the bran, more precisely in the aleurone and sub-aleurone layer (Fig. 5).



**Figure 5** Location of  $\beta$ -glucan in oat grain (oatsandhealth.org)

The content of dietary fibre and  $\beta$ -glucan varies between different oat products. Typical concentrations (% of dry weight) are presented below (Table 6).

#### Therapeutic attributes of oats against diabetes

**Table 6** Concentration of dietary fibre and  $\beta$ -glucan in different oat products


	Dietary fibre (%)	Beta-glucan (%)
Oat endosperm flour	5 - 10	1 - 3
Whole grain products	10 - 12	4 - 5
Conventional oat bran products	15 - 20	8 - 12
Oat bran concentrates	20 - 35	15 - 22
Beta-glucan isolates	80 - 100	Up to 80

## and its associated metabolic disorders

### Oat against type 2 diabetes

Insulin resistance, the main component and causative factor of metabolic syndrome associated with type 2 diabetes is an essential target in the therapeutic approach of oat  $\beta$ -glucans. The dietary intake of total as well as soluble and insoluble fiber is inversely associated with insulin resistance, supporting evidence that a high intake of dietary fiber is associated with enhanced insulin sensitivity (Galisteo 2008; Ylonen et al. 2003). Oat  $\beta$ -glucans have been used in several clinical trials to reduce glucose level in blood.

Oats act against diabetes and cause the lowering of blood glucose level through many ways (Fig. 6). Oat  $\beta$ -glucans favor increased glucose uptake into skeletal muscle and improves insulin sensitivity by delaying stomach emptying because of high viscosity and stability with pH (Butt et al. 2008) so that dietary glucose is absorbed more gradually (Kiho et al. 1995). The comparative reduction of glucagon response after the oat extracts may be partially responsible for the reduction in glucose concentrations (Judith et al. 1995). Another possible mechanism for oat  $\beta$ -glucans to reduce blood glucose level is mediated by signal pathway through PI3K/Akt activation.



**Figure 6** Mechanism of action of oats against diabetes

Studies showed that oat  $\beta$ -glucan (~5gm per day) decrease the glycemic response upto 50% after ingestion of 35 gm carbohydrate (Tappy et al. 1996) and in a 50 g carbohydrate portion each gram of  $\beta$ -glucan reduces the GI by 4 units (Jenkins et al. 2002). It has been also shown that oat bran flour was more effective than oat bran crisp explained by the three times higher  $\beta$ -glucan content in oat bran flour (Saris 2003; Tapola et al. 2005) against type 2 diabetes. After ingestion of the oat (bran flour or crisp), the blood glucose levels were lower at 15, 30, and 45 min but higher at 90 min after 12.5 g glucose loading (Tapola et al. 2005). These changes reduce the feeling of hunger caused by rapid decrease in blood glucose (Ludwig 2003; Saris 2003). Thus,  $\beta$ -glucans decrease appetite and reduce food intake. Zerm et al recently also reported that oatmeal diet days may improve insulin resistance in patients with type 2 diabetes mellitus (Zerm et al. 2013).

### Mechanism of action against type 2 diabetes

Decreased PI3K/Akt activity has been shown to play a key role in the pathogenesis of type 2 diabetes.  $\beta$ -glucans have been demonstrated to increase PI3K/Akt through several receptors (Hsu et al. 2002; Chen and Seviour 2007). These receptors stimulated by  $\beta$ -glucans include Dectin-1, complement receptor 3, lactosylceramide, scavenger and toll like receptors; each induces specific signal pathways. The interaction between  $\beta$ -glucans and their receptors has been studied by Brown (2006) and Chen and Seviour (2007). The administration of  $\beta$ -glucans could restore decreased PI3K in diabetes as mushroom extract have been shown to activate PI3K pathway (Li et al. 2006; Lee et al. 2008). These  $\beta$ -glucans can bind to dectin-1 receptor to stimulate the signal pathway (Underhill et al. 2005; Trinidad et al. 2006; Brown 2006; Olsson and Sundler 2007). Another  $\beta$ -glucan lentinan binds to scavenger receptors which can also activate PI3K pathway (Rice et al. 2002; Mineo et al. 2003).


A study in stroke-prone spontaneously hypertensive rats suggests that dietary fibre (mainly soluble)

supplementation effectively prevents insulin resistance by increasing skeletal muscle plasma membrane content of the insulin-responsive glucose transporter type (GLUT-4) by a mechanism different from the activation of the phosphatidylinositol-3 4 series of fatty acids stimulate peroxisome proliferator activated receptor PPAR $\gamma$ , whose activation has been reported to increase GLUT-4 content in adipocytes (Park et al. 1998). Therefore, it has been suggested and proved that short chain fatty acids (SCFAs) such as propionic and butyric acids, which result from anaerobic fermentation by bacteria of soluble fermentable dietary fibre in the colon (Cummings et al. 1987), increase muscle GLUT-4 via PPAR $\gamma$  (Songs et al. 2000).

also reported that oat bran high fibre diet significantly reduces plasma cholesterol level in hypercholesterolemic rats. Recently it was reported that instant oatmeal consumed daily for 6 weeks significantly increased fiber intake and decreased major risk factors for CVD in Chinese adults with hypercholesterolemia (Jhang et al. 2012). Rabey stated that oat bran appeared more efficient than barley bran in lowering the lipid profile levels in hypercholesterolemic rats (Rabey et al. 2013).

#### Mechanism of action against dyslipidemia

The mechanism of oat bran soluble fibres to lower LDL (Fig. 7) is considered to be mediated by bile acids binding property of  $\beta$ -glucans. Therefore  $\beta$ -glucans increase exclusion of bile acids (Marlett 1997;



**Figure 7** Action of Oat against dyslipidemia

#### Oat against dyslipidemia

$\beta$ -glucans have been shown to decrease LDL cholesterol and increase HDL to alleviate possibly dyslipidemia and reduce CVD (Anderson 1995; Reyna-Villasmil et al. 2007; Kapur et al. 2008). The required dose of  $\beta$ -glucan is 0.75 g/serving in food. Oats were first found to have a cholesterol-lowering effect and the active component was identified as  $\beta$ -glucans (Kerckhoffs et al. 2002). On the basis of numerous (Butt et al. 2008) clinical studies, the US Food and Drug Administration (FDA) permitted the use of a claim that oat soluble fibre has the ability to reduce the risk of heart disease. Oats reduced both serum total cholesterol and LDL cholesterol compared with control (Davidson et al. 1991; Van Horn et al. 1991; Gerhardt and Gallo 1998; Kerckhoffs et al. 2003; Karmally et al. 2005; Naumann et al. 2006; Reyna-Villasmil et al. 2007; Theuwissen and Mensink 2007). In 20 hypercholesterolemic male patients, oat bran was shown to be better than wheat bran in lowering cholesterol (Anderson et al. 1991). Usman et al (2010)

Ellegard and Anderson 2007) and this in turn activates cholesterol 7 $\alpha$ -hydroxylase and up-regulates low-density lipoprotein receptor (LDLR) and thus increase the transport of LDL into hepatocytes and the conversion of cholesterol into bile acids (Nilsson et al. 2007). It has been reported that every 1% reduction in LDL is associated with a decreased risk for CHD of 1% to 3% (Reynolds, 2000). In addition, oat  $\beta$ -glucans have also been shown to be fermented by human fecal micro biota to produce short-chain fatty acids such as acetate, propionate and butyrate and after absorption into the portal vein both acetate and propionate inhibit hepatic cholesterol synthesis (Wright et al. 1990). This in turn produces the hypocholesteolemic effect (Han et al. 2004; Držikova et al. 2005; Almingier and Eklund-Jonsson 2008; Hughes et al. 2008). The mechanism of oat bran soluble fibres to lower LDL is considered to be mediated by bile acids binding property of  $\beta$ -glucans. Therefore  $\beta$ -glucans increase exclusion of bile acids (Marlett 1997; Ellegard and

Anderson 2007) and this in turn activates cholesterol 7 $\alpha$ -hydroxylase and up-regulates low-density lipoprotein receptor (LDLR) and thus increase the transport of LDL into hepatocytes and the conversion of cholesterol into bile acids (Nilsson et al. 2007).


It has been reported that every 1% reduction in LDL is associated with a decreased risk for CHD of 1% to 3% (Reynolds 2000). In addition, oat  $\beta$ -glucans have also been shown to be fermented by human fecal micro biota to produce short-chain fatty acids such as acetate, propionate and butyrate and after absorption into the portal vein both acetate and propionate inhibit hepatic cholesterol synthesis (Wright et al. 1990). This in turn produces the hypocholesteolemic effect (Han et al. 2004; Drzikova et al. 2005; Alminger and Eklund- Jonsson 2008; Hughes et al. 2008).

### Oat against hypertension

A clinical trial with food containing oat  $\beta$ -glucans showed reduced blood pressure in the subjects with body mass index above medians (31.5 kg/m<sup>2</sup>) (Maki et al. 2007). This is further demonstrated by another trial that also showed effective role of oat against hypertension (He et al. 2004). However, no pure  $\beta$ -glucan has been used for these studies and thus it is difficult to identify which components are effective (Chen and Raymond 2008). In a randomized, double-blind, placebo-controlled trial in 110 participants aged 30 to 65 years who had untreated but higher than optimal blood pressure or Stage 1 hypertension (He et al. 2004), the effects of supplemented soluble fibre intake from oat bran on blood pressure for 12 weeks were analyzed. It was found that both systolic and diastolic blood pressure was significantly reduced in the study participants who consumed a high-fibre diet (mean reduction by 3.4 and 2.2 mmHg, respectively).

### Mechanism of action against hypertension

Increased body weight is a strong risk factor for hypertension (Neter et al. 2003). Several mechanisms for a potential effect of dietary fibre intake on blood pressure have been hypothesized. Dietary fibre has numerous effects on the digestion and absorption of



**Figure 8** Oat actions against Proinflammation

foods. Insulin resistance and its concomitant compensatory hyperinsulinemia have been suggested as major underlying pathogenesis mechanisms for the development of hypertension (Ferrannini et al. 1987). In this way, effectiveness of both soluble and insoluble fibres in reducing insulin resistance and insulin levels in both diabetic and healthy persons (McKeown et al. 2002, King et al. 2005, Qi et al. 2005) would contribute to treating or preventing hypertension. Intake of dietary fibre has also been shown to efficiently reduce body weight (Rigaud et al. 1990) and weight gain (Liu et al. 2003).

### Oat against proinflammatory state

There are interesting reports about the effects of dietary fibre intake on the acute-phase response marker CRP and the inflammatory cytokines IL-6, IL-18 and TNF- $\alpha$  whose levels are found abnormally elevated in the type 2 diabetes. Some studies have suggested an association between consumption of dietary fibre and the anti-inflammatory adiponectin. A study in diabetic women (Qi et al. 2006) indicated that after adjustment for age, body mass index (BMI), lifestyle and dietary covariates, intake of cereal fibres was associated with significantly decreasing trends of CRP (C reactive protein) and TNF- $\alpha$  receptor 2 among women with type 2 diabetes.

### Mechanism of action against proinflammatory state

Oat intake effect favoring weight loss could be involved as most weight loss studies have shown that decreasing body weight was related to decreases in CRP (Dietrich et al. 2005), TNF- $\alpha$  (Dandona et al. 1998) and to increases in adiponectin circulating levels of obese women (Esposito et al. 2003). In a study, King (2005) suggested that dietary fibre decreases lipid oxidation, which in turn is associated with decreased inflammation. These anti-inflammatory effects of dietary fibre could also be attributed, in the case of fermentable fibres, to the production of butyrate (Fig. 8). Clinical trials suggest that butyrate has anti-inflammatory properties in pathologies such as inflammatory bowel disease (Patz et al. 1996), and it has been described for anti-inflammatory activity in different human cellular types such as macrophages and monocytes (Segain et al. 2000).

### Oat against vascular injury

Vascular injury may contribute to the pathogenesis of CVD (Liuba and Pesonen 2005; Kibos et al. 2007) associated with type 2 diabetes. Dietary fibres may increase collagen synthesis (Wei et al. 2002). The study has shown that  $\beta$ -glucan can help in healing of wound in db/db mice (Berdal et al. 2007).

### Oat based product

Oats usage in human foods has increased as information on oats beneficial nutritional properties has come to light. The authorization of the heart health claim for oats in the United States by the FDA is especially

significant. Additionally, recent investigations on the health implications of minor oat constituents such as avenanthramides have raised hopes that the nutritional benefits of oats in human diets may go well beyond those currently recognized. There are many non-fermented and fermented oat based products are available in the market (Table 7).

Before oats are used to produce oat products, the hulls are removed, leaving the oat grout. Traditional commercial non-fermented oat products include rolled oats (whole-grain flakes), steel-cut grouts, quick oat

mainly due to lack of acceptability with an appearance, texture and sensory quality mainly the beany flavor associated with these products. The sensory attributes play pivotal role in consumer acceptance of novel functional foods. Once the healthful effects of such foods are validated and established through systematically designed studies using small animal models and human trials, their acceptability can be increased by flavoring these products using fruits, various artificial sugars and by modifying their texture using various stabilizers.

**Table 7** Fermented oats based products

Fermented oat products	Manufacturer	
Oatly	CeBa Food AB, Lund	Enzymatic hydrolysis of oat kernals, milk-free product
Oatwell	CreaNutrition AG, Switzerland Swedish oat fibre AB, Sweden	Oat bran
Primaliv™ Oatrim	Skåne mejerier, Malmö, Sweden ConAgra, Quaker Oats, Röhne-Poulenc	Yoghurt containing oat fibres Enzymatic hydrolysis of oat flour
YOSA®	Bioferme, Finland	fermented with a combination of probiotic bacteria i.e. <i>Bifidobacterium lactis</i> BB-12™ and <i>Lactobacillus acidophilus</i> LA-5™
Velle	Velle Oats, St. Petersburg, Russia	Oat Fermentation with Lactic acid bacteria
Oat based fermented product	Martensson <i>et al</i> , 2001	Fermented with commercial yoghurt starter cultures
Oat Bio Lacto	Bekers <i>et al</i> , 2001	Fermented with probiotic lactic acid bacteria i.e. <i>Lactobacillus acidophilus</i> T <sub>20</sub> and <i>Bifidobacteria</i> and commercial starter culture ABT-1
Oat based probiotic drink	Angelov <i>et al</i> , 2006	Fermented with Lactic acid bacteria <i>Lactobacillus plantarum</i>

flakes, baby oat flakes, instant oats, oat bran, and oat flour.

### Conclusion

Due to increased cost and side-effects of medication, consumer's interest is snowballing towards dietary modifications for improvement of health. It is a well documented fact that oats are effective against common disorders of lifestyle *in vogue*. Oats which earlier were grown as forage and fodder are a unique combinations of proteins, fats, dietary fibres, minerals, vitamins and alkaloid polyphenols (avenanthramides). They act as good substrate for the growth of pro-biotic lactic acid bacteria which enhance their potential to improve the health and well being of consumers. Though lactic acid fermentation of oats is an ancient processing method yet most of the oat based products available in market are non-fermented and even fermented ones are invariably non dairy based. Nevertheless, limited information is available regarding the development of functional food product having oats and pro-biotic in a completely milk-based product

### References

- Alam S M, Adams W A (1979) Effect of soil pH on the growth and mineral content of oats. *Pak J Sci Ind Res* 22: 147-151
- Alminger M, Eklund-Jonsson C (2008) Whole-grain cereal products based on a high- fibre barley or oat genotype lower post-prandial glucose and insulin responses in healthy humans. *Euop J Nutri* 47: 294-300
- Åman P, Rimsten L, Andersson R (2004) Molecular weight distribution of  $\beta$ -glucan in oat-based foods. *Cereal Chem* 81: 356-360
- Anderson J W (1995) Dietary fibre, complex carbohydrate and coronary artery disease. *Can J Cardiol* 11: 55G-62G
- Anderson J W, Gilinsky N H, Deakins D A, Smith S F, Oneal D S, Dillon D W, Oeltgen P R (1991) Lipid responses of hypercholesterolemic men to oat-bran and wheat-bran intake. *Am J Clin Nutri* 54: 678-683
- Angelov A, Gotcheva V, Kuncheva R, Hristozova T (2006) Development of a new oat-based probiotic drink. *Inter J Food Microbiol* 112(1): 75-80

- Anttila H, Sontag-Strohm T, Salovaara H (2004) Viscosity of beta-glucan in oat products. *Agri Food Sci* 13: 80-87
- Asp N G, Mattsson B, Onning G (1992) Variation in dietary fibre,  $\beta$ -glucan, starch, protein, fat and hull content of oats grown in Sweden 1987-1989) *Europ J Clin Nutr* 46: 31-37
- Bekers M, Marauska M, Laukevics J, Grube M, Vigants A, Karklina D, Skudra L, Viesturset U (2001) Oats and fat-free milk based functional food product. *Food Biotechnol* 15: 1 – 12
- Berdal M, Appelbom H I, Eikrem J H, Lund A, Zykova S, Busund L T, Seljelid R, Jenssen T (2007) Aminated  $\beta$ -1,3-Dglucan improves wound healing in diabetic db/db mice. *Wound Repair Regen* 15: 825-832
- Brand T S, Merwe J P (1996) Naked oats (*Avena nuda*) as a substitute for maize in diets for weanling and grower-finisher pigs. *Anim Feed Sci Technol* 57: 139-147
- Brown G D (2006) Dectin-1: a signalling non-TLR pattern-recognition receptor. *Nature Review Immunol* 6: 33-43
- Brunner B R, Freed R D (1994) Oat grain  $\beta$ -glucan content as affected by nitrogen level, location, and year. *Crop Sci* 34: 473-476
- Butt M S, Nadeem M T, Khan M K I, Shabir R (2008) Oat : unique among the cereals. *European J Nutrition*, 47: 68-79.
- Caldwell E F, Pomeranz Y (1973) Industrial uses of cereals: Oats. *Industrial Uses of Cereals*. Y. Pomeranz, ed. Am Assoc Cereal Chem, St. Paul, MN. 393-411
- Chen J, Raymond K (2008) B-glucans in the treatment of diabetes and associated cardiovascular risks. *Vascu Health Risk Manag*, 4(6): 1265-1272
- Chen J, Seviour R (2007) Medicinal importance of fungal  $\beta$ -(1-3) and (1-6)-glucans. *Mycological Res* 111: 635-52
- Coffman F A (1961) Oats and oats improvement. *Am society Agronomy Madison, Wisconsin*, 8: 263-329
- Collins F W (1989) Oat phenolics: Avenanthramides, novel substituted N- cinnamoylanthranilate alkaloids from oat groats and hulls. *J Agri Food Chem* 37: 60-66
- Collins F W (1986) Oat phenolics: structure, occurrence and function. *Oats: Chem and technology*. Webster, F.H (ed), Am Asso Cereal Chemists, St Paul, Minnesota, 227-295
- Dandona P, Weinstock R, Thusu K, Abdel-Rahman E, Aljada A, Wadden T (1998) Tumor necrosis factor- $\alpha$  in sera of obese patients: fall with weight loss. *J Clini Endocrinol Meta* 83: 2907-2910
- Davidson M H, Dugan L D, Burns J H, Bova J, Story k, Drennan K B (1991) The hypocholesterolemic effects of  $\beta$ -glucan in oatmeal and oat bran. A dose-controlled study. *JAM Medical Association* 265: 1833-1839
- Dietrich M, Jialal I (2005) The effect of weight loss on a stable biomarker of inflammation, C-reactive protein. *Nutri Reviews* 63: 22-28
- Dimberg L H, Gissén C, Nilsson J (2005) Phenolic compounds in oat grains (*Avena sativa* L.) grown in conventional and organic systems. *Am bio* 34: 331-337
- Drzikova B, Dongowski G, Gebhardt E, Habel A (2005) The composition of dietary fibre rich extrudates from oat affects bile acid binding and fermentation in vitro. *Food Chem* 90: 181-192
- Ellegard L, Andersson H (2007) Oat bran rapidly increases bile acid excretion and bile acid synthesis: an ileostomy study. *European. J Clinical Nutri* 61: 938-945
- Esposito K, Pontillo A, Di Palo C, Giugliano F, Masella M, Marfella R, Giugliano D (2003) Effect of weight loss and lifestyle changes on vascular inflammatory markers in obese women. *J Am Med Assoc* 289: 1799-1804
- Ferrannini E, Buzzigoli G, Bonadonna R, Giorico M A, Oleggini M, Graziadei L, Pedrinelli R, Brandi L, & Bevilacqua S (1987) Insulin resistance in essential hypertension. *The New Eng J Medicine* 317: 350-357
- Frolych W, Wahlgren M, Drankenberg T (1998) Studies on phytase activity in oats and wheat using  $^{31}\text{P}$ -NMR spectroscopy. *J Cereal Sci* 8: 47-53
- Galisteo M, Duarte J, Zarzuelo A (2008) Effects of dietary fibres on disturbances clustered in the metabolic syndrome. *J Nutritional BioChem* 19: 71-84
- Gerhardt A L, Gallo N B (1998) Full-fat rice bran and oat bran similarly reduce hypercholesterolemia in humans. *J Nutrition* 128: 865-869
- Gudmunsson M, Eliasson A C (1989) Some physicochemical properties of oat starches extracted from varieties with different oil content. *Acta Agri Scandinavia* 39: 101-111
- Hampshire J (2004) Variation in the content of nutrients in oats and its relevance for the production of cereal products. *Proceedings of the 7th International Oat Conference*. Helsinki, Finland. Peltonen-Sainio, P., Topi-Hulmi, M. (eds.) Jokioinen: MTT Agrifood Research Finland, Agrifood Research Reports 51: 126
- Han K H, Sekikawa M, Shimada K, Sasaki K, Ohba K, Fukushima M (2004) Resistant starch fraction prepared from kintoki bean affects gene expression of genes associated with cholesterol metabolism in rats. *Experimental Biol Med*. (Maywood) 229: 787-792
- He J, Streiffer R H, Muntner P, Krousel-Wood M A, Whelton P K (2004) Effect of dietary fibre intake on blood pressure: a randomized, double-blind, placebo-controlled trial. *J Hypertension* 22: 73-80
- Hostettmann K A, Marston A (1995) *Saponins. Chem and Pharmacology of Natural Products*, Cambridge, UK: Cambridge University Press
- Hsu M J, Lee S S, Lin W W (2002) Polysaccharide purified from *Ganoderma lucidum* inhibits spontaneous and Fas-mediated apoptosis in human neutrophils through activation of the phosphatidylinositol 3 kinase/Akt signaling pathway. *J Leukocyte Biology* 72: 207-216
- Hughes S A, Shewry P R, Gibson G R, McCleary B V, Rastall R A (2008) In vitro fermentation of oat and barley derived  $\beta$ -glucans by human faecal microbiota. *FEMS Microbiol Ecol* 64: 482-493
- Jacobs D R, Marquart L, Slavin J, Kushi L H (1998b) Whole grain intake may reduce the risk of ischemic heart disease death in postmenopausal women: The Iowa Women's Health Study. *Am J Clin Nutr* 68: 248-257
- Jacobs D R, Marquart L, Slavin J, Kushi L H (1998a) Whole-grain intake and cancer: an expanded review and meta-analysis. *Nutri Cancer* 30: 85-96
- Jacobs D R, Meyer K A, Kushi L H, Folsom A R (1998) Whole-grain intake may reduce the risk of ischemic heart disease death in postmenopausal women: the Iowa Women's Health Study. *Am J Clin Nutr* 68: 248-257

- Jenkins A L, Jenkins D J, Zdravkovic U (2002) Depression of the glycemic index by high levels of  $\beta$ -glucan fiber in two functional foods tested in type 2 diabetes. *Europ J Clinical Nutrition* 56: 622–628
- Judith H, Daniel J S, Behall K M (1995) Diets containing soluble oat extracts improve glucose and insulin responses of moderately hypercholesterolemic men and women. *Am J Clin Nutri* 61: 379–384
- Kapur N K, Ashen D, Blumenthal R S (2008) High density lipoprotein cholesterol: an evolving target of therapy in the management of cardiovascular disease. *Vascular Health Risk Manag* 4: 39–57
- Karmally W, Montez M G, Palmas W, Martinez W, Branstetter A, Ramakrishna R, Holleran S F, Haffner S M, Ginsberg H N (2005) Cholesterol-lowering benefits of oat-containing cereal in Hispanic Ams. *J Am Dietetic Assoc* 105: 967–970
- Kerckhoffs D A, Brouns F, Hornstra G, Mensink R P (2002) Effects on the human serum lipoprotein profile of  $\beta$ -glucan, soy protein and isoflavones, plant sterols and stanols, garlic and tocotrienols. *J Nutrition* 132: 2494–2505
- Kerckhoffs D A, Hornstra G, Mensink R P (2003) Cholesterol-lowering effect of  $\beta$ -glucan from oat bran in mildly hypercholesterolemic subjects may decrease when  $\beta$ -glucan is incorporated into bread and cookies. *Am J Clin Nutri* 78: 221–217
- Kibos A, Campeanu A, Tintoiu I (2007) Pathophysiology of coronary artery in-stent restenosis. *Acute Card Care* 9: 111–119
- Kiho T, Sobue S, Ukai S (1995) Polysaccharides in fungi: antidiabetic activity of an acidic polysaccharide from the fruiting bodies of *Tremella aurantia*. *Biological Pharma Bulle* 18: 1627–1629
- King D (2005) Dietary fibre, inflammation, and cardiovascular disease. *Mol Nutri Food Res* 49: 594–600
- Laszity R (1998) Oat grain—a wonderful reservoir of natural nutrients and biologically active substances. *Food Reviews Intern* 14: 19–199
- Lee J Y, Kim J Y, Lee Y G, Rhee M H, Hong E K, Cho J Y (2008) Molecular mechanism of macrophage activation by exopolysaccharides from liquid culture of *Lentinus edodes*. *J Microbiol Biotechnol* 18: 355–364
- Li B, Allendorf D J, Hansen R, Marroquin J, Ding C, Cramer D, Yan J (2006) Yeast  $\beta$ -glucan amplifies phagocyte killing of iC3b-opsonized tumor cells via complement receptor 3-Syk-phosphatidylinositol 3-kinase pathway. *J Immunol* 177: 1661–1669
- Lia A, Andersson H, Mekki N, Juhel C, Senft M, Lairon D (1997) Postprandial lipemia in relation to sterol and fat excretion in ileostomy subjects given oatbran and wheat test meals. *Am J Clin Nutri* 66: 357–365
- Liu S, Willett W C, Manson J E, Hu F B, Rosner B, Colditz G (2003) Relation between changes in intakes of dietary fibre and grain products and changes in weight and development of obesity among middle-aged women. *Am J Clin Nutri* 78: 920–927
- Liuba P, Pesonen E (2005) Infection and early atherosclerosis: does the evidence support causation. *Acta Paediatrica* 94: 643–51
- Lockhart H B, Hurt D H (1986) Nutrition in oats. In: *Oats: Chem and technology*. (F.H. Webster ed.) *Am Asso Cereal Chem* 297–308
- Lolas G M, Palambidis N, Markakis P (1976) *Am Association of Cereal Chemists* 53: 867–870
- Ludwig D S (2003) Dietary glycemic index and the regulation of body weight. *Lipids* 38: 117–121
- Lyly M, Salmenkallio-Marttila M, Suortti T, Autio K, Poutanen K, Lahteenmaki L (2004) The sensory characteristics and rheological properties of soups containing oat and barley beta-glucan before and after freezing. *Lebensm-Wiss Technology* 37: 749–761
- Maki K C, Galant R, Samuel P, Tesser J, Witchger M S, Ribaya-Mercado J D, Blumberg J B, Geohas J (2007) Effects of consuming foods containing oat  $\beta$ -glucan on blood pressure, carbohydrate metabolism and biomarkers of oxidative stress in men and women with elevated blood pressure. *European J Clinical Nutrition* 61: 786–795
- Manthey F A, Hareland G A, Huseby D J (1999) Soluble and insoluble dietary fiber content and composition in oat. *Cereal Chem* 76: 417–420.
- Marlett J A (1997) Sites and mechanisms for the hypocholesterolemic actions of soluble dietary fiber sources. *Advances in Exp Med and Biol* 427: 109–121
- Martensson O, Andersson C, Andersson K, Oste R, Holst O (2001) Formulation of an oat-based fermented product and its comparison with yoghurt. *Jr Sci of Food and Agr* 81: 1314–1321
- McKeown N M, Meigs J B, Liu S, Wilson P W, Jacques P F (2002) Wholegrain intake is favourably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study. *Am J Clin Nutri* 76: 390–398
- Miller S S, Wood P I, Pietrzak L N, Fulcher R G (1993) Mixed Linkage beta-Glucan, Protein Content, and Kernel Weight in *Avena* Species. *Cereal Chem* 70: 231–233
- Miller G A, Youngs V L, Oplinger E S (1980) Environmental and cultivar effects on oat phytic acid concentration. *Cereal Chem* 57: 189–191
- Miller S S, Fulcher R G (1995) Oat endosperm cell walls: II. Hot-water solubilization and enzymatic digestion of the wall. *Cereal Chem* 72: 428–432
- Miller S S, Fulcher R G, Sen A, Arnason J T (1995) Oat endosperm cell walls: Isolation, composition, and comparison with other tissues. *Cereal Chem* 72: 421–427
- Mineo C, Yuhanna I S, Quon M J, Shaul P W (2003) High density lipoprotein induced endothelial nitric-oxide synthase activation is mediated by Akt and MAP kinases. *J Biological Chem* 278: 9142–9149
- Mirmoghaddae L, Kadivar M, Shahedi M (2009) Effects of cross-linking and acetylation on oat starch properties. *Food Chem* 116: 709–713
- Morrison W R (1978) Cereal lipids. *Advances in Cereal Sci and Technology* 2: 221–348
- Morrissey J P, Osbourn A E (1999) Fungal resistance to plant antibiotics as a mechanism of pathogenesis. *Microbiology and Molecular Biology Reviews* 63: 708–724
- Naumann E, Van-Rees A B, Onning G, Oste R, Wydra M, Mensink R P (2006)  $\beta$ -glucan incorporated into a fruit drink effectively lowers serum LDL cholesterol concentrations. *Am J Clin Nutri* 83: 601–615

- Nefer J E, Stam B E, Kok F J, Grobbee D E, Geleijnse J M (2003) Influence of weight reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 42: 878–884
- Nilsson L M, Abrahamsson A, Sahlin S, Gustafsson U, Angelin B, Parini P, Einarsson C (2007) Bile acids and lipoprotein metabolism: effects of cholestyramine and chenodeoxycholic acid on human hepatic mRNA expression. *Biochem and Biophys Res Commun* 357: 707–711
- Olsson S, Sundler R (2007) The macrophage  $\beta$ -glucan receptor mediates arachidonate release induced by zymosan: essential role for Src family kinases. *Mol Immunol* 44: 1509–1515.
- Park K S, Ciaraldi T P, Lindgren K, Abrams-Carter L, Mudaliar S, Nikoulina S E (1998) Troglitazone effects on gene expression in human skeletal muscle of type II diabetes involve up-regulation of peroxisome proliferator-activated receptor- $\gamma$ . *J Clinical Endocrin Metabol* 83: 2830–2835
- Paton D, (1986) Oat starch: Physical, chemical, and structural properties. *Oats: Chem and Technology*. Am Association of Cereal Chem 93-120
- Patz J, Jacobsohn W Z, Gottschalk-Sabag S, Zeides S, Braverman D Z (1996) Treatment of refractory distal ulcerative colitis with short chain fatty acid enemas. *Am J Gastroent* 91: 731–734
- Peltonen-Sainio P, Kontturi M, Rajala A (2004) Impact dehulling oat grain to improve quality of on-farm produced feed. I. Hullability and associated changes in nutritive value and energy content. *Agricultural and Food Sci* 13: 18–28
- Peterson D M, Wood D F (1997) Composition and structure of high-oil oat. *J Cereal Sci* 26: 121–128
- Peterson D M (1995) Oat tocots: Concentration and stability in oat products and distribution within the kernel. *Cereal Chem* 72: 21-24
- Peterson D M (2004) Oat – a multifunctional grain. *Agri food Res Rep* 51: 21-26
- Petkov K, Biel W, Kowieska A, Jaskowska I (2001) The composition and nutritive value of naked oat grain (*Avena sativa* var. *nuda*). *J Animal and Feed Scis* 10 (2): 303–307
- Qi L, Rimm E, Liu S, Rifai N, Hu F B (2005) Dietary glycemic index, glycemic load, cereal fibre, and plasma adiponectin concentration in diabetic men. *Diabetes Care* 28: 1022–1028
- Qi L, Van Dam R M, Liu S, Franz M, Mantzoros C, Hu F B (2006) Wholegrain, bran, and cereal fibre intakes and markers of systemic inflammation in diabetic women. *Diabetes Care* 29: 207–211
- Rabey H, Al-seeni M, Amer H (2013) Efficiency of Barley bran and oat bran in ameliorating blood lipid profile and the adverse histological changes in hypercholesterolemic male rats. *Bio Med Res Int* 2013: 1-10
- Reyna-Villasmil N, Bermudez-Pirela V, Mengual-Moreno E, Arias N, Cano-Ponce C, Leal-Gonzalez E, Souki A, Inglett G E, Israili Z H, Hernández-Hernández R, Valasco M, Arraiz N (2007) Oat-derived  $\beta$ -glucan significantly improves HDLC and diminishes LDLC and non-HDL cholesterol in overweight individuals with mild hypercholesterolemia. *Am J Therap* 14: 203–212
- Reynolds H R, Quite E, Hunninghake D B (2000) Whole grain oat cereal lowers serum lipids. *Topics Clin Nutri* 15: 74-83
- Rice P J, Kelley J L, Kogan G, Ensley H E, Kalbfleisch J H, Browder I W, Williams D L (2002) Human monocyte scavenger receptors are pattern recognition receptors for (1–3)- $\beta$ -D-glucans. *J Leukocyte Biol* 72: 140–146
- Rigaud D, Rytting K R, Angel L A, Apfelbaum M (1990) Overweight treated with energy restriction and a dietary fibre supplement: a 6-month randomized, double-blind, placebo-controlled trial. *Int J Obesity* 14: 763–769
- Robbins G S, Pomeranz Y, Briggles L W (1971) Amino acid composition of oat groats. *J Agri and Food Chem* 19: 536–539
- Robert L S, Matlashewski G J, Adeli K, Nosolillo C, Altosaar I (1983) Electrophoretic and developmental characteristic of oat globulin in cultivars of different protein content. *Cereal Chem* 60: 231-234
- Saris W H (2003) Glycemic carbohydrate and body weight regulation. *Nutri Reviews* 61: S10–S16
- Schipper H, Frey K J (1991) Three recurrent selection regimes to increase oat-oil content of oat. *Crop Sci* 31: 1505-1510
- Schricket D J (1986) Oats production, value and use. *Oats: Chem and technol* 1-11
- Segain J P, Raingeard De La, Bletiere D, Bourreille A, Leray V, Gervois N, Rosales C, Ferrier L, Bonnet C, Blottière H M, Galmiche J P (2000) Butyrate inhibits inflammatory responses through NF kappa B inhibition: implications for Crohn's disease. *Gut* 47: 397–403
- Slavin J, Marquart L, Jacobs D (2000) Consumption of whole grain foods and decreased risk of cancer: Proposed mechanisms. *Cereal Foods World* 45: 54–58
- Sobotka W, Flis M, Antoszkiewicz Z, Lipinski K, Zdunczyk Z (2012) Effect of oat by-product antioxidants and vitamin E on the oxidative stability of pork from pigs fed diets supplemented with linseed oil. *Arch Anim Nutr* 66(1): 27-38
- Song Y J, Sawamura M, Ikeda K, Igawa S, Yamori Y (2000) Soluble dietary fibre improves insulin sensitivity by increasing muscle GLUT-4 content in stroke-prone spontaneously hypertensive rats. *Clin and Exp Pharmacol and Physiol* 27: 41–45
- Tapola N, Karvonen H, Niskanen L, Mikola M, Sarkkinen E (2005) Glycemic Response of Oat-bran products in type 2 diabetic patients. *Nutrition, Metabolism and Cardiovascular diseases* 15(4): 255-261
- Tappy L, Gugolz E, Wursch P (1996) Effects of breakfast cereals containing various amounts of  $\beta$ -glucan fibers on plasma glucose and insulin responses in NIDDM subjects. *Diabetes Care* 19: 831–834
- Theuwissen E, Mensink R P (2007) Simultaneous intake of  $\beta$ -glucan and plant stanol esters affects lipid metabolism in slightly hypercholesterolemic subjects. *J Nutrition* 137: 583–588
- Thompson L U (1994) Antioxidants and hormone-mediated health benefits of whole grains. *Critical Reviews in Food Sci and Nutri* 34: 473–497
- Trinidad A G, De-la Puerta M L, Fernandez N, Bayón Y, Crespo M S, Alonso A (2006) Coupling of C3b1 to IgG inhibits the tyrosine phosphorylation signaling cascade downstream

- Syk and reduces cytokine induction in monocytes. *J Leukocyte Biology* 79: 1073–1082
- Underhill D M, Rossnagle E, Lowell C A, Simmons R M (2005) Dectin-1 activates Syk tyrosine kinase in a dynamic subset of macrophages for reactive oxygen production. *Blood* 106: 2543–2550
- Usman S, Nazir S, Ali S, Nasreen Z, Najim A (2010) Determination of biochemical composition of *Avena sativa* (Oat) and to estimate the effect of high fibre diet on hypercholesteromic rats. *Bangladesh Res Publ J* 4: 312–319
- Van-Horn L, Moag-Stahlberg A, Liu K A, Ballew C, Ruth K, Hughes R, Stamler J (1991) Effects on serum lipids of adding instant oats to usual Am diets. *Am J Pub Health* 81: 183–188
- Wei D, Zhang L, Williams D L, Browder I W (2002) Glucan stimulates human dermal fibroblast collagen biosynthesis through a nuclear factor-1 dependent mechanism. *Wound Repair and Regeneration* 10: 161–168
- Wood P J (1993) Physicochemical characteristics and physiological properties of oat (1→3),(1→4)-β-D-glucan. *Am Association of Cereal Chem* 83:112
- Woods M N, Gorbach S L (2001) Influence of fiber on the ecology of the intestinal flora. *CRC Handbook of Dietary Fiber in Human Nutri* CRC Press 361–370
- Wright R W, Anderson J W, Bridges S R (1990) Propionate inhibits hepatic lipid synthesis. *Proceeding from the Society for Exp Biol and Med* 195: 26–29
- Ylonen K, Saloranta C, Kronberg-Kippila C, Groop L, Aro A, Virtanen M (2003) Associations of dietary fibre with glucose metabolism in non-diabetic relatives of subjects with type 2 diabetes. *Diabetes care* 26: 1979–1985
- Youngs V L (1986) *Oat: Chem and Technology*, AACC, St. Paul
- Zerm R, Helbrecht B, Jecht M, Hein A, Millet E, Girke M, Kroz m (2013) Oatmeal diet days may improve insulin resistance in patients with type 2 diabetes mellitus. *Forsch Complement med* 20: 465–468
- Zhang J, Li L, Song P, Wang C, Man Q, Meng L, Cai J, Kurilich A (2012) Randomized controlled trial of oatmeal consumption versus noodle consumption on blood lipids of urban Chinese adults with hypercholesterolemia. *Nutri J* 11: 54