



The application of predictive modelling of species distribution to biodiversity conservation

Jon Paul Rodríguez^{1*}, Lluís Brotons^{2,3}, Javier Bustamante⁴ and Javier Seoane⁵

¹Centro de Ecología, Instituto Venezolano de Investigaciones Científicas, Apdo. 21827, Caracas 1020-A, Venezuela, ²Centre Tecnològic Forestal de Catalunya, Àrea de Biodiversitat, Pujada del Seminari s/n, 25280 Solsona, Spain, ³Institut Català d'Ornitologia, Museu de Zoologia, Passeig Picasso s/n, 08003 Barcelona, Catalonia, Spain, ⁴Department of Applied Biology, Estación Biológica de Doñana, Consejo Superior de Investigaciones Científicas, Avda. María Luisa S/N, 41013 Sevilla, Spain, ⁵Departamento Interuniversitario de Ecología, Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Madrid, Spain

Keywords

Biodiversity, conservation biology, conservation planning, ecological data, environmental cartography, global change, invasive species, modelling, reserve design, surveys.

*Correspondence: Jon Paul Rodríguez, Centro de Ecología, Instituto Venezolano de Investigaciones Científicas, Apdo. 21827, Caracas 1020-A, Venezuela. E-mail: jonpaul@ivic.ve

INTRODUCTION

Predictive modelling of species geographical distributions is a thriving ecological and biogeographical discipline. Major advances in its conceptual foundation and applications have taken place recently, as well as the delineation of the outstanding challenges still to be met (Araújo & Guisán, 2006; Guisan *et al.*, 2006; Moisen *et al.*, 2006; accompanying papers). In this article, we focus on the application of predictive distribution modelling to biodiversity conservation. We begin by outlining a series of 'real-life' conservation problems that have been addressed with predictive modelling. Next, we consider the need to integrate these techniques with the development of systematic baseline data sets and models, in order to effectively monitor biodiversity change. The following section considers the availability of remotely sensed ecological data for use in predictive modelling, and introduces the readers to their major on-line sources. These general remarks provide a backdrop for a series of articles collected in this issue of *Diversity and Distributions* that emerged from a workshop on *Predictive Modelling of Species Distributions, New Tools for the XXI Century*, carried out under the auspices of the Universidad Internacional de Andalucía, sede Antonio Machado in Baeza, Spain, on 2–4 November 2005. The final section of this article provides an introduction to the collection of papers.

APPLICATIONS TO 'REAL-LIFE' PROBLEMS

Keeping up-to-date with the conceptual and technical advances of predictive modelling of species distribution is a challenge in itself, because it comprises a rapidly evolving set of versatile tools with a wide range of applications (Guisan & Zimmermann, 2000; Guisan & Thuiller, 2005). Predictive models have been successfully used in a variety of conservation biology studies, with the main purpose of addressing pressing conservation problems (e.g.

Brotons *et al.*, 2004; Linkie *et al.*, 2006). Confronted with extremely limited data on the distribution, abundance, and dynamics of most species on Earth (Rodríguez, 2003; Mace *et al.*, 2005), models allow for the extrapolation of relatively few field samples and museum records to the entire potential range of a species, thus creating predicted distribution maps (e.g. Guisan & Zimmermann, 2000; Heglund, 2002; Anderson *et al.*, 2003; Guisan & Thuiller, 2005). To carry out this task, detailed and reliable environmental data are needed to establish the species–environment relationships required to estimate species distribution outside sampling sites. The collection of field data using environmentally stratified sampling designs assures that the range of environmental variation is covered, and improves these predictions (Austin & Heyligers, 1991; Margules & Austin, 1991; Hortal *et al.*, 2001; Hortal & Lobo, 2005; J.R. Ferrer-Paris & J.P. Rodríguez, unpubl. data). Once data on species distribution have been collected in the field or derived from museum collections, and adequate environmental models have been fitted to the data, it is possible to generate predictive distribution maps. A number of potential applications of these generated maps rapidly arise in the field of biodiversity conservation and planning (provided that beyond statistical evaluations, also managers and planners are satisfied with the predictions provided by modellers). It is possible, for example, to assess the representation of species in nature reserve networks, evaluate their response to global climate change (of both the species and the reserves), quantify the impact of expected land cover change, and evaluate the influence of expanding exotic species, among other applications.

Growth and potential of predictive modelling techniques and approaches is boosting the applications of biodiversity mapping in recent years. A good example of structure construction of a discipline that has grown in parallel to biodiversity mapping is reserve design. Inspired by pioneering work carried out in Australia in the 1980s (Kirkpatrick, 1983), the Gap Analysis Program

(GAP) in the USA provided a major thrust to the application of predicted modelling to conservation (Scott *et al.*, 1993; Jennings, 2000; Peterson & Kluza, 2003). This programme has the main objective of creating and updating maps of the predicted distribution of native vertebrate species in the USA. By collating information from field research programmes and modelling their distribution using a variety of tools, GAP has become a powerful source of basic cartographic information for a number of planning and management decisions at the local, state, and national scale (Pearlstone *et al.*, 2002; Noon *et al.*, 2003). After generating predicted ranges, the next step in the process consists of overlaying these distributions with the geographical extent of the protected areas network, thus identifying existing 'gaps' — species that are not represented in the network — and providing potential locations for the future expansion or creation of new protected areas. Gap analysis has been proven highly successful and has become a very popular conservation tool. Its use is not limited to predicting geographical distributions, but also takes advantage of other sources of data on species ranges as well (e.g. De Klerk *et al.*, 2004; Rodrigues *et al.*, 2004).

The ample literature on reserve selection has been primarily devoted to optimizing the number, size, location, and cost of reserve networks, in order to assure that they include the largest possible fraction of biodiversity within the smallest amount of protected land and, ultimately, identify priority areas for conservation (Margules *et al.*, 2002; Wilson *et al.*, 2006; Alagador & Cerdeira, in press). Predictive modelling has enriched the discussion by including habitat suitability as an important characteristic to consider in the design of a network of reserves, because habitat suitability has been shown to correlate with the persistence of populations (Araújo *et al.*, 2002; Cabeza *et al.*, 2004). In an analogous way, habitat suitability has been incorporated into metapopulation models, which are primarily concerned with patch shape, size, and isolation (Akçakaya & Atwood, 1997; Kramer-Schadt *et al.*, 2005).

Regardless of how well the current reserve networks represent extant biodiversity, however, they are all threatened by global change, a series of complex forces that combine both climate change and land-use change (Araújo, 2004; Thomas *et al.*, 2004; Whittaker *et al.*, 2005; Araújo & Rahbek, 2006; Araújo *et al.*, 2006; Root & Schneider, 2006; Schlesinger, 2006; Sutherland, 2006; Thuiller *et al.*, 2006a,b). A number of studies, many based on habitat modelling, have raised the concern that species distributions may shift in response to climate change (mainly towards higher latitudes and altitudes), so that some may be forced to abandon reserves that were designed to protect them, and move to unprotected areas — if they can. The problem is that the current distribution of a particular species may be constrained by unsuitable surrounding habitats (e.g. perturbed areas, topographical barriers) or there may be a physical limit to their range, as is the case of species that live on mountains and cannot continue to shift upwards indefinitely (Dirnbock *et al.*, 2003; Thuiller *et al.*, 2006a). In fact, extensive modelling exercises predict an overall loss of biodiversity over the next 50 years as a result of climate change, in particular for mountain species (Bakkenes *et al.*, 2002; Thomas *et al.*, 2004). These predictions have been challenged,

however, because the results tend to vary with the technique applied (Thuiller *et al.*, 2004; Pearson *et al.*, 2006), and broad-scale models do not capture local interactions between organisms and their environment (Hampe, 2004), such as dispersal, interspecific interactions, or the existence of refuges at a finer spatial resolution than the climate change scenarios on which the models are based (Post *et al.*, 1999; Araújo & Pearson, 2005; Pearson, 2006). Although most studies are pessimistic about the extent of future range contractions, this is still an ongoing debate that relies heavily on habitat modelling.

Another promising but less explored line of research is the use of predictive habitat models to forecast conflicts between human activities and biodiversity conservation, and to try to minimize these conflicts. This is the case of the assessment of the impact of land-use changes linked to evolving agricultural practices. In Europe, for example, the simultaneous abandonment of traditional agriculture (areas that harbour several endangered species) and the intensification of agriculture on more profitable lands have led to a net decrease of biodiversity (Donald *et al.*, 2002; Newton, 2004). Environmental and agricultural policies tend to aim at different targets, and predictive models have been helpful for decision-makers to identify areas where these opposing forces may clash (Brotons *et al.*, 2004; Nogues-Bravo & Agirre, 2006; Seoane *et al.*, 2006). Similarly, maps that predict the likelihood of conflicts between carnivores and humans (due to livestock predation) have also been developed, guiding planners towards geographical locations where predation events are less likely to occur (Treves *et al.*, 2004). Another common conflict arises when human infrastructures, such as airports, roads, and power lines, are in the path of the regular movement of animals. For example, large birds pose a threat to airplanes because of bird strikes, particularly during the migratory season when birds gather in large flocks (<http://ecogrid.sara.nl/bambas/distribution/index.php>). Collision with power lines and road kills have been linked to the local decline of endangered species (Ferrer & Janss, 1999; Guzmán *et al.*, 2004), while some studies have modelled higher risk areas where corrective actions should be prioritized (Garthe & Huppopp, 2004; Malo *et al.*, 2004). Again, predictive models stand out as a useful tool for land-use planners seeking to make better decisions about biodiversity management and conservation.

Invasive species are one of the major drivers of global change, threatening numerous other species with extinction (Vitousek *et al.*, 1996; Chapin *et al.*, 2000). Biologists and agronomists initially began to search for traits that made species more aggressive colonizers, and for characteristics of ecosystems that made them more susceptible and vulnerable to invaders, in order to predict, mitigate, and perhaps prevent the negative consequences of biological invaders on native taxa (Crawley, 1986, 1987; D'Antonio & Vitousek, 1992; Crawley *et al.*, 1996; Mack, 1996; Veltman *et al.*, 1996; Williamson & Fitter, 1996; Williamson, 1999; Mack *et al.*, 2000; Huston, 2004; Richardson, 2004; Callaway & Maron, 2006). More recently, predictive habitat models have been used to map areas of high risk of being invaded (or that are currently more affected) both regionally (Mercado-Silva *et al.*, 2006; Muñoz & Real, 2006) and globally (Thuiller *et al.*, 2005). The

resulting picture can hardly be more worrying: six global biodiversity hotspots are highly susceptible of being invaded (Thuiller *et al.*, 2005).

On a relatively less ambitious scale, there are a number of additional applied questions that are being addressed with predictive models. Ecologists and field biologists, for example, are keen to use predictive models to generate potential habitat maps to plan for the reintroduction of species, and to evaluate reintroduction schemes underway (Mladenoff *et al.*, 1999; Schadt *et al.*, 2002; Hirzel *et al.*, 2004; Pellet *et al.*, 2004). Although reintroducing species into a portion of their historical range tends to be more successful than placing them elsewhere (Griffith *et al.*, 1989), the 'original' habitat is not always available (in fact this may be the cause of the disappearance of the species), and it may be necessary to look for new sites. But even when the original range is still a viable option, maps of potential suitable habitat help to recognize barriers to the dispersal of released individuals, to identify isolated areas unlikely to be colonized naturally, and to quantify the carrying capacity of the managed territory. These maps have also helped to detect species behavioural responses to fragmentation that could be used as early warnings of human perturbation (Laiolo & Tella, 2005).

BASELINE MONITORING AND SPATIAL MODELLING OF SPECIES DISTRIBUTION

Conservation, planning, and species management need basic monitoring data in order to gather critical information about where a species occurs and how its abundance changes through time (Underhill & Gibbons, 2002). Basic monitoring is aimed at sampling populations in order to describe their distribution in space (Donald & Fuller, 1998) and time (Bibby *et al.*, 2000).

Monitoring of species distribution has a strong link to predictive, habitat-based modelling techniques because both have the explicit objective of producing a spatial representation of a species' range. In fact, the development of predictive, habitat based models has been possible due the large amounts of data accumulated in large-scale distribution monitoring programmes such as atlases (Osborne & Tigar, 1992; Tobalske & Tobalske, 1999; Araújo & Pearson, 2005). But predictive distribution modelling has not only grown due to the availability of monitoring data. More recently, these techniques have started to play a key role in producing basic quantitative information on species distributions, allowing atlases move beyond their classic 'black and white' maps derived from grid mapping of presence-absence data. These new methodological approaches also offer testable predictive models that allow a more adequate representation of species distributions in a context of recognized but often unquantified uncertainty (Elith *et al.*, 2002; Barry & Elith, 2006).

Predictive models have already been applied to a number of atlases, for example, mammals (Hausser, 1995) and birds (Estrada *et al.*, 2004). Since atlases are a primary source of species distribution data for planning and species management conducted by governmental agencies, non-governmental organizations (NGOs), academia, and the private sector, the progressive inclusion of predictive habitat modelling will allow for the production of

an increasing number of detailed and testable maps, therefore increasing their potential impact and reliability.

For a number of conspicuous species, monitoring often involves counts to estimate their abundance in a given area. Population size is a fundamental ecological parameter, often required by local and international organizations and legislations, to assess the conservation status of a species, on the grounds that it is inversely correlated with the probability of extinction (IUCN, 2001; O'Grady *et al.*, 2004). However, the development of a standard sampling scheme to correctly make inferences from the numbers observed in the field to the entire study area is sometimes unfeasible. For example, studies on sea mammals commonly resort to the affordable technique of sampling from fishing or recreational boats (so-called platforms of opportunity), whose bearings are neither random nor systematic, and consequently any design-based estimate of abundance may be biased (Williams *et al.*, 2006). In contrast, model-based estimates are able to tackle the problem and have the further advantage that they provide maps at a higher spatial resolution (map units in model-based schemes are normally smaller than strata in design-based ones), as well as potential insights into the variables affecting abundance (Clarke *et al.*, 2003; Hedley & Buckland, 2004). These spatial modelling methods applied to field data have a large number of applications to conservation (see for example <http://www.ruwpa.st-and.ac.uk/Research/SpatialModelling/index.htm>).

Another promising application of habitat-based predictive modelling techniques to basic monitoring is less obvious. Typically, long-term monitoring programmes aim at detecting changes in population sizes. However, since they are generally based on a spatial sample of locations, these data also have the potential for producing species maps through interpolation or habitat modelling techniques (Sauer *et al.*, 1995; J.R. Ferrer-Paris & J.P. Rodríguez, unpubl. data). Recent attempts of using this kind of data have offered promising results as mapping tools (Carrascal *et al.*, 2006; Brotons *et al.*, 2007), including the incorporation of sampling biases via modern Bayesian estimation techniques (Thogmartin *et al.*, 2004, 2006).

Habitat models have another immediate link to monitoring efforts, since model predictions may help fill gaps in unsurveyed areas (as in atlas works) or to identify promising areas for future sampling, such as, for example, potential breeding nuclei for species in places yet to be explored (e.g. Seoane *et al.*, 2003; Jarnevich *et al.*, 2006; Brotons *et al.*, 2007). These spatial models have been shown to be an improvement on traditional atlas and reference works (Bustamante & Seoane, 2004; Bini *et al.*, 2006). Along the same line, drawing maps for sibling species whose identification relies on molecular characteristics is particularly demanding, and building habitat models with unequivocal genetic information is a cost-effective way to depict distributions (Real *et al.*, 2005).

The potential increase in the application of monitoring data to spatial mapping of species distribution suggests a trend towards more blurred boundaries between the objectives that these two activities cover. A promising approach is the integration of spatial mapping, including predictive modelling techniques, into basic, temporal monitoring of population trends in order to generate useful information on both changes in numbers and spatial

distribution of the monitored species (Travaini *et al.*, 2007). This may allow, for instance, the development of spatial alerts linked to temporal alerts, indicating worrying changes in species distribution and abundance (Greenwood, 2003). Traditional mapping projects, such as atlases, appear in this context as a fundamental complement, offering a detailed spatial picture of species distributions over a relatively small temporal scale.

In areas where sampling is expensive due to lack of observers or poor communications, representative sampling covering existing environmental variability offers an optimal allocation of limiting resources for monitoring (Hortal *et al.*, 2001; J.R. Ferrer-Paris & J.P. Rodríguez, unpubl. data). Such a sampling strategy is known to allow a potential representative estimation of species distributions through predictive modelling (Faith & Walker, 1996). In addition, such data may stand as baseline information for future sampling aimed at both better estimation of species distribution patterns or changes in population numbers or occurrence rates.

We think that the evidence shows that the use of predictive modelling of species distribution has been enormously useful and that conservation would benefit from a more explicit and structured integration of these techniques into basic monitoring programmes. We encourage governmental agencies, NGOs, academia, and the private sector, to develop basic monitoring programmes that incorporate as one of their cornerstones habitat modelling approaches to create and update species distributions.

A fundamental limitation that still exists, however, is that current biodiversity monitoring efforts are extremely limited in terms of their spatial and taxonomic coverage, survey methods are not standardized, and data sets collected at different scales lack integration (Pereira & Cooper, 2006). Worldwide monitoring initiatives such as the Global Amphibian Assessment (Stuart *et al.*, 2004) and the Biodiversity Intactness Index (Scholes & Biggs, 2005), rely heavily on expert opinion rather than on field data, museum collections, or predicted distribution modelling (Mace, 2005). For only the better-known taxonomic groups, such as birds and mammals, are there maps of their geographical distributions (Patterson *et al.*, 2003; Ridgely *et al.*, 2003), and only in a small number better-known locations is there information on their relative abundance and temporal dynamics (e.g. Gibbons *et al.*, 1993; Sauer *et al.*, 2005). The challenge is particularly large in the tropics, which harbour the majority of the world's species, but lack the financial resources and personnel for conservation (Rodríguez, 2003; Rodríguez *et al.*, 2005). The design, implementation, and funding of systematic biodiversity surveys at the scale of entire tropical countries, which minimize the costs and time required to perform them, apply standardized methods, quantify both distribution and abundance, and generate data that are adequate for species distribution modelling, are a major gap in global biodiversity monitoring that needs to be addressed.

AVAILABILITY OF ENVIRONMENTAL CARTOGRAPHY

Using species distribution models to generate maps relies to a great extent on the availability of environmental cartography. There can be a strong correlation between an environmental




Figure 1 Cumulative number of Earth observation remote sensing missions on satellite platforms (Donnio, 2006).

variable and the distribution of a species, but if this variable is not available as a map, it cannot be applied to predict the distribution of the species. Traditionally, the availability of environmental cartography has been very dependent on the economic development of a country. Also, even if environmental maps exist, there has been a wide variability in the access to those data, especially if they were considered to hold commercial or strategic value. During the last two decades several changes in the *status quo* have quickly improved access to high-quality environmental cartography. Earth observation remote sensing mission has increased steadily since 1972 (Fig. 1), and sensors on those satellites provide environmental data at a rate never imagined before. Although some data sets still need to be purchased at fairly high costs, others are readily available (see Appendix S1 in Supplementary Material). Some require the users to demonstrate a scientific pursuit (rather than a commercial one), but most can be freely downloaded. Other sources of data, such as global climate maps based on modelling of records from meteorological stations are also accessible on the Internet (e.g. Worldclim, <http://www.worldclim.org/>, Hijmans *et al.*, 2005). Although the spatial and temporal resolution of most data sets are adequate for species distribution modelling for both terrestrial and marine organisms, they have not been widely used. Notable exceptions include the work of Munoz *et al.* (2004), in which data of the NASA Wind Scatterometer were used to model landmass connectivity for plant dispersal in the southern hemisphere, and Travaini *et al.* (2007), in which Spot-Vegetation and Shuttle Radar Topographic Mission data were used to model guanaco distribution. Species distribution models, coupled with wider availability of environmental cartography, and relatively inexpensive and user-friendly geographical information systems (GIS), will surely result in a significant and rapid change in our knowledge of the distribution and abundance of biodiversity in the decades to come.

INTRODUCTION TO THE INVITED PAPERS

This special volume includes both methodological and applied works. Among those of a methodological nature, Ferrier *et al.*

(2007) introduce the technique of Generalized Dissimilarity Modelling (GDM), a new statistical method focused on modelling the distribution of biotic communities based on compositional dissimilarity. Ferrier *et al.* (2007) describe for the first time the technique in detail and present several examples of its use. Although originally designed to predict distributions of communities, it is a flexible approach that can also be accommodated to model individual species. Variations of the method can be used to predict the spatial distribution of phylogenetic dissimilarity, to regionalize areas based on species distribution patterns, or to include information on barriers to dispersal in the modelling. Elith & Leathwick (2007) explore the use of Multivariate Adaptive Regression Splines (MARS) to model presence-only data. MARS constitutes an extension of the better-known generalized additive models (GAMs), directed at modelling communities instead of individual species. Both GDM and MARS are examples of new techniques directed at modelling species groups to solve the common problem of species lacking enough information to be modelled as individual entities (Pearce & Boyce, 2006).

Two other new techniques: Geographically Weighted Regression (GWR) and Varying Coefficient Modelling (VCM) are explored by Osborne *et al.* (2007) to address the problem of non-stationarity in ecological systems. These new methods are complementary to global techniques, and provide details and habitat associations at local scales that global methods miss (Foody, 2005; Jetz *et al.*, 2005). Guisan *et al.* (2007) also consider aspects of scale, and explore the effect of grain size in species distribution models to try to answer the question of whether there is an optimal grain size. They present the first extensive test of the effect of coarsening the resolution of environmental variables in species distribution models, and show that, although there is a general trend towards degradation of model performance with resolution coarsening, it does not affect equally all data set, species, or modelling techniques. In their analyses, a new modelling technique, Boosted Regression Trees (BRT), ranks as the best among the 10 they used (including building environmental envelopes and regression methods). The papers of Elith & Leathwick (2007) and Guisan *et al.* (2007) consider the important aspect of building distribution models with presence-only data from herbaria and museum collections, often the only source of information on historical distribution changes and sometimes an important source of information for rare species.

A very important aspect considered in this special issue is the practical application of species distribution models. Travaini *et al.* (2007) provide an example on how new technologies can be integrated to reduce the costs of mapping species distributions. New technologies include statistical modelling techniques, as well as the use of freely available remote sensing data, accurate georeferencing of species observations with global positioning systems (GPS) and laser range-finders, optimal data gathering using personal digital assistants (PDAs) and customized software, and the use of GIS for extrapolation and map production. Their results show that new technologies constitute very promising tools for large and remote regions that lack accurate species distribution maps. Even in areas where apparently reliable distribution data exist, species distribution modelling techniques have

been shown to provide more accurate maps (Bustamante & Seoane, 2004). The article by Brotons *et al.* (2007) highlights the potential role of species distribution modelling techniques to improve and update our knowledge of species ranges using data from long-term monitoring programmes, which tend to be common in developed countries. Anadón *et al.* (2007) provide a practical example on the use of a hierarchical species distribution model to identify the causes of decline of an endangered species. Richardson & Thuiller (2007) use climate matching models to successfully identify species that may become potential invaders, and provide a technique to potentially design preventive measures for some species. As they emphasize, the most cost-effective way of reducing future problems is to prevent the introduction of high-risk invasive species.

REFERENCES

- Akçakaya, H.R. & Atwood, J.L. (1997) A habitat-based meta-population model of the California gnatcatcher. *Conservation Biology*, **11**, 422–434.
- Alagador, D. & Cerdeira, J.O. (in press) Designing spatially-explicit reserve networks in the presence of mandatory sites. *Biological Conservation*.
- Anadón, J.D., Giménez, A.M., Martínez, J.A.P. & Esteve, M.A. (2007) Assessing changes in habitat quality due to land use changes in the spur-thighed tortoise *Testudo graeca* using hierarchical predictive habitat models. *Diversity and Distributions*, **13**.
- Anderson, R., Lew, D. & Peterson, A. (2003) Evaluating predictive models of species' distributions: criteria for selecting optimal models. *Ecological Modelling*, **162**, 211–232.
- Araújo, M.B. (2004) Matching species with reserves — uncertainties from using data at different resolutions. *Biological Conservation*, **118**, 533–538.
- Araújo, M.B. & Guisán, A. (2006) Five (or so) challenges for species distribution modelling. *Journal of Biogeography*, **33**, 1677–1688.
- Araújo, M.B. & Pearson, R.G. (2005) Equilibrium of species' distributions with climate. *Ecography*, **28**, 693–695.
- Araújo, M.B. & Rahbek, C. (2006) How does climate change affect biodiversity? *Science*, **313**, 1396–1397.
- Araújo, M.B., Williams, P.H. & Fuller, R.J. (2002) Dynamics of extinction and the selection of nature reserves. *Proceedings of the Royal Society of London Series B, Biological Sciences*, **269**, 1971–1980.
- Araújo, M.B., Thuiller, W. & Pearson, R.G. (2006) Climate warming and the decline of amphibians and reptiles in Europe. *Journal of Biogeography*, **33**, 1712–1728.
- Austin, M.P. & Heyligers, P.C. (1991) New approach to vegetation survey design: gradsect sampling. *Nature conservation: cost effective biological surveys and data analysis* (ed. by C.R. Margules and M.P. Austin), pp. 31–36. CSIRO — Commonwealth Scientific and Industrial Research Organization, Canberra, Australia.
- Bakkenes, M., Alkemade, J.R.M., Ihle, F., Leemans, R. & Latour, J.B. (2002) Assessing effects of forecasted climate change on

- the diversity and distribution of European higher plants for 2050. *Global Change Biology*, **8**, 390–407.
- Barry, S. & Elith, J. (2006) Error and uncertainty in habitat models. *Journal of Applied Ecology*, **43**, 413–423.
- Bibby, J.C., Burgess, N.D., Hill, D.A. & Mustoe, S.H. (2000) *Bird census techniques*. 2nd edn. Academic Press, London.
- Bini, L.M., Diniz, J.A.F., Rangel, T., Bastos, R.P. & Pinto, M.P. (2006) Challenging Wallacean and Linnean shortfalls: knowledge gradients and conservation planning in a biodiversity hotspot. *Diversity and Distributions*, **12**, 475–482.
- Brotons, L., Herrando, S. & Pla, M. (2007) Updating bird species distribution at large spatial scales: applications of habitat modelling to data from long term monitoring programs. *Diversity and Distributions*, **13**.
- Brotons, L., Mañosa, S. & Estrada, J. (2004) Modelling the effects of irrigation schemes on the distribution of steppe birds in Mediterranean farmland. *Biodiversity and Conservation*, **13**, 1039–1058.
- Bustamante, J. & Seoane, J. (2004) Predicting the distribution of four species of raptors (Aves: Accipitridae) in southern Spain: statistical models work better than existing maps. *Journal of Biogeography*, **31**, 295–306.
- Cabeza, M., Araujo, M.B., Wilson, R.J., Thomas, C.D., Cowley, M.J.R. & Moilanen, A. (2004) Combining probabilities of occurrence with spatial reserve design. *Journal of Applied Ecology*, **41**, 252–262.
- Callaway, R.M. & Maron, J.L. (2006) What have exotic plant invasions taught us over the past 20 years? *Trends in Ecology & Evolution*, **21**, 369–374.
- Carrascal, L., Seoane, J., Palomino, D., Alonso, C.L. & Lobo, J. (2006) Species-specific features affect the ability of census-derived models to map winter avian distribution. *Ecological Research*, **21**, 681–691.
- Chapin, F.S. III, Zavaleta, E.S., Eviner, V.T., Naylor, R.L., Vitousek, P.M., Reynolds, H.L., Hooper, D.U., Lavorel, S., Sala, O.E., Hobbie, S.E., Mack, M.C. & Díaz, S. (2000) Consequences of changing biodiversity. *Nature*, **405**, 234–242.
- Clarke, E.D., Spear, L.B., McCracken, M.L., Marques, F.F.C., Borchers, D.L., Buckland, S.T. & Ainley, D.G. (2003) Validating the use of generalized additive models and at-sea surveys to estimate size and temporal trends of seabird populations. *Journal of Applied Ecology*, **40**, 278–292.
- Crawley, M.J. (1986) The population biology of invaders. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences*, **314**, 711–731.
- Crawley, M.J. (1987) What makes a community invisable? *Colonization, succession and stability* (ed. by A.J. Gray, M.J. Crawley and P.J. Edwards), pp. 429–453. Blackwell Scientific Publications, Oxford.
- Crawley, M.J., Harvey, P.H. & Purvis, A. (1996) Comparative ecology of the native and alien floras of the British Isles. *Philosophical Transactions of the Royal Society of London Series B, Biological Sciences*, **351**, 1251–1259.
- D'Antonio, C.M. & Vitousek, P.M. (1992) Biological invasions by exotic grasses, the grass/fire cycle, and global change. *Annual Review of Ecology and Systematics*, **23**, 63–87.
- De Klerk, H.M., Fjeldsa, J., Blyth, S. & Burgess, N.D. (2004) Gaps in the protected area network for threatened Afrotropical birds. *Biological Conservation*, **117**, 529–537.
- Dirnbock, T., Dullinger, S. & Grabherr, G. (2003) A regional impact assessment of climate and land-use change on alpine vegetation. *Journal of Biogeography*, **30**, 401–417.
- Donald, P.F. & Fuller, R.J. (1998) Ornithological atlas data: a review of uses and limitations. *Bird Study*, **45**, 129–145.
- Donald, P.F., Pisano, G., Rayment, M.D. & Pain, D.J. (2002) The Common Agricultural Policy, EU enlargement and the conservation of Europe's farmland birds. *Agriculture Ecosystems and Environment*, **89**, 167–182.
- Donnio, J.-P. (2006) *The satellite encyclopedia*. V0.74.001, 54th edn (http://www.tbs-satellite.com/tse/online/mis-teledeccion_res.html, accessed on 20 September 2006).
- Elith, J., Burgman, M.A. & Regan, H.M. (2002) Mapping epistemic uncertainties and vague concepts in predictions of species distribution. *Ecological Modelling*, **157**, 313–329.
- Elith, J. & Leathwick, J. (2007) Predicting species distributions from museum and herbarium records using multiresponse models fitted with multivariate adaptive regression splines. *Diversity and Distributions*, **13**.
- Estrada, J., Pedrocchi, V., Brotons, L. & Herrando, S. (2004) *Catalan breeding bird atlas (1999–2002)*. Lynx Edicions, Barcelona.
- Faith, D.P. & Walker, P.A. (1996) Environmental diversity: on the best-possible use of surrogate data for assessing the relative biodiversity of sets of areas. *Biodiversity and Conservation*, **5**, 399–415.
- Ferrer, M. & Janss, G.F.E. (1999) *Birds and power lines: collision, electrocution and breeding*. Servicios Informativos Ambientales/Quercus, Madrid.
- Ferrier, S., Manion, G., Elith, J. & Richardson, K. (2007) Using generalised dissimilarity modelling to analyse and predict patterns of beta-diversity in regional biodiversity assessment. *Diversity and Distributions*, **13**.
- Foody, G.M. (2005) Clarifications on local and global data analysis. *Global Ecology and Biogeography*, **14**, 99–100.
- Garthe, S. & Huppopp, O. (2004) Scaling possible adverse effects of marine wind farms on seabirds: developing and applying a vulnerability index. *Journal of Applied Ecology*, **41**, 724–734.
- Gibbons, D.W., Reid, J.B. & Chapman, R.A. (1993) *The new atlas of breeding birds in Britain and Ireland: 1988–1991*. T & AD Poyser Ltd., London, UK.
- Greenwood, J.J.D. (2003) The monitoring of British breeding birds: a success story for conservation science? *Science of the Total Environment*, **310**, 221–230.
- Griffith, B., Scott, J.M., Carpenter, J.W. & Reed, C. (1989) Translocations as a species conservation tool: status and strategy. *Science*, **245**, 477–480.
- Guisan, A., Graham, C.H., Elith, J., Huettmann, F. & Group, N.S.D.M. (2007) Sensitivity of predictive species distribution models to change in grain size. *Diversity and Distributions*, **13**.
- Guisan, A., Lehmann, A., Ferrier, S., Austin, M., Overton, J.M.C., Aspinall, R. & Hastie, T. (2006) Making better biogeographical

- predictions of species' distributions. *Journal of Applied Ecology*, **43**, 386–392.
- Guisan, A. & Thuiller, W. (2005) Predicting species distribution: offering more than simple habitat models. *Ecology Letters*, **8**, 993–1009.
- Guisan, A. & Zimmermann, N.E. (2000) Predictive habitat distribution models in ecology. *Ecological Modelling*, **135**, 147–186.
- Guzmán, J.N., García González, F.J., Garrote, G., Pérez de Ayala, R. & Iglesias, C. (2004) *El lince ibérico (Lynx pardinus) en España y Portugal*. Ministerio de Medio Ambiente, Madrid.
- Hampe, A. (2004) Bioclimate envelope models: what they detect and what they hide. *Global Ecology and Biogeography*, **13**, 469–471.
- Hausser, J. (1995) *Mammifères de la Suisse*. Birkhäuser-Verlag, Basel.
- Hedley, S.L. & Buckland, S.T. (2004) Spatial models for line transect sampling. *Journal of Agricultural Biological and Environmental Statistics*, **9**, 181–199.
- Heglund, P.J. (2002) Foundations of species-environment relations. *Predicting species occurrences: issues of accuracy and scale* (ed. by J.M. Scott, P.J. Heglund, M.L. Morrison, J.B. Haufler, M.G. Raphael, W.A. Wall and F.B. Samson), pp. 35–41. Island Press, Washington, D.C.
- Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**, 1965–1978.
- Hirzel, A.H., Posse, B., Oggier, P.A., Crettenand, Y., Glenz, C. & Arlettaz, R. (2004) Ecological requirements of reintroduced species and the implications for release policy: the case of the bearded vulture. *Journal of Applied Ecology*, **41**, 1103–1116.
- Hortal, J. & Lobo, J.M. (2005) An ED-based protocol for the optimal sampling of biodiversity. *Biodiversity and Conservation*, **14**, 2913–2947.
- Hortal, J., Lobo, J.M. & Martin-Piera, F. (2001) Forecasting insect species richness scores in poorly surveyed territories: the case of the Portuguese dung beetles (Col. Scarabaeinae). *Biodiversity and Conservation*, **10**, 1343–1367.
- Huston, M.A. (2004) Management strategies for plant invasions: manipulating productivity, disturbance, and competition. *Diversity and Distributions*, **10**, 167–178.
- IUCN (2001) *IUCN red list categories and criteria: version 3.1*. Species Survival Commission, World Conservation Union (IUCN), Gland, Switzerland and Cambridge, UK.
- Jarnevich, C.S., Stohlgren, T.J., Barnett, D. & Kartesz, J. (2006) Filling in the gaps: modelling native species richness and invasions using spatially incomplete data. *Diversity and Distributions*, **12**, 511–520.
- Jennings, M.D. (2000) Gap analysis: concepts, methods, and recent results. *Landscape Ecology*, **15**, 5–20.
- Jetz, W., Rahbek, C. & Lichstein, J.W. (2005) Local and global approaches to spatial data analysis in ecology. *Global Ecology and Biogeography*, **14**, 97–98.
- Kirkpatrick, J.B. (1983) An iterative method for establishing priorities for the selection of nature reserves: an example from Tasmania. *Biological Conservation*, **25**, 127–134.
- Kramer-Schadt, S., Revilla, E. & Wiegand, T. (2005) Lynx reintroductions in fragmented landscapes of Germany: projects with a future or misunderstood wildlife conservation? *Biological Conservation*, **125**, 169–182.
- Laiolo, P. & Tella, J.L. (2005) Habitat fragmentation affects culture transmission: patterns of song matching in Dupont's lark. *Journal of Applied Ecology*, **42**, 1183–1193.
- Linkie, M., Chapron, G., Martyr, D.J., Holden, J. & Leader-Williams, N. (2006) Assessing the viability of tiger subpopulations in a fragmented landscape. *Journal of Applied Ecology*, **43**, 576–586.
- Mace, G.M. (2005) Biodiversity — an index of intactness. *Nature*, **434**, 32–33.
- Mace, G., Masundire, H., Baillie, J., Ricketts, T., Brooks, T., Hoffmann, M., Stuart, S., Balmford, A., Purvis, A., Reyers, B., Wang, J., Revenga, C., Kennedy, E., Naeem, S., Alkemade, R., Allnutt, T., Bakarr, M., Bond, W., Chanson, J., Cox, N., Fonseca, G., Hilton-Taylor, C., Loucks, C., Rodrigues, A., Sechrest, W., Stattersfield, Janse van Rensburg, B. & Whiteman, C. (2005) Biodiversity. *Ecosystems and human well-being: current state and trends*, Vol. 1. *Findings of the condition and trends working group* (ed. by R. Hassan, R. Scholes and N. Ash), pp. 77–122. Island Press, Washington, D.C.
- Mack, R.N. (1996) Predicting the identity and fate of plant invaders: emergent and emerging approaches. *Biological Conservation*, **78**, 107–121.
- Mack, R.N., Simberloff, D., Lonsdale, W.M., Evans, H., Clout, M. & Bazzaz, F. (2000) Biotic invasions: causes, epidemiology, global consequences and control. *Issues in Ecology*, **5**, 1–20.
- Malo, J.E., Suarez, F. & Diez, A. (2004) Can we mitigate animal-vehicle accidents using predictive models? *Journal of Applied Ecology*, **41**, 701–710.
- Margules, C.R. & Austin, M.P. (1991) *Nature conservation: cost effective biological surveys and data analysis*. CSIRO — Commonwealth Scientific and Industrial Research Organization, Canberra, Australia.
- Margules, C.R., Pressey, R.L. & Williams, P.H. (2002) Representing biodiversity: data and procedures for identifying priority areas for conservation. *Journal of Biosciences*, **27**, 309–326.
- Mercado-Silva, N., Olden, J.D., Maxted, J.T., Hrabik, T.R. & Zanden, M.J.V. (2006) Forecasting the spread of invasive rainbow smelt in the Laurentian Great Lakes Region of North America. *Conservation Biology*, **20**, 1740–1749.
- Mladenoff, D.J., Sickley, T.A. & Wydeven, A.P. (1999) Predicting gray wolf landscape recolonization: logistic regression models vs. new field data. *Ecological Applications*, **9**, 37–44.
- Moisen, G.G., Edwards, J.T.C. & Osborne, P.E. (2006) Further advances in predicting species distributions. *Ecological Modelling*, **199**, 129–131.
- Munoz, J., Felicísimo, A.M., Cabezas, F., Burgaz, A.R. & Martínez, I. (2004) Wind as a long-distance dispersal vehicle in the Southern Hemisphere. *Science*, **304**, 1144–1147.

- Muñoz, A.R. & Real, R. (2006) Assessing the potential range expansion of the exotic monk parakeet in Spain. *Diversity and Distributions*, **12**, 656–665.
- Newton, I. (2004) The recent declines of farmland bird populations in Britain: an appraisal of causal factors and conservation actions. *Ibis*, **146**, 579–600.
- Nogues-Bravo, D. & Agirre, A. (2006) Modelling the spatial location of Dupont lark *Chersophilus duponti* during the breeding session in the Ablitas Natura Network area (Navarra). *Ardeola*, **53**, 55–68.
- Noon, B.R., Murphy, D.D., Beissinger, S.R., Shaffer, M.L. & Dellasala, D. (2003) Conservation planning for US National Forests: conducting comprehensive biodiversity assessments. *Bioscience*, **53**, 1217–1220.
- O'Grady, J.J., Burgman, M.A., Keith, D.A., Master, L.L., Andelman, S.J., Brook, B.W., Hammerson, G.A., Regan, T. & Frankham, R. (2004) Correlations among extinction risks assessed by different systems of threatened species categorization. *Conservation Biology*, **18**, 1624–1635.
- Osborne, P.E., Foody, G.M. & Suárez-Seoane, S. (2007) Non-stationarity and local approaches to modelling the distributions of wildlife. *Diversity and Distributions*, **13**.
- Osborne, P.E. & Tigar, B.J. (1992) Interpreting bird atlas data using logistic-models — an example from Lesotho, Southern Africa. *Journal of Applied Ecology*, **29**, 55–62.
- Patterson, B.D., Ceballos, G., Sechrest, W., Tognelli, M.F., Brooks, T., Luna, L., Ortega, P., Salazar, I. & Young, B.E. (2003) *Digital distribution maps of the mammals of the Western Hemisphere*, Version 1.0. NatureServe, Arlington, Virginia.
- Pearce, J.L. & Boyce, M.S. (2006) Modelling distribution and abundance with presence-only data. *Journal of Applied Ecology*, **43**, 405–412.
- Pearlstine, L.G., Smith, S.E., Brandt, L.A., Allen, C.R., Kitchens, W.M. & Stenberg, J. (2002) Assessing state-wide biodiversity in the Florida Gap analysis project. *Journal of Environmental Management*, **66**, 127–144.
- Pearson, R.G. (2006) Climate change and the migration capacity of species. *Trends in Ecology & Evolution*, **21**, 111–113.
- Pearson, R.G., Thuiller, W., Araújo, M.B., Martinez-Meyer, E., Brotons, L., McClean, C., Miles, L., Segurado, P., Dawson, T.P. & Lees, D.C. (2006) Model-based uncertainty in species range prediction. *Journal of Biogeography*, **33**, 1704–1711.
- Pellet, J., Guisan, A. & Perrin, N. (2004) A concentric analysis of the impact of urbanization on the threatened European tree frog in an agricultural landscape. *Conservation Biology*, **18**, 1599–1606.
- Pereira, H.M. & Cooper, H.D. (2006) Towards the global monitoring of biodiversity change. *Trends in Ecology & Evolution*, **21**, 123–129.
- Peterson, A.T. & Kluza, D. (2003) New distributional modelling approaches for gap analysis. *Animal Conservation*, **6**, 47–54.
- Post, E., Peterson, R.O., Stenseth, N.C. & McLaren, B.E. (1999) Ecosystem consequences of wolf behavioural response to climate. *Nature*, **401**, 905–907.
- Real, R., Barbosa, A.M., Martinez-Solano, I. & Garcia-Paris, M. (2005) Distinguishing the distributions of two cryptic frogs (Anura: Discoglossidae) using molecular data and environmental modeling. *Canadian Journal of Zoology-Revue Canadienne de Zoologie*, **83**, 536–545.
- Richardson, D.M. (2004) Plant invasion ecology — dispatches from the front line. *Diversity and Distributions*, **10**, 315–319.
- Richardson, D.M. & Thuiller, W. (2007) Home away from home — objective mapping of high-risk source areas for plant introductions. *Diversity and Distributions*, **13**.
- Ridgely, R.S., Allnutt, T.F.T., Brooks, T., McNicol, D.K., Mehlman, D.W., Young, B.E. & Zook, J.R. (2003) *Digital distribution maps of the birds of the Western Hemisphere*, Version 1.0. NatureServe, Arlington, Virginia.
- Rodrigues, A.S.L., Akcakaya, H.R., Andelman, S.J., Bakarr, M.I., Boitani, L., Brooks, T.M., Chanson, J.S., Fishpool, L.D.C., Da Fonseca, G.A.B., Gaston, K.J., Hoffmann, M., Marquet, P.A., Pilgrim, J.D., Pressey, R.L., Schipper, J., Sechrest, W., Stuart, S.N., Underhill, L.G., Waller, R.W., Watts, M.E.J. & Yan, X. (2004) Global gap analysis: Priority regions for expanding the global protected-area network. *Bioscience*, **54**, 1092–1100.
- Rodríguez, J.P. (2003) Challenges and opportunities for surveying and monitoring tropical biodiversity — a response to Danielsen *et al.* *Oryx*, **37**, 411–411.
- Rodríguez, J.P., Simonetti, J.A., Premoli, A. & Marini, M.A. (2005) Conservation in Austral and Neotropical America: building scientific capacity equal to the challenges. *Conservation Biology*, **19**, 969–972.
- Root, T.L. & Schneider, S.H. (2006) Conservation and climate change: the challenges ahead. *Conservation Biology*, **20**, 706–708.
- Sauer, J.R., Hines, J.E. & Fallon, J. (2005) *The North American breeding bird survey, results and analysis 1966–2005*, Version 6.2.2005. USGS Patuxent Wildlife Research Center, Laurel, Maryland.
- Sauer, J.R., Pendleton, G.W. & Orsillo, S. (1995) Mapping of bird distributions from point count surveys. *Monitoring bird populations by point counts* (ed. by C.J. Ralph, J.R. Sauer and S. Droege), pp. 151–160. USDA Forest Service, Pacific Southwest Research Station, General Technical Report PSW-GTR-149, Albany, California.
- Schadt, S., Knauer, F., Kaczensky, P., Revilla, E., Wiegand, T. & Trepl, L. (2002) Rule-based assessment of suitable habitat and patch connectivity for the Eurasian lynx. *Ecological Applications*, **12**, 1469–1483.
- Schlesinger, W.H. (2006) Global change ecology. *Trends in Ecology & Evolution*, **21**, 348–351.
- Scholes, R.J. & Biggs, R. (2005) A biodiversity intactness index. *Nature*, **434**, 45–49.
- Scott, J.M., Davis, F., Csuti, B., Noss, R., Butterfield, B., Groves, C., Anderson, H., Caicco, S., D'Erchia, F., Edwards, T.C. Jr, Ulliman, J. & Wright, R.G. (1993) Gap analysis: a geographic approach to protection of biological diversity. *Wildlife Monographs*, **123**, 1–41.
- Seoane, J., Justríbó, J.H., García, F., Retamar, J., Rabadán, C. & Atienza, J.C. (2006) Habitat-suitability modelling to assess the effects of land-use changes on Dupont's lark *Chersophilus*

- duponti*: a case study in the Layna Important Bird Area. *Biological Conservation*, **128**, 67–78.
- Seoane, J., Viñuela, J., Díaz-Delgado, R. & Bustamante, J. (2003) The effects of land use and climate on red kite distribution in the Iberian peninsula. *Biological Conservation*, **111**, 401–414.
- Stuart, S.N., Chanson, J.S., Cox, N.A., Young, B.E., Rodrigues, A.S.L., Fischman, D.L. & Waller, R.W. (2004) Status and trends of amphibian declines and extinctions worldwide. *Science*, **306**, 1783–1786.
- Sutherland, W.J. (2006) Predicting the ecological consequences of environmental change: a review of the methods. *Journal of Applied Ecology*, **43**, 599–616.
- Thogmartin, W.E., Knutson, M.G. & Sauer, J.R. (2006) Predicting regional abundance of rare grassland birds with a hierarchical spatial count model. *Condor*, **108**, 25–46.
- Thogmartin, W.E., Sauer, J.R. & Knutson, M.G. (2004) A hierarchical spatial model of avian abundance with application to Cerulean Warblers. *Ecological Applications*, **14**, 1766–1779.
- Thomas, C.D., Cameron, A., Green, R.E., Bakkenes, M., Beaumont, L.J., Collingham, Y.C., Erasmus, B.F.N., Ferreira de Siqueira, M., Grainger, A., Hannah, L., Hughes, L., Huntley, B., Van Jaarsveld, A.S., Midgley, G.F., Miles, L., Ortega-Huerta, M.A., Peterson, A.T., Phillips, O.L. & Williams, S.E. (2004) Extinction risk from climate change. *Nature*, **427**, 145–148.
- Thuiller, W., Araujo, M.B., Pearson, R.G., Whittaker, R.J., Brotons, L. & Lavorel, S. (2004) Uncertainty in predictions of extinction risk. *Nature*, **430**, 10.1038/nature02716.
- Thuiller, W., Broennimann, O., Hughes, G., Alkemade, J.R.M., Midgley, G.F. & Corsi, F. (2006a) Vulnerability of African mammals to anthropogenic climate change under conservative land transformation assumptions. *Global Change Biology*, **12**, 424–440.
- Thuiller, W., Lavorel, S., Sykes, M.T. & Araujo, M.B. (2006b) Using niche-based modelling to assess the impact of climate change on tree functional diversity in Europe. *Diversity and Distributions*, **12**, 49–60.
- Thuiller, W., Richardson, D.M., Pysek, P., Midgley, G.F., Hughes, G.O. & Rouget, M. (2005) Niche-based modelling as a tool for predicting the risk of alien plant invasions at a global scale. *Global Change Biology*, **11**, 2234–2250.
- Tobalske, C. & Tobalske, B.W. (1999) Using atlas data to model the distribution of woodpecker species in the Jura, France. *Condor*, **101**, 472–483.
- Travaini, A., Bustamante, J., Rodríguez, A., Zapata, S., Procopio, D., Pedrana, J. & Martínez-Peck, R. (2007) An integrated framework to map animal distributions in large and remote regions. *Diversity and Distributions*, **13**.
- Treves, A., Naughton-Treves, L., Harper, E.K., Mladenoff, D.J., Rose, R.A., Sickley, T.A. & Wydeven, A.P. (2004) Predicting human-carnivore conflict: a spatial model derived from 25 years of data on wolf predation on livestock. *Conservation Biology*, **18**, 114–125.
- Underhill, L.G. & Gibbons, D.W. (2002) Mapping and monitoring bird populations: their conservation uses. *Conserving bird biodiversity: general principles and their application* (ed. by K. Norris and D.J. Pain), pp. 34–60. Cambridge University Press, Cambridge.
- Veltman, C.J., Nee, S. & Crawley, M.J. (1996) Correlates of introduction success in exotic New Zealand birds. *The American Naturalist*, **147**, 542–557.
- Vitousek, P.M., D'Antonio, C.M., Loope, L.L. & Westbrooks, R. (1996) Biological invasions as global environmental change. *The American Scientist*, **84**, 468–478.
- Whittaker, R.J., Araujo, M.B., Paul, J., Ladle, R.J., Watson, J.E.M. & Willis, K.J. (2005) Conservation biogeography: assessment and prospect. *Diversity and Distributions*, **11**, 3–23.
- Williams, R., Hedley, S.L. & Hammond, P.S. (2006) Modeling distribution and abundance of Antarctic baleen whales using ships of opportunity. *Ecology Society*, **11**.
- Williamson, M. (1999) Invasions. *Ecography*, **22**, 5–12.
- Williamson, M. & Fitter, A. (1996) The varying success of invaders. *Ecology*, **77**, 1661–1666.
- Wilson, K.A., McBride, M.F., Bode, M. & Possingham, H.P. (2006) Prioritizing global conservation efforts. *Nature*, **440**, 337–340.

SUPPLEMENTARY MATERIAL

The following supplementary material is available for this article:

Appendix S1 Environmental predictors of global extent accessible through the Internet.

This material is available as part of the online article from:

<http://www.blackwell-synergy.com/doi/abs/10.1111/j.1472-4642.2007.00356.x>

(This link will take you to the article abstract).

Please note: Blackwell Publishing are not responsible for the content or functionality of any supplementary materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.