

Met Office. 2014. UK seasonal weather summary: winter 2013/2014. *Weather* **69**: 99.

Neilson BT, Chapra SC, Stevens DK et al. 2010. Two-zone transient storage modeling using temperature and solute data with multiobjective calibration: 1. Temperature. *Water Resour. Res.* **46**: W12520.

Orr HG, Simpson GL, des Clers S et al. 2014. Detecting changing river temperatures in England and Wales. *Hydrol. Processes*. doi:10.1002/hyp.10181

Parry S, Marsh T, Kendon M. 2013. 2012: from drought to floods in England and Wales. *Weather* **68**: 268–274.

Prudhomme C, Young A, Watts G et al. 2012. The drying up of Britain? A national estimate of changes in seasonal river flows from 11 Regional Climate Model simulations. *Hydrol. Processes* **26**: 1115–1118.

Punzet M, Voß F, Voß A et al. 2012. A global approach to assess the potential impact of climate change on stream water temperatures and related in-stream first order decay rates. *J. Hydrometeorol.* **13**: 1052–1065.

Sabouri F, Gharabaghi B, Mahboubi AA et al. 2013. Impervious surfaces and sewer pipe effects on stormwater runoff temperature. *J. Hydrol.* **502**: 10–17.

Smith K, Lavis ME. 1975. Environmental influences on the temperature of a small upland stream. *Oikos* **26**: 228–236.

Subehi L, Fukushima T, Onda Y et al. 2010. Analysis of stream water temperature changes during rainfall events in forested watersheds. *Limnology* **11**: 115–124.

Toffolon M, Siviglia A, Zolezzi G. 2010. Thermal wave dynamics in rivers affected by hydropeaking. *Water Resour. Res.* **46**: W08536.

Torgersen CE, Faux RN, McIntosh BA et al. 2001. Airborne thermal remote sensing for water temperature assessment in rivers and streams. *Remote Sens. Environ.* **76**: 386–398.

Trenberth KE. 2011. Changes in precipitation with climate change. *Clim. Res.* **47**: 123–138.

Webb BW. 1995. Regulation and thermal regime in a Devon river system, in *Sediment and Water Quality in River Catchments*. Foster IDL, Gurnell AM, Webb BW (eds). John Wiley & Sons: Chichester, UK, pp 65–94.

Webb BW, Walling DE. 1985. Temporal variation of river water temperatures in a Devon river system. *Hydrol. Sci. J.* **30**: 449–464.

Webb BW, Hannah DM, Moore RD et al. 2008. Recent advances in stream and river temperature research. *Hydrol. Processes* **22**: 902–918.

Westhoff MC, Bogaard TA, Savenije HHG. 2010. Quantifying the effect of in-stream rock clasts on the retardation of

heat along a stream. *Adv. Water Resour.* **33**: 1417–1425.

Westhoff MC, Gooseff MN, Bogaard TA et al. 2011. Quantifying hyporheic exchange at high spatial resolution using natural temperature variations along a first order stream. *Water Resour. Res.* **47**: W10508.

Wilby RL, Orr H, Watts G et al. 2010. Evidence needed to manage freshwater ecosystems in a changing climate: turning adaptation principles into practice. *Sci. Total Environ.* **408**: 4150–4164.

Wilby RL, Johnson MF, Toone JA. 2012. The Loughborough University Temperature Network (LUTEN): rationale and analysis of stream temperature variations. *Proceedings of Earth Systems Engineering 2012: Systems Engineering for Sustainable Adaptation to Global Change*, Newcastle, UK.

Wilby RL, Johnson MF, Toone JA. 2014. Nocturnal river water temperatures: spatial and temporal controls. *Sci. Total Environ.* **482–483**: 157–173.

Zolezzi G, Siviglia A, Toffolon M et al. 2011. Thermo-peaking in Alpine streams: event characterization and time scales. *Ecohydrology* **4**: 564–576.

Correspondence to: Robert L. Wilby
r.l.wilby@lboro.ac.uk

© 2015 Royal Meteorological Society
doi:10.1002/wea.2435

Non-classic extratropical cyclones on Met Office sea-level pressure charts: double cold and warm fronts

**Katy C. Mulqueen¹
and David M. Schultz²**

¹*School of Earth, Atmospheric and Environmental Sciences, University of Manchester*

²*Centre for Atmospheric Science, School of Earth, Atmospheric and Environmental Sciences, University of Manchester*

Introduction

The evolution of extratropical cyclones has traditionally been described in the context of the Norwegian cyclone model, where a developing cyclone grows on a stationary

front, developing single cold and warm fronts, then occluding (Figure 1(a)). First described by Bjerknes (1919) and Bjerknes and Solberg (1922), the Norwegian cyclone model provided a context for scientists and forecasters to describe the evolution of cyclones and led to the modernisation of meteorology in the first half of the twentieth century (e.g. Namias, 1980; 1983; Newton and Rodebush Newton, 1994). The model has withstood the test of time, with relatively few modifications. One modification is that of the occlusion process as the wrap-up of the air masses involved rather than the catch-up of the warm front by the cold front (Schultz and Vaughan, 2011).

Not all cyclones undergo an evolution similar to the Norwegian cyclone model,

however. In the 1980s, field programmes targeting rapidly developing marine extratropical cyclones and mesoscale numerical model simulations showed a slightly different evolution (Shapiro and Keyser, 1990). Instead of a narrowing warm sector during occlusion, as in the Norwegian cyclone model, the Shapiro–Keyser cyclone model exhibited a warm front nearly perpendicular to the cold front in a frontal T-bone, a bent-back front and an eventual warm seclusion (Figure 1(b)). Schultz *et al.* (1998) and Schultz and Zhang (2007) showed that cyclones in large-scale diffluence tended to form Norwegian cyclones and cyclones in large-scale confluence tended to form Shapiro–Keyser cyclones: the lower friction over water also facilitates the development

of Shapiro–Keyser characteristics (Hines and Mechoso, 1993).

Over the central and eastern USA, other types of cyclone structures and evolutions are possible, including lee troughs and arctic fronts associated with lee cyclones

(Keshishian *et al.*, 1994; Steenburgh and Mass, 1994) warm-sector rainbands caused by cold fronts aloft (Hobbs *et al.*, 1990; 1996; Locatelli *et al.*, 2002) and cyclones with multiple warm fronts (Metz *et al.*, 2004). Multiple fronts have also been documented

in idealised modelling studies (Hoskins *et al.*, 1984). In the UK, Miles (1962) and Browning and Monk (1982) recognised the importance of prefrontal troughs and upper-level fronts in the split-front model (also known as a katafront), and multiple fronts have also been documented (Young, 2014).

Consideration of the Met Office sea-level pressure charts for the North Atlantic Ocean and western Europe on a regular basis leads to the realisation that many cyclones possess frontal structures and evolutions inconsistent with the classic Norwegian cyclone model, specifically cyclones with an array of fronts (Figure 2). For example, the cyclone in Figure 2(a) consists of two warm sectors with two cold and two warm fronts, with a pre-warm-front trough and post-cold-front trough. The Met Office defines a trough as the following: *An elongated area of relatively low surface pressure. The troughs marked on weather charts may also represent an area of low thickness (thickness trough), or a perturbation in the upper troposphere (upper trough).* All are associated with increasing cloud and risk of precipitation (<http://www.metoffice.gov.uk/guide/weather/symbols>). In Figure 2(b) the cyclone is occluded with a bent-back front, an upper cold front, and warm and cold fronts to the north of the cyclone centre. In Figure 2(c), a cyclone with an occluded front, two warm fronts and three cold fronts is connected to an occluded cyclone to its northeast. In Figure 2(d), a cyclone with a bent-back occluded front is associated with three other warm sectors, each with its own cold and warm fronts, and one even having another bent-back front.

Perhaps an initial reaction is to dismiss these ‘octopus cyclones’ (because of their many radiating fronts from the low centre) as too inconsistent with the Norwegian cyclone model, too inappropriate, too difficult to understand, or unclassifiable (Mass, 1991). Such reactions though reduce the users’ confidence in the Met Office sea-level pressure charts as a useful tool for understanding the weather. For example, Prichard (2006) lamented the lack of consistency of features analysed on Met Office charts. Rather than dismiss these analyses, our research is aimed at understanding these non-classic structures taken at face-value from the analyses. As an initial step, this article focuses on a class of cyclones from these charts: those with two warm fronts, two cold fronts, or both two warm fronts and two cold fronts (hereafter all three are classified as *double-front cyclones*). Figure 2(a) epitomises these types of cyclones in their most simple form.

Such double fronts have been observed occasionally in the literature. Even the Norwegians recognised the existence of multiple fronts (Bjerknes, 1930, his figures 3, 5, 10 and A3–A5). Browning *et al.* (1995)




Figure 1. Two types of cyclone conceptual models: (a) Norwegian cyclone model, (b) Shapiro–Keyser cyclone model. (top) Lower-tropospheric (e.g. 850hPa) geopotential height and fronts, and (bottom) lower-tropospheric potential temperature. The stages in the respective cyclone evolutions are separated by approximately 6–24h, and the frontal symbols are conventional. The characteristic scale of the cyclones based on the distance from the geopotential height minimum, denoted by L, to the outermost geopotential height contour in stage IV is 1000km. (Figures and caption from Schultz and Vaughan, 2011.)




Figure 2. Examples of non-classic multiple-front cyclones from Met Office sea-level pressure charts: (a) 0000 UTC on 26 January 2013, (b) 0000 UTC on 18 November 2012, (c) 1800 UTC on 27 November 2012, and (d) 0000 UTC on 15 January 2013. (Images courtesy of wetter3.de and Crown copyright.)




Figure 3. Diagrams representing double-front extratropical cyclones on Met Office sea-level pressure charts. (a) A double warm-front cyclone with a secondary warm front located ahead of the primary warm front. (b) A double warm-front cyclone with a secondary warm front located behind the primary warm front. (c) A double cold-front cyclone with a secondary cold front located ahead of the primary cold front. (d) A double cold-front cyclone with a secondary cold front located behind the primary cold front. (e) A double front cyclone with both a secondary warm front and a secondary cold front.

studied a cyclone with two cold fronts during the FRONTS 92 field programme. Parker (1998, his figure 1, from T. Hewson) showed a schematic of a frontal wave inside a cyclone's warm sector that yielded a cyclone with two warm and two cold fronts. Schultz (2005) discussed prefrontal troughs and wind shifts, features that could be represented as fronts on surface charts.

The questions we want to examine in this study are how common these non-classic cyclones are and how they evolve. We will examine four cool seasons of Met Office sea-level pressure charts, classifying them as single-front cyclones or double-front cyclones. We will produce a classification scheme for double-front cyclones and describe common structures and evolutions. This study has three goals. First, we aim to provide some guidance for interpreting Met Office sea-level pressure charts with non-classic cyclone structures. Second, we wish to bring a greater appreciation to the variety of cyclone structures and evolutions present in the atmosphere. Third, we want to show that the Met Office sea-level pressure charts are more sensible than they may seem at first from a cursory glance at a single chart.

Data and methods

Our data are the six-hourly sea-level pressure charts for the North Atlantic Ocean and western Europe produced by the UK Met Office and archived at wetter3.de. Production of the charts at the Met Office begins with background fields from a previous model run, typically 6h forecasts

from the Met Office global forecast model. In the next step, fronts are initially hand drawn, although automated fronts derived from the 850hPa wet-bulb potential temperature field (Hewson, 1998; Hewson and Tittley, 2010) are available to the forecaster to supplement their analyses. The fields referenced by the analyst are partly subjective, but they always include mean sea-level pressure and 850hPa wet-bulb potential temperature. Other commonly used fields include the model's explicitly resolved precipitation rate and parameterised convective-precipitation rate, relative humidity at one or more lower-tropospheric levels, 1000–500hPa thickness and surface dewpoint temperature. The analyst generally looks for discrepancies compared with observations. Where these clearly exist, the analyst will follow the observations not the model fields, but for the vast majority of the time, the model is followed. The analyst is also able to follow the recent history of particular ship and buoy observations, which can be helpful when discrepancies arise.

In this study, we examine the cool-season months October to March because cyclones and fronts are more abundant and better defined in the cool season. The four most recent cool seasons were examined: 2010–2011, 2011–2012, 2012–2013 and 2013–2014.

A three-step process was used to identify cyclones. In the first step, we considered just the existence of cyclones. A cyclone was selected for consideration if it had one or more closed isobars, had a central sea-level pressure lower than 995hPa, was

located within the North Atlantic region (40°N–70°N, 50°W–10°E), and was isolated from other cyclones (i.e. no double-barrelled low-pressure centres). This process yielded 217 cyclones in our dataset.

In the second step, the 217 cyclones were classified into either single-front or double-front cyclones. Single-front cyclones had a single warm front, single cold front, and a possible occluded front throughout their time within the domain described above. Nominally, these could be Norwegian cyclones or Shapiro–Keyser cyclones, or even another type of evolution. Double-front cyclones had either a double warm front, double cold front, or both, at some point in their evolution within the domain. There were 94 single-front cyclones and 123 double-front cyclones.

The basic structures of these 123 double-front cyclones are displayed in Figure 3. A double-front cyclone consisted of a cyclone with connected warm and cold fronts passing through the centre of the cyclone (the primary fronts), with an additional front or fronts nearby the cyclone (the secondary front or fronts). The 123 double-front cyclones consisted of 63 cyclones (51% of the 123) with double warm fronts (Figure 3(a) and (b)), 37 (30%) cyclones with double cold fronts (Figure 3(c) and (d)), and 23 (19%) cyclones with both double warm and cold fronts (Figure 3(e)).

In the third selection step we considered the evolution of the double-fronted cyclones. From the 123 cyclones above, a double-front cyclone was further selected if its evolution on the Met Office sea-level pressure charts took place within the domain (40°N–70°N, 50°W–10°E), if the secondary front and the cyclone were less than 200km apart, the secondary front was at least 500km in length and the secondary front was present for at least 6h (appears on two maps 6h apart). As four sea-level pressure charts are produced, at 0000, 0600, 1200 and 1800 UTC each day, 6h is the minimum duration of a secondary front – and hence the minimum period of time during which the cyclone maintained its double-front structure. (Strictly speaking, a feature that existed on two consecutive maps 6h apart, but not three consecutive maps, could have had a lifetime of at least 6h, but not more than 18h. For the purposes of this paper, we say that a feature appearing on two consecutive 6h maps had a duration of 6h, although we recognise this as a lower limit.)

Those cyclones remaining after the third selection step above were also examined for common evolutionary sequences. A double-front cyclone meeting the criteria above was considered classifiable if it was possible to understand how the double front formed based on the evolution displayed on the sea-level pressure charts.

Through this process, it became evident that double-front cyclones underwent a small number of classifiable evolutions, allowing for the construction of a classification scheme. A cyclone was considered unclassifiable if its double-front evolution could not be easily recognised from the sea-level pressure charts. Doswell (1991) expressed the importance of having an unclassifiable category, stating that it *leaves open the possibility that someone might find a way to classify them in the future*. With these additional criteria, the number of classifiable cyclones was 53 (43% of the 123 cyclones) and the number of unclassifiable cyclones was 70 (57%).

Climatology

The distribution of the 123 double-front cyclones shows only a slight preference for some cool-season months over others (Figure 4). For example, a relative maximum of 26 events (6.5 events per month) occurs in January compared with a relative minimum of 16 events (4 events per month) in November. Given that only four cool seasons were examined, the possibility of the small sample size of the classifiable cyclones producing a different result for a different set of years because of interannual variability precludes further analysis of this graph.




Figure 4. A stacked bar chart showing the annual cycle of 53 classifiable and 70 unclassifiable double-front extratropical cyclones from the four cool seasons (October–March) of 2010–2011 to 2013–2014.




Figure 5. Duration (in hours) of secondary fronts in 53 double-front cyclones.

Nevertheless, Figure 4 does show that double-front cyclones on Met Office sea-level pressure charts are relatively common, averaging about five per cool-season month. Indeed, within our dataset of cyclones, double-front cyclones are more commonly analysed than Norwegian cyclones on the Met Office charts.

The distribution of the durations of the secondary fronts in the 53 double-front cyclones is plotted in Figure 5. Although some double-front cyclones existed for only 6h, others existed for as many as 54h (Figure 5). The mode of the distribution is 12h, with 15 out of 53 cases occurring for this length of time. The mean is 18.7h, and the median is 18h. Thus, double fronts were not isolated features that appeared for a limited period of time only, but were often long-lived and persistently analysed structural features of the cyclones.

Classification

Our classification scheme produced four broad categories of double-front cyclones at some point in their evolution within our analysis domain: A, B, C and D. Category A cyclones started out with both a double cold and double warm front within our analysis domain (Figure 6): in our dataset,

43 cyclones (81.1% of 53) were classified as category A. Five different evolutions were identified within category A, four of which involved the loss of a front, presumably by merger or frontolysis. Type A1 represents cyclones with two cold fronts where the secondary cold front was located ahead of the primary cold front: 2/53 (3.8%) cyclones formed by this evolution. Type A2 also had two cold fronts, but the secondary cold front was located behind the primary cold front: they were the second most common type of evolution, with 16/53 (30.2%) cyclones. Next, A3 cyclones had two warm fronts where the secondary warm front was located ahead of the primary warm front: they were the most common type of evolution, with 18/53 (34.0%) cyclones. The A4 cyclones also had two warm fronts, but the secondary warm front was located behind the primary warm front: 3/53 (5.7%) cyclones formed by this evolution. Finally, A5 represented double-front cyclones with both a double warm and a double cold front throughout the evolution of the cyclone: four (7.5%) cyclones underwent this evolution.

Category B cyclones started out as a classic Norwegian cyclone with a single cold front and a single warm front, but with an occluded front to the north that became emplaced on the cyclone, producing either a double cold- or double warm-front cyclone (Figure 7). This evolution is similar to a conceptual model proposed by Metz *et al.* (2004, their figure 7) for the formation of multiple warm-front-like baroclinic zones within extratropical cyclones east of the Rocky Mountains: attachment of a northern baroclinic zone to a pre-existing cyclone. In our dataset, four cyclones (7.5% of 53) were classified as category B. Type B1 evolved into a cyclone characterised by only a double cold front (2/53, 3.8%), and B2 evolved into a cyclone characterised by only a double warm front (2/53, 3.8%) (Figure 7). Interestingly, this evolution did not produce any cyclones with both a double cold front and a double warm front (although the small sample size of four cyclones could be a factor).

Category C cyclones began life as a cyclone with both a surface warm front




Figure 6. Schematic diagrams for the five possible evolutions of category A cyclones and the percentage (rounded to one decimal place) in red of the 53 cyclones fitting that behaviour.

and an upper-level warm front (Figure 8). At some point in the cyclone's evolution, the analyst changed the upper-level warm front to a surface warm front, producing a cyclone with two analysed warm fronts (C1). In our dataset, three cyclones (5.7% of 53) were classified as category C.

Finally, the fourth classification, category D cyclones, had a trough behind the primary cold front. At a later time in the evolution of the cyclone, that trough became analysed as a secondary cold front (Figure 9). In our dataset, three cyclones (5.7% of 53) were classified as category D.

To summarise, the most common evolutions were those in the A classification, specifically types A2 and A3. These two evolutions involve either the weakening of a warm front (A2) or a cold front (A3), leaving a secondary cold or warm front outside of the warm sector (B1, B2, C1, D1). Evolutions leaving fronts within the warm sector (A1, A4 and A5) were relatively uncommon.




Figure 7. Schematic diagrams for the two possible evolutions of category B cyclones and the percentage (rounded to one decimal place) in red of the 53 cyclones fitting that behaviour.




Figure 8. Schematic diagram for the evolution of category C cyclones and the percentage (rounded to one decimal place) in red of the 53 cyclones fitting that behaviour.




Figure 9. Schematic diagram for the evolution of category D cyclones and the percentage (rounded to one decimal place) in red of the 53 cyclones fitting that behaviour. The solid line represents a trough (see text for definition).

Discussion

The natural question to ask is where the double-front structure that began the evolution in category A came from. Examination of these events indicated that in many cases the innermost cold and warm fronts to the warm sector resulted from the attachment of an open-wave warm sector at a lower latitude (perhaps associated with the subtropical anticyclone) to a more eastwardly mobile extratropical cyclone at a higher latitude. Figure 10 illustrates one of these evolutions of a cyclone in our dataset classified as type A2. As the mobile cyclone

passed and deepened from 992 to 960hPa in 24h, it attached to the southernmost 1002hPa open-wave cyclone and became a double cold-front and double warm-front cyclone (Figure 10). This evolution is similar to the conceptual model proposed by Metz *et al.* (2004, their figure 6). As the cyclone neared the UK, the leading warm front lost its identity, probably due to a merger with the leading warm front (Figure 10(b) and (c)).

Another question is why these charts had such an abundance of analysed double fronts compared with charts produced by other agencies (National Oceanic and Atmospheric Administration, Deutsche Wetterdienst). We suggest that one reason may be the use of 850hPa wet-bulb potential temperature (θ_w). As gradients in θ_w result from gradients in both potential temperature and humidity mixing ratio, some 'fronts' may actually have been moisture gradients. Sanders and Doswell (1995), Sanders (1999), Schultz (2009, p. 355) and Schultz and Blumen (2015) have discussed the importance of analysing fronts as a function of temperature. Although a mask for potential-temperature gradients was available for the forecasters (Hewson and Tittley, 2010), comparing some analysed cases with gridded-model output indicated that this might have been a factor for some cases, but not all. Indeed, global frontal climatologies produced using the same θ_w scheme have identified 'fronts' in the tropics and subtropics (Berry *et al.*, 2011) that lay along moisture gradients, not temperature gradients (G. Berry, pers. comm.). A more thorough analysis of this issue awaits further investigation.

Although this preliminary study has presented these evolutions and described the most common ones, it does not describe why these double-front cyclones happen. That remains for future research.

Conclusion

The purpose of this study was to examine how common non-classic double-front cyclones are and how they form. Analysing Met Office sea-level pressure charts from four cool seasons between 2010–2011 and 2013–2014 identified 123 double-front cyclones (<995hPa), exceeding the 94 single-front cyclones (which include cyclones adhering to the Norwegian conceptual model). The 123 double-front cyclones were re-examined, specifically focusing on their evolution. This additional scrutiny led to a dataset of 53 double-front cyclones where the evolution could be classified into one of four categories. Category A (cyclones with two warm fronts and two cold fronts) dominated the dataset with 81.1% of the 53 double-front cyclones. Types A2 and A3 were the most dominant evolutions, with 30.2% and 34.0% of 53 double-front




Figure 10. Example from Met Office sea-level pressure charts of the evolution of a double-front cyclone classified as type A2, with two warm fronts and two cold fronts, that loses one warm front: (a) 1200 UTC on 11 December 2011, (b) 0000 UTC on 12 December 2011 and (c) 1200 UTC on 12 December 2011. (Images courtesy of wetter3.de and Crown copyright.).

cyclones forming from these evolutions, respectively. Specifically, type A2 cyclones evolved to a cyclone with a secondary cold front, and A3 cyclones evolved to a cyclone with a secondary warm front.

This paper highlights the relative abundance of double-front cyclones in reality and suggests that such cyclones are not rare or a result of errors by the analysts. Instead, they are common structures with recurring classifiable patterns that are displayed on published Met Office sea-level pressure charts. Moreover, they are more common than single-front cyclones. Therefore, forecasters and researchers should be aware that such double-front cyclones can occur and usually relate to sensible weather on the ground. Although the single-front conceptual models (i.e. Norwegian and Shapiro–Keyser conceptual models) continue to provide a useful framework for many cyclones, this paper adds to a growing body of literature demonstrating that important structural and evolutionary differences exist among real-world cyclones.

Acknowledgements

This article arose from Mulqueen's third-year undergraduate dissertation at the University of Manchester. We thank the two anonymous reviewers and Editor Jim Galvin for their comments that improved the manuscript. Partial funding for Schultz comes from grant NE/I005234/1 from the UK Natural Environment Research Council (NERC) to the Diabatic Influences on Mesoscale Structures in Extratropical Storms (DIAMET) project at the University of Manchester.

References

- Berry G, Reeder MJ, Jakob C.** 2011. A global climatology of fronts. *Geophys. Res. Lett.* **38**: L04809.
- Bjerknes J.** 1919. On the structure of moving cyclones. *Geofys. Publ.* **1**(2): 1–8.
- Bjerknes J.** 1930. Practical examples of polar-front analysis over the British Isles in 1925–1926. *Geophys. Mem.* **5**(10): 1–21 and 28 pp. of figures.
- Bjerknes J, Solberg H.** 1922. Life cycle of cyclones and the polar front theory of atmospheric circulation. *Geofys. Publ.* **3**(1): 3–18.
- Browning KA, Monk GA.** 1982. A simple model for the synoptic analysis of cold fronts. *Q. J. R. Meteorol. Soc.* **108**: 435–452.
- Browning KA, Clough SA, Davitt CSA et al.** 1995. Observations of the meso-scale sub-structure in the cold air of

a developing frontal cyclone. *Q. J. R. Meteorol. Soc.* **121**: 1229–1254.

- Doswell CA III.** 1991. Comments on "Mesoscale convective patterns of the southern High Plains". *Bull. Am. Meteorol. Soc.* **72**: 389–390.
- Hewson TD.** 1998. Objective fronts. *Meteorol. Appl.* **5**: 37–65.
- Hewson TD, Tittley HA.** 2010. Objective identification, typing and tracking of the complete life-cycles of cyclonic features at high spatial resolution. *Meteorol. Appl.* **17**: 355–381.
- Hines KM, Mechoso CR.** 1993. Influence of surface drag on the evolution of fronts. *Mon. Weather Rev.* **121**: 1152–1175.
- Hobbs PV, Locatelli JD, Martin JE.** 1990. Cold fronts aloft and the forecasting of precipitation and severe weather east of the Rocky Mountains. *Weather and Forecasting* **5**: 613–626.
- Hobbs PV, Locatelli JD, Martin JE.** 1996. A new conceptual model for cyclones generated in the lee of the Rocky Mountains. *Bull. Am. Meteorol. Soc.* **77**: 1169–1178.
- Hoskins BJ, Caetano Neto E, Cho H-R.** 1984. The formation of multiple fronts. *Q. J. R. Meteorol. Soc.* **110**: 881–896.
- Keshishian LG, Bosart LF, Bracken WE.** 1994. Inverted troughs and cyclogenesis over interior North America: a limited regional climatology and case studies. *Mon. Weather Rev.* **122**: 565–607.
- Locatelli JD, Schwartz RD, Stoelinga MT et al.** 2002. Norwegian-type and cold front aloft-type cyclones east of the Rocky Mountains. *Weather and Forecasting* **17**: 66–82.
- Mass CF.** 1991. Synoptic frontal analysis: time for a reassessment? *Bull. Am. Meteorol. Soc.* **72**: 348–363.
- Metz ND, Schultz DM, Johns RH.** 2004. Extratropical cyclones with multiple warm-front-like baroclinic zones and their relationship to severe convective storms. *Weather and Forecasting* **19**: 907–916.
- Miles MK.** 1962. Wind, temperature and humidity distribution at some cold fronts over SE England. *Q. J. R. Meteorol. Soc.* **88**: 286–300.
- Namias J.** 1980. The early influence of the Bergen School on synoptic meteorology in the United States. *Pure Appl. Geophys.* **119**: 491–500.
- Namias J.** 1983. The history of polar front and air mass concepts in the United States — an eyewitness account. *Bull. Am. Meteorol. Soc.* **64**: 734–755.
- Newton C, Rodebush Newton H.** 1994. The Bergen School concepts come to America, in *The Life Cyclones of Extratropical Cyclones*. Shapiro MA, Grønås S (eds). American Meteorological Society: Boston, MA, pp 41–59.

Parker DJ. 1998. Secondary frontal waves in the North Atlantic region: a dynamical perspective of current ideas. *Q. J. R. Meteorol. Soc.* **124**: 829–856.

- Prichard B.** 2006. Synoptic analysis. *Weather* **61**: 322.
- Sanders F.** 1999. A proposed method of surface map analysis. *Mon. Weather Rev.* **127**: 945–955.
- Sanders F, Doswell CA III.** 1995. A case for detailed surface analysis. *Bull. Am. Meteorol. Soc.* **76**: 505–521.
- Schultz DM.** 2005. A review of cold fronts with prefrontal troughs and wind shifts. *Mon. Weather Rev.* **133**: 2449–2472.
- Schultz DM.** 2009. *Eloquent Science: A Practical Guide to Becoming a Better Writer, Speaker, and Atmospheric Scientist*. American Meteorological Society: Boston, MA, 412 pp.
- Schultz DM, Blumen W.** 2015. Fronts, in *Encyclopedia of Atmospheric Sciences*, Volume 5, 2nd Edition. North GR, Pyle J, Zhang F (eds). Elsevier: London, pp 337–343.
- Schultz DM, Vaughan G.** 2011. Occluded fronts and the occlusion process: a fresh look at conventional wisdom. *Bull. Am. Meteorol. Soc.* **92**: 443–466, ES19–ES20.
- Schultz DM, Zhang F.** 2007. Baroclinic development within zonally-varying flows. *Q. J. R. Meteorol. Soc.* **133**: 1101–1112.
- Schultz DM, Keyser D, Bosart LF.** 1998. The effect of large-scale flow on low-level frontal structure and evolution in mid-latitude cyclones. *Mon. Weather Rev.* **126**: 1767–1791.
- Shapiro MA, Keyser D.** 1990. Fronts, jet streams and the tropopause, in *Extratropical Cyclones: The Erik Palmén Memorial Volume*. Newton CW, Holopainen EO (eds). American Meteorological Society: Boston, MA, pp 167–191.
- Steenburgh WJ, Mass CF.** 1994. The structure and evolution of a simulated Rocky Mountain lee trough. *Mon. Weather Rev.* **122**: 2740–2761.
- Young MV.** 2014. An unusual case of line convection – the birth of a classic cold front over the UK. *Weather* **69**: 259–263.

Correspondence to: David M. Schultz
David.Schultz@manchester.ac.uk

© 2015 The Authors. *Weather* published by John Wiley & Sons Ltd on behalf of Royal Meteorological Society.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

doi:10.1002/wea.2463