

---

# Measurements of K- shell production cross-section and fluorescence yield for Y element

Hamza Tunç<sup>1</sup>, Rafet Yilmaz<sup>2,\*</sup>, Abdullah Ozkartal<sup>2</sup>, Turgay Ozmen<sup>1</sup>

<sup>1</sup>Institute of Sciences, Yuzuncu Yıl University, Van, Turkey

<sup>2</sup>Department of Physics, Faculty of Sciences and Arts, Yuzuncu Yıl University, Van, Turkey

## Email address:

hamza-tunc23@hotmail.com (H. Tunç), ryilmaz@yyu.edu.tr (R. Yilmaz), abduallahozkartal@yyu.edu.tr (A. Ozkartal), turgay333@hotmail.com (T. Ozmen)

## To cite this article:

Hamza Tunç, Rafet Yilmaz, Abdullah Ozkartal, Turgay Ozmen. Measurements of *K- shell* Production Cross-Section and Fluorescence Yield for Y Element. *American Journal of Physics and Applications*. Vol. 3, No. 1, 2015, pp. 1-5. doi: 10.11648/j.ajpa.20150301.11

---

**Abstract:** *K- shell* X- ray production cross-section have been measured for Y element. Measurements have been carried out at 16.896 keV excitation energy using secondary source. *K* X-rays emitted by samples have been counted by a Si(Li) detector with 160 eV resolution at 5.9 keV. The values of *K- shell* fluorescence yield has been evaluated for Y element. The results obtained for fluorescence yield and *K* X- ray production cross-section has been compared with the theoretically calculated values and other available semiempirical fits values.

**Keywords:** X-Ray Fluorescence, Fluorescence Yields, Cross-Sections

---

## 1. Introduction

*K* X-ray production cross-sections ( $\sigma_{K\alpha}$ ) and fluorescence yields ( $\omega_K$ ) are important in a variety of fields such as atomic physics, nuclear physics, space physics, plasma physics, elemental analysis by X- ray emission technique, medical research, industrial processing. The de-excitation of an atom with an inner-shell *K* vacancy can proceed either by emission of an X-ray photon or by the ejection of Auger electrons. The de-excitation of an atomic shell is characterized by these fluorescence yields and is defined as the probability that a vacancy in the *K*-shell is filled through a radiative transition.

*K* X-ray production cross-sections and fluorescence yields for different elements have been investigated for many years. Earlier experimental *K* X-ray production cross-sections have been measured [1,2]. *K* X-ray production cross-sections have been determined theoretically for all the elements at energies ranging from 10 to 60 keV [3]. *K*-shell X-ray production cross-sections and fluorescence yields have been investigated for different elements [4,5,6].

*K* shell fluorescence yields for different elements have been investigated for many years. Bambynek et al. have fitted their collection of selected most reliable experimental values in the  $13 \geq Z \geq 92$  range [7]. Krause present a table of  $\omega_K$  adopted values for elements  $5 \geq Z \geq 110$  by using all

theoretical and experimental data on the parameter contributing to the *K*-shell fluorescence yields [8]. Hubbel et al. have compiled more recent experimental values [9]. Theoretical values of  $\omega_K$  were obtained using the Hartree-Fock-Slater model [10]. Kostroun et al. present computations for elements in the range  $10 \geq Z \geq 70$  [11]. *K*-shell fluorescence yields have been investigated for different elements [12,13,14]. *K* shell X-ray production cross-sections and fluorescence yields for Cr, Mn, Fe and Co elements have been measured using secondary source by Yilmaz [15]. There is scarcely experimental *K*-shell fluorescence yield data for <sup>39</sup>Y.

The purpose of this study is to examine in low energy the *K* X-ray production cross-section of <sup>39</sup>Y. Measurements of fluorescence yields for elements at various photoionization energies are important because of their widespread use.

In the present study, measurements have been carried out at 16.896 keV. *K* shell has been excited by the *K* X-rays of the secondary source excited by 59.5 keV photons emitted by the primary source. *K- shell* fluorescence yields were obtain by using the measured cross-sections, the theoretical photoionization cross-sections, and fractional X-ray emission rates. The results obtained for *K* X-ray production cross-sections and fluorescence yields are compared with the theoretically calculated values and other available

semiempirical values.

## 2. Experimental

The geometry and the shielding arrangement of the experimental set-up employed in the present work are as shown in Fig.1. Also, it is available in our previous study [16]. As shown in Fig. 1., the sample (target) was placed at a 45° angle with respect to the direct beam from secondary source excited by 59.5 keV photons emitted by the primary source.




Fig . 1. Experimental set-up used for measurements of production cross-section and yields.

The samples were excited by the *K* X-rays of secondary source excited at 59.5 keV  $\gamma$ - rays from a <sup>241</sup>Am point source. The *K* X-ray spectra from target was recorded by a calibrated Si(Li) X-ray spectrometer (full-width at half-maximum (FWHM) =160 eV at 5.96 keV, active area = 12.5 mm<sup>2</sup>, sensitivity depth = 3.5 cm, Be window thickness = 12.5 $\mu$ m) coupled to a Nuclear Data MCA system (ND66B) consisting of a 4096-channel analyzer and spectroscopy amplifier. The net peak areas of the *K* X-rays of each target were determined after background subtraction, escape-peak corrections [17]. The secondary excitation source was pure Nb (99.99%). The excitation energy was taken as average of *K* <sub>$\alpha$</sub>  and *K* <sub>$\beta$</sub>  X-ray energies. For Nb, weighted averages *K* <sub>$\alpha$</sub> , *K* <sub>$\beta$</sub> , *K* <sub>$\alpha\beta$</sub>  energies are 16.584 keV, 18.661 KeV and 16.896 keV, respectively [18].




Fig. 2. A typical *K* X-ray spectrum for <sup>39</sup>Y.

## 3. Data Analysis

The *K*-shell X-ray production cross-section ( $\sigma_{K\alpha}$ ) required for the determination of *K* shell fluorescence yield was evaluated by measuring the characteristic *K* X- ray intensities for <sup>39</sup>Y. The experimental *K* X-ray production cross-sections

were measured using the relation.

$$\sigma_{Ki} = \frac{N_{Ki}}{I_0 G \epsilon_{Ki} \beta t} \tag{1}$$

where  $N_{Ki}$  ( $i = \alpha, \beta$ ) is the number of counts per unit time under the corresponding photopeak. A typical *K* X-ray

spectrum for  $^{39}\text{Y}$  is shown in Fig.2.  $I_0$  is the intensity of exciting radiation,  $G$  is the geometrical factor,  $\epsilon_{Ki}$  is the detector efficiency for the  $K_i$  X-rays,  $t$  is the mass of the sample in  $\text{g}/\text{cm}^2$  and  $\beta$  is the self-absorption correction factor for the incident photons and emitted  $K$  X-ray photons.  $\beta$  is the calculated by using the relation[15],

$$\beta = \frac{1 - \exp[-(\mu_1 / \sin \theta + \mu_2 / \sin \theta)t]}{(\mu_1 / \sin \theta + \mu_2 / \sin \theta)t} \quad (2)$$

where  $\mu_1$  and  $\mu_2$  are the absorption coefficients ( $\text{cm}^2/\text{g}$ ) of incident photons and emitted characteristic X-Rays, respectively [9]. The angle of incident photons and emitted X-rays, with respect to the normal at the surface of the sample,

$\theta$  was equal to  $45^\circ$  in the present set-up. In the present study, as shown in Fig 3, the values of the factors  $I_0\epsilon G$ , which contain terms related to the incident photon flux, geometrical factor and the efficiency of the X-ray detector, were determined by collecting the  $K$  X-ray spectra of thin samples of Fe, Ga, Se, Sr and Zr, in the same geometry in which the  $K$  X-ray fluorescence cross-sections were measured and using the equation [19].

$$I_0 G \epsilon_{Ki} = \frac{N_{Ki}}{\sigma_{Ki} \beta t} \quad (3)$$

where  $N_{Ki}$ ,  $\beta$  and  $t$  are as in Eq.(1).




Fig. 3.  $I_0\epsilon G$  vs.  $K$  X-ray energy.

#### 4. Theoretical Method

The theoretical values of  $K$  X-ray production cross-section  $\sigma_{K\alpha}$  has been calculated using the following equation [5],

$$\sigma_{K\alpha} = \sigma_K^p(E) \omega_K f_{K\alpha} \quad (4)$$

Where  $\sigma_K^p(E)$  is the  $K$ -shell photoionization cross-section for the given element at excitation energy  $E$ ,  $\omega_K$  is the  $K$ -shell fluorescence yield and  $f_{K\alpha}$  is fractional X-ray emission rate for  $K_\alpha$  X-rays and are defined as

$$f_{K\alpha} = (1 + I_{K\beta} / I_{K\alpha})^{-1} \quad (5)$$

where  $I_{K\beta} / I_{K\alpha}$  is the  $K_\beta$  to  $K_\alpha$  X-ray intensity ratio. In the present calculations, the values of  $\sigma_K^p(E)$  were taken from Scofield [20] based on Hartree-Slater potential theory, and the

values of  $\omega_K$  were taken from the tables of Krause[8].

$I_{K\beta} / I_{K\alpha}$  values based on relativistic Hartree-Slater potential theory were used for the evaluation of theoretical  $K$  X-ray cross-sections [21].

5. Results and Discussion  
The experimental values of  $K$  shell X-ray production cross-section for  $^{39}\text{Y}$  at 16.896 keV is listed in Table 1 together with the theoretical value obtained using Eq.(4). The overall error in the present measurements is estimated to be <8%. This error is the quadrature sum of the uncertainties in the different parameters used to deduce  $K$  X-ray production cross-sections, namely, the error in the area evaluation under the  $K_\alpha$  and  $K_\beta$  X-ray peak (2%), in the self-absorption correction factor ratio (<1%), the product  $I_0\epsilon G$ (5%) and the other error in the non-uniform thickness(<1%). In this work, in order to reduce the absorption, thin sample was also used as the target. In order to reduce the statistical error, the spectra was recorded and about 6000 counts were collected under the  $K_\alpha$  peak. It can be seen from Table 1 that our measured value is in good

agreement, within the experimental uncertainties, with the calculated theoretical value.. The agreement between the

presently measured *K* shell X-ray production cross-section and theoretical prediction is within % 3 for <sup>39</sup>Y element.

**Table 1.** Experimental and theoretical  $K_{\alpha}$  X- ray cross-sections (barns/atom) for <sup>39</sup>Y.

Element	Excitation energy (keV)	Present Work	Theoretical predictions
<sup>39</sup> Y	16.896	6002±468	6187.863

The present measured value of *K*-shell fluoresans yield ( $\omega_K$ ), deduced using Eq.(4) for <sup>39</sup>Y is compared with the other available calculated values, and semiempirical fits in Tables 2,3. In this study,our experimental result agree to the experimental result agree to <3.9% for <sup>39</sup>Y element with The

*K*-shell fluorescence yields calculated using a semiempirical expression by Hubbell et al.[9] , Bambynek et al., [7] and Krause et al. [8]. The present measurement is in good agreement with the semiempirical values deduced by Hubbell et al.[9], with 0.4% (Table 3).

**Table 2.** Comparison of the present *K* shell fluorescence yield ( $\omega_K$ ) and theoretical values for <sup>39</sup>Y.

Element	Present work	Walters [10]	Kostroun [11]
<sup>39</sup> Y	0.688±0.053	0.742	0.722

**Table 3.** Present experimental result and semiempirical fits values of  $\omega_K$  for <sup>39</sup>Yb

Element	Present work	Bambynek [7]	Hubbell [9]	Krause [8]
<sup>39</sup> Y	0.688±0.053	0.7155	0.685	0.710

As a result, the present agreement between the theoretical and present experimental values lead to the conclusion that the data presented here will benefit for analytical purpose and satisfactory for many other applications employing the fundamental parameter approach. Also, secondary source are quite useful in low energy studies.

## References

- [1] Garg, L., Mehta, D., Kumar, S., Mangal, P.C., Trehan, P.N., 1985. Energy dependence of photon-induced  $K_{\alpha}$  and  $K_{\beta}$  X-ray fluorescence cross-sections for some elements with  $20 \leq Z \leq 56$ . X-ray Spectrom. 14, 165-169.
- [2] Rao. D. V., Cesereo, R. and Gigante, G. E. (1993d). Photon-excited K X ray fluorescence cross-sections for some light elements in the range 20-60 keV. X-ray Spectrom. 22, 406.
- [3] Krause, M.O., Nester, C. V., Sparks Jr., C.J., Ricci, E., 1978. X-ray fluorescence cross-sections for K and L X-rays of elements. Oak Ridge national laboratory, Report ORNL 5399.
- [4] Kumar, S., Singh, S., Mehta, D., Singh, N., Mangal, P.C., Trehan, P. N., 1987. Measurement of K-shell fluorescence cross-sections for some elements with  $23 \leq Z \leq 55$  in the energy range 8-60 keV, X- ray Spectrom. 16, 203-206.
- [5] Durak, R., Erzenoğlu S., Kurucu Y. and Şahin Y., 1998. Measurement of photon-induced K X-ray production cross-sections for some elements with  $40 \leq Z \leq 70$  at 122 keV. Radiation Physics and Chemistry.51, 45-48.
- [6] Özdemir, Y., Durak, R., Öz. E., 2002. K shell X- ray fluorescence cross-sections and fluorescence yields in some medium-Z elements. Radiation Physics and Chemistry. 65, 199-204.
- [7] Bambynek, W., Crasemann, B., Fink, R. W., Freund, H.U., Mark, H., Swift, C.D., Price, R.E., Rao, P.V., 1972. X-ray fluorescence yields, Auger and Coster-Kronig transitions probabilities. Rev. Mod.phys. 44, 716-813
- [8] Krause, M.O., 1979. Atomic radiative and radiationless yields for K and L shell. J.Phys.Chem. Ref.Data 8, 307-327
- [9] Hubbell, J.H., and Seltzer, S. M., 1995. Tables of X-ray mass attenuation coefficients and mass energy absorption coefficients 1 keV to 20 MeV for elements Z=1 to 92 and 48 additional substance of dosimetric interest. NISTIR 5632.
- [10] Walters, D.L., Bhalla, C. P., 1971. Nonrelativistik Auger rates, X-rays rates, and fluorescence yields for the K shell. Phys. Rev. A 3, 1919-1927.
- [11] Kostroun, V.O., Chen, M. H., Crasemann, B., 1971. Atomic radiation transition probabilities to the is state and theoretical K- shell fluorescence yields. Phys. Rev. A 3, 533-545.
- [12] Arora, S.K., Allawadhi, K.L., Sood, B.S., 1981, Measurement of K-shell fluorescence in elements  $28 \leq Z \leq 53$ , Physica B, C 111C, 71-75.
- [13] Pious, J.K., Balakrishna, K.M., Lingappa, N., Siddappa, K., 1992. Total K fluorescence yields for Fe, Cu, Zn, Ge and Mo, J. Phys. B:Atom. Mol. Opt. Phys. 24, 1155-1160.
- [14] Şimşek, Ö., Yılmaz, D., Karagöz, M. Ertuğrul, 2002, Measurement of K-shell fluorescence cross sections and K shell fluorescence yields for the atomic region  $22 \leq Z \leq 64$  by 59.5 keV photons. Journal of Radioanalytical and Nuclear Chemistry 253, 143-147.
- [15] Yılmaz, R., 2014. Measurements of K shell X-ray production cross-sections and fluorescence yields for Cr, Mn, Fe and Co elements. Radiation Physics and Chemistry. 96, 1-4.

- [16] Yilmaz, R., 2012. Measurements of  $L_{2,3}$  subshells X-ray production cross-sections and average  $L_{2,3}$  subshells fluorescence yield values of some elements in the atomic number range  $70 \leq Z \leq 78$ . Radiation Physics and Chemistry. 81,1539-1542
- [17] Öz E., 2006. Determination of emission probabilities of Auger electrons and K-L shell radiative vacancy transfer probabilities for 17 elements from Mn to Mo at 59.5 keV. Journal of Quantitative Spectroscopy & Radiative Transfer 97, 41-50
- [18] Storm, E., Israel, I., 1970. Photons cross-sections from 1 keV to 100 MeV for elements  $Z=1$  to  $Z=100$ . Nuclear Data Tables A 7, 565-681.
- [19] Durak, R., Özdemir, Y., 2001. Measurement of K-shell fluorescence cross-sections and yields of 14 elements in the atomic number range  $25 \leq Z \leq 47$  using photoionization. Radiation Physics and Chemistry. 61, 19-25.
- [20] Scofield, J.H., 1973. Theoretical photoionisation cross-sections from 1 to 1500 keV. Report UCRL 51326, Lawrence Livermore Laboratory, Livermore, CA.
- [21] Scofield, J.H., 1974. Relativistic Hartree-Slater values for K and L X-ray emission rates. Atom. Data Nucl. Data Tables 14, 121-137.