

Spontaneous Rupture of the Stomach in a Healthy Adult Man After Sodium Bicarbonate Ingestion

MICHAEL R. MASTRANGELO, M.D.; and EDWARD W. MOORE, M.D.

Medical College of Virginia, Virginia Commonwealth University; Richmond, Virginia.

SPONTANEOUS RUPTURE of the stomach is a rare event that usually is fatal. We report the case of a patient in

whom rupture occurred within 1 minute after ingestion of a label-recommended dose of sodium bicarbonate. The patient had surgery and survived. This is the fifth such case reported in the literature.

A 31-year-old white man was apparently in excellent health before the present illness. He had no history of peptic ulcer disease, abdominal trauma, or surgery. At approximately 2000 h on 25 June 1981, the patient began eating dinner at a Mexican restaurant. He had two margaritas, an order of nachos, and a large combination plate. Because of the size of the meal, he did not have dessert but did drink a cup of coffee. After arriving home at about 2200 h, the patient noted an uncomfortable feeling of fullness in the abdomen. Because he could not find any Alka-Seltzer (Miles Laboratories, Elkhart, Indiana), he went to the refrigerator and removed a box of baking soda, a product he remembered could be used for indigestion (although he had not taken it before). He placed $\frac{1}{2}$ teaspoon of baking soda in $\frac{1}{2}$ glass of water and quickly drank the solution.

Within 1 minute after drinking the bicarbonate, the patient had severe abdominal pain without nausea or retching. Believing that he could induce vomiting to relieve the pain, he went to the bathroom to attempt to vomit. By this time, however, the pain was so intense that he could no longer stand erect, so he lay down on the floor. He then noted that his abdomen was distended, and he released his belt from the buckle. He called for help from his fiancée, who immediately recognized his dire condition and called the rescue squad. The patient was rushed to the hospital. At no time before entering the hospital did the patient eructate, retch, or vomit.

At a local hospital, the patient was believed to have either a Boerhaave rupture of the esophagus or perforated peptic ulcer and was taken to the operating room for exploratory laparotomy. At operation, the peritoneal cavity was noted to contain a large amount of coffee-ground material containing beans and other food particles. After vigorous irrigation, a 5-cm linear perforation was noted along the lesser curvature of the stomach, beginning near the esophagogastric junction. The stomach was noted to contain a "massive" amount of retained food. No evidence of ulceration or other abnormality was seen, and the perforation was oversewn in two layers. Histologic examination of the margins of the involved stomach wall revealed "full thickness hemorrhage and focal necrosis, but relatively little attendant acute inflammation."

The patient tolerated the procedure well, but his postoperative course was complicated by the development of a right subphrenic abscess, subhepatic abscess, and respiratory failure believed secondary to adult respiratory distress syndrome. After 21 days in the hospital, he was discharged and has continued to do well.

"Spontaneous" rupture of the stomach is an uncommon condition—only 71 cases have been reported in the English literature. Six of these were associated with antacid ingestion (1-6); only four reports specified sodium bicarbonate as the antacid (1-4).

Presumably, the rupture is caused by increasing intragastric pressure secondary to distention of the stomach by any combination of solid, liquid, or gas. In addition to the previous cases of gastric rupture associated with sodium bicarbonate ingestion, several cases in the literature suggest that fermentation might have occurred in the stomach (7, 8), producing large quantities of gas leading to gastric distention and rupture. This pathogenesis is similar to that in the present case—ingestion of sodium bicarbonate resulted in the rapid production of enough carbon dioxide gas to rupture a stomach already distended with food.

Spontaneous rupture of the stomach is associated with a high mortality rate. In Albo and colleagues' series (9), there was an 85% mortality rate; however, 26 of 44 pa-

tients did not undergo surgery, and all of the them died. Of the 18 patients who did have surgery, there was a 65% mortality rate. In the 30 cases reported since Albo and colleagues' series, all patients had surgery, and 11 survived, producing a 63% mortality rate.

We present the fifth case of spontaneous rupture of the stomach associated with sodium bicarbonate ingestion. Our patient was uncomfortable until he had ingested sodium bicarbonate, at which time he became acutely ill. We believe that his initial discomfort was due to "overdistention" of the stomach and that his acutely ill state was due to gastric rupture secondary to release of carbon dioxide by the reaction of sodium bicarbonate with gastric acid. (The kinetics of this reaction will be discussed in a later report.) Because of the high mortality rate associated with this lesion and because of the availability of effective antacids without gas production as a side effect, we believe that the use of sodium bicarbonate should be discontinued for the treatment of heartburn or other gastrointestinal symptoms.

ACKNOWLEDGMENTS: The authors thank Dr. Robert Wagner of the Arlington Hospital, Arlington, Virginia, for his cooperation in obtaining clinical and pathological material pertinent to the case.

Grant support: by Research Grant #AM 32130 from the National Institutes of Health. Dr. Mastrangelo is the recipient of an NIH Research Fellowship Award (#AM 07244) from the National Institute of Arthritis, Diabetes, Digestive, and Kidney Diseases.

► Requests for reprints should be addressed to Edward W. Moore, M.D.; Box 711, MCV Station, Medical College of Virginia, 12th & Broad Streets; Richmond, VA 23298.

REFERENCES

1. BRUNO MS, GRIER WRN, OBER WB. Spontaneous laceration and rupture of esophagus and stomach. *Arch Intern Med.* 1963;112:574-83.
2. LEMMON WT, PASCHAL GW. Rupture of the stomach following ingestion of sodium bicarbonate. *Ann Surg.* 1941;114:997-1003.
3. MURDFIELD P. Rupture of the stomach from sodium bicarbonate. *Klin Wochenschr.* 1926;5:1613. (Abstract in *JAMA.* 1926;87:1692-3 and *Ann Surg.* 1929;89:247-63.)
4. ZER M, CHAIMOFF C, DINTSMAN M, TIKVA P. Spontaneous rupture of the stomach following ingestion of sodium bicarbonate. *Arch Surg.* 1970;101:532-3.
5. KREGEL LA. Spontaneous rupture of the stomach. *Am J Surg.* 1955;21:505-7.
6. SHALDON C. Spontaneous rupture of the stomach. *Br J Surg.* 1961;48:640-1.
7. FIESTER RF, ZINN LD. Spontaneous gastric rupture during pregnancy (case report). *W Va Med J.* 1975;71:110.
8. FLEMING LB, HORTON JAG, WAGGET J. Spontaneous rupture of the stomach: a rare complication of pyloric stenosis. *Br J Surg.* 1966;53:384-7.
9. ALBO R, DE LORIMIER AA, SILEN W. Spontaneous rupture of the stomach in the adult. *Surgery.* 1963;53:797-805.

©1984 American College of Physicians