

Pseudorandom Generators Hard for k -DNF Resolution and Polynomial Calculus Resolution

Alexander A. Razborov *

May 28, 2014

Abstract

A pseudorandom generator $G_n : \{0, 1\}^n \rightarrow \{0, 1\}^m$ is *hard* for a propositional proof system P if (roughly speaking) P can not efficiently prove the statement $G_n(x_1, \dots, x_n) \neq b$ for *any* string $b \in \{0, 1\}^m$. We present a function ($m \geq 2^{n^{\Omega(1)}}$) generator which is hard for $\text{Res}(\epsilon \log n)$; here $\text{Res}(k)$ is the propositional proof system that extends Resolution by allowing k -DNFs instead of clauses.

As a direct consequence of this result, we show that whenever $t \geq n^2$, every $\text{Res}(\epsilon \log t)$ proof of the principle $\neg \text{Circuit}_t(f_n)$ (asserting that the circuit size of a Boolean function f_n in n variables is greater than t) must have size $\exp(t^{\Omega(1)})$. In particular, $\text{Res}(\log \log N)$ ($N \sim 2^n$ is the overall number of propositional variables) does not possess efficient proofs of $\mathbf{NP} \not\subseteq \mathbf{P}/\text{poly}$. Similar results hold also for the system PCR (the natural common extension of Polynomial Calculus and Resolution) when the characteristic of the ground field is different from 2.

As a by-product, we also improve on the small restriction switching lemma from [SBI04] by removing a square root from the final bound. This in particular implies that the (moderately) weak pigeonhole principle PHP_n^{2n} is hard for $\text{Res}(\epsilon \log n / \log \log n)$.

*Institute for Advanced Study, Princeton, US, on leave from Steklov Mathematical Institute, Moscow, Russia. Supported by The State of New Jersey and by the RFBR grant 02-02-01290. Current address: University of Chicago, US, razborov@cs.uchicago.edu.

1. Introduction

Propositional proof complexity is an area of study that has seen a rapid development over the last decade. It plays as important a role in the theory of feasible proofs as the role played by the complexity of Boolean circuits in the theory of efficient computations. And in most cases the basic question of propositional proof complexity boils down to this. Given a mathematical statement encoded as a propositional tautology ϕ and a class of admissible mathematical proofs formalized as a propositional proof system P , what is the minimal possible complexity of a P -proof of ϕ ?

1.1. General overview

For most “interesting” propositional proof systems P one can easily define the accompanying (non-uniform) complexity class \mathcal{C}_P typically consisting of functions computable by lines allowed in efficient P -proofs. This correspondence leads to a classification of propositional proof systems that is somewhat imprecise and potentially (and hopefully) time-dependent but nonetheless very instructive. Namely, we call a propositional proof system P *weak* if we (currently) know how to prove super-polynomial lower bounds for the accompanying circuit class \mathcal{C}_P and *strong* otherwise.

There is a steady progress in studying the complexity of proofs in weak proof systems surveyed e.g. in [Urq95, Kra95, Raz96, BP01, Pud98, Raz02a].

For strong proof systems the current situation is by far more miserable. Although there are no rigorous results along these lines (and, moreover, this feeling is not universal – see e.g. [Kra04]), the empirical evidence strongly suggests that lower bounds for a proof system P are even harder to attain than computational lower bounds for the companion class \mathcal{C}_P . Therefore, with our current understanding, we can not apparently hope to show lower bounds for systems like Frege or Extended Frege without first making a major breakthrough in complexity theory.

A more accessible task which (in the author’s opinion) is almost as interesting would be to show at least that proof complexity lower bounds are *at most* as hard as comparable problems in the computational world. Let us (informally) identify this task as

PROVING LOWER BOUNDS FOR STRONG PROOF SYSTEMS P
 LIKE FREGE OR EXTENDED FREGE MODULO ANY HARDNESS
 ASSUMPTION IN THE PURELY COMPUTATIONAL WORLD, HOW-
 EVER STRONG BUT STILL NATURAL AND BELIEVABLE

This task will be referred to as *proving conditional lower bounds* (for the proof system P). It should be remarked in this respect that $\mathbf{NP} \neq \text{co-NP}$ implies lower bounds for *any* propositional proof system whatsoever. Therefore, *purely computational* above refers to the demand that the assumption itself should speak only about computations and should not attempt to restrict the power of proofs even in a disguised form.

One extremely exciting and, in a sense, model approach to this task was gradually developed in the sequence of papers [Kra94, Raz95b, BPR97, Kra97a, Pud97] and finally became known as the *Efficient Interpolation Property* (EIP in what follows). EIP was shown to be true for some weak proof systems and it was also remarked that for every proof system (be it weak or strong) EIP implies conditional lower bounds. Unfortunately, it turned out rather soon [KP98, BPR00] that neither Frege nor Extended Frege have Efficient Interpolation modulo (somewhat ironically) hardness assumptions of the same sort that are needed to prove conditional lower bounds for proof systems *with* EIP.

This omnipresent hardness assumption is nothing other than the existence of pseudorandom generators (arbitrary or specific) which also turns out to be the main primitive of the modern cryptography. After the (apparent) failure of the efficient interpolation approach, it was independently proposed in [Kra01a, ABRW00] to employ pseudorandom generators for proving conditional lower bounds in a more direct manner. On the conceptual level, a mapping $G_n : \{0, 1\}^n \rightarrow \{0, 1\}^m$, where $m > n$, is called *hard* for a propositional proof system P if P can not efficiently prove the (properly encoded) statement $G_n(x_1, \dots, x_n) \neq b$ for *any*¹ string $b \in \{0, 1\}^m$. Since $m > n$, for at least half of all b 's this statement is a tautology. Therefore, conditional lower bounds for a proof system P follow from the following task:

¹The reader wondering whether it might be more natural to weaken here the condition “for any string b ” to “for some string b ” or perhaps to “for some explicit string b ” is referred to an extensive discussion of this issue in [ABRW00, Section 1] where the concept was introduced.

PROVE THAT FOR A REASONABLE CLASS OF MAPPINGS $G_n : \{0, 1\}^n \rightarrow \{0, 1\}^m$ WITH $m > n$, THEIR HARDNESS IN ANY REASONABLE COMPUTATIONAL OR COMBINATORIAL SENSE IMPLIES HARDNESS FOR P .

this will be referred to as the *generator approach* (to conditional lower bounds for P).

The generator approach certainly does not work for *arbitrary* mappings G_n believed to be pseudorandom generators in the standard sense of [Yao82], and specific counterexamples almost immediately follow from the results in [KP98] on the limitations of EIP. On the positive side, it was observed in [ABRW00] that for proof systems P with EIP there is an easy and general way of *converting any* pseudorandom generator that is *computationally* hard (in the standard sense) into a pseudorandom generator that is *hard for P* . No such *general* transformation is known for a single non-trivial proof system without EIP.

Thus, it is vital for the generator approach that we somehow restrict the class of mappings for which one hopes to trade computational hardness for proof complexity hardness. Along these lines [ABRW00] specifically proposed to consider the class of Nisan-Wigderson generators (concrete results from that paper as well as from later improvements [AR03, Kra04] will be reviewed after we are done with this general overview). Some arguments advocating this choice (as opposed to other classical cryptographic constructions) were presented in [ABRW00] and in the introduction to [Raz04b]: the principle expressing hardness of NW-generators is tightly related to such familiar personages in proof complexity as the pigeonhole principle and Tseitin tautologies.

Elaborating on these arguments, we can further remark that the NW-generator is in a sense the quintessence of the very idea of “local consistency”. Namely, the information contained in the output bits of a NW-generator is “local” (in the sense that every output bit depends only on a “small” subset of input bits which are “nearly independent” for different output bits), and it is locally consistent to such an extent that no interesting conclusion about the global behaviour of the generator can be obtained by an “easy” analysis of this local knowledge. This simple methodology is behind a great deal of lower bounds existing in proof complexity for weak proof systems, and it is also behind the efficient interpolation property.

This methodology also has a very clean mathematical meaning. A (non-

existent) falsifying assignment to the tautology ϕ corresponds to a manifold with given local properties. The proof system P tries to argue that no such manifold may exist using tools at its disposal. And we (lower bounds provers) try to fool it by feeding into the potential proof something which looks like a (non-existing) manifold to such an extent that P can not discern the difference. See [Raz98] for (apparently) the cleanest implementation of this intuitive scheme.

Anyway, the moving forces that make the Nisan-Wigderson generator work in the computational world are of so general nature that we are ready to spell out the formal conjectures that the generator approach *always* works for Nisan-Wigderson generators. More specifically (assuming that the constructions are based on combinatorial designs with the same parameters as in the seminal paper [NW94]),

Conjecture 1. *Any NW-generator based on any poly-time function that is hard on average for $\mathbf{NC}^1/poly$, is hard for the Frege proof system.*

Conjecture 2. *Any NW-generator based on any function in $\mathbf{NP} \cap co-\mathbf{NP}$ that is hard on average for $\mathbf{P}/poly$ (e.g. $B(f^{-1}(r))$, where $f(x)$ is any one-way permutation and $B(x)$ its hard-core bit), is hard for Extended Frege.*

The suggestion to use Nisan-Wigderson generators for lower bounds in proof complexity has been recently re-iterated in [Kra04]. That paper also proposes a paradigm similar in spirit to the construction from [Gol11] in the context of computational complexity: hardness of the resulting mapping should depend on the randomness of the base functions rather than their complexity. So far all known results on the hardness of NW-generators for weak proof systems have not directly appealed to the randomness and used instead specific combinatorial properties of the base functions. But of course it remains to be seen yet which of the two paradigms (if any) will turn out more fruitful in the long run.

The task of proving lower bounds (even conditional) for strong proof systems is, in the author's opinion, extremely interesting and well justified in its own right (whereas the popular motivation that this should be regarded as an intermediate step in approaching the \mathbf{NP} vs. $co-\mathbf{NP}$ problem looks, again in the author's opinion, more of a speculation). One venerable way, however, to make this study even more interesting is to look at the proof complexity of

statements whose *validity* is also not known, and whose importance stretches well beyond any proof-theoretical studies.

To that end, [Raz95a] proposed² to study the proof complexity of the principle $\neg\text{Circuit}_t(f_n)$ expressing that the circuit size of the Boolean function f_n in n variables, given as its truth-table, is lower bounded by $t = t(n)$ (thus, e.g. $\mathbf{NP} \not\subseteq \mathbf{P}/\text{poly}$ is essentially equivalent to the validity of $\neg\text{Circuit}_{t(n)}(\text{SAT}_n)$ for some $t(n) \geq n^{\omega(1)}$). [Raz95a] put forward the thesis (so far not refuted) that *all existing* proofs of lower bounds for restricted classes of circuits and for explicit functions translate to Extended Frege proofs (often to much weaker proof systems) of size $2^{O(n)}$. This makes the question of the efficient provability of the original principle $\neg\text{Circuit}_t(f_n)$ for *general* circuits even more intriguing.

The connection of this question to the generator approach above is the same as in the context of Natural Proofs [RR97]. Namely, if we have a function pseudorandom generator $G_n : \{0, 1\}^{t_0} \rightarrow \{0, 1\}^{2^n}$ that is hard for a proof system P , such that the associated predicate $G(x)_y$ ($x \in \{0, 1\}^{t_0}$, $y \in \{0, 1\}^n$) can be computed by a size t circuit, then for every fixed seed $x \in \{0, 1\}^{t_0}$, the Boolean function with the truth-table $G(x)$ is also computed by a size t circuit. Since for any given f_n the system P can not efficiently refute that f_n is different even from the functions $G(x)$ in the image of the generator G , it is not capable of efficient proofs of $\neg\text{Circuit}_t(f_n)$. In plain words,

TO SHOW THAT P DOES NOT HAVE EFFICIENT PROOFS OF THE FORMULA $\neg\text{Circuit}_t(f_n)$, IT SUFFICES TO DESIGN A SUFFICIENTLY CONSTRUCTIVE PSEUDORANDOM GENERATOR HARD FOR P AND SUCH THAT THE NUMBER OF OUTPUT BITS, AS A FUNCTION OF THE NUMBER OF INPUT BITS, IS AS LARGE AS POSSIBLE.

The larger number of output bits we can manage, the smaller are the parameters t_0, t (relatively to n) and the stronger is the result. In particular, in order to conclude the efficient unprovability of $\mathbf{NP} \not\subseteq \mathbf{P}/\text{poly}$ (or, for that matter, that any function f_n is not in \mathbf{P}/poly), one needs a generator that

²[Raz95a] dealt with provability of Σ_0^b -statements in the theories of Bounded Arithmetic, which is the uniform counterpart of propositional proof complexity. At the suggestion of Jan Krajíček, in later papers it was recast in more convenient framework of propositional proof complexity.

stretches n bits to 2^{n^ϵ} bits; such generators are commonly known as *function pseudorandom generators*.

The tight connection between pseudorandom generators and the tautologies $\neg\text{Circuit}_t(f_n)$ has been also fruitfully exploited from more structural point of view. We already remarked above that $\mathbf{NP} \neq \text{co-NP}$ implies the existence of hard tautologies for any propositional proof system, but this does not give any clue as to what these hard tautologies actually are. [IKW02] proved that under the assumption $\mathbf{NEXP} \not\subseteq \mathbf{P}/\text{poly}$, it is the specific tautologies $\neg\text{Circuit}_t(f_n)$ (for any f_n whatsoever) that are hard for any proof system. R. Impagliazzo (see a footnote in [Kra04, Section 1]) and independently M. Alekhnovich recently observed that the same conclusion holds under the assumption $\mathbf{BPP} \not\subseteq \mathbf{NP}$ (stronger than $\mathbf{NP} \neq \text{co-NP}$). Although none of these two assumptions looks particularly plausible (and none of them is “purely computational” either), this still serves as another indication of the “distinguished” character of the tautologies $\neg\text{Circuit}_t(f_n)$.

The project of proving lower bounds for stronger and stronger classes of circuits until we arrive at $\mathbf{P} \neq \mathbf{NP}$ has met a solid obstacle in the form of Natural Proofs [RR97]. The project of proving lower bounds for stronger and stronger proof systems until we arrive at the (Extended) Frege proof system is restricted by the empirical observation that this task is even harder than the previous one. Given this gloomy background, fulfilling the generator approach for strong proof systems is certainly not an easy task, and the progress in this direction is much slower than originally hoped. However, it seems that we currently do not know of any general reasons (like those for the two previous projects formulated above) making us suspect that this task is unfeasible and/or requires an entirely different view of the subject.

1.2. Previous results and our contributions

In all known partial results along the generator approach, Nisan-Wigderson generators play the central role. Hardness results for generators of this kind are determined by combinatorial properties of the underlying set system, conditions imposed on the base functions, and by the specific way their computation is encoded as a propositional tautology. Disregarding for the moment all these technical issues, on the conceptual level [ABRW00] proved that the NW-generator is hard for Resolution but only when the complexity is measured by *width*. Another result from [ABRW00] says that the *Nisan*

generator (that is, the partial case of the NW-generator in which all base functions must be linear mod 2) is hard for Polynomial Calculus (PC in what follows) over fields \mathbb{F} with $\text{char}(\mathbb{F}) \neq 2$. The latter result was further extended and generalized in [AR03] to show that the Nisan-Wigderson generator is hard for PC over any field.

Results for resolution width and polynomial calculus degree are applicable to function generators stretching n bits to 2^{n^ϵ} bits. As long as the *size* complexity measures are concerned, [ABRW00] exhibited hard Nisan-Wigderson generators for the system PCR (that is a natural common extension of Polynomial Calculus and Resolution) but only when $m \leq o(n^2)$. This poor input/output ratio hindered their potential application to proving that $\mathbf{NP} \not\subseteq \mathbf{P}/\text{poly}$ is hard even for Resolution, and this was established by somewhat different methods in [Raz04a, Raz04b].

A prominent *general* way for enhancing the I/O performance of pseudorandom generators in proof complexity has been recently proposed in [Kra04]. Like the classical constructions in computational complexity [Yao82, GGM86], it is very natural to try to achieve this goal by composing the given generator with itself. Unfortunately, it is far from clear whether hardness in the context of proof complexity is preserved under composition. [Kra04] proposed a way around this difficulty by showing that it is indeed the case if hardness is replaced by a stronger notion of *s-iterability* (the latter, in turn, being a variant of a similar notion of freeness earlier introduced in [Kra01b]). As a first application of this approach, [Kra04] showed that one particular construction of the Nisan generator from [ABRW00] can be iterated with itself once, thus giving a pseudorandom generator with $m = n^{3-\epsilon}$ output bits that is hard for Resolution.

In the current paper we continue this line of research. Let $\text{Res}(k)$ be the propositional proof system that extends Resolution by allowing as its lines arbitrary k -DNFs. Our first main result (Theorem 2.7) exhibits Nisan generators that are hard for $\text{Res}(k)$ and stretch n input bits to as many as $n^{(\epsilon \log n)/k}$ output bits. A relatively easy modification of this argument for $k = 1$ shows that this generator (from n to $n^{\epsilon \log n}$ bits) is hard not only for Resolution but also for its extension PCR (Theorem 2.18)³. These results were proved independently of [Kra04]; the proof method uses the resolution width/PC degree bounds from [ABRW00] cited above in combination with

³Here and in the rest of introduction we implicitly assume that $\text{char}(\mathbb{F}) \neq 2$.

the machinery from the recent paper [SBI04] based upon the so-called *small restriction switching lemma*. In order to bring these two together, we also introduce a special kind of random restrictions specifically tailored to deal with Nisan generators.

Then, using a very simple reduction, we show that our generator is not only hard for $\text{Res}(k)$ and PCR but it is in fact $\exp(n^{\Omega(1)})$ -iterable for these systems (Theorems 2.10, 2.19). According to the paradigm from [Kra04], this implies that if we compose this generator with itself as many as $\exp(n^{\Omega(1)})$ times, the resulting mapping will still be hard for $\text{Res}(k)/\text{PCR}$. Applying in particular the classical GGM-construction [GGM86], we get in this way a function generator $G_n : \{0, 1\}^n \rightarrow \{0, 1\}^{2^{n^\epsilon}}$ that is hard for $\text{Res}(\epsilon \log n)$, $\epsilon > 0$ a sufficiently small constant (Theorem 2.12), and for PCR. As we discussed in Section 1.1, this implies that neither $\text{Res}(\log \log N)$ (where $N \sim 2^n$ is the overall number of propositional variables) nor PCR possess efficient proofs of $\mathbf{NP} \not\subseteq \mathbf{P}/\text{poly}$ (Theorems 2.13, 2.20), which are the first results of this kind for *any* propositional proof system that (most likely) does not have the Efficient Interpolation Property.

In the course of this work we were able to get a quadratic (in k) improvement of the small restriction switching lemma from [SBI04] based on Janson inequality (Lemma 4.4). This in particular implies that the moderately weak pigeonhole principle $PHP_n^{2^n}$ is exponentially hard for $\text{Res}(k)$ when $k \leq (\epsilon \log n) / \log \log n$ (Theorem 2.15), as opposed to $k \leq \epsilon \sqrt{\log n / \log \log n}$ in [SBI04] that in turn was an improvement on the previous papers [BT88] ($k = 1$, i.e. resolution) and [ABE02] ($k = 2$).

Finally we prove a miscellaneous result about the complexity of Nisan generators themselves that indicates the sensitivity of their proof complexity behaviour with respect to the choice of encoding. Namely, we show how to modify the (natural and reasonable) encoding of the Nisan generator used in the rest of the paper in such a way that this generator becomes hard for PCR even in the functional case $m = 2^{n^\epsilon}$ (Theorem 2.21). This encoding, however, is in a sense very bad: it does not seem to allow any reduction to $\neg \text{Circuit}_t(f_n)$, and the proof method apparently fails already for $\text{Res}(2)$.

2. Definitions and statements of our results

In this section we typically confine ourselves to defining only those notions that are needed for stating our main results. Auxiliary concepts needed for

their proofs will normally appear in respective places.

Let x be a *propositional variable*, i.e. a variable that ranges over the set $\{0, 1\}$. A *literal* of x is either x (denoted sometimes as x^1) or \bar{x} (denoted sometimes as x^0). A *clause* [*term*] is either a constant 0 or 1 (corresponding to FALSE and TRUE, respectively) or a disjunction [conjunction, respectively] of literals. A *CNF* [*DNF*] is a conjunction of clauses [disjunction of terms], often specified as the set of all participating clauses [terms, respectively]. Accordingly, a clause/term/CNF/DNF is a sub-clause/sub-term/sub-CNF/sub-DNF of another clause/term/CNF/DNF if every literal/literal/clause/term appearing in the first, appears also in the second. A clause/term/CNF/DNF is *monotone* if it does not contain occurrences of negated literals \bar{x} .

For a Boolean function f [a propositional formula F], let $Vars(f)$ [$Vars(F)$] be the set of its essential variables [the set of variables explicitly occurring in F , respectively]. The *width* of a clause C [of a term t] is defined as $w(C) \stackrel{\text{def}}{=} |Vars(C)|$ [$w(t) \stackrel{\text{def}}{=} |Vars(t)|$, respectively]. A *k-CNF* [*k-DNF*] is a CNF [DNF] in which all clauses [terms, respectively] are of width at most k . $|F|$ is the number of terms in a DNF F .

An *assignment* to a Boolean function f [a propositional formula F] is a mapping $\alpha : Vars(f) \rightarrow \{0, 1\}$ [$\alpha : Vars(F) \rightarrow \{0, 1\}$, respectively]. A *restriction of f* [of F] that, depending on the context, will be sometimes called a *partial assignment* is a mapping $\rho : Vars(f) \rightarrow \{0, 1, \star\}$ [$\rho : Vars(F) \rightarrow \{0, 1, \star\}$, respectively]. We let $\text{sup}(\rho) \stackrel{\text{def}}{=} \rho^{-1}(\{0, 1\})$ denote the set of assigned variables. The *restriction of a function f* [of a formula F] by ρ , denoted $f|_\rho$ [$F|_\rho$], is the Boolean function [propositional formula] obtained from f [from F , respectively] by setting the value of each $x_i \in \text{sup}(\rho)$ to $\rho(x_i)$, and leaving each $x_i \notin \text{sup}(\rho)$ unassigned. In the case of propositional formulas we as usual assume that simplifications are performed only when a sub-formula has become explicitly constant. A *variable substitution* of variables in V_1 by variables in V_2 is a mapping ρ that takes variables in V_1 to either propositional constants or *literals* of variables in V_2 (thus, restrictions are viewed as a special case of variable substitutions with $V_2 = V_1$). Variable substitutions ρ of variables in $Vars(f)$ or $Vars(F)$ also naturally act on the Boolean function f or propositional formula F , and, as before, we denote the result of this action by $f|_\rho$, $F|_\rho$.

For an integer n , let $[n] \stackrel{\text{def}}{=} \{1, 2, \dots, n\}$. Whenever we use probabilistic methods, random variables will always appear in the bold face, including

deterministic parameters they depend on, if any. We extensively use the Ω -notation (customary in complexity theory) that is opposite to the ordinary O -notation. For example, given two functions f, g with values in the set of non-negative reals, $f \geq \Omega(g)$ means that there exists an absolute constant $\epsilon > 0$ such that $f \geq \epsilon g$ for any specification of the parameters occurring in f, g .

Definition 2.1 ([Kra01a, ABRW00]) Let $m > n$, C be a Boolean circuit with n inputs x_1, \dots, x_n and m outputs, and $b \in \{0, 1\}^m$ be an arbitrary Boolean vector. For every computational gate v of the circuit C we introduce a special *extension variable* y_v , and when v is the j th input gate, we identify y_v with the corresponding propositional variable x_j . Let $Vars_C \stackrel{\text{def}}{=} \{x_1, \dots, x_n\} \cup \{y_v \mid v \text{ a computational gate of } C\}$ ⁴.

By $\tau(C, b)$ we denote the CNF in the variables $Vars_C$ that expresses the fact “ $C(x_1, \dots, x_n) = b$ ” and consists of the following clauses:

1. $y_{v_1}^{\bar{\epsilon}_1} \vee \dots \vee y_{v_d}^{\bar{\epsilon}_d} \vee y_v^{\pi(\epsilon_1, \dots, \epsilon_d)}$ whenever $v := \pi(v_1, \dots, v_d)$ is an instruction of C of arity d and $\epsilon \in \{0, 1\}^d$ is an arbitrary vector;
2. $y_{v_i}^{b_i}$ when v_i is the i th output gate of C , $i \in [m]$.

Remark 1 In this paper we will be mostly interested in the case when the circuit C is *linear*, that is consists only of linear instructions $v := v_1 \oplus v_2$ of arity 2. Moreover, the circuits considered will be read-once circuits in which variables are added one at a time. However, even with these restrictions there remains an ambiguity about the order in which the variables are introduced, and this does become important for weak proof systems like those considered in this paper. The next definition makes this formal.

For A an $m \times n$ 0-1 matrix and $i \in [m]$, let $J_i(A) \stackrel{\text{def}}{=} \{j \in [n] \mid a_{ij} = 1\}$ and $X_i(A) \stackrel{\text{def}}{=} \{x_j \mid a_{ij} = 1\}$ be the corresponding set of propositional variables. The set system $J_i(A)$ provides an alternative (and often more convenient) way to represent the matrix A .

Definition 2.2 An *ordering* \leq of an $m \times n$ 0-1 matrix A is a tuple (\leq_1, \dots, \leq_m) , where \leq_i is a linear ordering of the set $J_i(A)$. Given any ordering

⁴although $Vars_C$ is in a one-to-one correspondence with the set of all gates, we prefer to draw a clear distinction between inputs and computational gates

\leq , let C_1, \dots, C_m be the (single-output) circuits naturally computing the parity functions $\bigoplus \{x_j \mid x_j \in X_i(A)\}$ according to this ordering. That is:

- gates of C_i have the form v_Σ^i , where Σ is a non-empty \leq_i -initial segment of $J_i(A)$;
- when x_j is the minimal element of $J_i(A)$, $v_{\{x_j\}}^i$ is the input gate x_j ;
- instructions have the form $v_{\Sigma \cup \{x_j\}}^i := v_\Sigma^i \oplus x_j$, where Σ is a proper initial segment of $J_i(A)$ and j is the minimal element in $J_i(A) \setminus \Sigma$;
- $v_{J_i(A)}^i$ is the output gate.

Denote by $C_{A, \leq}$ the m -output linear circuit which is a disconnected union of C_1, \dots, C_m (that is, these circuits do not have any gates in common except for input gates). Let $\text{Vars}_{\leq}(A) \stackrel{\text{def}}{=} \text{Vars}_{C_{A, \leq}}$ and $\tau_{\leq}(A, b) \stackrel{\text{def}}{=} \tau(C_{A, \leq}, b)$.

Definition 2.3 For a set of rows $I \subseteq [m]$ in the matrix A , we define its *boundary* $\partial_A(I)$ as the set of all $j \in [n]$ (called *boundary elements*) such that $\{a_{ij} \mid i \in I\}$ contains exactly one 1. We say that A is an (r, d) -*lossless expander* if

$$\forall I \subseteq [m] (|I| \leq r \Rightarrow \sum_{i \in I} |J_i(A)| - |\partial_A(I)| \leq d \cdot |I|). \quad (1)$$

Remark 2 Another useful way to interpret the expansion property (1) in this definition is to say that rows in I have at most d *non-boundary* elements on average. In the regular case, that is when all $J_i(A)$ have the same cardinality s , (r, d) -lossless expanders are exactly (r, s, c) -expanders in the terminology of [ABRW00] for $c = s - d$. We introduce this new definition mainly to stress that it is in fact the difference between s and c that matters, and for most applications we need not know s (and in fact even do not need regularity). Also, the “ordinary” lossless expanders recently constructed in [CRVW02] correspond to the case $d = \epsilon s$ for a small constant $\epsilon > 0$, whence our choice of the name.

Despite recent progress, no explicit constructions of (r, d) -lossless expanders with the parameters sufficient for our purposes are currently known. Fortunately, we will be satisfied with the following simple non-constructive bound.

Definition 2.4 $\mathbf{A}_{m,n}$ is a random $m \times n$ 0-1 matrix in which all entries are independent and $\mathbf{P}[\mathbf{a}_{ij} = 1] = n^{-2/3}$.

Theorem 2.5 Let $m \leq 2^{n^\epsilon}$, where $\epsilon > 0$ is a sufficiently small constant. Then $\mathbf{A}_{m,n}$ is an $(n^{\Omega(1)}, O(\frac{\log m}{\log n}))$ -lossless expander with probability $\geq 1 - O(1/m)$.

The proof of Theorem 2.5 is straightforward, it is deferred to Section 7.

Definition 2.6 $\text{Res}(k)$ is the propositional proof system whose lines are k -DNFs, whose only axioms are $\ell \vee \bar{\ell}$ (ℓ a literal) and whose inference rules are given below (F, G are k -DNFs, $1 \leq w \leq k$ and ℓ, ℓ_i are literals).

$$\frac{F}{F \vee \ell} \text{ (WEAKENING)} \qquad \frac{F \vee \ell_1 \ \dots \ F \vee \ell_w}{F \vee (\bigwedge_{i=1}^w \ell_i)} \text{ (AND-INTRODUCTION)}$$

$$\frac{F \vee (\bigwedge_{i=1}^w \ell_i)}{F \vee \ell_i} \text{ (AND-ELIMINATION)} \qquad \frac{F \vee (\bigwedge_{i=1}^w \ell_i) \quad G \vee \bigvee_{i=1}^w \bar{\ell}_i}{F \vee G} \text{ (CUT)}$$

A $\text{Res}(k)$ refutation of a set of k -DNFs is a $\text{Res}(k)$ proof of 0 from this set. In particular, a $\text{Res}(k)$ refutation of a CNF τ is a $\text{Res}(k)$ refutation of the set of clauses τ consists of.

We define the *size* of a $\text{Res}(k)$ proof as the number of *lines* in it. Note that since in this paper we deal exclusively with lower bounds, this makes our results only stronger (as opposed to measuring the complexity by the number of bits).

Note that any variable substitution ρ takes a $\text{Res}(k)$ refutation of a CNF τ into a $\text{Res}(k)$ refutation of $\tau|_\rho$. This in particular implies that the minimal size of a $\text{Res}(k)$ refutation of τ is at least the same size for $\tau|_\rho$.

Finally, the case $k = 1$ corresponds to *Resolution*: this system operates with clauses and has the only inference rule

$$\frac{F \vee x \quad G \vee \bar{x}}{F \vee G}$$

called *resolution rule* (the weakening rule does not change the power of Resolution and is usually, often implicitly, also assumed for convenience). The *width* of a resolution proof is the maximal width of a clause occurring in this proof.

Now we are ready to formulate our main result for Nisan generators. For the sake of definiteness, all algorithms in this paper are assumed to be base 2.

Theorem 2.7 *Let A be an $m \times n$ (r, d) -lossless expander, and assume that*

$$\min_{i \in [m]} |J_i(A)| \geq Cd(k + \log m) \quad (2)$$

for a sufficiently large constant $C > 0$. Let \leq be an arbitrary ordering of A and $b \in \{0, 1\}^m$ be an arbitrary vector. Then every $\text{Res}(k)$ refutation of $\tau_{\leq}(A, b)$ must be of size $\geq \exp(r/2^{O(kd)})$.

Combining Theorems 2.7 and 2.5, we in particular get

Corollary 2.8 *Let m, n, k be parameters such that*

$$m \leq n^{(\epsilon \log n)/k}, \quad (3)$$

where $\epsilon > 0$ is a sufficiently small constant. Then with probability $1 - O(1/m)$ the following holds. For every ordering \leq of $\mathbf{A}_{m,n}$ and every $b \in \{0, 1\}^m$, every $\text{Res}(k)$ -refutation of $\tau_{\leq}(\mathbf{A}_{m,n}, b)$ must have size $\geq \exp(n^{\Omega(1)})$.

Definition 2.9 ([Kra04]) For an n -input m -output circuit C and a vector z_1, \dots, z_m of propositional variables, let $\tau_C(x_1, \dots, x_n, \vec{y}, z_1, \dots, z_m)$ be defined in the same way as $\tau(C, b)$ (see Definition 2.1) with the difference that the “output axioms” $y_{v_i}^{b_i}$ get replaced by the two clauses $y_{v_i} \vee \bar{z}_i, \bar{y}_{v_i} \vee z_i$ (thus, $\tau(C, b)$ is the same as $\tau_C(\vec{x}, \vec{y}, b)$).

For a proof system P and an integer S , the circuit C is S -iterable in P if for every CNF of the form

$$\bigwedge_{\nu=1}^H \tau_C(x_1^{(\nu)}, \dots, x_n^{(\nu)}, \vec{y}^{(\nu)}, q_1^{(\nu)}, \dots, q_m^{(\nu)}), \quad (4)$$

every refutation of this CNF in the system P must have size $\geq S$. Here $x_j^{(\nu)}, \vec{y}^{(\nu)}$ are pairwise disjoint tuples of variables, and every $q_i^{(\nu)}$ is either a Boolean constant or belongs to the set $\{x_j^{(\mu)} \mid j \in [n], \mu < \nu\}$.

Since $\tau(C, b)$ itself has the form (4) (with $H = 1$), iterability is stronger than just hardness. As the proof of the following theorem will show, for the particular case of NW-generators they are not very much apart from each other.

Theorem 2.10 *Let A be an $m \times n$ (r, d) -lossless expander, and assume that*

$$s \stackrel{\text{def}}{=} \min_{i \in [m]} |J_i(A)| \geq Cd(k + \log m) \quad (5)$$

for a sufficiently large constant $C > 0$. Then for every ordering \leq of the matrix A , the circuit $C_{A, \leq}$ from Definition 2.2 is $\exp(\min\{\Omega(s/d), r/2^{O(kd)}\})$ -iterable in $\text{Res}(k)$.

As in [Kra04], this in particular implies that composing the generator $C_{A, \leq}$ with itself preserves hardness. We will note here only one particular iteration protocol corresponding to the classical construction from [GGM86].

Definition 2.11 Let C be a Boolean circuit (over an arbitrary basis) with n inputs and $2n$ outputs, and $h \geq 1$ be an integer. The circuit C^h with n inputs and $2^h n$ outputs is constructed as follows. For every binary string u with $|u| \leq h - 1$ we prepare an isomorphic copy C_u of C . The inputs of C^h are the inputs of C_Λ (Λ the empty string), and the outputs of C^h are the outputs of the circuits C_u with $|u| = h - 1$. Finally, for every u with $|u| < h - 1$, the first n outputs of C_u are identified with the inputs of C_{u*0} , and the last n output bits are identified with inputs of C_{u*1} (in everything else these circuits are completely independent).

Theorem 2.12 *Let \leq be an arbitrary ordering of an $(2n \times n)$ (r, d) -lossless expander A , and assume that*

$$s \stackrel{\text{def}}{=} \min_{i \in [2n]} |J_i(A)| \geq Cd(k + \log n)$$

for a sufficiently large constant $C > 0$. Let $h \geq 1$ be an arbitrary integer, and $b \in \{0, 1\}^{2^h n}$ be an arbitrary vector. Then every $\text{Res}(k)$ refutation of $\tau(C_{A, \leq}^h, b)$ must have size $\geq \exp(\min\{\Omega(s/d), r/2^{O(kd)}\})$.

For a Boolean function f_n in n variables⁵ and $t \leq 2^n$, denote by $\text{Circuit}_t(f_n)$ an $O(1)$ -CNF of size $2^{O(n)}$ encoding the description of a size- t fan-in 2 Boolean

⁵Note that the “intended meaning” of n in Theorems 2.13, 2.20 is completely different from all other results

circuit over the standard basis $\{\neg, \wedge, \vee\}$ for computing f_n . We will recall an exact definition in Section 8; for the time being let us just observe that proving that the circuit size of f_n is greater than t is tantamount to showing that $\text{Circuit}_t(f_n)$ is unsatisfiable.

Theorem 2.13 *Let f_n be any Boolean function in n variables, and $n^2 \leq t \leq 2^n$. Then every $\text{Res}(\epsilon \log t)$ refutation of $\text{Circuit}_t(f_n)$ ($\epsilon > 0$ a sufficiently small constant) must have size $\exp(t^{\Omega(1)})$.*

Let us now consider the pigeonhole principle.

Definition 2.14 Assume $m > n$. The (negation of the) *onto pigeonhole principle* is the unsatisfiable CNF in the variables $\{x_{ij} \mid i \in [m], j \in [n]\}$ denoted by $\neg\text{onto} - \text{PHP}_n^m$ that is the conjunction of the following clauses:

$$\begin{aligned} Q_i &\stackrel{\text{def}}{=} \bigvee_{j=1}^n x_{ij} \quad (i \in [m]); \\ Q_{i_1, i_2; j} &\stackrel{\text{def}}{=} (\bar{x}_{i_1 j} \vee \bar{x}_{i_2 j}) \quad (i_1 \neq i_2 \in [m], j \in [n]); \\ Q_j &\stackrel{\text{def}}{=} \bigvee_{i=1}^m x_{ij} \quad (j \in [n]) \end{aligned}$$

(the prefix “onto” refers to the presence of the last group of axioms).

[SBI04] proved that $\neg\text{onto} - \text{PHP}_n^{2n}$ is exponentially hard to refute in $\text{Res}(k)$ as long as $k \leq \sqrt{\log n / \log \log n}$. We improve this as follows:

Theorem 2.15 *Every $\text{Res}(k)$ refutation of $\neg\text{onto} - \text{PHP}_n^{2n}$ must have size $\exp(n/(\log n)^{O(k)})$.*

This bound is exponential up to $k = (\epsilon \log n) / \log \log n$.

Let us now recall another extension of Resolution that is of more algebraic flavour.

Definition 2.16 ([CEI96]) Let \mathbb{F} be a fixed field. *Polynomial Calculus* (PC for short) is the proof system whose lines are polynomials $f \in \mathbb{F}[x_1, \dots, x_n]$ (a polynomial f is interpreted as the polynomial equation $f = 0$). It has polynomials $x_i^2 - x_i$ ($i \in [n]$) as *default axioms* and has two inference rules:

$$\frac{f}{\alpha f + \beta g}; \quad \alpha, \beta \in \mathbb{F} \quad (\text{SCALAR ADDITION});$$

$$\frac{f}{x_i \cdot f} \text{ (VARIABLE MULTIPLICATION).}$$

Definition 2.17 ([ABRW02]) Let again \mathbb{F} be a fixed field. *Polynomial Calculus with Resolution* (PCR) is the proof system whose lines are polynomials from $F[x_1, \dots, x_n, \bar{x}_1, \dots, \bar{x}_n]$, where $\bar{x}_1, \dots, \bar{x}_n$ are treated as new formal variables. PCR has all default axioms and inference rules of PC (including, of course, those that involve new variables \bar{x}_i), plus additional default axioms $x_i + \bar{x}_i = 1$ ($i \in [n]$).

For a clause C , denote by Γ_C the monomial

$$\Gamma_C \stackrel{\text{def}}{=} \prod_{\bar{x} \in C} x \cdot \prod_{x \in C} \bar{x}, \quad (6)$$

and for a CNF τ , let $\Gamma_\tau \stackrel{\text{def}}{=} \{\Gamma_C \mid C \in \tau\}$ (note that τ is unsatisfiable if and only if the polynomials Γ_τ have no common root in \mathbb{F} satisfying all default axioms of PCR). A *PCR refutation of a CNF τ* is a PCR proof of the contradiction $1=0$ from Γ_τ .

The *degree* of a PCR proof is defined as the maximal degree of a polynomial appearing in it, and its *size* is the number of different *monomials* in this proof.

Remark 3 PC and PCR are equivalent with respect to the degree measure (via the linear transformation $\bar{x}_i \mapsto 1 - x_i$). Also note that we measure the size of PCR proofs differently from Definition 2.6; namely, by the number of monomials. We do not know if our results still hold if the size is measured by the number of lines.

All our lower bounds for Res(1) (= Resolution) generalize to PCR over any field \mathbb{F} with $\text{char}(\mathbb{F}) \neq 2$. That is,

Theorem 2.18 *Let A be an $m \times n$ (r, d) -lossless expander, and assume that*

$$\min_{i \in [m]} |J_i(A)| \geq Cd \log m \quad (7)$$

for a sufficiently large constant $C > 0$. Let \leq be an arbitrary ordering of A , $b \in \{0, 1\}^m$ be an arbitrary vector, and \mathbb{F} be an arbitrary field with $\text{char}(\mathbb{F}) \neq 2$. Then every PCR refutation of $\tau_{\leq}(A, b)$ over the field \mathbb{F} must be of size $\geq \exp(r/2^{O(d)})$.

Theorem 2.19 *Let A be an $m \times n$ (r, d) -lossless expander such that (7) holds, and let $s \stackrel{\text{def}}{=} \min_{i \in [m]} |J_i(A)|$. Then for every ordering \leq of the matrix A and for every field \mathbb{F} with $\text{char}(\mathbb{F}) \neq 2$, the circuit $C_{A, \leq}$ is $\exp(\min\{\Omega(s/d), r/2^{O(d)}\})$ -iterable in PCR over \mathbb{F} .*

Let $\text{Circuit}_t^\oplus(f_n)$ be defined in the same way as $\text{Circuit}_t(f_n)$, with the exception that besides the standard connectives $\{\neg, \wedge, \vee\}$, the encoded circuit also allows PARITY gates of fan-in 2.

Theorem 2.20 *Let f_n be any Boolean function in n variables, $n^2 \leq t \leq 2^n$, and \mathbb{F} be any field with $\text{char}(\mathbb{F}) \neq 2$. Then every PCR refutation of $\text{Circuit}_t^\oplus(f_n)$ over \mathbb{F} must have size $\exp(t^{\Omega(1)})$.*

We conclude with one miscellaneous result about the hardness of Nisan generators that is specific to PCR. For these generators we still do not know how to get more than $n^{\epsilon \log n}$ output bits even in case of Resolution. We now show how to make a *function* (that is, with $m = 2^{n^\epsilon}$ bits) Nisan generator hard for PCR at the expense of spoiling its encoding.

Every ordering j_1, \dots, j_d of a finite set J can be also viewed as a *cyclic order*, that is as an injective mapping $\alpha : J \rightarrow S^1$ into the unit circle S^1 given by $\alpha(j_\nu) = \nu/d$. A *cyclic interval* is a set of the form $\alpha^{-1}(A)$, where $A \subseteq S^1$ is an arc.

Fix a tuple $\leq = (\leq_1, \dots, \leq_m)$ of orderings of the sets $(J_1(A), \dots, J_m(A))$ and let

$$\text{Vars}_{\leq}^{\text{Cycl}}(A) \stackrel{\text{def}}{=} \{x_1, \dots, x_n\} \cup \left\{ y_\Delta^i \mid i \in [m], \Delta \text{ a cyclic interval in } J_i(A) \right. \\ \left. \text{such that } \Delta = J_i(A) \text{ or } \frac{|J_i(A)|}{3} - 1 \leq |\Delta| \leq \frac{2|J_i(A)|}{3} + 1 \right\}.$$

Let $\tau_{\leq}^{\text{Cycl}}(A, b)$ be the 3-CNF consisting of the clauses that result from expanding those constraints of the form

$$y_{\Delta \cup \{j\}}^i \equiv y_\Delta^i \oplus x_j \quad (i \in [m], j \text{ adjacent to } \Delta); \quad (8)$$

$$y_{\Delta_1 \cup \Delta_2}^i \equiv y_{\Delta_1}^i \oplus y_{\Delta_2}^i \quad (i \in [m], \Delta_1, \Delta_2 \text{ disjoint and adjacent}); \quad (9)$$

$$y_{J_i(A)}^i \equiv b_i. \quad (10)$$

in which all variables belong to $\text{Vars}_{\leq}^{\text{Cycl}}(A)$ (that is, $\Delta, \Delta \cup \{j\}, \Delta_1, \Delta_2$ and $\Delta_1 \cup \Delta_2$ must obey the size bound in its definition).

Let us verify that $\tau_{\leq}^{Cycl}(A, b)$ is indeed a complete encoding of the linear system $AX = b$ (this is not immediately clear due to the presence of the constraints on $|\Delta|$).

Fact 1 *The system of \mathbb{F}_2 -linear equations $AX = b$ is consistent if and only if $\tau_{\leq}^{Cycl}(A, b)$ is satisfiable.*

Proof. The “only if” part is obvious. For the opposite direction, we only have to show that the x -part of every satisfying assignment for $\tau_{\leq}^{Cycl}(A, b)$ also satisfies the system $AX = b$.

For any given $i \in [m]$ fix an arbitrary partition $J_i(A) = \Delta_{i,1} \dot{\cup} \Delta_{i,2} \dot{\cup} \Delta_{i,3}$ into three cyclic intervals of almost equal sizes: $\frac{|J_i(A)|}{3} - 1 \leq |\Delta_{i\nu}| \leq \frac{|J_i(A)|}{3} + 1$. Then (9) (applied twice) and (10) imply that in every satisfying assignment we have $y_{\Delta_{i,1}}^i \oplus y_{\Delta_{i,2}}^i \oplus y_{\Delta_{i,3}}^i = b_i$. On the other hand, by summing up appropriate axioms (8) (with $j \in \Delta_{i\nu}$), we obtain $y_{\Delta_{i\nu'}}^i \oplus \bigoplus_{j \in \Delta_{i\nu}} x_j = y_{\Delta_{i\nu} \cup \Delta_{i\nu'}}^i$ ($\nu' \neq \nu$) and then, using again (9), $\bigoplus_{j \in \Delta_{i\nu}} x_j = y_{\Delta_{i\nu}}^i$ for every $\nu \in [3]$. The statement follows. ■

Theorem 2.21 *Let A be an $m \times n$ (r, d) -lossless expander such that $\min_{i \in [m]} |J_i(A)| \geq Cd \log m$ for a sufficiently large constant $C > 0$, \leq be an arbitrary ordering of A , $b \in \{0, 1\}^m$ be an arbitrary vector, and \mathbb{F} any field with $\text{char}(\mathbb{F}) \neq 2$. Then every PCR refutation of $\tau_{\leq}^{Cycl}(A, b)$ over the field \mathbb{F} must have size $\geq \exp(r/d^{O(1)})$.*

The bound of this theorem is as good as we might hope. The encoding itself, however, is not very useful: it does not correspond to any circuit, and, moreover, the proof of this theorem seems to completely break apart for $\text{Res}(2)$.

3. Preliminaries

In this section we begin the proof of Theorem 2.7 by presenting some known results in the form adapted to our purposes.

Recall several definitions from [ABRW00].

Definition 3.1 A Boolean function f is ℓ -robust if every restriction ρ such that $f|_{\rho} = \text{const}$, satisfies $|\text{sup}(\rho)| \geq \ell$.

Definition 3.2 Let A be an $m \times n$ 0-1 matrix. For every Boolean function f with the property $\exists i \in [m](\text{Vars}(f) \subseteq X_i(A))$ we introduce a new *extension variable* y_f . Let $\text{Vars}(A)$ be the set of all these variables.

Given Boolean functions $\vec{g} = (g_1, \dots, g_m)$ such that $\text{Vars}(g_i) \subseteq X_i(A)$, we denote by $\tau(A, \vec{g})$ the CNF in the variables $\text{Vars}(A)$ that consists of those clauses $y_{f_1}^{\epsilon_1} \vee \dots \vee y_{f_w}^{\epsilon_w}$ for which there exists $i \in [m]$ such that

$$\text{Vars}(f_1) \cup \dots \cup \text{Vars}(f_w) \subseteq X_i(A)$$

and

$$g_i \leq f_1^{\epsilon_1} \vee \dots \vee f_w^{\epsilon_w}.$$

We now have the following (minor) generalization of [ABRW00, Theorem 3.1].

Theorem 3.3 *Let A be an (r, d) -lossless expander of size $m \times n$, and g_1, \dots, g_m be ℓ -robust functions with $\text{Vars}(g_i) \subseteq X_i(A)$, where $\ell \geq d + 1$. Then every resolution refutation of $\tau(A, \vec{g})$ must have width $> \frac{r(\ell-d)}{2\ell}$.*

Proof. The only difference from [ABRW00, Theorem 3.1] is that we now use (r, d) -lossless expanders instead of (r, s, c) -expanders with $s - c = d$. The proof goes the same way, and we only remark on the changes to be made to its text. All these changes pertain to the proof of [ABRW00, Claim 3.3].

First, the bound on the size of $\partial_A(I)$ now becomes $|\partial_A(I)| \geq \sum_{i \in I} |J_i(A)| - d \cdot |I|$ (as opposed to $|\partial_A(I)| \geq c \cdot |I|$ in [ABRW00]). Next, the proof of the crucial inequality $|J_{i_1}(A) \cap \partial_A(I)| \leq s - \ell$ in [ABRW00] actually shows $|J_{i_1}(A) \cap \partial_A(I)| \leq |J_{i_1}(A)| - \ell$. Last, the final calculation now looks like

$$\begin{aligned} \sum_{i \in I} |J_i(A)| - d \cdot |I| &\leq |\partial_A(I)| \leq \sum_{i \in I_0} |J_i(A)| + \sum_{i \in I_1} (|J_i(A)| - \ell) \\ &= \sum_{i \in I} |J_i(A)| - \ell \cdot |I_1| \leq \sum_{i \in I} |J_i(A)| - \ell \cdot (|I| - w(C)) \end{aligned}$$

which implies our bound $w(C) > \frac{r(\ell-d)}{2\ell}$ since $|I| > r/2$. ■

We will only need the following special case of this.

Corollary 3.4 *Let A be an (r, d) -lossless expander of size $m \times n$ such that $|J_i(A)| \geq 2d$ for all $i \in [m]$. Then for every ordering \leq and every $b \in \{0, 1\}^m$, every resolution refutation of $\tau_{\leq}(A, b)$ must have width $> r/4$.*

Proof. In Theorem 3.3, let g_i be the function $\bigoplus_{j \in J_i(A)} x_j = b_i$ and $\ell := 2d$. Since $\tau_{\leq}(A, b)$ is a sub-CNF of $\tau(A, \vec{g})$ (for any ordering \leq), the result follows. ■

Definition 3.5 ([SBI04]) A *decision tree* is a rooted binary tree such that every internal node is labelled with a variable, the edges leaving this node correspond to whether the variable is set to 0 or 1, and the leaves are labelled with either 0 or 1. As usual, we assume that on every given path no variable appears more than once. Then every path from the root to a leaf may be viewed as a partial assignment, and this assignment, in turn, will be sometimes identified with the corresponding leaf. For a decision tree T and $\epsilon \in \{0, 1\}$, we write the set of paths (partial assignments) that lead from the root to a leaf labelled ϵ as $Br_{\epsilon}(T)$. We say that a decision tree T *strongly represents a DNF* F if for every $\pi \in Br_0(T)$ and for all $t \in F$, $t|_{\pi} = 0$ and for every $\pi \in Br_1(T)$, there exists $t \in F$ such that $t|_{\pi} = 1$. Let the *representation height* of F , $h(F)$ be the minimum height of a decision tree strongly representing F .

Proposition 3.6 ([SBI04]) *Let τ be an h -CNF, P be a $Res(k)$ refutation of τ , and let ρ be a partial assignment so that for every line F of P , $h(F|_{\rho}) \leq h$. Then $\tau|_{\rho}$ has a resolution refutation of width $\leq kh$.*

Remark 4 One small technicality is that the system $Res(k)$, as defined in [SBI04], does not automatically include the axioms $\ell \vee \bar{\ell}$. Therefore, formally speaking, an application of their result only implies that $\tau^*|_{\rho}$ has a resolution refutation of the required width, where τ^* is obtained from τ by adding these trivial axioms. It is, however, obvious that the latter can be eliminated from any resolution proof without increasing its width.

Finally, we recall a combinatorial inequality originally proved in [Jan90] and further generalized in [AS08, Section 8.1].

Definition 3.7 For propositional variables x_1, \dots, x_n and probabilities $p_1, \dots, p_n \in [0, 1]$, denote by $\mathbf{a}_{\vec{p}}$ a random assignment that independently assigns every variable x_i to 1 with probability p_i , and to 0 with probability $(1 - p_i)$.

Proposition 3.8 *Let t_1, \dots, t_H be monotone terms (not necessarily distinct) in the variables x_1, \dots, x_n , and $p_1, \dots, p_n \in [0, 1]$. Let*

$$\kappa \stackrel{\text{def}}{=} \sum_{\alpha \in [H]} \mathbf{P}[t_\alpha(\mathbf{a}_{\vec{p}}) = 1] \quad (11)$$

be the expectation of the number of terms satisfied by $\mathbf{a}_{\vec{p}}$, and let

$$\Delta \stackrel{\text{def}}{=} \sum_{\substack{\alpha \neq \beta \in [H] \\ \text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta) \neq \emptyset}} \mathbf{P}[t_\alpha(\mathbf{a}_{\vec{p}}) = t_\beta(\mathbf{a}_{\vec{p}}) = 1].$$

Then

$$\mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\vec{p}}) = 0 \right] \leq e^{-\kappa + \frac{\Delta}{2}}.$$

Proof. The notation of [AS08, Section 8.1] corresponds to ours as follows: Ω is $\{x_1, \dots, x_n\}$, I is $[H]$, $A_i := \text{Vars}(t_i)$ ($i \in [H]$), and B_i is the event $t_i(\mathbf{a}_{\vec{p}}) = 1$; we also renamed μ to κ . After this translation, our proposition is exactly the second inequality in [AS08, Theorem 8.1.1]. ■

Remark 5 We will actually need another version of this inequality ([AS08, Theorem 8.1.2]), useful when $\Delta \gg \kappa$. We, however, will need to dig into the proof of that theorem rather deeply. For this reason we do not formulate the corresponding general statement here, but rather incorporate it as an ad hoc argument in the proof of Lemma 4.4.

4. Small restriction switching lemma

In this section we give a quadratic improvement on the original version of this lemma from [SBI04]. In our main application, the underlying random restriction will not act totally independently on different variables, but at least it will have some “weak local independence” property. We will be able to capture this property in the main statement so that the proof will not become more complicated than for truly independent restrictions, but this will require several auxiliary definitions.

Definition 4.1 We say that a (deterministic) restriction ρ' is a *sub-restriction* of another restriction ρ if $\text{sup}(\rho') \subseteq \text{sup}(\rho)$ and ρ, ρ' coincide on $\text{sup}(\rho')$. For

random restrictions ρ, ρ', ρ' is a *sub-restriction* of ρ if there exists a set Ω , a random variable $\omega \in \Omega$, and functions π, π' from Ω to the set of all restrictions such that:

1. ρ has the same distribution as $\pi(\omega)$, and ρ' has the same distribution as $\pi'(\omega)$;
2. for every individual $\omega \in \Omega$, $\pi'(\omega)$ is a sub-restriction of $\pi(\omega)$.

Clearly, if ρ' is a sub-restriction of ρ then for any terms t_1, \dots, t_H we have the inequality $\mathbf{P}\left[\bigvee_{\alpha \in [H]} (t_\alpha | \rho' \equiv 1)\right] \leq \mathbf{P}\left[\bigvee_{\alpha \in [H]} (t_\alpha | \rho \equiv 1)\right]$.

Definition 4.2 A *weight function* is any function $\mu : \{x_1, \dots, x_n\} \longrightarrow \mathbb{Z}^+$ from variables to strictly positive integers. The *weight* $\mu(V)$ of a set of variables V is defined as $\mu(V) \stackrel{\text{def}}{=} \sum_{x \in V} \mu(x)$, and the *weight of a term* t is $\mu(t) \stackrel{\text{def}}{=} \mu(\text{Vars}(t))$. We will denote by μ_{triv} the *trivial weight function* identically equal to 1; in this case the weight of a term is equal to its width. For arbitrary weight function μ we have this in one direction: $\mu(t) \geq w(t)$. A DNF F is a *weighted K -DNF* (w.r.t. a weight function μ) if all terms $t \in F$ have weight $\leq K$.

Finally, we define the amount of “weak local independence” needed to carry out the proof of our switching lemma. It is similar to the ordinary r -wise independence with one important change. Namely, we do not demand that on small sets of variables our random restriction behaves *exactly* as the genuine i.i.d restriction. We only require that it can be obtained from the latter by assigning more variables if necessary.

Definition 4.3 Let μ be a weight function, and let $p \in [0, 1]$. For a set of variables X , let $\rho_{\mu, X, p}$ be the random restriction of these variables that independently assigns every $x \in X$ to 0,1 with probability $p^{\mu(x)}/2$, and leaves it unassigned with probability $(1 - p^{\mu(x)})$. Given an integer r , say that a random restriction ρ is (r, μ, p) -*independent* if for every subset X of variables with $|X| \leq r$, $\rho_{\mu, X, p}$ is a sub-restriction of $\rho|_X$.

Thus, if ρ is (r, μ, p) -independent, then for any set of terms t_1, \dots, t_H such that $X \stackrel{\text{def}}{=} \bigcup_{i=1}^H \text{Vars}(t_i)$ has size at most r , we have

$$\mathbf{P}\left[\bigvee_{\alpha \in [H]} (t_\alpha | \rho_{\mu, X, p} \equiv 1)\right] \leq \mathbf{P}\left[\bigvee_{\alpha \in [H]} (t_\alpha | \rho \equiv 1)\right] \quad (12)$$

(and, as a matter of fact, this is the only property of (r, μ, p) -independent restrictions we will need).

The rest of this section is devoted to the proof of the following lemma.

Lemma 4.4 *Let μ be a weight function, and F be a weighted K -DNF w.r.t. μ . Suppose that ρ is a random (r, μ, p) -independent restriction. Then for every $h \leq r$,*

$$\mathbf{P}[h(F|\rho) > h] \leq \exp(-h(p/2)^{O(K)}). \quad (13)$$

Proof. We begin with the case when F is monotone, and in that case we are going to prove by induction on K that

$$\mathbf{P}[h(F|\rho) > h] \leq e^{-h(p\epsilon)^{2K}}, \quad (14)$$

where $\epsilon > 0$ is a constant chosen sufficiently small so that the arguments for Cases 1 and 2 below work⁶. This clearly implies (13).

Base $K = 0$ is obvious since F is a constant and $h(F|\rho) = 0$ with probability 1.

Inductive step. Let F be a non-trivial monotone weighted K -DNF, and assume that (14) is established for all weighted K' -DNF with $K' < K$ (and for all integers h). We define a numerical invariant $\delta(F)$ that represents, up to a scaling factor, the optimal value of the parameter Δ in Proposition 3.8 when we attempt to apply it to the formula F . Further analysis will be sharply divided according to whether $\delta(F)$ is small or large (cf. [SBI04, Theorem 3]).

First, for a set of variables V we let

$$\delta(V) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } V = \emptyset \\ (2/p)^{\mu(V)} & \text{otherwise.} \end{cases}$$

Next, given a random term $\mathbf{t} \in F$ (viewed, for the duration of this proof, as a probability distribution on the set of all terms of F), we define

$$\delta(\mathbf{t}) \stackrel{\text{def}}{=} \mathbf{E}[\delta(\text{Vars}(\mathbf{t}) \cap \text{Vars}(\mathbf{t}'))],$$

⁶The factor 2 in (14) can be removed by using a slightly more sophisticated analysis in Case 2 below, but we do not need that precision in what follows.

where \mathbf{t}' is an independent copy of \mathbf{t} . Finally, let

$$\delta(F) \stackrel{\text{def}}{=} \min_{\mathbf{t}} \delta(\mathbf{t}),$$

where the minimum is taken over all random terms $\mathbf{t} \in F$. (This minimum exists since the space of all probability distributions on terms of F is compact, and the function $\delta(\mathbf{t})$ is continuous.) Let

$$s \stackrel{\text{def}}{=} \lceil 2h(p\epsilon)^{2K} \rceil \quad (15)$$

(so that the right-hand side of (14) is roughly $e^{-s/2}$). Consider two cases.

Case 1: $\delta(F) \leq s^{-1}$. Let $\mathbf{t} \in F$ be the random term for which $\delta(\mathbf{t}) \leq s^{-1}$. Arguing as in the proof of [AS08, Theorem 8.1.2], let $\mathbf{t}_1, \dots, \mathbf{t}_H$ be independent samples from F according to the distribution of \mathbf{t} , where

$$H \stackrel{\text{def}}{=} \lceil s(2/p)^K \rceil. \quad (16)$$

Then

$$\mathbf{E} \left[\sum_{\alpha \neq \beta \in [H]} \delta(\text{Vars}(\mathbf{t}_\alpha) \cap \text{Vars}(\mathbf{t}_\beta)) \right] = H(H-1)\delta(\mathbf{t}) \leq H(H-1)s^{-1},$$

and we fix any particular sampling t_1, \dots, t_H for which

$$\sum_{\alpha \neq \beta \in [H]} \delta(\text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta)) \leq H(H-1)s^{-1}. \quad (17)$$

The terms t_1, \dots, t_H altogether contain at most HK variables, and $HK \leq h \leq r$ as long as the constant ϵ is small enough. Therefore, since ρ is (r, μ, p) -independent, (12) implies

$$\mathbf{P}[h(F|\rho) > h] \leq \mathbf{P} \left[\bigwedge_{\alpha \in [H]} (t_\alpha|_\rho \neq 1) \right] \leq \mathbf{P} \left[\bigwedge_{\alpha \in [H]} (t_\alpha|_{\rho_{\mu,p}} \neq 1) \right]. \quad (18)$$

Furthermore, since the terms t_α are monotone, $t_\alpha|_{\rho_{\mu,p}} \equiv 1$ if and only if $t_\alpha(\mathbf{a}) = 1$, where \mathbf{a} is the random (total) assignment obtained from $\rho_{\mu,p}$ by additionally assigning to 0 all unassigned variables. Now, \mathbf{a} has the same

distribution as the assignment $\mathbf{a}_{\tilde{p}}$ from Definition 3.7 for the vector of probabilities \tilde{p} given by $\tilde{p}_i \stackrel{\text{def}}{=} p^{\mu(x_i)}/2$. Let $p_i \stackrel{\text{def}}{=} (p/2)^{\mu(x_i)}$; then $p_i \leq \tilde{p}_i$ and, since the event $\bigvee_{\alpha \in [H]} t_\alpha(a) = 0$ is anti-monotone in a , we get

$$\mathbf{P} \left[\bigwedge_{\alpha \in [H]} (t_\alpha |_{\rho_{\mu,p}} \neq 1) \right] = \mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\tilde{p}}) = 0 \right] \leq \mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\tilde{p}}) = 0 \right]. \quad (19)$$

We are going to upper bound $\mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\tilde{p}}) = 0 \right]$, given (17).

For every $\alpha \in [H]$ such that $\mu(t_\alpha) < K$ introduce a new auxiliary variable y_α with weight $K - \mu(t_\alpha)$, and replace t_α with $t_\alpha \wedge y_\alpha$. This operation does not change the value of the sum $\sum_{\alpha \neq \beta \in [H]} \delta(\text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta))$ in (17), and $\mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\tilde{p}}) = 0 \right]$ may only increase. Therefore, we may assume w.l.o.g. that all terms in F have weight *exactly* K .

And now we apply Proposition 3.8. All events $\mathbf{P}[t_\alpha(\mathbf{a}_{\tilde{p}}) = 1]$ have the same probability $(p/2)^K$; therefore, the quantity κ given by (11) is equal to $H(p/2)^K$. Also, whenever $\text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta) \neq \emptyset$, we have

$$\mathbf{P}[t_\alpha(\mathbf{a}_{\tilde{p}}) = t_\beta(\mathbf{a}_{\tilde{p}}) = 1] = (p/2)^{\mu(t_\alpha \wedge t_\beta)} = (p/2)^{2K} \delta(\text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta));$$

therefore,

$$\Delta = (p/2)^{2K} \sum_{\alpha \neq \beta \in [H]} \delta(\text{Vars}(t_\alpha) \cap \text{Vars}(t_\beta)) \leq (p/2)^{2K} H(H-1)s^{-1} \leq H(p/2)^K = \kappa$$

by (17) and (16). Applying Proposition 3.8, we get

$$\mathbf{P} \left[\bigwedge_{\alpha \in [H]} t_\alpha(\mathbf{a}_{\tilde{p}}) = 0 \right] \leq e^{-\frac{H(p/2)^K}{2}} \leq e^{-s/2}$$

which, along with (18) and (19), implies (14).

Case 2: $\delta(F) \geq s^{-1}$. Note first that

$$\begin{aligned} \delta(\mathbf{t}) &= \mathbf{E}[\delta(\text{Vars}(\mathbf{t}) \cap \text{Vars}(\mathbf{t}'))] = \sum_V \delta(V) \cdot \mathbf{P}[\text{Vars}(\mathbf{t}) \cap \text{Vars}(\mathbf{t}') = V] \\ &\leq \sum_V \delta(V) \cdot \mathbf{P}[V \subseteq \text{Vars}(\mathbf{t}) \wedge V \subseteq \text{Vars}(\mathbf{t}')] = \sum_{V \neq \emptyset} (2/p)^{\mu(V)} \mathbf{P}[V \subseteq \text{Vars}(\mathbf{t})]^2; \end{aligned}$$

therefore, we also have

$$\min_{\mathbf{t}} \sum_{V \neq \emptyset} (2/p)^{\mu(V)} \mathbf{P}[V \subseteq \text{Vars}(\mathbf{t})]^2 \geq s^{-1}.$$

Let \mathbf{t} be the random term on which the quadratic form $\sum_{V \neq \emptyset} (2/p)^{\mu(V)} \mathbf{P}[V \subseteq \text{Vars}(\mathbf{t})]^2$ in the variables $p_t \stackrel{\text{def}}{=} \mathbf{P}[\mathbf{t} = t]$ describing the density function of the associated distribution attains its minimal value $\delta \geq s^{-1}$. Denoting further $\mathbf{P}[V \subseteq \text{Vars}(\mathbf{t})]$ (viewed as a linear form in the variables p_t) by p_V we have

$$\sum_{\substack{t \in F \\ p_t \neq 0}} \frac{\partial(\sum_{V \neq \emptyset} (2/p)^{\mu(V)} p_V^2)}{\partial p_t} p_t = 2 \sum_{V \neq \emptyset} (2/p)^{\mu(V)} p_V^2 = 2\delta.$$

Since $\sum_{\substack{t \in F \\ p_t \neq 0}} p_t = 1$, this implies the existence of some $t_0 \in F$ with $p_{t_0} \neq 0$ and such that

$$\frac{\partial(\sum_{V \neq \emptyset} (2/p)^{\mu(V)} p_V^2)}{\partial p_{t_0}} = 2 \sum_{\substack{V \neq \emptyset \\ V \subseteq \text{Vars}(t_0)}} (2/p)^{\mu(V)} p_V \geq 2\delta.$$

Therefore, for any term $t \in F$ (even when $p_t = 0$!) we also have $\sum_{\substack{V \neq \emptyset \\ V \subseteq \text{Vars}(t)}} (2/p)^{\mu(V)} p_V \geq \delta$ since otherwise, by setting p_{t_0} to 0 and p_t to $p_t + p_{t_0}$, we would have obtained contradiction with the assumption that \mathbf{t} minimizes the quadratic form $\sum_{V \neq \emptyset} (2/p)^{\mu(V)} \mathbf{P}[V \subseteq \text{Vars}(\mathbf{t})]^2$.

Summarizing the above argument, we have found coefficients $p_V \geq 0$ (V a non-empty set of variables) and $\delta \geq s^{-1}$ such that

$$\sum_{V \neq \emptyset} (2/p)^{\mu(V)} p_V^2 = \delta; \tag{20}$$

$$\forall t \in F \left(\sum_{\substack{V \neq \emptyset \\ V \subseteq \text{Vars}(t)}} (2/p)^{\mu(V)} p_V \geq \delta \right). \tag{21}$$

These are the only properties of p_V s we will need in the rest of the proof; in particular, at this point we can forget their interpretation as certain probabilities.

Let us order all non-empty sets of variables V in decreasing order with respect to the coefficient p_V : $p_{V_1} \geq p_{V_2} \geq p_{V_3} \geq \dots$. Assume w.l.o.g. that the variables x_1, \dots, x_n are enumerated in the order in which they appear for the first time in the sequence V_1, V_2, V_3, \dots (equivalently, we require that every union $V_1 \cup V_2 \cup \dots \cup V_H$ should be an initial segment in the set of variables ordered according to their indices). For a non-empty V , denote by $i(V)$ the *maximal* index i for which $x_i \in V$.

Now we classify terms $t \in F$ as follows. Represent a given term $t \in F$ in the form $t = x_{i_1} \wedge x_{i_2} \wedge \dots \wedge x_{i_w}$ ($i_1 < i_2 < \dots < i_w$). For a non-empty

$V \subseteq Vars(t)$, let $\mu_t(V) \stackrel{\text{def}}{=} \mu(x_{i_1} \wedge x_{i_2} \wedge \dots \wedge x_{i(V)})$ (in other words, this is the weight of the minimal initial segment in $Vars(t)$ containing V). We split the sum in (21) with respect to the value of $\mu_t(V)$, and classify t according to which part is large enough. Formally, let

$$F_\mu \stackrel{\text{def}}{=} \left\{ t \in F \mid \sum_{\substack{\mu_t(V)=\mu \\ V \subseteq Vars(t)}} p_V \geq \delta(p/4)^\mu \right\};$$

let us see that $F = \bigvee_{\mu=1}^K F_\mu$. Assume for the sake of contradiction that for some $t \in F$ and all real μ we have

$$\sum_{\substack{\mu_t(V)=\mu \\ V \subseteq Vars(t)}} p_V < \delta(p/4)^\mu < \delta(p/2)^\mu.$$

Summing these inequalities with coefficients $(2/p)^\mu$, we would get

$$\sum_{V \subseteq Vars(t)} p_V (2/p)^{\mu_t(V)} < \delta.$$

that would be in contradiction with (21) since clearly $\mu_t(V) \geq \mu(V)$ for all t, V . This contradiction shows that $t \in F_\mu$ for at least one μ , that is indeed $F = \bigvee_{\mu=1}^K F_\mu$.

Then, noting that $\mu_t(V) \geq \mu(V)$ for all t, V , we have $F = \bigvee_{\mu=1}^K F_\mu$. This clearly implies $h(F|_\rho) \leq \sum_{\mu=1}^K h(F_\mu|_\rho)$ for every restriction ρ (the concatenation of decision trees strongly representing $F_\mu|_\rho$ for $\mu = 1, \dots, K$, strongly represents $F|_\rho$). Thus, we obtain

$$\mathbf{P}[h(F|_\rho) > h] \leq \sum_{\mu=1}^K \mathbf{P}[h(F_\mu|_\rho) > h2^{-\mu}], \quad (22)$$

and we treat every term in the right-hand side separately.

So, let us fix μ , $1 \leq \mu \leq K$, and let us fix $t \in F_\mu$; $t = x_{i_1} \wedge x_{i_2} \wedge \dots \wedge x_{i_w}$ ($i_1 < i_2 < \dots < i_w$). Let d be the index for which $\mu(x_{i_1} \wedge x_{i_2} \wedge \dots \wedge x_{i_d}) = \mu$. Then $d \leq \mu$, which implies that there are at most 2^μ subsets $V \subseteq t$ with $\mu_t(V) = \mu$. Consulting the definition of F_μ , we see that there exists a particular $V \subseteq Vars(t)$ with $\mu_t(V) = \mu$ such that $p_V \geq \delta(p/8)^\mu$.

Let this particular V have rank ℓ in our enumeration of all non-empty subsets. We are going to upper bound $\sum_{\nu=1}^\ell |V_\nu|$. Notice that, according to

the choice of this enumeration, $p_{V_\nu} \geq p_V$ for all $\nu \in [\ell]$. Therefore,

$$\sum_{\nu=1}^{\ell} (2/p)^{\mu(V_\nu)} p_{V_\nu}^2 \geq p_V^2 \cdot \sum_{\nu=1}^{\ell} (2/p)^{\mu(V_\nu)} \geq \delta^2 (p^2/64)^\mu \sum_{\nu=1}^{\ell} \mu(V_\nu) \geq \delta^2 (p^2/64)^\mu \sum_{\nu=1}^{\ell} |V_\nu|.$$

Comparing this with (20), we get

$$\sum_{\nu=1}^{\ell} |V_\nu| \leq \delta^{-1} (64/p^2)^\mu \leq s (64/p^2)^\mu. \quad (23)$$

The conclusion (23) holds for every $t \in F_\mu$. Therefore, if we define ℓ_μ as the maximal value ℓ for which the bound (23) holds, then for *every* $t \in F_\mu$, $V_1 \cup \dots \cup V_{\ell_\mu}$ will contain some $V \subseteq \text{Vars}(t)$ with $\mu_t(V) = \mu$. Since $V_1 \cup \dots \cup V_{\ell_\mu}$ is an initial segment in $\{x_1, \dots, x_n\}$, this implies that $\mu((V_1 \cup \dots \cup V_{\ell_\mu}) \cap \text{Vars}(t)) \geq \mu$.

Let now T_μ be the (oblivious) decision tree that queries all variables in $V_1 \cup \dots \cup V_{\ell_\mu}$ (in an arbitrary order). Our analysis implies that for every leaf π of T_μ , $(F_\mu)|_\pi$ is a monotone weighted $(K - \mu)$ -DNF. Therefore, we may apply the inductive assumption (14) and conclude that for each leaf π ,

$$\mathbf{P} \left[h((F_\mu)|_\pi) > \frac{1}{2} h 2^{-\mu} \right] \leq e^{-\frac{1}{2} h 2^{-\mu} (p\epsilon)^{2(K-\mu)}}.$$

By the definition of ℓ_μ and (15), the height of T_μ does not exceed $h(p\epsilon)^{2K} \cdot \{O(1/p^2)\}^\mu$, which is at most $\frac{1}{4} h 2^{-\mu} (p\epsilon)^{2(K-\mu)}$, as long as ϵ is small enough. Summing over all leaves of T_μ ,

$$\mathbf{P} \left[\exists \pi \in \{0, 1\}^{V_1 \cup \dots \cup V_{\ell_\mu}} \left(h((F_\mu)|_\pi) > \frac{1}{2} h 2^{-\mu} \right) \right] \leq e^{-\frac{1}{4} h 2^{-\mu} (p\epsilon)^{2(K-\mu)}}.$$

Noting that the height of the tree $(T_\mu)|_\rho$ never exceeds $\frac{1}{2} h 2^{-\mu}$, the latter event is logically implied by $h(F_\mu)|_\rho > h 2^{-\mu}$. Comparing with (22),

$$\mathbf{P}[h(F)|_\rho > h] \leq \sum_{\mu=1}^K e^{-\frac{1}{4} h 2^{-\mu} (p\epsilon)^{2(K-\mu)}} \leq \sum_{\mu=1}^K e^{-h(p\epsilon)^{2K} \cdot \frac{1}{4} \left(\frac{1}{2\epsilon^2}\right)^\mu},$$

which is at most $e^{-h(p\epsilon)^{2K}}$, as long as ϵ is small enough.

This proves (14) in Case 2, completes the inductive step, and the proof of Lemma 4.4 for monotone F .

In order to extend this lemma to the general case, let us note first that, by symmetry, we automatically have it when the DNF F is *pseudo-monotone*, defined as those DNF that can be turned into monotone by negating some variables. Now, every DNF F (interpreted as a set of terms) has a straightforward *fractional* cover by pseudo-monotone DNFs of acceptable size; that is, it possesses a random pseudo-monotone sub-DNF \mathbf{G} that contains every individual term $t \in F$ with sufficiently large probability. This \mathbf{G} is generated simply by picking a (total) assignment \mathbf{a} at random and including those terms that satisfy it. If it were an ordinary cover, we would have been trivially done. The following general lemma (that will also be repeatedly used in both our applications of Lemma 4.4) shows how to handle the more general fractional case.

Lemma 4.5 *Let F be a k -DNF, and \mathbf{G} be a random sub-DNF such that*

$$\forall t \in F (\mathbf{P}[t \in \mathbf{G}] \geq \epsilon), \quad (24)$$

ϵ an arbitrary parameter. Let ρ be a random restriction such that for every fixed G from the support of \mathbf{G} ,

$$\mathbf{P}[h(G|\rho) > h] \leq \delta,$$

δ another parameter. Then

$$\mathbf{P} \left[h(F|\rho) > h \left(\frac{2k}{\epsilon} + k + 1 \right) \right] \leq 2\delta/\epsilon.$$

Proof of Lemma 4.5. Arguing by averaging over the distribution of \mathbf{G} , we get $\mathbf{P}[h(\mathbf{G}|\rho) > h] \leq \delta$, where \mathbf{G} and ρ are assumed independent. By Markov's inequality,

$$\mathbf{P}_\rho[\mathbf{P}_\mathbf{G}[h(\mathbf{G}|\rho) > h] \geq \epsilon/2] \leq 2\delta/\epsilon.$$

Therefore, we only have to show that for every individual restriction ρ ,

$$\mathbf{P}[h(\mathbf{G}|\rho) > h] < \epsilon/2 \quad (25)$$

logically implies $h(F|\rho) \leq h \left(\frac{2k}{\epsilon} + k + 1 \right)$.

This is done by an adaptation of the ‘‘block sensitivity’’ method [Nis91, BBC⁺01] to our setting. Assume that (25) holds. We want to construct a

decision tree of height $\leq h \left(\frac{2k}{\epsilon} + k + 1 \right)$ strongly representing $F|_\rho$. We begin with a recursive construction of a sequence of decision trees $T_0, T_1, \dots, T_\ell, \dots$ that goes as follows.

T_0 is the trivial tree of height 0. In order to construct $T_{\ell+1}$ from T_ℓ , examine one by one all leaves π of T_ℓ . If either $t|_\pi = 0$ for all $t \in F|_\rho$, or $t|_\pi = 1$ for some $t \in F|_\rho$, we leave π alone. Otherwise, pick up an arbitrary non-trivial term $t_{\ell+1} \in (F|_\rho)|_\pi$, and append to the leaf π the oblivious decision tree querying all variables in $\text{Vars}(t_{\ell+1}) \setminus (\text{sup}(\rho) \cup \text{sup}(\pi))$. Repeating this procedure for all leaves π of T_ℓ , we get $T_{\ell+1}$.

We terminate this construction after $s \stackrel{\text{def}}{=} \lceil (2h/\epsilon) \rceil$ steps. The only leaves π of T_s that still may violate Definition 3.5 are those for which the procedure of appending a new tree was repeated all s times before we arrived at π . Let us fix any such leaf π , and let t_1, t_2, \dots, t_s be the terms picked by our algorithm along the path to π .

By (24),

$$\mathbf{E}[|\{t_1, t_2, \dots, t_s\} \cap \mathbf{G}|] \geq s \cdot \epsilon.$$

On the other hand, denoting by **Bad** the indicator function of the event $h(\mathbf{G}|_\rho) > h$, (25) implies

$$\mathbf{E}[|\{t_1, t_2, \dots, t_s\} \cap \mathbf{G}| \cdot \mathbf{Bad}] \leq s \cdot \mathbf{E}[\mathbf{Bad}] < s \cdot \frac{\epsilon}{2}.$$

Therefore, $\mathbf{E}[|\{t_1, t_2, \dots, t_s\} \cap \mathbf{G}| \cdot (1 - \mathbf{Bad})] > \frac{s\epsilon}{2} \geq h$; pick up a particular sub-DNF G that contains $(h + 1)$ terms $t_{\alpha_1}, \dots, t_{\alpha_{h+1}}$ from the list $\{t_1, t_2, \dots, t_s\}$ (and possibly some other terms) such that $h(G|_\rho) \leq h$. Let T_π be the decision tree of height $\leq h$ strongly representing $G|_\rho$. We complete out construction by appending the tree T_π to the leaf π (all vertices asking questions from $\text{sup}(\pi)$ are contracted in the process), and repeating this for all leaves π that still violate Definition 3.5.

Clearly, the final tree constructed in this way has height at most $sk + h \leq h \left(\frac{2k}{\epsilon} + k + 1 \right)$, and we claim that it strongly represents $F|_\rho$. For that we only have to check the leaves of the form $\pi * \sigma$, where π is a problematic leaf of T_s , and σ is a leaf of T_π such that π and σ are consistent.

For every $\nu \in [h + 1]$, let $V_\nu \stackrel{\text{def}}{=} \text{Vars}(t_{\alpha_\nu}) \setminus \bigcup_{\alpha < \alpha_\nu} \text{Vars}(t_\alpha)$. Then V_ν are disjoint and non-empty; therefore, since $|\text{sup}(\sigma)| \leq h$, there exists at least one $\nu_0 \in [h + 1]$ such that $V_{\nu_0} \cap |\text{sup}(\sigma)| = \emptyset$. Now, the values of π at the variables V_{ν_0} can be changed in such a way that for the resulting partial assignment π' we have $t_{\alpha_{\nu_0}}|_{\pi'} = 1$. $V_{\nu_0} \cap |\text{sup}(\sigma)| = \emptyset$ implies that π' and

σ are still consistent, hence $t_{\alpha_{\nu_0}}|_{\sigma} \not\equiv 0$ and, since $t_{\alpha_{\nu_0}} \in G|_{\rho}$, $\sigma \in Br_0(T_{\pi})$ is impossible. Thus, $\sigma \in Br_1(T_{\pi})$ which implies that there exists a term $t \in G|_{\rho} \subseteq F|_{\rho}$ such that $t|_{\sigma} = 1$, and the more $t|_{\pi^*\sigma} = 1$.

We have constructed (under the assumption (25)) a tree of height $\leq h\left(\frac{2k}{\epsilon} + k + 1\right)$ strongly representing $F|_{\rho}$. This completes the proof of Lemma 4.5. ■

Let us now finish the proof of Lemma 4.4 for arbitrary weighted K -DNF F . For every (total) assignment a to F , let F_a be the set of all terms in F satisfied by a . Then F_a is a pseudo-monotone sub-DNF of F ; therefore, we already have for it the required bound (13). Next, if \mathbf{a} is picked completely at random, then for every $t \in F$, $\mathbf{P}[t \in F_{\mathbf{a}}] = 2^{-w(t)} \geq 2^{-K}$. Thus, we may apply Lemma 4.5 with $k := K$, $\mathbf{G} := F_{\mathbf{a}}$, $\epsilon := 2^{-K}$, $h := \frac{h}{2K2^{K+K+1}}$ and $\delta := \exp\left(-\frac{h}{2K2^{K+K+1}}(p/2)^{O(K)}\right) \leq \exp(-h(p/2)^{O(K)})$, and this completes the proof of Lemma 4.4 in the general situation. ■

5. Hardness of Nisan generator for $\text{Res}(k)$

In this section we prove Theorem 2.7. The proof is technically involved, and for that reason is split into a chain of auxiliary claims. They will be assembled together only at the end of the section, although some informal intuition as to what we are doing will be provided as we go along.

Let us fix for the rest of the section an $m \times n$ (r, d) -lossless expander A , its ordering \leq , $b \in \{0, 1\}^m$ and an integer $k \geq 1$ such that (2) holds. Without loss of generality we can assume that $r \geq k$ (since otherwise the bound is trivial) and that $d \geq 1$ (since otherwise $\tau_{\leq}(A, b)$ is consistent and hence does not possess any refutations whatsoever). The overall strategy of our proof for the first time appeared in [BP96] and since that has become a standard tool in proof complexity. Namely, we want to design a random partial assignment ρ of the variables $\text{Vars}_{\leq}(A)$ that has the following two properties:

Height-reduction for every fixed k -DNF F , $h(F|_{\rho})$ is small with high probability;

Width-preservation with high probability every *resolution* refutation of $\tau_{\leq}(A, b)|_{\rho}$ must have large width.

Now, if a small size $\text{Res}(k)$ refutation P of $\tau_{\leq}(A, b)$ existed, then with high probability $h(F|_{\rho})$ would be small for *every* $F \in P$ by the **Height-reduction** property, and we could apply Proposition 3.6. Its conclusion, however, would be in immediate contradiction with the **Width-preservation** property.

We begin realizing this plan with **Width-preservation** (mostly because this part is by far easier), and we will show that large width of resolution refutations of $\tau_{\leq}(A, b)|_{\rho}$ is implied by a simple combinatorial property of ρ .

Some conventions on notation. In the rest of the paper we will be abbreviating the extension variables $y_{v_{\Sigma}}^i$ (where Σ is a non-empty \leq_i -initial segment of $J_i(A)$) by y_{Σ}^i . For technical reasons it will be also convenient to introduce the variables y_{\emptyset}^i , along with the axioms \bar{y}_{\emptyset}^i . Likewise, it will be convenient *not* to identify $y_{\{j\}}^i$ with x_j and with each other (as required in the general Definition 2.1), but introduce instead new axioms $y_{\{j\}}^i \vee \bar{x}_j$, $\bar{y}_{\{j\}}^i \vee x_j$. These conventions imply that all variables in the vectors \vec{x}, \vec{y}^i ($i \in [m]$) are pairwise distinct, and the axioms of $\tau_{\leq}(A, b)$ become of particularly symmetric form

$$y_{\emptyset}^i = 0, \quad y_{\Sigma \cup \{j\}}^i = y_{\Sigma}^i \oplus x_j, \quad y_{J_i(A)}^i = b_i \quad (i \in [m]) \quad (26)$$

(more precisely, they are clauses of width ≤ 3 resulting from the straightforward CNF expansion of these linear equations mod 2).

Definition 5.1 A restriction ρ of the variables $\text{Vars}_{\leq}(A)$ is *closed* if $\rho(y_{\emptyset}^i) = 0$ and $\rho(y_{J_i(A)}^i) = b_i$ for all $i \in [m]$. Let $J_x(\rho) \stackrel{\text{def}}{=} \{j \in [n] \mid x_j \in \text{sup}(\rho)\}$. ρ is *sparse* if it is closed and for every $i \in [m]$ and every two different initial segments $\Sigma \subset \Sigma'$ in $J_i(A)$ such that $y_{\Sigma}^i, y_{\Sigma'}^i \in \text{sup}(\rho)$, we have $|(\Sigma' \setminus \Sigma) \setminus J_x(\rho)| \geq 2d$.

Claim 5.2 *If ρ is sparse then every resolution refutation of $\tau_{\leq}(A, b)|_{\rho}$ must be of width $> r/4$.*

Proof. Given a closed restriction ρ of the variables $\text{Vars}_{\leq}(A)$, define the matrix $A|_{\rho}$ as follows. For every row i of the original matrix A let $\emptyset = \Sigma_0^i \subset \Sigma_1^i \subset \dots \subset \Sigma_{s_i}^i = J_i(A)$ be the complete list of those initial segments Σ in $J_i(A)$ for which $y_{\Sigma}^i \in \text{sup}(\rho)$. Then the rows of the matrix $A|_{\rho}$ are, by definition, indexed by the pairs (i, ν) ($i \in [m], \nu \in [s_i]$), its columns are indexed by $[n] \setminus J_x(\rho)$ (i.e., by those x -variables that are left unassigned by ρ), and the underlying set system is described as $J_{i,\nu}(A|_{\rho}) \stackrel{\text{def}}{=} (\Sigma_{\nu}^i \setminus \Sigma_{\nu-1}^i) \setminus J_x(\rho)$.

Note that $A|_\rho$ satisfies all assumptions of Corollary 3.4. Indeed, the bound $|J_{i,\nu}(A|_\rho)| \geq 2d$ is exactly the definition of sparseness. Next, given any set $\{(i_1, \nu_1), \dots, (i_\ell, \nu_\ell)\}$ of rows in the matrix $A|_\rho$ with $\ell \leq r$, applying the expansion property (1) for the original matrix A to the set $\{i_1, \dots, i_\ell\}$ gives us

$$\sum_{i \in \{i_1, \dots, i_\ell\}} |J_i(A)| - |\partial_A(\{i_1, \dots, i_\ell\})| \leq d \cdot |\{i_1, \dots, i_\ell\}| \leq d \cdot \ell. \quad (27)$$

On the other hand, it is easy to see that

$$\left. \begin{aligned} & \sum_{\alpha=1}^{\ell} |J_{i_\alpha, \nu_\alpha}(A|_\rho)| - |\partial_{A|_\rho}(\{(i_1, \nu_1), \dots, (i_\ell, \nu_\ell)\})| \\ & \leq \sum_{i \in \{i_1, \dots, i_\ell\}} |J_i(A)| - |\partial_A(\{i_1, \dots, i_\ell\})|. \end{aligned} \right\} \quad (28)$$

Indeed, let $\ell_j \stackrel{\text{def}}{=} |\{\alpha \in [\ell] \mid j \in J_{i_\alpha, \nu_\alpha}(A|_\rho)\}|$, then the left-hand side of (28) can be equivalently re-written as $\sum_{j=1}^n f(\ell_j)$, where $f(x) \stackrel{\text{def}}{=} \begin{cases} 0, & x \leq 1 \\ x, & x \geq 2 \end{cases}$ is a *non-decreasing* function. Likewise, the right-hand side is equal to $\sum_{j=1}^n f(\widehat{\ell}_j)$, where $\widehat{\ell}_j \stackrel{\text{def}}{=} |\{i \in \{i_1, \dots, i_\ell\} \mid j \in J_i(A)\}|$. But $\ell_j \leq \widehat{\ell}_j$ since for every fixed i the sets $\{J_{i_\alpha, \nu_\alpha}(A|_\rho) \mid i_\alpha = i\}$ are mutually disjoint, whence (28) follows.

Now, (27) and (28) imply that $A|_\rho$ is an (r, d) -lossless expander and therefore all assumptions of Corollary 3.4 are fulfilled.

We now define (in a natural way) the assignment $b|_\rho$ of the rows of the matrix $A|_\rho$ by letting

$$(b|_\rho)|_{i,\nu} \stackrel{\text{def}}{=} \rho(y_{\Sigma_{\nu-1}^i}^i) \oplus \rho(y_{\Sigma_\nu^i}^i) \oplus \bigoplus_{j \in (\Sigma_\nu^i \setminus \Sigma_{\nu-1}^i) \cap J_x(\rho)} \rho(x_j).$$

Finally, let $\leq|_\rho$ be the ordering of the matrix $A|_\rho$, where $(\leq|_\rho)|_{i,\nu}$ is the restriction of \leq_i onto $J_{i,\nu}(A|_\rho)$.

We extend the restriction ρ to a variable substitution ρ' of variables in $\text{Vars}_{\leq}(A)$ by variables in $\text{Vars}_{\leq|_\rho}(A|_\rho)$ defining it outside of $\text{sup}(\rho)$ as follows. All $x_j \notin \text{sup}(\rho)$ are simply left alone: $\rho'(x_j) \stackrel{\text{def}}{=} x_j$. For every $y_\Sigma^i \notin \text{sup}(\rho)$ we

identify the index ν such that $\Sigma_{\nu-1}^i \subset \Sigma \subset \Sigma_\nu^i$, and let

$$\rho'(y_\Sigma^i) \stackrel{\text{def}}{=} y_{(\Sigma \setminus \Sigma_{\nu-1}^i) \setminus J_x(\rho)}^{(i,\nu)} \oplus \rho(y_{\Sigma_{\nu-1}^i}^i) \oplus \bigoplus_{j \in (\Sigma \setminus \Sigma_{\nu-1}^i) \cap J_x(\rho)} \rho(x_j)$$

(cf. the definition of $b|_\rho$). It is straightforward to check that this extension of the original restriction ρ takes every equation in (26) either to $0=0$ or to an equation of the same form corresponding to $\tau_{\leq|\rho'}(A|_{\rho'}, b|_{\rho'})$, and therefore maps every resolution refutation P of $\tau_{\leq}(A, b)|_\rho$ to a resolution refutation $P|_{\rho'}$ of $\tau_{\leq|\rho'}(A|_{\rho'}, b|_{\rho'})$. By Corollary 3.4 (applied to $(A|_{\rho'}, \leq|\rho', b|_{\rho'})$), $P|_{\rho'}$ must contain a clause of width $> r/4$. Hence its pre-image in P also has width $> r/4$. Claim 5.2 is proved. ■

Our second (and much more complicated) goal is to design a distribution on sparse restrictions that fulfils the **Height-reduction** property. After a short thinking it becomes clear that in full generality this is impossible. Namely, if a term t contains a couple of variables $y_\Sigma^i, y_{\Sigma'}^i$ with Σ, Σ' close to each other than *no* sparse restriction whatsoever can set t to 1, which completely ruins the whole argument. Thus, our most immediate task is to get rid of such “nasty” terms.

More notation. For a term t in the variables $Vars_{\leq}(A)$, $J_x(t) \stackrel{\text{def}}{=} \{j \in [n] \mid x_j \in Vars(t)\}$ and $\text{dom}(t) \stackrel{\text{def}}{=} \{i \in [m] \mid \exists \Sigma (y_\Sigma^i \in Vars(t))\}$. For a DNF F , $\text{dom}(F) \stackrel{\text{def}}{=} \bigcup_{t \in F} \text{dom}(t)$. For a non-empty \leq_i -initial segment Σ , let $r(\Sigma)$ be its right end (maximal element). For uniformity of notation, we let $r(\emptyset) \stackrel{\text{def}}{=} \text{nil}_i$, where nil_i is an imaginary element with $\text{nil}_i <_i j$ for all $j \in J_i(A)$. In this notation, the difference $\Sigma' \setminus \Sigma$ of two initial segments $\Sigma \subseteq \Sigma'$ coincides with the *interval* $(r(\Sigma), r(\Sigma'))$. For subsets $L, R \subseteq [n]$, $\text{Conv}(L, R)$ is the minimal interval containing them both.

Definition 5.3 Let t be a term in the variables $Vars_{\leq}(A)$. For every $i \in \text{dom}(t)$ list in the form $\Sigma_1^i \subset \Sigma_2^i \subset \dots \subset \Sigma_{k_i}^i$ all initial segments Σ such that $y_\Sigma^i \in Vars(t)$. Say that t is *protected* if there exists a system of subsets $L_{i\nu}, R_{i\nu} \subseteq J_i(A)$ ($i \in \text{dom}(t)$, $\nu \in [k_i]$) such that:

1. $L_{i,1} <_i r(\Sigma_1^i) <_i R_{i,1} <_i L_{i,2} <_i r(\Sigma_2^i) <_i \dots <_i r(\Sigma_{k_i}^i) <_i R_{i,k_i}$;
2. $|L_{i\nu}|, |R_{i\nu}| = 3d$;
3. $\sum_{i \in \text{dom}(t)} \sum_{\nu \in [k_i]} |\text{Conv}(L_{i\nu}, R_{i\nu})| \leq 20kd$;

4. all $L_{i\nu}, R_{i\nu}$ are disjoint with $J_x(t)$.

The pair $(L_{i\nu}, R_{i\nu})$ (often also written in the form $(L_{i\Sigma}, R_{i\Sigma})$, where $\Sigma = \Sigma_\nu^i$), will be called a *protection of the variable* $y_{\Sigma_\nu^i}^i \in \text{Vars}(t)$, and the entire set system $\{L_{i\nu}, R_{i\nu}\}$ will be called a *protection of the term* t (protections are, of course, not necessarily uniquely defined). A DNF F is *protected* if all terms $t \in F$ have this property.

We now show that every term t with $w(t) \leq k$ can be replaced by a protected DNF that will be denoted by $R(t)$. For that purpose we will pick in a special way several ‘‘anchor’’ variables y_Σ^i ; those will be left intact. All other y -variables will be expanded as linear forms in x -variables modulo our knowledge of the value of a nearby anchor variable (see Definition 5.6 for details), and $R(t)$ will then be the naive DNF-expansion of the resulting $\wedge \oplus$ -circuit.

Definition 5.4 Let t be a term of width $\leq k$ in the variables $\text{Vars}_{\leq}(A)$. Denote by t' the term obtained from t by appending to it the literals $\bar{y}_\emptyset^i \wedge (y_{J_i(A)}^i)^{b_i}$ for every $i \in \text{dom}(t)$ (if t contains either y_\emptyset^i or $(y_{J_i(A)}^i)^{\bar{b}_i}$ for some $i \in [m]$, we immediately abort the construction and let $R(t) \stackrel{\text{def}}{=} 0$).

For a fixed $i \in \text{dom}(t)$, list all initial segments Σ with $y_\Sigma^i \in \text{Vars}(t')$: $\emptyset = \Sigma_0^i \subset \Sigma_1^i \subset \dots \subset \Sigma_{k_i+1}^i = J_i(A)$, and let $r_{i\nu} \stackrel{\text{def}}{=} r(\Sigma_\nu^i)$. Let

$$\text{Ker}_i(t) \stackrel{\text{def}}{=} J_i(A) \cap (J_x(t) \cup \bigcup_{i' \in \text{dom}(t) \setminus \{i\}} J_{i'}(A)). \quad (29)$$

Construct the graph G_i on $\{0, 1, \dots, k_i + 1\}$ by connecting $(\nu - 1)$ with ν if $|(r_{i,\nu-1}, r_{i\nu}] \setminus \text{Ker}_i(t)| < 6d$, and let $\Gamma_{i,0}, \Gamma_{i,1}, \dots, \Gamma_{i,\ell_i}$ be the connected components of this graph. $\Gamma_{i\alpha}$ is an interval in $\{0, 1, \dots, k_i + 1\}$; let $\Gamma_{i\alpha} = [\nu_{i\alpha}^\ell, \nu_{i\alpha}^r]$, and $\hat{\Gamma}_{i\alpha} \stackrel{\text{def}}{=} (r_{i,\nu_{i\alpha}^\ell}, r_{i,\nu_{i\alpha}^r}]$ be the corresponding interval in $J_i(A)$.

Claim 5.5 a)

$$\sum_{i \in \text{dom}(t)} |\text{Ker}_i(t) \cup \bigcup_{\alpha=0}^{\ell_i} \hat{\Gamma}_{i\alpha}| \leq 14kd; \quad (30)$$

b) $\ell_i \geq 1$ for every $i \in [m]$ (that is, y_\emptyset^i and $y_{J_i(A)}^i$ are not in the same connected component).

Proof. a). We have

$$\begin{aligned}
\text{Ker}_i(t) \cup \bigcup_{\alpha=0}^{\ell_i} \hat{\Gamma}_{i\alpha} &= \text{Ker}_i(t) \cup \bigcup_{(\nu-1, \nu) \in G_i} (r_{i, \nu-1}, r_{i\nu}] \\
&= \text{Ker}_i(t) \cup \bigcup_{(\nu-1, \nu) \in G_i} ((r_{i, \nu-1}, r_{i\nu}] \setminus \text{Ker}_i(t)) \\
&= (J_i(A) \cap \bigcup_{i' \in \text{dom}(t) \setminus \{i\}} J_{i'}(A)) \\
&\quad \cup (J_i(A) \cap (J_x(t) \setminus \bigcup_{i' \in \text{dom}(t) \setminus \{i\}} J_{i'}(A))) \\
&\quad \cup \bigcup_{(\nu-1, \nu) \in G_i} ((r_{i, \nu-1}, r_{i\nu}] \setminus \text{Ker}_i(t)).
\end{aligned}$$

Accordingly,

$$\left. \begin{aligned}
&\sum_{i \in \text{dom}(t)} |\text{Ker}_i(t) \cup \bigcup_{\alpha=0}^{\ell_i} \hat{\Gamma}_{i\alpha}| \\
&\leq \sum_{i \in \text{dom}(t)} |J_i(A) \cap \bigcup_{i' \in \text{dom}(t) \setminus \{i\}} J_{i'}(A)| \\
&\quad + \sum_{i \in \text{dom}(t)} |J_i(A) \cap (J_x(t) \setminus \bigcup_{i' \in \text{dom}(t) \setminus \{i\}} J_{i'}(A))| \\
&\quad + \sum_{i \in \text{dom}(t)} \sum_{(\nu-1, \nu) \in G_i} |(r_{i, \nu-1}, r_{i\nu}] \setminus \text{Ker}_i(t)|.
\end{aligned} \right\} (31)$$

The first summand in the right-hand side is bounded by kd due to the expansion property (1). Sets appearing in the second summand are disjoint, therefore, it is bounded by $|J_x(t)| \leq k$. Finally, $|(r_{i, \nu-1}, r_{i\nu}] \setminus \text{Ker}_i(t)| < 6d$ for every $i \in \text{dom}(t)$ and $(\nu - 1, \nu) \in G_i$, and also $\sum_{i \in \text{dom}(t)} |G_i| \leq \sum_{i \in \text{dom}(t)} (k_i + 1) \leq 2k$. Therefore, the third summand is bounded by $12kd$. (30) follows.

Part b) of this claim immediately follows from part a) and the bound (2). ■

Definition 5.6 (construction of $R(t)$ continued) Fix now one representative $\nu_{i\alpha}$ in every connected component $\Gamma_{i\alpha}$ such that $\nu_{i,0} = 0$ and $\nu_{i\ell_i} = k_i + 1$ (Claim 5.5 b) guarantees that this is possible): these will be the “anchor”

variables mentioned before Definition 5.4, and they will serve as reference points for variables $y_{\Sigma_\nu}^i$ with $\nu \in \Gamma_\alpha$. Let ϵ_α^i be the sign with which the anchor variable $y_{\Sigma_{\nu_{i\alpha}}}^i$ appears in t' . Then, since both t' and $R(t)$ assert that $y_{\Sigma_{\nu_{i\alpha}}}^i = \epsilon_\alpha^i$, modulo this fact we can replace y -variables corresponding to all others $\nu \neq \nu_{i\alpha}$ in $\Gamma_{i\alpha}$ as a linear form in x -variables only.

Formally, we define the expression $\tilde{R}(t)$ as the result of replacing in the term t' all variables $y_{\Sigma_\nu}^i$ with $\nu \in \Gamma_{i\alpha}$ and $\nu \neq \nu_{i\alpha}$ by the following linear forms:

$$y_{\Sigma_\nu}^i \mapsto \bigoplus \left\{ x_j \mid j \in \Sigma_\nu^i \Delta \Sigma_{\nu_{i\alpha}}^i \right\} \oplus \epsilon_\alpha^i. \quad (32)$$

Finally, we let $R(t)$ be the straightforward DNF expansion of $\tilde{R}(t)$, in which we also remove all ‘‘cosmetic’’ literals $\bar{y}_\emptyset^i, (y_{J_i(A)}^i)^{b_i}$ inserted there at the beginning of the construction.

Claim 5.7 a) $R(t)$ is a protected DNF such that $|Vars(R(t))| \leq O(kd)$.

b) There exist $Res(O(kd))$ inferences of $R(t)$ from $t, \tau_{\leq}(A, b)$ and, vice versa, of t from $R(t), \tau_{\leq}(A, b)$ that have size $\exp(O(kd))$ and contain at most $O(kd)$ variables.

Proof. a). The bound on $|Vars(R(t))|$ follows from the construction and Claim 5.5 a). We protect terms in $R(t)$ as follows. Let $y_{\Sigma_{\nu_{i\alpha}}}^i \in Vars(R(t))$ (note that $\Sigma_{\nu_{i\alpha}}^i \notin \{\emptyset, J_i(A)\}$). Protect this variable by the sets $L_{i\alpha}, R_{i\alpha}$, where $L_{i\alpha}$ consists of (3d) right-most points in $(r_{\nu_{i\alpha}^\ell - 1}, r_{\nu_{i\alpha}^\ell}] \setminus \text{Ker}_i(t)$, and $R_{i\alpha}$ consists of (3d) left-most points in $(r_{\nu_{i\alpha}^r}, r_{\nu_{i\alpha}^r + 1}] \setminus \text{Ker}_i(t)$. Let us check that all requirements of Definition 5.3 are fulfilled.

Since $(\nu_{i\alpha}^\ell - 1, \nu_{i\alpha}^\ell), (\nu_{i\alpha}^r, \nu_{i\alpha}^r + 1) \notin G_i$, the sets $(r_{\nu_{i\alpha}^\ell - 1}, r_{\nu_{i\alpha}^\ell}] \setminus \text{Ker}_i(t)$ and $(r_{\nu_{i\alpha}^r}, r_{\nu_{i\alpha}^r + 1}] \setminus \text{Ker}_i(t)$ have cardinality $\geq 6d$ each, which implies 1).

2) is obvious.

For 3) note that the intervals $\text{Conv}(L_{i\alpha}, R_{i\alpha})$ are disjoint for every fixed i , and that $\text{Conv}(L_{i\alpha}, R_{i\alpha}) \subseteq \text{Ker}_i(t) \cup L_{i\alpha} \cup \hat{\Gamma}_{i\alpha} \cup R_{i\alpha}$. Hence $\sum_{i \in \text{dom}(t)} \sum_\alpha |\text{Conv}(L_{i\alpha}, R_{i\alpha})| = \sum_{i \in \text{dom}(t)} |\bigcup_\alpha \text{Conv}(L_{i\alpha}, R_{i\alpha})|$; applying Claim 5.5 a) along with the obvious estimate $\sum_{i \in \text{dom}(t)} |\bigcup_\alpha (L_{i\alpha} \cup R_{i\alpha})| \leq 6kd$, we get the required bound.

Finally, 4) immediately follows from the construction. Indeed, given any fixed protection $(L_{i_0\alpha_0}, R_{i_0\alpha_0})$, all new x -variables introduced in $J_{i_0}(A)$ by the substitution (32) either belong to $\text{Ker}_{i_0}(t)$, or belong to $\bigcup_\alpha \hat{\Gamma}_{i_0\alpha}$. Both sets are disjoint with $L_{i_0\alpha_0}, R_{i_0\alpha_0}$.

b). As we explained inside Definition 5.6, the axioms from (26) pertaining to the same connected component of G_i semantically imply the equivalence of both sides in (32) in the presence of the literal $(y_{\Sigma_{\nu_{i\alpha}}}^i)^{\epsilon_\alpha}$, contain the axioms \bar{y}_0^i , $(y_{J_i(A)}^i)^{b_i}$, and thus semantically imply $t \equiv R(t)$. Now we only have to refer to the implicational completeness of $\text{Res}(k)$ and to the well-known fact that everything provable in this theory (and, in fact, even in Resolution) also has a proof of size at most exponential in the number of variables. ■

Remark 6 It is worth noting that the construction of $R(t)$ in fact gives yet another property crucial for the argument: the protections $L_{i\alpha}, R_{i\alpha}$ are pairwise disjoint (even when i varies). This property, however, will be re-established in the proof of Claim 5.13 anyway, and for this reason is not included in Definition 5.3.

Claim 5.8 *Assume that $\tau_{\leq}(A, b)$ has a $\text{Res}(k)$ refutation of size S . Then it also has an $\text{Res}(O(kd))$ refutation of size $S \cdot \exp(O(kd))$ in which all lines are of the form $F \vee F'$, where F is a protected $O(kd)$ -DNF, and $|\text{Vars}(F')| \leq O(kd)$.*

Proof. For a k -DNF F , let $R(F) \stackrel{\text{def}}{=} \bigvee_{t \in F} R(t)$. Claim 5.7 b) implies that every axiom $C \in \tau_{\leq}(A, b)$ (of width ≤ 3) gets converted to a DNF $R(C)$ that has an inference P from $\tau_{\leq}(A, b)$ of size $\exp(O(kd))$ and with $|\text{Vars}(P)| \leq O(kd)$. Also, the image of any inference rule in Definition 2.6 is admissible which can be seen by decoding and then encoding back the principal formula. For example, we simulate the R -image of AND-introduction as follows:

$$\frac{\frac{R(F) \vee R(\ell_1)}{R(F) \vee \ell_1} \quad \dots \quad \frac{R(F) \vee R(\ell_w)}{R(F) \vee \ell_w}}{R(F) \vee \bigwedge_{\nu=1}^w \ell_\nu} \quad \frac{\quad}{R(F) \vee R(\bigwedge_{\nu=1}^w \ell_\nu)}$$

In order to do this encoding/decoding, we simply use the inferences from Claim 5.7 b) weakened by $R(F)$. Since weakening does not change the number of lines, this induced inference has all the required properties. ■

The claims proved so far will allow us to concentrate for the purpose of **Height-reduction** on protected DNFs, and we now define a distribution on

sparse restrictions that, with an overwhelming probability, will reduce the representation height of any such DNF.

Definition 5.9 The random restriction ρ of the variables $Vars_{\leq}(A)$ is constructed as follows. Pick a subset $\mathbf{J} \subseteq [n]$ completely at random. Then for every $i \in [m]$ independently apply the following construction.

Pick a random set of variables $\tilde{\mathbf{Y}}^i \subseteq \{y_{\Sigma}^i \mid \Sigma \text{ an initial segment of } J_i(A)\}$ by including there the cosmetic variables y_{\emptyset}^i and $y_{J_i(A)}^i$ with probability 1, and every other y_{Σ}^i with probability $1/(2d)$, independently of each other. Say that $y_{\Sigma}^i, y_{\Sigma'}^i \in \tilde{\mathbf{Y}}^i$ with $\Sigma \subset \Sigma'$ *collide* if $|(\Sigma' \setminus \Sigma) \setminus \mathbf{J}| < 2d$. Remove from $\tilde{\mathbf{Y}}^i$ all variables $y_{\Sigma}^i \notin \{y_{\emptyset}^i, y_{J_i(A)}^i\}$ that collide with at least one other variable in $\tilde{\mathbf{Y}}^i$ (possibly, with y_{\emptyset}^i or $y_{J_i(A)}^i$). Let $\mathbf{Y}^i \subseteq \tilde{\mathbf{Y}}^i$ be the resulting set.

ρ , by definition, assigns y_{\emptyset}^i to 0, assigns $y_{J_i(A)}^i$ to b_i and assigns all other variables in $\{x_j \mid j \in \mathbf{J}\} \cup \bigcup_{i=1}^m \mathbf{Y}^i$ at random (and independently of each other).

Claim 5.10 $\mathbf{P}[\rho \text{ is sparse}] \geq 1/2$.

Proof. By inspecting definitions, ρ may be not sparse only in the “pathological” case when for some $i \in [m]$ we have $\mathbf{Y}^i = \{y_{\emptyset}^i, y_{J_i(A)}^i\}$. We bound the probability of this bad event separately for every $i \in [m]$.

By the bound (2), we may choose $s \geq \frac{C \log m}{9}$ disjoint intervals $\Delta_1, \dots, \Delta_s$ in $J_i(A)$, of length $9d$ each. Subdivide every Δ_{ν} into three sub-intervals $\Delta_{\nu}^{\ell}, \Delta_{\nu}^c, \Delta_{\nu}^r$, where $|\Delta_{\nu}^{\ell}| = |\Delta_{\nu}^r| = 4d$ and $|\Delta_{\nu}^c| = d$. Then the following events:

- $|\Delta_{\nu}^{\ell} \cap \mathbf{J}| \leq 2d, |\Delta_{\nu}^r \cap \mathbf{J}| \leq 2d;$
- $\tilde{\mathbf{Y}}^i$ contains exactly one variable y_{Σ}^i with $r(\Sigma) \in \Delta_{\nu}^c$ and no variables with $r(\Sigma) \in \Delta_{\nu}^{\ell} \cup \Delta_{\nu}^r$

have probability $\Omega(1)$ each, are independent and logically imply that the variable $y_{\Sigma}^i \in \tilde{\mathbf{Y}}^i$ with $r(\Sigma) \in \Delta_{\nu}^c$ does not collide with any other variable and hence stays in \mathbf{Y}^i . Therefore, the probability that this happens for at least one $\nu \in [s]$ (and, in particular, $\mathbf{Y}^i \neq \{y_{\emptyset}^i, y_{J_i(A)}^i\}$) is at least $1 - 1/(2m)$, provided the constant C in (2) is large enough. By the union bound, this implies $\mathbf{P}[\forall i \in [m](\mathbf{Y}^i \neq \{y_{\emptyset}^i, y_{J_i(A)}^i\})] \geq 1/2$. ■

We are going to apply Lemma 4.4 to show that for any protected DNF, $h(F|_{\rho})$ is small with high probability. For that we need to know that the

restriction ρ , when restricted to the set of variables $\text{Vars}(F)$, satisfies the weak independence property from Definition 4.3. The intuitive reason why it should be the case is already suggested by the proof of Claim 5.10 (the role of the “wings” $\Delta_\nu^\ell, \Delta_\nu^r$ in that proof will be played by protections). The major problem is, of course, that protections need not be disjoint, may be inconsistent for different occurrences of the same variable etc. We circumvent this by showing that F has a relatively small fractional cover by sub-DNFs for which these problems already do not occur, whereupon we will apply Lemma 4.5.

More notation. Given a protected DNF F , we fix once and for all protections $\{L_{i\Sigma}(t), R_{i\Sigma}(t)\}$ for every $t \in F$. Let also $P_{i\Sigma}(t) \stackrel{\text{def}}{=} L_{i\Sigma}(t) \cup R_{i\Sigma}(t)$.

Definition 5.11 A protected DNF F is *weakly regular* if the following three properties hold:

1. the sets $\bigcup_{t \in F} J_x(t)$ and $\bigcup_{t \in F} \bigcup_{i, \Sigma} P_{i\Sigma}(t)$ are disjoint;
2. for every $y_\Sigma^i \in \text{Vars}(F)$, the protection $(L_{i\Sigma}(t), R_{i\Sigma}(t))$ does not actually depend on the term t (and henceforth will be denoted simply by $(L_{i\Sigma}, R_{i\Sigma})$);
3. for every **fixed** $i \in [m]$, intervals $\text{Conv}(L_{i\Sigma}, R_{i\Sigma})$, where Σ runs over all \leq_i -initial segments with $y_\Sigma^i \in \text{Vars}(F)$ are pairwise disjoint.

Remark 7 The adjective “weakly” refers to the fact that we do not require the protections $P_{i\Sigma}$ to be disjoint for *different* i . Recall (Remark 6) that this disjointness property automatically follows from our construction for protections in any *fixed* term $t \in F$. This, however, seems to be of absolutely no help whatsoever for the uniform version, when t varies. The only way to enforce the disjointness of protections uniformly we know of additionally requires $|F|$ to be small, and for that reason will be incorporated in the proof of Claim 5.13.

Claim 5.12 *For every protected $O(kd)$ -DNF F , there exists a random weakly regular sub-DNF $\mathbf{G} \subseteq F$ such that $\min_{t \in F} \mathbf{P}[t \in \mathbf{G}] \geq \exp(-O(kd))$.*

Proof. The proof consists of three independent steps, and at every step we enforce one property required in Definition 5.11.

Step 1. For a colouring $\chi : [n] \rightarrow \{0, 1\}$, let

$$G_\chi \stackrel{\text{def}}{=} \left\{ t \in F \mid \chi(J_x(t)) \equiv 0 \wedge \chi\left(\bigcup_{i,\Sigma} P_{i\Sigma}(t)\right) \equiv 1 \right\}.$$

Then for every χ , G_χ has property 1) in Definition 5.11. Let $\chi : [n] \rightarrow \{0, 1\}$ be picked completely at random. Then, due to property 3) in Definition 5.3, $\mathbf{P}[t \in G_\chi] \geq \exp(-O(kd))$ for every particular $t \in F$.

Step 2. The idea behind enforcing property 2) in Definition 5.11 is similar to Step 1, but calculations get substantially more involved. For any system of protections $\vec{P} = (P_{i\Sigma} \mid y_\Sigma^i \in \text{Vars}(F))$, the sub-DNF $G_{\vec{P}} \stackrel{\text{def}}{=} \{t \in F \mid \forall y_\Sigma^i \in \text{Vars}(t) (P_{i\Sigma}(t) = P_{i\Sigma})\}$ has the required property 2). We only need to construct a random system \vec{P} in such a way that

$$\min_{t \in F} \mathbf{P}[t \in G_{\vec{P}}] \geq \exp(-O(kd)). \quad (33)$$

All $P_{i\Sigma}$ will be mutually independent. In order to construct $P_{i\Sigma}$ for a fixed variable $y_\Sigma^i \in \text{Vars}(F)$, pick first an integer $\mu_{i\Sigma} \geq 6d$ according to the distribution $\mathbf{P}[\mu_{i\Sigma} = \mu] = 2^{6d-\mu-1}$ ($\mu \geq 6d$). Next, let $\Delta_{i\Sigma}$ be the interval of length $2\mu_{i\Sigma}$ centered at $r(\Sigma)$ (if $|\Sigma| < \mu_{i\Sigma}$ or $|J_i(A) \setminus \Sigma| < \mu_{i\Sigma}$, then we abort the construction and output $P_{i\Sigma}$ arbitrarily). Finally, pick $P_{i\Sigma}$ as a random subset of $\Delta_{i\Sigma}$ that has cardinality $6d$.

Consider an arbitrary $t \in F$, and let $\mu_{i\Sigma}(t) \stackrel{\text{def}}{=} |\text{Conv}(L_{i\Sigma}(t), R_{i\Sigma}(t))|$. Then $\mathbf{P}[\mu_{i\Sigma} = \mu_{i\Sigma}(t)] \geq \exp(-O(\mu_{i\Sigma}(t)))$, and $\mathbf{P}[P_{i\Sigma} = P_{i\Sigma}(t) \mid \mu_{i\Sigma} = \mu_{i\Sigma}(t)] \geq \binom{2\mu_{i\Sigma}(t)}{6d}^{-1} \geq \left(\frac{\mu_{i\Sigma}(t)}{d}\right)^{-O(d)}$. Combining these bounds together, we get $\mathbf{P}[P_{i\Sigma} = P_{i\Sigma}(t)] \geq \exp(-O(\mu_{i\Sigma}(t))) \cdot \left(\frac{\mu_{i\Sigma}(t)}{d}\right)^{-O(d)}$. Multiplying over all $y_\Sigma^i \in \text{Vars}(t)$,

$$\mathbf{P}[t \in G_{\vec{P}}] \geq \prod_{y_\Sigma^i \in \text{Vars}(t)} \left\{ \exp(-O(\mu_{i\Sigma}(t))) \cdot \left(\frac{\mu_{i\Sigma}(t)}{d}\right)^{-O(d)} \right\}.$$

Since $\sum_{y_\Sigma^i \in \text{Vars}(t)} \mu_{i\Sigma}(t) \leq O(kd)$ by property 3) in Definition 5.3, the first term in this product is bounded from below by $\exp(-O(kd))$. In order to bound the second term, we apply the arithmetic-geometric mean inequality. Denoting the number of y -variables in t by w (which, due to properties 2), 3) in Definition 5.3, is bounded by $O(k)$), we have the calculation

$$\prod_{y_\Sigma^i \in \text{Vars}(t)} \left(\frac{\mu_{i\Sigma}(t)}{d}\right)^{-O(d)} \geq \left(\frac{\sum_{y_\Sigma^i \in \text{Vars}(t)} \mu_{i\Sigma}(t)}{dw}\right)^{-O(dw)} \geq (O(k/w))^{-O(dw)} \geq \exp(-O(kd)).$$

This proves (33) and concludes the analysis at Step 2.

Step 3. For this step we additionally assume that F already satisfies the consistency property 2) in Definition 5.11, and let \vec{P} be the corresponding system of protections of the variables $y_\Sigma^i \in \text{Vars}(F)$. Choose $\tilde{\mathbf{Y}}^i$ according to Definition 5.9. Say that $y_\Sigma^i, y_{\Sigma'}^i \in \text{Vars}(F) \cap \tilde{\mathbf{Y}}^i$ \vec{P} -collide if $\text{Conv}(L_{i\Sigma}, R_{i\Sigma}) \cap \text{Conv}(L_{i\Sigma'}, R_{i\Sigma'}) \neq \emptyset$. For every \vec{P} -colliding pair $y_\Sigma^i, y_{\Sigma'}^i$, identify arbitrarily any one point j in the above intersection, and remove from $\tilde{\mathbf{Y}}^i$ that variable $y_{\Sigma_0}^i$ ($\Sigma_0 \in \{\Sigma, \Sigma'\}$) for which the \leq_i -distance between $r(\Sigma_0)$ and j is larger (both y_Σ^i and $y_{\Sigma'}^i$ if these distances are equal). Let $\mathbf{Y}^i(\vec{P})$ be the set of remaining variables.

Clearly, the sub-DNF $G_{\mathbf{Y}(\vec{P})} \stackrel{\text{def}}{=} \{t \in F \mid \text{Vars}(t) \subseteq \{x_1, \dots, x_n\} \cup \bigcup_{i \in \text{dom}(t)} \mathbf{Y}^i(\vec{P})\}$ has the required property 3). Further, every variable y_Σ^i may be removed from $\tilde{\mathbf{Y}}^i$ only if it \vec{P} -collides with some $y_{\Sigma'}^i$ such that the \leq_i -distance between $r(\Sigma)$ and $r(\Sigma')$ does not exceed $2|\text{Conv}(L_{i\Sigma}, R_{i\Sigma})|$.

Now, for every particular term $t \in F$ mark all those variables $y_{\Sigma'}^i$ in $\text{Vars}(F)$ for which the above bound on the \leq_i -distance holds for at least one $y_\Sigma^i \in \text{Vars}(t)$. Property 3) in Definition 5.3 implies that we altogether have marked at most $O(kd)$ variables. Therefore, the event “for every marked variable $y_\Sigma^i, y_{\Sigma'}^i \in \bigcup_{i \in \text{dom}(t)} \tilde{\mathbf{Y}}^i$ if and only if $y_\Sigma^i \in \text{Vars}(t)$ ” has probability $\geq \exp(-O(kd))$. On the other hand, since variables in $\text{Vars}(t)$ never \vec{P} -collide with each other (due to property 1) in Definition 5.3), this event logically implies that $\text{Vars}(t) \subseteq \{x_1, \dots, x_n\} \cup \bigcup_{i \in \text{dom}(t)} \mathbf{Y}^i(\vec{P})$. Thus, $\mathbf{P}[t \in G_{\mathbf{Y}(\vec{P})}] \geq \exp(-O(kd))$, which completes the analysis of Step 3.

Combining things together, the random sub-DNF $G_\chi \cap G_{\vec{P}} \cap G_{\mathbf{Y}(\vec{P})}$ (where χ, \vec{P} and the auxiliary random variables $\tilde{\mathbf{Y}}^i$ are independent) has all the required properties. This completes the proof of Claim 5.12. ■

We assume that any weakly regular DNF F is weighted according to the following weight function μ_F : $\mu_F(x_j) \stackrel{\text{def}}{=} 1$ and $\mu_F(y_\Sigma^i) \stackrel{\text{def}}{=} |\text{Conv}(L_{i\Sigma}, R_{i\Sigma})|$.

Claim 5.13 *There exists an absolute constant $p > 0$ such that for every weakly regular protected DNF F , the random restriction ρ restricted to the variables $\text{Vars}(F)$ is (r, μ_F, p) -independent.*

Proof. Let $Z \subseteq \text{Vars}(F)$ be a set of variables with $|Z| \leq r$. Denote by I the set of all $i \in [m]$ for which Z contains at least one variable y_Σ^i . Clearly,

$|I| \leq r$. Therefore, by the expansion property (1), there exists $i_0 \in I$ such that

$$|J_{i_0}(A) \cap \bigcup_{i \in I \setminus \{i_0\}} J_i(A)| \leq d. \quad (34)$$

This implies that for every Σ with $y_{\Sigma}^{i_0} \in \text{Vars}(F)$ there exist $L'_{i_0\Sigma} \subseteq L_{i_0\Sigma}$ and $R'_{i_0\Sigma} \subseteq R_{i_0\Sigma}$ of size $2d$ each that are disjoint with $\bigcup_{i \in I \setminus \{i_0\}} J_i(A)$. In particular, these $L'_{i_0\Sigma}, R'_{i_0\Sigma}$ are disjoint not only between themselves (by property 3) in Definition 5.11), but also with any other $P_{i\Sigma}$ when $i \neq i_0$. Next, choose $i_1 \in I \setminus \{i_0\}$ such that $|J_{i_1}(A) \cap \bigcup_{i \in I \setminus \{i_0, i_1\}} J_i(A)| \leq d$ and repeat this procedure until we find pairwise disjoint subsets $L'_{i\Sigma} \subseteq L_{i\Sigma}$, $R'_{i\Sigma} \subseteq R_{i\Sigma}$ for all $y_{\Sigma}^i \in Z$. Let as before $P'_{i\Sigma} \stackrel{\text{def}}{=} L'_{i\Sigma} \cup R'_{i\Sigma}$.

Now, the space Ω required in Definition 4.1 consists of three parts: $\Omega = \Omega_1 \times \Omega_2 \times \Omega_3$, and $\omega = (\omega_1, \omega_2, \omega_3)$, where $\omega_i \in \Omega_i$ are independent. The role of ω_1 is played by the random set $\{x_j \mid j \in \mathbf{J}\} \cup \bigcup_{i=1}^m \tilde{\mathbf{Y}}^i$ from Definition 5.9, and ω_2 consists of additional independent Bernoulli variables used for assigning $\{x_j \mid j \in \mathbf{J}\} \cup \bigcup_{i=1}^m \mathbf{Y}^i$. π depends only on ω_1, ω_2 and represents the construction of the random variable ρ from that definition.

Next, we construct the second required mapping $\pi'(\omega_1, \omega_2, \omega_3)$ as follows. Let $J \stackrel{\text{def}}{=} \{j \mid x_j \in \omega_1\}$ and let \tilde{Y}^i consist of all $y_{\Sigma}^i \in \omega_1$. First, define from J and \tilde{Y}^i random subsets $\mathbf{J}' \subseteq J$ and $(\mathbf{Y}^i)' \subseteq \tilde{Y}^i \cap Z$ (the third random variable ω_3 accounts for the extra randomness used in this construction). Let $j \in \mathbf{J}'$ with probability $2p$, independently for all $j \in J \cap Z$. If $y_{\Sigma}^i \in \tilde{Y}^i \cap Z$ and either $J \cap P'_{i\Sigma} \neq \emptyset$ or \tilde{Y}^i contains any other $y_{\Sigma'}^i$ with $r(\Sigma') \in \text{Conv}(L'_{i\Sigma}, R'_{i\Sigma})$, then $y_{\Sigma}^i \notin (\mathbf{Y}^i)'$. Otherwise, y_{Σ}^i is included into $(\mathbf{Y}^i)'$ with some probability $p_{i\Sigma}$ to be specified later, independently for all variables y_{Σ}^i (and independently of \mathbf{J}'). Finally, $\pi'(\omega_1, \omega_2, \omega_3)$ assigns variables in $\{x_j \in Z \mid j \in \mathbf{J}'\} \cup \bigcup_{i \in I} (\mathbf{Y}^i)'$ using ω_2 .

$y_{\Sigma}^i \in (\mathbf{Y}^i)'$ can take place only if $J \cap P'_{i\Sigma} = \emptyset$ and \tilde{Y}^i does not contain any $y_{\Sigma'}^i$ with $r(\Sigma') \in \text{Conv}(L'_{i\Sigma}, R'_{i\Sigma})$ other than y_{Σ}^i . This implies that y_{Σ}^i can not collide with any other variable in \tilde{Y}^i . Therefore, $(\mathbf{Y}^i)' \subseteq \mathbf{Y}^i$, and $\pi'(\omega_1, \omega_2, \omega_3)$ is a sub-restriction of $\pi(\omega_1, \omega_2) \approx \rho$.

We are only left to show that for a suitable choice of the probabilities $p_{i\Sigma}$, $\pi'(\omega_1, \omega_2, \omega_3)$ will be equidistributed with $\rho_{\mu_F, Z, p}$.

The properties listed in Definition 5.11, along with the disjointness property of $L'_{i\Sigma}, R'_{i\Sigma}$ ensured at the beginning of our proof, imply that the facts $j \in \mathbf{J}'$, $y_{\Sigma}^i \in (\mathbf{Y}^i)'$ ($x_j, y_{\Sigma}^i \in Z$) depend on the behaviour of J, \tilde{Y}^i on pairwise different variables. Therefore, if we randomize also over $\mathbf{J}, \tilde{\mathbf{Y}}^i$,

all these events are independent of each other. Denote by A the following event: “ $\mathbf{J} \cap P'_{i\Sigma} = \emptyset$ and $y_\Sigma^i \in \tilde{\mathbf{Y}}^i$ and $\tilde{\mathbf{Y}}^i$ does not contain any y_Σ^i with $r(\Sigma') \in \text{Conv}(L'_{i\Sigma}, R'_{i\Sigma})$ other than y_Σ^i ”: this is exactly the pre-requisite for including y_Σ^i into $(\mathbf{Y}^i)'$. The last remark implies that the three parts of this event are independent and hence we conclude

$$\mathbf{P}[A] = 2^{-4d} \frac{1}{2d} \left(1 - \frac{1}{2d}\right)^{|\text{Conv}(L'_{i\Sigma}, R'_{i\Sigma})| - 1} \geq 2^{-4d} \cdot \frac{1}{2d} \left(1 - \frac{1}{2d}\right)^{\mu(y_\Sigma^i) - 1} \geq \exp(-O(\mu(y_\Sigma^i))),$$

where for the last estimate we used the obvious bound $\mu(y_\Sigma^i) \geq 6d$. Hence

$$\mathbf{P}[y_\Sigma^i \in (\mathbf{Y}^i)'] = \mathbf{P}[A] \cdot \mathbf{P}[y_\Sigma^i \in (\mathbf{Y}^i)' | A] \geq \exp(-O(\mu(y_\Sigma^i))) p_{i\Sigma}. \quad (35)$$

Therefore, if $p > 0$ is small enough, the probabilities $p_{i\Sigma} \leq 1$ can be chosen in such a way that $\mathbf{P}[y_\Sigma^i \in (\mathbf{Y}^i)'] = p^{\mu(y_\Sigma^i)}$, and, clearly, $\mathbf{P}[j \in \mathbf{J}'] = p$ for $x_j \in Z$. Thus, $\pi'(\omega_1, \omega_2, \omega_3)$ has the same distribution as $\rho_{\mu_F, Z, p}$, which completes the proof of Claim 5.13. ■

Now we have at our disposal all tools necessary for proving the following switching lemma for protected DNFs.

Claim 5.14 *Let F be a protected $O(kd)$ -DNF. Then for every $h \leq r$,*

$$\mathbf{P}[h(F|_\rho) \geq h] \leq \exp(-h/2^{O(kd)}).$$

Proof. Assume first that F is weakly regular, and let $\mu = \mu_F$. Then property 3) in Definition 5.3 implies that F is even a *weighted* $O(kd)$ -DNF. With this remark, our claim for weakly regular F immediately follows from Lemma 4.4 and Claim 5.13.

In order to generalize this to the case of arbitrary protected $O(kd)$ -DNF, we use Lemma 4.5 in combination with Claim 5.12, just in the same way as at the end of Section 4. ■

Now we can easily finish the proof of Theorem 2.7. Let P be a $\text{Res}(k)$ refutation of $\tau_{\leq}(A, b)$ that has size S . We need to prove that $S \geq \exp(r/2^{O(kd)})$.

Applying Claim 5.8, for some $K = O(kd)$ we get a $\text{Res}(K)$ refutation P' of $\tau_{\leq}(A, b)$ that has size $\leq S2^K$, and in which every line has the form $F \vee F'$, where F is a protected K -DNF and $|Vars(F')| \leq K$. By Claims 5.10, 5.2,

$$\mathbf{P}[\text{every resolution refutation of } \tau_{\leq}(A, b)|_\rho \text{ has width } > r/4] \geq 1/2. \quad (36)$$

Comparing this with Proposition 3.6 ($k := K$ and $h := r/(4K)$), we get

$$\mathbf{P}[\exists G \in P'(h(G|_\rho) > r/(4K))] \geq 1/2. \quad (37)$$

On the other hand,

$$\mathbf{P}[\exists G \in P'(h(G|_\rho) > r/(4K))] \leq S2^K \cdot \max_{G \in P'} \mathbf{P}[h(G|_\rho) > r/(4K)], \quad (38)$$

and we treat every line $G \in P'$ individually. Let $G = F \vee F'$, where F is a protected K -DNF and $|Vars(F')| \leq K$. Obviously, $h(F'|_\rho) \leq K$, and we can assume w.l.o.g. that $K < r/(8K)$ (otherwise, the bound we are proving becomes trivial). Thus,

$$\mathbf{P}[h(G|_\rho) > r/(4K)] \leq \mathbf{P}[h(F|_\rho) > r/(8K)]. \quad (39)$$

Applying Claim 5.14 (with $h := r/(8K) - 1$), we find

$$\mathbf{P}[h(F|_\rho) > r/(8K)] \leq \exp(-r/2^{O(kd)}). \quad (40)$$

Theorem 2.7 follows by comparing (40), (39) and (38) with (37).

6. Stretching the number of output bits

In this section we prove Theorems 2.10, 2.12.

Proof of Theorem 2.10. Let A be an $m \times n$ (r, d)-lossless expander such that the bound (5) holds, and let \leq be an arbitrary ordering of A . Let S be the minimal size of a $\text{Res}(k)$ refutation P of any CNF having the form

$$\bigwedge_{\nu=1}^H \tau_{C_{\leq, A}}(x_1^{(\nu)}, \dots, x_n^{(\nu)}, \bar{y}^{(\nu)}, q_1^{(\nu)}, \dots, q_m^{(\nu)}), \quad (41)$$

from Definition 2.9. First we remark that we may assume w.l.o.g. that $H \leq S$.

Indeed, P may contain at most S axioms; therefore, there exist at most S indices $\nu \in [H]$ for which P contains an axiom from $\tau_{C_{\leq, A}}(x_1^{(\nu)}, \dots, x_n^{(\nu)}, \bar{y}^{(\nu)}, q_1^{(\nu)}, \dots, q_m^{(\nu)})$. Remove from (41) all other conjunctive terms; then P will still be a $\text{Res}(k)$ refutation of the resulting sub-CNF. This sub-CNF itself has the form (41), with the only exception that some $q_i^{(\nu)}$ may be equal to variables not appearing in the lists $\bar{x}^{(\nu)}, \bar{y}^{(\nu)}$. Substituting them arbitrarily with Boolean

constants 0,1, we get a CNF of the form (41) with $H \leq S$ that still has a $\text{Res}(k)$ refutation of size S .

Note that (41) encodes the system of linear equations

$$\bigoplus_{j \in J_i(A)} x_j^{(\nu)} = q_i^{(\nu)} \quad (\nu \in [H], i \in [m]).$$

Let us transfer those $q_i^{(\nu)}$ which actually are the variables from the list $\{x_j^{(\mu)} \mid j \in [n], \mu < \nu\}$ to the left-hand side. Then we get a linear system with constant right-hand side, and it also has the form $\hat{A}X = \hat{b}$ for some matrix \hat{A} and vector \hat{b} . It will turn out that \hat{A} has almost as good expansion properties as A itself, and we will apply to it Theorem 2.7.

In order to formalize this intuition, let A^H be the *direct sum* of H copies of A . That is, rows of A^H are indexed by $[H] \times [m]$, columns are indexed by $[H] \times [n]$ and $a_{(\nu,i),(\mu,j)}^H \stackrel{\text{def}}{=} \begin{cases} a_{ij} & \text{if } \nu = \mu \\ 0 & \text{if } \nu \neq \mu \end{cases}$. Let \hat{A} be obtained from

A^H by setting additionally $\hat{a}_{(\nu,i),(\mu,j)} := 1$ whenever $q_i^{(\nu)} = x_j^{(\mu)}$. Let also $\hat{b}_{(\nu,i)} \stackrel{\text{def}}{=} \begin{cases} q_i^{(\nu)} & \text{if } q_i^{(\nu)} \text{ is a Boolean constant} \\ 0 & \text{if } q_i^{(\nu)} \text{ is a variable} \end{cases}$. Finally, we order the rows of

A^H according to the ordering \leq of the matrix A . Whenever we add in \hat{A} the new 1-entry (μ, j) to the row (ν, i) (that is, when $q_i^{(\nu)} = x_j^{(\mu)}$), we declare it to be the *largest* element in $J_{(i,\nu)}(\hat{A})$. Denote the resulting ordering of \hat{A} by $\hat{\leq}$.

Given the above intuition, it is straightforward to check that there exists a (naturally defined) variable substitution that takes every clause in (41) into a clause having a resolution inference from $\tau_{\hat{\leq}}(\hat{A}, \hat{b})$ of size $O(1)$. This implies that $\tau_{\hat{\leq}}(\hat{A}, \hat{b})$ has a $\text{Res}(k)$ refutation of size $O(S)$.

On the other hand, it is easy to see that the direct sum of (r, d) -lossless expanders is still an (r, d) -lossless expander. Also, if we append at most one 1-entry per row to an (r, d) -lossless expander, we come up with an $(r, d+1)$ -lossless expander. Applying Theorem 2.7, we see that either the condition (2) is violated for \hat{A} or we have $S \geq \exp(r/2^{O(kd)})$ (in which case we are done). It remains to note that in the first case we have $s \leq Cd(k + \log(mH))$ which implies (if the constant in (5) is twice as large as the constant in (2)) that $H \geq \exp(\Omega(s/d))$. Since $S \geq H$, Theorem 2.10 follows in this case as well. ■

Proof of Theorem 2.12. (cf. [Kra04]) Given any Boolean circuit C with n inputs and $2n$ outputs and any $b \in \{0, 1\}^{2^h \cdot n}$, $\tau(C^h, b)$ can be expressed in the form (4) as $\bigwedge_{|u| \leq h-1} \tau_C(x_1^{(u)}, \dots, x_n^{(u)}, \bar{y}^{(u)}, q_1^{(u)}, \dots, q_{2n}^{(u)})$, where

$$q_i^{(u)} \stackrel{\text{def}}{=} \begin{cases} x_i^{(u*0)} & \text{if } |u| < h-1, i \leq n \\ x_{n-i}^{(u*1)} & \text{if } |u| < h-1, i \geq n+1 \\ b_{(u*0),i} & \text{if } |u| = h-1, i \leq n \\ b_{(u*1),i} & \text{if } |u| = h-1, i \geq n+1. \end{cases}$$

Now Theorem 2.12 immediately follows from Theorem 2.10. ■

7. Random matrices have good expansion properties

The statements of this sort have been re-appearing in the literature at a steady rate beginning from [Pin73]. We, however, have not been able to find any particular source handling the matter in the generality needed for our purposes. Thus, we prove Theorem 2.5 from the scratch (this is not hard anyway).

Let

$$r \stackrel{\text{def}}{=} n^\delta, \quad d \stackrel{\text{def}}{=} C \frac{\log m}{\log n},$$

where δ is a sufficiently small and C a sufficiently large constant; assume for simplicity that d is even. Note that $d \leq n^\epsilon$, where ϵ is the constant from the statement of the theorem. We claim that if a set I of rows with $|I| = \ell \leq r$ in the matrix $\mathbf{A}_{m,n}$ violates the expansion property (1), then there exists a set of $\ell d/2$ columns J such that $\mathbf{A}_{m,n}$ contains at least ℓd ones in the rectangle $I \times J$. Indeed, let $J \stackrel{\text{def}}{=} \{j \in [n] \mid |\{i \in I \mid j \in J_i(A)\}| \geq 2\} (= \bigcup_{i \in I} J_i(A) \setminus \partial_A(I))$. The bound ℓd on the number of ones in $I \times J$ follows from our assumption that I violates (1). Hence, if $|J| \leq \ell d/2$, we are done (add $\ell d/2 - |J|$ columns to J arbitrarily). Otherwise, its arbitrary subset of cardinality exactly $\ell d/2$ would do. Calling a rectangle $I \times J$ such that $|I| \leq r$, $|J| = \frac{d}{2}|I|$ and $I \times J$ contains at least $d|I|$ ones *dense*, it remains to show that the probability of existence of at least one dense rectangle is $O(1/m)$.

For any fixed value of ℓ , there exist at most m^ℓ choices of I with $|I| = \ell$, at most $n^{\ell d/2}$ choices of J with $|J| = \ell d/2$ and at most $(\ell d)^{O(\ell d)}$ choices of ℓd positions in $I \times J$. Therefore, the probability that for given ℓ there exists at least one dense $\ell \times (\ell d/2)$ rectangle does not exceed

$$m^\ell n^{\ell d/2} (\ell d)^{O(\ell d)} (n^{-2/3})^{\ell d} \leq m^\ell \left(\frac{(\ell d)^{O(1)}}{n} \right)^{\ell d/6}.$$

Next, this is $\leq m^\ell n^{-\Omega(\ell d)}$ provided the constants ϵ, δ are small enough which, in turn, is $\leq m^{-2\ell}$ provided the constant C in the definition of d is large enough.

This is the bound on the probability that $\mathbf{A}_{m,n}$ contains a dense rectangle $I \times J$ with $|I| = \ell$. Therefore, the probability that $\mathbf{A}_{m,n}$ is not an (r, d) -lossless expander is bounded by $\sum_{\ell=1}^{\infty} m^{-2\ell} \leq O(1/m)$. Theorem 2.5 is proved.

8. Unprovability of circuit lower bounds by small $\text{Res}(k)$ proofs

The general idea toward extracting Theorem 2.13 from a pseudorandom function generator was extensively discussed in Introduction. We are, however, dealing with a rather weak proof system and, moreover, Definition 2.2 severely restricts the choice of the encoding for the circuit $C_{A,\leq}^h$. Thus, we should be careful in checking that the natural reduction can be indeed carried over with the limited tools at our disposal (cf. the previous arguments of this sort in [Raz98, Raz04a, Raz04b, Kra04]).

Let f_n be a Boolean function in n variables and $n^2 \leq t \leq 2^n$. We begin with reproducing the formal definition of the CNF $\text{Circuit}_t(f_n)$ from [Raz98, Raz04b].

First, we list all variables of $\text{Circuit}_t(f_n)$ (some of them have peculiar long names like $\text{InputType}'_\nu(v)$), along with their intended meaning:

$$\begin{aligned} y_{av} \ (a \in \{0, 1\}^n, v \in [t]) & \quad \text{the Boolean value computed at} \\ & \quad \text{the computational node } v \text{ on the input string } a; \\ y_{a\nu v} \ (a \in \{0, 1\}^n, \nu \in \{1, 2\}, v \in [t]) & \quad \text{the value on } a \text{ brought to } v \text{ by} \\ & \quad \text{the } \nu\text{'s input to } v; \end{aligned}$$

- $Fanin(v)$ – this is 0 if v is NOT-gate and 1 if v is AND-gate or OR-gate;
- $Type(v)$ – when $Fanin(v) = 1$, this is 0 if v is AND-gate and 1 if v is OR-gate;
- $InputType_\nu(v)$ – this is 0 if ν 's input to v is a constant or a variable and 1 if it is one of the previous computational gates;
- $InputType'_\nu(v)$ – when $InputType_\nu(v) = 0$, this is 0 if ν 's input to v is a constant, and 1 if it is a variable;
- $InputType''_\nu(v)$ – when $InputType_\nu(v) = InputType'_\nu(v) = 0$, this equals the ν 's input to v ;
- $InputVar_\nu(v, i)$ ($i \in [n]$) – when $InputType_\nu(v) = 0$, $InputType'_\nu(v) = 1$, this is 1 iff ν 's input to v is the i th variable;
- $INPUTVAR_\nu(v, i)$ – equals $\bigvee_{i' \leq i} InputVar_\nu(v, i')$, introduced to keep bottom fan-in bounded;
- $InputNode_\nu(v, v')$ ($v' < v$) – when $InputType_\nu(v) = 1$, this is 1 iff ν 's input to v is the previous gate v' ;
- $INPUTNODE_\nu(v, v')$ – analogously to $INPUTVAR_\nu(v, i)$.

$Circuit_t(f_n)$ is the conjunction of (conjunctive normal forms equivalent to)

the following axioms:

$$\begin{aligned}
& \neg \text{InputType}_\nu(v) \wedge \neg \text{InputType}'_\nu(v) \longrightarrow (y_{av} \equiv \text{InputType}''_\nu(v)); \\
& \neg \text{InputType}_\nu(v) \wedge \text{InputType}'_\nu(v) \longrightarrow \neg(\text{InputVar}_\nu(v, i) \wedge \\
& \quad \text{InputVar}_\nu(v, i')) \quad (i \neq i'); \\
& \neg \text{InputType}_\nu(v) \wedge \text{InputType}'_\nu(v) \longrightarrow (\text{INPUTVAR}_\nu(v, i) \equiv \\
& \quad (\text{INPUTVAR}_\nu(v, i-1) \vee \text{InputVar}_\nu(v, i))) \\
& \quad (\text{INPUTVAR}_\nu(v, 0) \stackrel{\text{def}}{=} 0); \\
& \neg \text{InputType}_\nu(v) \wedge \text{InputType}'_\nu(v) \longrightarrow \text{INPUTVAR}_\nu(v, n); \\
& \neg \text{InputType}_\nu(v) \wedge \text{InputType}'_\nu(v) \wedge \text{InputVar}_\nu(v, i) \longrightarrow (y_{av} \equiv a_i); \\
& \text{the analogous group of axioms for } \text{InputNode}; \\
& \neg \text{Fanin}(v) \longrightarrow (y_{av} \equiv \neg y_{a1v}); \\
& \text{Fanin}(v) \wedge \neg \text{Type}(v) \longrightarrow (y_{av} \equiv (y_{a1v} \wedge y_{a2v})); \\
& \text{Fanin}(v) \wedge \text{Type}(v) \longrightarrow (y_{av} \equiv (y_{a1v} \vee y_{a2v})); \\
& y_{at} \equiv f(a).
\end{aligned}$$

Let $t_0 \stackrel{\text{def}}{=} \delta \sqrt{t/n}$, δ a sufficiently small constant. Note that $t_0 \geq t^{\Omega(1)}$ (since $t \geq n^2$). Fix an arbitrary $(2t_0 \times t_0)$ ($t^{\Omega(1)}, O(1)$)-lossless expander A such that $|J_i(A)| \geq t^{\Omega(1)}$ for all $i \in [2t_0]$ (its existence follows from Theorem 2.5). Fix an arbitrary ordering \leq of A and consider the iterated circuit $C_{A, \leq}^{n+1}$ from Section 6. Let $b \in \{0, 1\}^{2^{n+1}t_0}$ be given by $b_{aj} \stackrel{\text{def}}{=} f_n(a_1, \dots, a_n)$ ($a \in \{0, 1\}^n$, $j \in [2t_0]$).

In order to prove Theorem 2.13, we are going to reduce $\tau(C_{A, \leq}^{n+1}, b)$ to $\text{Circuit}_t(f_n)$ (that is, substitute variables of $\text{Circuit}_t(f_n)$ by “simple” formulas in the variables of $\tau(C_{A, \leq}^{n+1}, b)$ so that “simple” (which in our context means $\text{Res}(k)$) refutations of $\text{Circuit}_t(f_n)$ get transformed into simple ($\text{Res}(2k)$) refutations of $\tau(C_{A, \leq}^{n+1}, b)$). For that purpose we will convert the circuit $C_{A, \leq}^{n+1}$ (with t_0 inputs and $2^{n+1}t_0$ outputs, naturally split into 2^n groups with $2t_0$ bits each, of which we will select one bit per group) to the (single-output) Boolean circuit $D_{n, \vec{x}}$ in n Boolean variables z_1, \dots, z_n parameterized by t_0 variables x_1, \dots, x_{t_0} . We will require that $D_{n, \vec{x}}(a_1, \dots, a_n)$ is equal to the a 'th selected bit in $C_{A, \leq}^{n+1}(x)$, and that the size of $D_{n, \vec{x}}$ is *polynomial* in n . We employ the same construction that was used for self-defeating Natural Proofs [RR97, Theorem 4.1]. But since we need to check that this transformation can be carried over in a rather weak proof system, we provide a few (tedious)

technical details. The new parameter ℓ below corresponds to the iteration level.

The skeleton of $D_{n,\bar{x}}$ consists of the gates $v[\ell, i, \Sigma]$, where $0 \leq \ell \leq n$, $i \in [2t_0]$ and Σ an initial segment in $J_i(A)$ computing Boolean functions $f_{\bar{x}}[\ell, i, \Sigma]$ in the variables z_1, \dots, z_n . The values $f_{\bar{x}}[\ell, i, \Sigma](a_1, \dots, a_n)$ of these functions are defined as follows. We take the circuit $(C_{A,\leq})_{a_1 \dots a_\ell}$ from Definition 2.11, look at the gate v_{Σ}^i in this circuit (that we will denote in what follows by $v_{\Sigma}^i(a_1 \dots a_\ell)$) and output as $f_{\bar{x}}[\ell, i, \Sigma](a_1, \dots, a_n)$ its value when $\tau(C_{A,\leq}^{m+1}, b)$ is fed with x_1, \dots, x_{t_0} (we also naturally let $f_{\bar{x}}[\ell, i, \emptyset](a_1, \dots, a_n) \stackrel{\text{def}}{=} 0$). Then these functions have the following recursive definition

$$\left. \begin{aligned} f_{\bar{x}}[\ell, i, \emptyset] &:= 0 \\ f_{\bar{x}}[0, i, \Sigma \cup \{j\}] &:= f_{\bar{x}}[0, i, \Sigma] \oplus x_j \\ f_{\bar{x}}[\ell, i, \Sigma \cup \{j\}] &:= f_{\bar{x}}[\ell, i, \Sigma] \oplus \{(\bar{z}_\ell \wedge f_{\bar{x}}[\ell - 1, j, J_j(A)]) \\ &\quad \vee (z_\ell \wedge f_{\bar{x}}[\ell - 1, j + t_0, J_{j+t_0}(A)])\}, \ell \geq 1. \end{aligned} \right\} \quad (42)$$

We define $D_{n,\bar{x}}$ as the circuit resulting from expanding these recursive definitions in the standard basis $\{\neg, \wedge, \vee\}$. We let the output gate of $D_{n,\bar{x}}$ be $v[n, 1, J_1(A)]$ (cf. the definition of b above).

$D_{n,\bar{x}}$ has size $O(t_0^2 n)$ which is at most t if the constant δ in the definition of t_0 is small enough. We begin constructing the required substitution ρ by assigning first all structural variables $Fanin(v), Type(v)$ etc., except for $InputType''_\nu(v)$, to appropriate Boolean constants describing the topology of $D_{n,\bar{x}}$. We also let $\rho(InputType''_\nu(v)) \stackrel{\text{def}}{=} x_j$ whenever v is the gate of $D_{n,\bar{x}}$ (necessarily resulting from an instruction in (42) of the second type) whose ν 's input is x_j . In the case $InputType''_\nu(v) = 1$ we let $\rho(y_{avv}) \stackrel{\text{def}}{=} \rho(y_{av'})$ ($\rho(y_{av'})$ themselves are yet to be defined), where (v', v) is the ν 's input leading to v in $D_{n,\bar{x}}$. If $\neg InputType''_\nu(v)$, then ν 's input to v is either one of the variables z_i or a known Boolean constant $\epsilon \in \{0, 1\}$ or one of the unknown constants x_j , and we let $\rho(y_{avv})$ be a_i, ϵ or x_j , respectively. Further, if the gate $v[\ell, i, \Sigma]$ explicitly appears in (42), then we let $\rho(y_{a,v[\ell,i,\Sigma]}) \stackrel{\text{def}}{=} y_{v_{\Sigma}^i}^i(a_1 \dots a_\ell)$ (which is the variable of $\tau(C_{A,\leq}^{m+1}, b)$).

We, however, can not extend ρ as a variable substitution to the remaining variables y_{av} , where v is an auxiliary gate resulting from expanding instructions in (42) in the standard basis $\{\neg, \wedge, \vee\}$. But even in this case we still

can let $\rho(y_{av})$ be a Boolean function of just two variables $y_{v_{\Sigma}^i(a_1 \dots a_{\ell})}$ and $y_{v_{j^*}^{j^*}(A)(a_1 \dots a_{\ell-1})}$, where $j^* = j$ (the right end of Σ) if $a_{\ell} = 0$ and $j + t_0$ if $a_{\ell} = 1$.

Summarizing the above argument, we have constructed a substitution ρ that takes every variable of $Circuit_t(f_n)$ to a Boolean function depending of at most two variables of $\tau(C_{A, \leq}^{m+1}, b)$. Let us extend this substitution to k -DNF formulas F by letting $\rho(F) \stackrel{\text{def}}{=} \bigvee_{t \in F} \widehat{\rho(t)}$, where $\widehat{\rho(t)}$ is the straightforward DNF expansion of $\rho(t)$. Clearly, $\rho(F)$ is a $2k$ -DNF, and the ρ -image of any inference rule of $\text{Res}(k)$ can be simulated in $\text{Res}(2k)$ by an inference of size $\exp(O(k))$. Further, given the “intended interpretation”, it is easy to check by inspection that for every axiom C of $Circuit_t(f_n)$, $\rho(C)$ has a $\text{Res}(2)$ inference of size $O(1)$ from $\tau(C_{A, \leq}^{m+1}, b)$.

These remarks imply that every $\text{Res}(k)$ refutation of $Circuit_t(f_n)$ of size S gives rise to a $\text{Res}(2k)$ refutation of $\tau(C_{A, \leq}^{m+1}, b)$ of size $S \cdot \exp(O(k))$. The proof of Theorem 2.13 is now completed by applying Theorem 2.12 (with $n := t_0$; $r, s \geq t^{\Omega(1)}$; $d \leq O(1)$; $h := n + 1$).

9. Pigeonhole principle

In this section we prove Theorem 2.15. Review first some more material from [SBI04].

Let $m = 2n$ (the same proof works with $m = (1 + \epsilon)n$, for any absolute constant $\epsilon > 0$). For a bipartite graph $G = ([m] \cup [n], E)$ ($E \subseteq [m] \times [n]$), let $\neg\text{onto} - \text{PHP}(G)$ be the CNF in the variables $\{x_{ij} \mid (i, j) \in E\}$ which is obtained from $\neg\text{onto} - \text{PHP}_n^m$ by the restriction assigning to 0 all variables $\{x_{ij} \mid (i, j) \notin E\}$. Let ρ^G be the random restriction constructed in the following way. Pick a random subset $\mathbf{J} \subseteq [n]$ by including there every $j \in [n]$ with probability $1/4$ independently of each other. For $j \in \mathbf{J}$ select uniformly (and independently for different j) one neighbour i_j of j in G , assign $x_{i_j j}$ to 1, and assign to 0 all other x_{ij} with $(i, j) \in E$ and $i \neq i_j$.

Proposition 9.1 ([SBI04]) *There exists a graph G of maximal degree $O(\log n)$ such that*

\mathbf{P} [every resolution refutation of $\neg\text{onto} - \text{PHP}(G)|_{\rho^G}$ has width $> n/24$] $\geq 1/2$.

Proof. This is essentially [SBI04, Lemma 18], with the only difference that we have additionally included the group of onto axioms Q_j . This does not affect its proof in any way. ■

Fix from now on any particular graph G with properties from Proposition 9.1, and let $\Delta \leq O(\log n)$ be its maximal degree. Since $\neg\text{onto} - \text{PHP}(G)$ is obtained from $\neg\text{onto} - \text{PHP}_n^m$ by a restriction, it is sufficient to prove the required bound for $\neg\text{onto} - \text{PHP}(G)$. The proof follows the pattern laid out in Section 5, although in the current case it is much simpler.

Definition 9.2 A term in the variables $\{x_{ij} \mid (i, j) \in E\}$ is *reduced* if it is monotone and does not contain any sub-term of the form $x_{ij} \wedge x_{i'j}$, $i \neq i'$. A DNF F is *reduced* if all terms $t \in F$ have this property.

Reduced DNFs will play the role of protected ones in Section 5. Let us mention for the record (we will need this in the proof of Claim 9.4) that every reduced DNF that mentions at most k pigeons contains at most $k\Delta$ variables and $O(\Delta^k)$ clauses.

Definition 9.3 For a term t , let us denote by $R(t)$ the reduced DNF that is constructed as follows. Let $\tilde{R}(t)$ be the result of replacing in the term t all negative literals \bar{x}_{ij} with $\bigvee \{x_{i'j} \mid (i', j) \in E \wedge i' \neq i\}$. Let $R(t)$ be the straightforward DNF expansion of $\tilde{R}(t)$, in which we remove all terms containing at least one sub-term of the form $x_{ij} \wedge x_{i'j}$ ($i' \neq i$).

Claim 9.4 *Let t be a term of width $\leq k$ in the variables $\{x_{ij} \mid (i, j) \in E\}$.*

- a) $R(t)$ is a reduced DNF.
- b) *There exist $\text{Res}(O(k))$ -inferences of $R(t)$ from $t, \neg\text{onto} - \text{PHP}(G)$ and, vice versa, of t from $R(t), \neg\text{onto} - \text{PHP}(G)$ that have size $\Delta^{O(k)}$ and contain at most $k\Delta$ variables.*

Proof. a) is obvious. For part b) we could have used the same reasoning based on implicational completeness as in the proof of Claim 5.7, but this would have led to an inference of size $\exp(O(k\Delta))$, at least a priori. We circumvent this by the following ad hoc hybrid-type argument.

Let $\{i_1, \dots, i_\ell\}$ ($\ell \leq k$) be an enumeration of all pigeons mentioned in t . For $0 \leq \nu \leq \ell$ split the term t as $t = t'_\nu \wedge t''_\nu$, where t'_ν is the part corresponding to the pigeons $\{i_1, \dots, i_\nu\}$, and t''_ν corresponds to the remaining

pigeons $\{i_{\nu+1}, \dots, i_\ell\}$. Let $R_\nu(t) \stackrel{\text{def}}{=} R(t'_\nu) \wedge t''_\nu$, so that $R_0(t) = t$ and $R_\ell(t) = R(t)$. We consequently infer in $\text{Res}(k)$ all equivalences $R_\nu(t) \equiv R_{\nu+1}(t)$ and then combine them together. Since by an earlier observation, every reduced DNF $R_\nu(t)$ contains only $O(\Delta^k)$ clauses, it can be done by an inference of size $\Delta^{O(k)}$. ■

Claim 9.5 *Assume that $\neg\text{onto-PHP}(G)$ has a $\text{Res}(k)$ refutation of size S . Then it also has an $\text{Res}(O(k))$ refutation of size $S \cdot \Delta^{O(k)}$ in which all lines are of the form $F \vee F'$, where F is a reduced k -DNF, and $|\text{Vars}(F')| \leq k\Delta$.*

Proof. As in the proof of Claim 5.8, for a DNF F let $R(F) \stackrel{\text{def}}{=} \bigvee_{t \in F} R(t)$. Notice that R does not change the monotone axioms Q_i, Q_j , and that $R(Q_{i_1, i_2; j}) = Q_j$. With these remarks in mind, the rest of the proof is identical to the proof of Claim 5.8. ■

Finally, let us prove the following PHP-oriented switching lemma.

Claim 9.6 *For any reduced k -DNF F and any parameter h ,*

$$\mathbf{P}[h(F|_\rho) > h] \leq \exp(-h/\Delta^{O(k)}).$$

Proof. Call a reduced DNF F *regular* if $\text{Vars}(F)$ does not contain any pair of variables $x_{ij}, x_{i'j}$ with $i \neq i'$. Then for every regular F , ρ^G acts independently on the variables from $\text{Vars}(F)$, and, moreover, $\mathbf{P}[\rho^G(x_{ij}) = 0], \mathbf{P}[\rho^G(x_{ij}) = 1] \geq \frac{1}{4\Delta}$. This implies that the restriction of ρ^G to the variables in $\text{Vars}(F)$ is $(\infty, \mu_{\text{triv}}, 1/(2\Delta))$ -independent. Applying Lemma 4.4, we prove our claim in the case F is regular.

For the general case we apply the same trick as before. Let the function $\theta : [n] \rightarrow [2n]$ be picked completely at random, and $G_\theta \stackrel{\text{def}}{=} \{t \in F \mid \forall x_{ij} \in \text{Vars}(t)(i = \theta(j))\}$. Then G_θ is regular, and $\forall t \in F(\mathbf{P}[t \in G_\theta] \geq \Delta^{-k})$. Now the proof is completed by applying Lemma 4.5. ■

The proof of Theorem 2.15 is completed in exactly the same way as the proof of Theorem 2.7 at the end of Section 5.

10. Polynomial Calculus with Resolution

Throughout this section we fix an arbitrary field \mathbb{F} with $\text{char}(\mathbb{F}) \neq 2$. First we need the following generalization of Corollary 3.4.

Corollary 10.1 *Let A be an (r, d) -lossless expander of size $m \times n$ such that $|J_i(A)| \geq 2d$ for all $i \in [m]$. Then for every ordering \leq and every $b \in \{0, 1\}^m$, every PCR refutation of $\tau_{\leq}(A, b)$ must have degree $> r/8$.*

Proof. Although this follows by the technique of [BGIP01] (cf. remark in [ABRW00] before Theorem 3.10), it is easier to apply more general result from [BSI10] that for our purposes can be stated as follows. If a CNF τ results from expanding a set of \mathbb{F}_2 -linear equations, then every PCR refutation of τ over \mathbb{F} (remember that $\text{char}(\mathbb{F}) \neq 2$) gives rise to a *resolution* refutation of τ whose width is at most twice as large as the degree of P .

Remark 8 Strictly speaking, [BSI10, Theorem 2.7] is formulated in terms of so-called *Gaussian refutations*, but it is a well-known fact that for systems of linear equations, Gaussian width and resolution width coincide. It is also worth mentioning here (we will need this observation in the proof of Claim 10.4 below) that the overhead factor of two comes from the following stronger property that is a by-product of their proof: for every clause C appearing in the resulting resolution refutation we have $\text{Vars}(C) \subseteq \text{Vars}(m_1) \cup \text{Vars}(m_2)$, where m_1 and m_2 are some monomials in the original PCR refutation.

Since $\tau_{\leq}(A, b)$ always has this “linear” form, Corollary 10.1 follows from Corollary 3.4. ■

Now the proofs of Theorems 2.18, 2.19 and 2.20 are more or less straightforward adaptation of the corresponding results for Resolution (= Res(1)).

Proof of Theorem 2.18. Monomials in the variables $x_1, \dots, x_n, \bar{x}_1, \dots, \bar{x}_n$ can be identified, via the transformation (6) and up to a multiplicative constant $\alpha \in \mathbb{F}^*$, with respective clauses. In this way, variable substitutions naturally act on polynomials from $\mathbb{F}[x_1, \dots, x_n, \bar{x}_1, \dots, \bar{x}_n]$ and take PCR inferences to PCR inferences of the same (or lesser) size. This remark, along with Corollary 10.1, implies the following analogue of Claim 5.2 (proved in exactly the same way):

Claim 10.2 *If ρ is sparse then every PCR refutation of $\tau_{\leq}(A, b)|_{\rho}$ must have degree $> r/8$.*

The analogue of (36) will thus be

$$\mathbf{P}[\text{every PCR refutation of } \tau_{\leq}(A, b)|_{\rho} \text{ has degree } > r/8] \geq 1/2. \quad (43)$$

Consider the mapping R from Definitions 5.4, 5.6 restricted to literals (R , in fact, is almost always identical, except for literals of those variables y_Σ^i for which either Σ or $J_i(A) \setminus \Sigma$ is very small). Let $R_{\mathbb{F}}$ denote the corresponding polynomial homomorphism over the field \mathbb{F} . $R_{\mathbb{F}}$ takes any PCR refutation P of $\tau_{\leq}(A, b)$ into another PCR refutation in which every line has the form $R_{\mathbb{F}}(f) \cdot f'$, where $f \in P$ and $\deg(f') \leq O(d)$. Note that unlike Claim 5.8, we do *not* make any conclusions about the *size* of the refutation $R_{\mathbb{F}}(P)$ (and in fact it *may* grow out of control).

Next, we remark that the proof of Claim 5.14 actually allows a finer analysis for protected $O(d)$ -DNF of the form $R(C)$, C a clause. Namely, assuming $k = 1$, we get

Claim 10.3 *For every clause C in the variables $\text{Vars}_{\leq}(A)$, and every $h \leq r$, either $|\text{Vars}(R(C))| \leq h$ or*

$$\mathbf{P}[R(C)|_{\rho} \neq 1] \leq \exp(-h/2^{O(d)})$$

(*unlike all previous claims of this sort, $R(C)|_{\rho} \neq 1$ here simply means that $R(C)|_{\rho}$ is not **semantically** equal to 1*).

Proof of Claim 10.3. Assume that $|\text{Vars}(R(C))| > h$. Applying Claim 5.14, we get that with the required probability $1 - \exp(-h/2^{O(d)})$ there exists a decision tree T_{ρ} of height $< h$ strongly representing $R(C)|_{\rho}$. Inspecting the proof of Lemma 4.4 for monotone F , we see that the tree T given by this construction actually has a stronger property. Namely, whenever $\pi \in Br_0(T)$, then not only $(t|_{\rho})|_{\pi} = 0$ for all $t \in F$ (as required by Definition 3.5), but in fact even $t|_{\pi} = 0$. The proof of Lemma 4.5 can be easily modified to preserve this property: we only have to go over terms $t \in F$ (rather than $t \in F|_{\rho}$) in the construction of the sequence $T_0, T_1, \dots, T_{\ell}, \dots$. Therefore, we may also assume that the tree $T|_{\rho}$ also has this stronger property.

$R(C)$, however, is the result of a DNF expansion of a disjunction of linear forms. Such DNFs can be set to 0 only by restrictions that assign *all* their variables. Therefore, since $|\text{Vars}(R(C))| > h$ and the height of $T|_{\rho}$ is at most h , we have $Br_0(T|_{\rho}) = \emptyset$. $R(C)|_{\rho} = 1$ (in the semantical sense) follows. ■

We now finish the proof of Theorem 2.18. Let P be a PCR refutation of $\tau_{\leq}(A, b)$. From (43) we in particular get

$$\mathbf{P}[\exists g \in R_{\mathbb{F}}(P)(\deg(g|_{\rho}) > r/8)] \geq 1/2.$$

As we remarked above, every line in the refutation $R_{\mathbb{F}}(P)$ has the form $R_{\mathbb{F}}(f) \cdot f'$ with $f \in P$ and $\deg(f') \leq O(d) \leq r/16$. Next, $\deg(R_{\mathbb{F}}(f)|_{\rho}) > r/16$ implies that there exists a monomial $\alpha \cdot \Gamma_C$ in f ($\alpha \in \mathbb{F}^*$) such that $|\text{Vars}(R(C))| > r/16$ and $R_{\mathbb{F}}(\Gamma_C)|_{\rho} \neq 0$. Applying Claim 10.3 (with $h := r/16$), we see that the probability of this event for every particular $\alpha \cdot \Gamma_C \in P$ is bounded by $\exp(-r/2^{O(d)})$. Theorem 2.18 now follows by the same calculation as at the end of Section 5. ■

All proofs in Section 6 hold for any proof system that is closed under variable substitutions. In particular, Theorem 2.19 (as well as the analogue of Theorem 2.12 for PCR not stated explicitly in Section 2) follows from Theorem 2.18 by the same proof.

The only problem with the proof of Theorem 2.13 is that the reduction ρ constructed in Section 8 is not a variable substitution (and, as we already remarked above in the proof of Theorem 2.18, the size of PCR proofs may blow up exponentially from applying such ρ). The only variables that create this problem are y_{av} , where v is an auxiliary gate resulting from expanding the instructions in (42). This time, however, our circuit $D_{n,\bar{x}}$ is also allowed to contain the parity gates; therefore, such auxiliary gates are confined to be one of $\bar{z}_\ell \wedge f[\ell-1, j, J_j(A)]$, $z_\ell \wedge f[\ell-1, j+t_0, J_{j+t_0}(A)]$, $(\bar{z}_\ell \wedge f[\ell-1, j, J_j(A)]) \vee (z_\ell \wedge f[\ell-1, j+t_0, J_{j+t_0}(A)])$ explicitly appearing in (42). It remains to notice that for every fixed $a \in \{0, 1\}^n$, every one of these three gates computes either 0 or one of the two functions $f[\ell-1, j, J_j(A)]$, $f[\ell-1, j+t_0, J_{j+t_0}(A)]$. Thus, at the expense of allowing \oplus -gates, ρ can be turned into a variable substitution, and the rest of the proof carries over to PCR without any further changes.

We now turn to the proof of our last result, Theorem 2.21, that does not seem to have any obvious analogue for $\text{Res}(k)$.

Let ρ be a restriction of the variables $\text{Vars}_{\leq}^{\text{Cycl}}(A)$. Denote by $E^i(\rho)$ the set of all endpoints of all cyclic intervals $\Delta \not\equiv J_i(A)$ with $y_\Delta^i \in \text{sup}(\rho)$. $E^i(\rho)$ defines a partition of $J_i(A)$ into cyclic intervals $\Delta_1^i(\rho), \dots, \Delta_{\ell_i(\rho)}^i(\rho)$ such that whenever $y_\Delta^i \in \text{sup}(\rho)$, Δ is a disjoint union of some of these intervals. Say that ρ is *consistent* if whenever $y_{\Delta(1)}^i, \dots, y_{\Delta(w)}^i \in \text{sup}(\rho)$, and $\Delta(1) \oplus \dots \oplus \Delta(w) = 0$, we have $\rho(\Delta(1)) \oplus \dots \oplus \rho(\Delta(w)) = 0$. Like in Section 5, let $J_x(\rho) \stackrel{\text{def}}{=} \{j \in [n] \mid x_j \in \text{sup}(\rho)\}$, and say that ρ is *sparse* if $|\Delta_\nu^i(\rho) \setminus J_x(\rho)| \geq 2d$ for every $i \in [m]$, $\nu \in [\ell_i(\rho)]$.

For a clause C in the variables $\text{Vars}_{\leq}^{\text{Cycl}}(A)$, let $J_x(C) \stackrel{\text{def}}{=} \{j \in [n] \mid x_j \in \text{Vars}(C)\}$ and $\text{dom}(C) \stackrel{\text{def}}{=} \{i \in [m] \mid \exists \Delta \neq J_i(A) (y_\Delta^i \in \text{Vars}(C))\}$. Call the quantity

$w_A(C) \stackrel{\text{def}}{=} |J_x(C)| + |\text{dom}(C)|$ the A -width of the clause C . The A -degree of a monomial $\alpha \cdot \Gamma_C$ ($\alpha \in \mathbb{F}^*$) is defined as $\deg_A(\alpha \cdot \Gamma_C) \stackrel{\text{def}}{=} w_A(C)$, and the A -degree of a polynomial is the maximal A -degree of a monomial occurring in it⁷. The A -width of a resolution proof [A -degree of a PCR proof] is the maximal A -width [A -degree] of a clause [polynomial, respectively] occurring in it.

Claim 10.4 *If ρ is consistent and sparse then every PCR refutation P of $\tau_{\leq}^{\text{Cycl}}(A, b)$ must contain a monomial m such that $\deg_A(m) > r/8$ and $m|_{\rho} \neq 0$.*

Proof. This is analogous to the proof of Claim 5.2, so we only remark the differences.

First, in the cyclic case the reduction ρ constructed in the proof of that claim is no longer a variable substitution: $\rho(y_{\Delta}^i)$ is in general the parity of two variables in $\text{Vars}_{\leq|\rho}^{\text{Cycl}}(A|_{\rho})$. If, however, we extend in a natural way the notions of A -width and A -degree to clauses/monomials in the variables $\text{Vars}_{\leq|\rho}^{\text{Cycl}}(A|_{\rho})$ (simply by ignoring the second superscript ν in $y_{\Delta}^{(i,\nu)}$), then it turns out that ρ still does not increase A -degree. Assuming (for the sake of contradiction) that no monomial m with the required properties exist, we conclude that $\deg_A(P|_{\rho}) \leq r/8$.

Next, we use the same argument from [BSI10] as in the proof of Corollary 10.1 to convert $P|_{\rho}$ into a resolution refutation of $\tau_{\leq|\rho}^{\text{Cycl}}(A|_{\rho}, b|_{\rho})$. As we already noticed above (see Remark 8), for every clause C in the resulting resolution refutation, $\text{Vars}(C) \subseteq \text{Vars}(m_1) \cup \text{Vars}(m_2)$, where m_1, m_2 are some monomials in the original PC refutation. Hence, the A -width of this resulting refutation of $\tau_{\leq|\rho}^{\text{Cycl}}(A|_{\rho}, b|_{\rho})$ is at most $r/4$.

Finally, when Corollary 3.4 is applied to the triple $(A|_{\rho}, \leq|_{\rho}, b|_{\rho})$, it can be generalized in two ways. First, $\tau_{\leq|\rho}(A|_{\rho}, b|_{\rho})$ can be replaced by $\tau_{\leq|\rho}^{\text{Cycl}}(A|_{\rho}, b|_{\rho})$ (since, like $\tau_{\leq|\rho}(A|_{\rho}, b|_{\rho})$, this is also a sub-CNF of $\tau(A|_{\rho}, \vec{g})$ for the same \vec{g}). Second, and this is more crucial, width can be replaced by A -width. This is also done by an easy adjustment of [ABRW00, Theorem 3.1] and Theorem 3.3. The only non-trivial thing to be remarked in this respect is that in the matrix $A|_{\rho}$ the expansion property holds for every set of rows $\{(i_1, \nu_1), \dots, (i_{\ell}, \nu_{\ell})\}$ whose projection $\{i_1, \dots, i_{\ell}\}$ onto the first coordinate

⁷Our notion of A -degree is slightly different from the one used in [ABRW00], mainly since the variables we consider here are automatically stratified w.r.t. rows $i \in [m]$

has size $\leq r$ (cf. the bound (27) in the proof of Claim 5.2), that is to every set of the form $\text{dom}(C)$ with $w_A(C) \leq r$.

The generalization of Corollary 3.4 obtained in this way gives the required contradiction with $\deg_A(P|\rho) \leq r/8$ and completes the proof of Claim 10.4. ■

Next, we, similarly to Definition 5.9, define a consistent random restriction ρ of the variables $\text{Vars}_{\leq}^{\text{Cycl}}(A)$. As before, ρ assigns x -variables completely at random (with $\mathbf{P}[\rho(x_j) = 0] = \mathbf{P}[\rho(x_j) = 1] = 1/4$). y -variables are assigned as follows. Pick a random subset of endpoints $\tilde{\mathbf{E}}^i$ by including there every endpoint with probability $(1/2d)$, independently of each other. $\tilde{\mathbf{E}}^i$ induces a partition of $J_i(A)$ into cyclic intervals $\tilde{\Delta}_1^i, \dots, \tilde{\Delta}_{\ell_i}^i$. We take all those $\tilde{\Delta}_\nu^i$ for which $|\tilde{\Delta}_\nu^i \setminus J_x(\rho)| < 2d$, and remove from $\tilde{\mathbf{E}}^i$ both endpoints of these intervals. Let \mathbf{E}^i be the resulting set of endpoints and $\Delta_1^i, \dots, \Delta_{\ell_i}^i$ be the corresponding partition into cyclic intervals. We pick at random Boolean values $\mathbf{b}_{i,1}, \dots, \mathbf{b}_{i,\ell_i}$ subjected to the only linear constraint $\bigoplus_{\nu=1}^{\ell_i} \mathbf{b}_{i,\nu} = b_i$, and assign in the natural way all those variables $y_\Delta^i \in \text{Vars}_{\leq}^{\text{Cycl}}(A)$ for which both endpoints of Δ are in \mathbf{E}^i . That is, if $\Delta = \bigcup_{\nu \in \Gamma} \Delta_\nu^i$ then $\rho(y_\Delta^i) \stackrel{\text{def}}{=} \bigoplus_{\nu \in \Gamma} \mathbf{b}_{i,\nu}$.

Clearly, ρ is consistent with probability 1.

Claim 10.5 $\mathbf{P}[\rho \text{ is sparse}] \geq 1/2$.

Proof. By exactly the same analysis as in the proof of Claim 5.10. ■

Lastly, we need the following simple version of Claim 10.3 (note the polynomial dependence on d – this is where our choice of the encoding will be used).

Claim 10.6 *Let C be any clause in the variables $\text{Vars}_{\leq}^{\text{Cycl}}(A)$ with $w_A(C) \leq r$. Then*

$$\mathbf{P}[C|\rho \neq 1] \leq \exp(-w_A(C)/d^{O(1)}).$$

Proof. We may assume w.l.o.g. that C consists either only of x -variables, or only of y -variables. In the first case the claim is obvious (since ρ acts on x -variables completely at random). In the second, for every $i \in \text{dom}(C)$ choose arbitrarily a cyclic interval Δ^i with $\frac{|J_i(A)|}{3} - 1 \leq |\Delta^i| \leq \frac{2|J_i(A)|}{3} + 1$ such that $y_{\Delta^i}^i \in \text{Vars}(C)$. Let j_1^i, j_2^i be its endpoints, and for $\alpha = 1, 2$, let $L_{i\alpha}, R_{i\alpha}$ be the two cyclic intervals, of length $5d$ each, with the endpoint j_α^i . Note that, due to the constraints on $|\Delta^i|$, we have $\text{Conv}(L_{i1}, R_{i1}) \cap \text{Conv}(L_{i2}, R_{i2}) = \emptyset$, and we use $L_{i\alpha}, R_{i\alpha}$ as we used protections in Section 5. The property of

weak regularity is immediate (C does not contain x -variables, and for every $i \in [m]$ contains at most one y_Δ^i), therefore we need neither the reduction operator R nor any analogue of Claim 5.12.

In the proof of Claim 5.13, the sub-protections $L'_{i\alpha}, R'_{i\alpha}$ will now have cardinalities $4d$ (as opposed to $2d$), and we can save by relaxing the requirement $J \cap P'_{i\alpha} \neq \emptyset$ in the definition of $(\mathbf{Y}^i)'$ to $|J \cap L'_{i\alpha}|, |J \cap R'_{i\alpha}| \leq 2d$. Then in (35) we have the better bound $\mathbf{P}[y_\Sigma^i \in (\mathbf{Y}^i)'] \geq \Omega(p_{i\Sigma}/d)$, and with the same argument we get that ρ restricted to $\text{Vars}(C)$ is $(r, \mu_{triv}, \Omega(1/d))$ -independent. Claim 10.6 follows. ■

Theorem 2.21 is immediately implied by Claims 10.4, 10.5 and 10.6.

11. Open problems

The central open problem in this area is obvious: construct pseudo-random generators that would be hard for as strong proof systems as possible and get as many pseudo-random output bits as possible. We complement this with several other (more minor) questions.

Does there exist a *function* pseudorandom generator of Nisan-Wigderson type which is hard for Resolution? Say, do there exist any A, \leq, b such that $m = 2^{n^\epsilon}$ and the minimal resolution refutation size of $\tau_{\leq}(A, b)$ is exponential in n ? The importance of this problem is, of course, greatly undermined by the iterability trick which allowed us to turn around it in Theorem 2.13. Still, this problem might be interesting in its own right. Also, it is not clear at the moment how general this trick will turn out, so after all a better understanding of the hardness of NW-generators themselves still may be useful in further research.

The next problem is of similar flavour. Even $n^{\Omega(\log n)}$ output bits we were able to get only for Nisan generators, that is, when the base functions are parity functions. Get more than quadratic number of output bits for a wider class of base functions. Suppose for example that $m = O(n^2)$, and the functions \vec{g} are picked at random. Is $\tau(\mathbf{A}_{mn}, \vec{g})$ hard for Resolution?

If the last problem is solved, then we might ask to extend the resulting bound to the system PCR over fields of characteristic 2 (this case is left completely open by the current paper).

Does PCR possess efficient proofs of $\mathbf{NP} \not\subseteq \mathbf{P/poly}$ when the latter class is defined by circuits over the standard basis $\{\neg, \wedge, \vee\}$? The natural attempt

to simply ignore the difficulty occurred in the proof of Theorem 2.20 leads to the system PCRes(2) which is a natural hybrid of PC and Res(2). Now, lower bounds for this system are known (see much more general result in [Kra97b]), but what we really need is a pseudo-random generator hard for it.

Last, but not the least, construct explicit lossless expanders (ideally, expanders with parameters close to those in Theorem 2.5). The importance of this last problem stretches of course well beyond proof complexity (see e.g. the impressive list of potential applications of expanders in [CRVW02]).

Acknowledgement

I am grateful to late Misha Alekhnovich, Jan Krajíček and Avi Wigderson for their interest, involvement and useful discussions at various stages of this ongoing project. I am very much indebted to both anonymous referees for the exceptionally thorough job they did on reading this technically involved text. The many remarks made by them have hopefully helped to significantly improve presentation in quite a few places.

References

- [ABE02] A. Atserias, M. L. Bonet, and J. L. Esteban. Lower bounds for the weak pigeonhole principle beyond resolution. *Information and Computation*, 176:136–152, 2002.
- [ABRW02] M. Alekhnovich, E. Ben-Sasson, A. Razborov, and A. Wigderson. Space complexity in propositional calculus. *SIAM Journal on Computing*, 31(4):1184–1211, 2002.
- [ABRW00] M. Alekhnovich, E. Ben-Sasson, A. Razborov, and A. Wigderson. Pseudorandom generators in propositional proof complexity. *SIAM Journal on Computing*, 34(1):67–88, 2004.
- [AR03] M. Alekhnovich and A. Razborov. Lower bounds for the polynomial calculus: non-binomial case. *Proceedings of the Steklov Institute of Mathematics*, 242:18–35, 2003.
- [AS08] N. Alon and J. Spencer. *The probabilistic method*. Wiley-Interscience, 3 edition, 2008.

- [BBC⁺01] R. Beals, H. Buhrman, R. Cleve, M. Mosca, and R. de Wolf. Quantum lower bounds by polynomials. *Journal of the ACM*, 48(4):778–797, 2001.
- [BGIP01] S. Buss, D. Grigoriev, R. Impagliazzo, and T. Pitassi. Linear gaps between degrees for the Polynomial Calculus modulo distinct primes. *Journal of Computer and System Sciences*, 62:267–289, 2001.
- [BSI10] E. Ben-Sasson and R. Impagliazzo. Random CNF’s are hard for the polynomial calculus. *Computational Complexity*, 19(4):501–519, 2010.
- [BP96] P. Beame and T. Pitassi. Simplified and improved resolution lower bounds. In *Proceedings of the 37th IEEE FOCS*, pages 274–282, 1996.
- [BP01] P. Beame and T. Pitassi. Propositional proof complexity: Past, present and future. In G. Paun, G. Rozenberg, and A. Salomaa, editors, *Current Trends in Theoretical Computer Science: Entering the 21st Century*, pages 42–70. World Scientific Publishing, 2001.
- [BPR97] M. Bonnet, T. Pitassi, and R. Raz. Lower bounds for cutting planes proofs with small coefficients. *Journal of Symbolic Logic*, 62(3):708–728, 1997.
- [BPR00] M. Bonnet, T. Pitassi, and R. Raz. On interpolation and automatization for Frege systems. *SIAM Journal on Computing*, 29(6):1939–1967, 2000.
- [BT88] S. Buss and G. Turán. Resolution proofs of generalized pigeon-hole principle. *Theoretical Computer Science*, 62:311–317, 1988.
- [CEI96] M. Clegg, J. Edmonds, and R. Impagliazzo. Using the Groebner basis algorithm to find proofs of unsatisfiability. In *Proceedings of the 28th ACM STOC*, pages 174–183, 1996.
- [CRVW02] M. Capalbo, O. Reingold, S. Vadhan, and A. Wigderson. Randomness conductors and constant-degree expansion beyond the

- degree/2 barrier. In *Proceedings of the 34th ACM Symposium on the Theory of Computing*, pages 659–668, 2002.
- [GGM86] O. Goldreich, S. Goldwasser, and S. Micali. How to construct random functions. *Journal of the ACM*, 33(4):792–807, 1986.
- [Gol11] O. Goldreich. Candidate one-way functions based on expander graphs. In *Studies in Complexity and Cryptography. Miscellanea on the Interplay between Randomness and Computation*, pages 76–87. Springer-Verlag, 2011.
- [IKW02] R. Impagliazzo, V. Kabanets, and A. Wigderson. In search of an easy witness: exponential time vs. probabilistic polynomial time. *Journal of Computer and System Sciences*, 65(4):672–694, 2002.
- [Jan90] S. Janson. Poisson approximation for large deviations. *Random Structures and Algorithms*, 1:221–230, 1990.
- [KP98] J. Krajíček and P. Pudlák. Some consequences of cryptographical conjectures for S_2^1 and EF . *Information and Computation*, 142:82–94, 1998.
- [Kra94] J. Krajíček. Lower bounds to the size of constant-depth propositional proofs. *Journal of Symbolic Logic*, 59(1):73–86, 1994.
- [Kra95] J. Krajíček. *Bounded arithmetic, propositional logic and complexity theory*. Cambridge University Press, 1995.
- [Kra97a] J. Krajíček. Interpolation theorems, lower bounds for proof systems and independence results for bounded arithmetic. *Journal of Symbolic Logic*, 62(2):457–486, 1997.
- [Kra97b] J. Krajíček. Lower bounds for a proof system with an exponential speed-up over constant-depth Frege systems and over polynomial calculus. In P. Ružička and I. Prívvara, editor, *Proceedings of the 22nd International Symposium on the Mathematical Foundations of Computer Science (Bratislava, August '97)*, *Lecture Notes in Computer Science* 1295, pages 85–90. Springer-Verlag, 1997.

- [Kra01a] J. Krajíček. On the weak pigeonhole principle. *Fundamenta Mathematicae*, 170(1-3):123–140, 2001.
- [Kra01b] J. Krajíček. Tautologies from pseudo-random generators. *Bulletin of Symbolic Logic*, 7(2):197–212, 2001.
- [Kra04] J. Krajíček. Dual weak pigeonhole principle, pseudo-surjective functions, and provability of circuit lower bounds. *Journal of Symbolic Logic*, 69(1):265–286, 2004.
- [Nis91] N. Nisan. CREW PRAMs and decision trees. *SIAM Journal on Computing*, 20(6):999–1007, 1991.
- [NW94] N. Nisan and A. Wigderson. Hardness vs. randomness. *Journal of Computer and System Sciences*, 49(2):149–167, October 1994.
- [Pin73] M. Pinsky. On the complexity of a concentrator. In *7th Annual Teletraffic Conference*, pages 318/1–318/4, Stockholm, 1973.
- [Pud97] P. Pudlák. Lower bounds for resolution and cutting planes proofs and monotone computations. *Journal of Symbolic Logic*, 62(3):981–998, 1997.
- [Pud98] P. Pudlák. The lengths of proofs. In S. Buss, editor, *Handbook of Proof Theory*, pages 547–637. Elsevier, 1998.
- [Raz04a] R. Raz. Resolution lower bounds for the weak pigeonhole principle. *Journal of the ACM*, 51(2):115–138, 2004.
- [Raz95a] A. Razborov. Bounded Arithmetic and lower bounds in Boolean complexity. In P. Clote and J. Remmel, editors, *Feasible Mathematics II. Progress in Computer Science and Applied Logic*, vol. 13, pages 344–386. Birkhäuser, 1995.
- [Raz95b] A. Razborov. Unprovability of lower bounds on the circuit size in certain fragments of bounded arithmetic. *Izvestiya of the RAN*, 59(1):201–224, 1995.
- [Raz96] A. Razborov. Lower bounds for propositional proofs and independence results in Bounded Arithmetic. In F. Meyer auf der

- Heide and B. Monien, editors, *Proceedings of the 23rd ICALP, Lecture Notes in Computer Science*, 1099, pages 48–62, New York/Berlin, 1996. Springer-Verlag.
- [Raz98] A. Razborov. Lower bounds for the polynomial calculus. *Computational Complexity*, 7:291–324, 1998.
- [Raz02a] A. Razborov. Proof complexity of pigeonhole principles. In *Proceedings of the 5th DLT, Lecture Notes in Computer Science*, 2295, pages 100–116, New York/Berlin, 2002. Springer-Verlag.
- [Raz04b] A. Razborov. Resolution lower bounds for perfect matching principles. *Journal of Computer and System Sciences*, 69(1):3–27, 2004.
- [RR97] A. Razborov and S. Rudich. Natural proofs. *Journal of Computer and System Sciences*, 55(1):24–35, 1997.
- [SBI04] N. Segerlind, S. Buss, and R. Impagliazzo. A switching lemma for small restrictions and lower bounds for k -DNF resolution. *SIAM Journal on Computing*, 33(5):1171–1200, 2004.
- [Urq95] A. Urquhart. The complexity of propositional proofs. *Bulletin of Symbolic Logic*, 1:425–467, 1995.
- [Yao82] A. Yao. Theory and applications of trapdoor functions. In *Proceedings of the 23rd IEEE FOCS*, pages 92–99, 1982.