

# Calcium-Activated Potassium Channels: Multiple Contributions to Neuronal Function

E. S. LOUISE FABER and PANKAJ SAH

Calcium-activated potassium channels are a large family of potassium channels that are found throughout the central nervous system and in many other cell types. These channels are activated by rises in cytosolic calcium largely in response to calcium influx via voltage-gated calcium channels that open during action potentials. Activation of these potassium channels is involved in the control of a number of physiological processes from the firing properties of neurons to the control of transmitter release. These channels form the target for modulation for a range of neurotransmitters and have been implicated in the pathogenesis of neurological and psychiatric disorders. Here the authors summarize the varieties of calcium-activated potassium channels present in central neurons and their defining molecular and biophysical properties. *NEUROSCIENTIST* 9(3):181–194, 2003. DOI: 10.1177/1073858403252673

**KEY WORDS** SK channel, AHP, sAHP, Accommodation

Calcium influx into cells has a variety of consequences that includes activation of second messenger systems, initiation of gene transcription, release of calcium from intracellular stores, and opening of calcium-dependent ion channels. Among the best studied of these events is the activation of calcium-dependent potassium channels. Initial indications that rises in cytosolic calcium could change plasmalemmal potassium permeability in response to changes in intracellular calcium concentrations were made by Gardos in red blood cells (Gardos 1958). The first identification of an ionic current activated by a rise in cytosolic calcium was made by Meech and Strumwasser who described a calcium-activated potassium current in snail neurons (Meech and Strumwasser 1970). It is now clear that such currents are present in a wide variety of cell types and are mediated by the opening of potassium-selective ion channels that are gated by a rise in intracellular calcium. These channels have a number of physiological roles ranging from secretion to the control of neuronal firing properties. In recent years, the functional roles of calcium-activated potassium channels have begun to be elucidated. In this review, we will discuss the molecular and physiological properties of calcium-activated potassium channels and the functions that have been attributed to these channels, including their role in memory and learning, aging and disease.

## Types of Calcium-Activated Potassium Channel

Three broad families of calcium-activated potassium channels have been identified, which can be separated on biophysical and pharmacological grounds. These

have been called BK, SK, and IK channels (Sah 1996; Vegara and others 1998).

### BK Channels

BK channels, also known as large conductance or maxiK channels (Marty 1981), were the first calcium-activated potassium channels to be identified. They were also the first type to be cloned from *Drosophila*, as the product of the *slowpoke* gene (Adelman and others 1992; Atkinson and others 1992). Subsequently, BK channels have been identified in a number of other species. These channels are highly potassium selective, have large single-channel conductances of 200 to 400 pS, and require both calcium and membrane depolarization for their activation (Marty 1981; McManus 1991). The calcium dependence of these channels is steeply dependent on the membrane potential (Cui and others 1997), the K<sub>d</sub> for calcium being in the micromolar range at resting membrane potentials (~–60 mV) but in the nanomolar range at depolarized potentials (+20 to +40 mV) (Latorre and others 1989). Interestingly, these channels can open in the absence of calcium, and it appears that the effects of calcium and membrane potential are independent processes, both of which enhance the open probability (Horrigan and Aldrich 2002). Although the voltage dependence of these channels is relatively weak when compared with purely voltage-dependent potassium channels, their voltage dependence allows these channels to operate as coincidence detectors and is of central importance to their physiological function.

BK channels open rapidly when calcium is raised on the cytosolic face and close rapidly upon removal of calcium. Several pharmacological blockers of BK channels are known. These include tetraethylammonium (TEA) in the low micromolar range (Blatz and Magleby 1987) and the two scorpion-derived peptides charybdotoxin and iberiotoxin (Galvez and others 1990). Recently, the

Division of Neuroscience, John Curtin School of Medical Research, Australian National University, Canberra, Australia

**Address correspondence to:** Pankaj Sah, Division of Neuroscience, John Curtin School of Medical Research, GPO Box 334, Canberra, ACT 2601, Australia (e-mail: pankaj.sah@anu.edu.au).

mycotoxins paxilline and penitrem A have also been shown to block BK channels (Knaus, McManus, and others 1994; Sanchez and McManus 1996; Strobaek and others 2000) (Fig. 2). Of these agents, iberiotoxin and paxilline are selective for BK channels whereas the other agents also block a number of other potassium channels. Some BK channels can also be specifically activated by dehydrosoyasaponin-1 (DHS-1), a compound isolated from a Ghanese medicinal herb (McManus and others 1993).

As with many other ion channels, two distinct subunits, named  $\alpha$  and  $\beta$ , have been associated with BK channels. The primary sequence of the pore-forming  $\alpha$  subunit shares significant homology with the pore regions of other members of the potassium channel family (Jan and Jan 1997). In agreement, the crystal structure of a homologous bacterial BK channel suggests that the ion-selective pores of BK channels share many properties with other potassium selective channels (Wei and others 1994; Meera and others 1997; Jiang and others 2002). However, unlike other voltage-dependent potassium channels, hydrophathy analysis suggests that the  $\alpha$  subunit has seven membrane-spanning domains (Fig. 1). In addition, an extra four hydrophobic segments, S7–S10, have been identified in the cytoplasmic carboxy terminal domain (Wei and others 1994; Meera and others 1997). The functions of these extra regions are not currently understood. Although BK channels are clearly activated by rises in cytoplasmic calcium, no defined calcium-binding domain has been identified. A series of negatively charged residues near the intracellular C-terminus referred to as the “calcium bowl” have been suggested to contain the calcium-binding site (Wei and others 1994; Schreiber and Salkoff 1997; Jiang and others 2002). However, removal of the C-terminus of BK channels does not disrupt calcium activation of BK channels, indicating that the calcium binds to a region outside of the calcium bowl (Piskorowski and Aldrich 2002). The exact location of the calcium-binding site on these channels still needs resolution.

Unlike other members of the potassium channel family, the pore-forming  $\alpha$  subunit of BK channels is coded for by a single gene—*slowpoke* (Adelman and others 1992). Diversity in the physiological properties of these channels is generated by alternative splicing of the *slowpoke* RNA (Atkinson and others 1992), by phosphorylation of the  $\alpha$  subunit (Toro and Stefani 1991; Reinhart and Levitan 1995; Sansom and others 1997), and by heteromeric assembly with the modulatory  $\beta$  subunit (Garcia-Calvo and others 1994; Meera and others 1997). The  $\beta$  subunit is composed of two putative transmembrane domains with a large extracellular loop (Fig. 1). Whereas only one  $\alpha$  subunit has been identified, three  $\beta$  subunits,  $\beta 1$ ,  $\beta 2/3$ , and  $\beta 4$ , have been cloned so far (Dworetzky and others 1994; Knaus, Folander, and others 1994; Tseng-Crank and others 1996; Brenner and others 2000; Meera and others 2000). The native channel is thought to comprise either an  $\alpha$  subunit alone as a tetramer (Shen and others 1994) or coassemble in combination with one to three  $\beta$  subunits (Adelman and

others 1992; Garcia-Calvo and others 1994; McManus and others 1995). The exact subunit composition of these channels in vivo, however, has not yet been determined.

Coassembly of the  $\alpha$  subunit with  $\beta$  subunits significantly modifies the pharmacology, voltage dependence, and kinetics of the assembled channel. Expression of the  $\beta$  subunit can shift the voltage dependence of activation to more negative membrane potentials and increases the calcium sensitivity (McManus and others 1995; Dworetzky and others 1996; Nimigeon and Magleby 1999; Wallner and others 1999). Channels containing the  $\alpha$  subunit in isolation or in combination with the  $\beta 1$  or  $\beta 4$  subunits produce sustained currents that do not inactivate (Wallner and others 1999; Meera and others 2000). In contrast, coexpression of the  $\alpha$  subunit with  $\beta 2/3$  subunits gives rise to channels that show rapid inactivation, thought to be mediated by blockade of the channel by the N terminus cytoplasmic “ball” (Wallner and others 1999; Xia and others 1999; Brenner and others 2000). Moreover, compared to when the  $\alpha$  subunit is expressed alone, coexpression of the  $\beta 2/3$  subunits with the  $\alpha$  subunit lowers the channel’s sensitivity to charybdotoxin and DHS-1 (Ding and others 1998; Wallner and others 1999; Xia and others 1999). Furthermore, recent electrophysiological studies have shown that coexpression of  $\alpha$  subunits with the neuronal  $\beta 4$  subunit confers insensitivity to iberiotoxin and charybdotoxin (Meera and others 2000) and slows the activation time of the channel.

The diversity in BK channel properties produced by posttranslational modification has clear physiological effects. For example, in the avian cochlea, differential splicing of BK channels is thought to confer the altered sensitivity of these channels to calcium and leads to the tonotopic organization of these cells (Navaratnam and others 1997; Rosenblatt and others 1997). Furthermore, the splice variants of BK channels, and thus their electrophysiological properties, are also controlled by hormonal status in adrenal chromaffin cells, indicating that the physiological properties of these channels are not fixed but can change with different physiological conditions (Xie and McCobb 1998; Lovell and McCobb 2001).

### SK Channels

Three types of SK channels have been cloned: SK1, SK2, and SK3 (Kohler and others 1996). SK channels have a single-channel conductance of 2–20 pS and are activated by rises in cytosolic calcium with half maximal activation in the 400–800 nM range (Blatz and Magleby 1986; Park 1994). Unlike BK channels, they are voltage insensitive and unaffected by low concentrations of TEA, charybdotoxin, or iberiotoxin. However, they are potently blocked by the bee venom apamin (Romey and others 1984; Blatz and Magleby 1986) (Fig. 3), tubocurarine, and quaternary salts of bicuculline (Johnson and Seutin 1997; Seutin and Johnson 1999). A new series of compounds that block SK channels include dequalinium

and a large set of related bis-quinolinium cyclophanes (Dunn 1994; Campos Rosa and others 2000; Chen and others 2000). Furthermore, 1-ethyl-2-benzimidazolone (EBIO) has been found to activate SK channels by altering their calcium sensitivity and open probability (Olesen and others 1994; Syme and others 2000; Pedarzani and others 2001) (Fig. 3). The activity of SK channels following action potentials leads to more long-lasting currents than those mediated by BK channels, owing to their higher affinity for calcium at hyperpolarized membrane potentials.

SK channels are widely expressed throughout the central nervous system but are also found in the periphery (Kohler and others 1996; Stocker and Pedarzani 2000; Sailer and others 2002). These channels are typical potassium channels with six putative transmembrane spanning regions (Fig. 1). Their primary structure shows approximately 60% sequence homology with each other, but SK channels only share homology with voltage-gated potassium channels in the pore region of the channel (Kohler and others 1996). Owing to the overall similarity in the transmembrane structure between SK channels and voltage-gated potassium channels (six putative transmembrane domains and cytoplasmic carboxy and amino terminals), it has been proposed that SK channels assemble as tetramers (Vegara and others 1998 #1556). However, direct evidence for this is lacking. Like BK channels, SK channels also do not have a clear calcium-binding domain. Instead, they are covalently linked to the calcium-binding protein calmodulin, and binding of calcium to calmodulin leads to a conformational change in the channel, which causes its opening (Xia and others 1998; Keen and others 1999; Schumacher and others 2001).

When expressed as homomultimers (Kohler and others 1996), SK channel subunits form ion channels that have functional characteristics typical of SK channels described in neurons, skeletal muscle (Romey and others 1984; Romey and Lazdunski 1984; Blatz and Magleby 1987), lymphocytes (Grissmer and others 1993), and adrenal chromaffin cells (Artalejo and others 1993; Park 1994). Thus, they respond rapidly to calcium applied to their cytoplasmic face and are voltage independent (Hirschberg and others 1998). Pharmacologically, SK2 and SK3 channels are very sensitive to blockade by apamin with  $IC_{50}$ s in the pM range (Kohler and others 1996; Ishii, Maylie, and others 1997). In contrast, homomeric hSK1 channels initially expressed in *Xenopus* oocytes were described as being apamin insensitive (Kohler and others 1996). However, when expressed in mammalian cell lines, they are blocked by apamin, suggesting that the sensitivity of these channels to apamin is partly dependent on the expression system used (Shah and Haylett 2000; Strobaek and others 2000). Interestingly, the SK1 gene has eight exons and undergoes extensive alternate splicing with over 30 predicted transcripts (Shmukler and others 2001). Of these, at least 20 transcripts have been detected in the brain (Shmukler and others 2001). Although the different predicted pro-

teins have yet to be detected, these findings suggest that there is a much larger diversity of SK channels than previously suspected.


### IK Channels

The third type of calcium-activated potassium channel has an intermediate single-channel conductance (20–100 pS) (Ishii, Silvia, and others 1997; Joiner and others 1997; Logsdon and others 1997). These channels have therefore been called intermediate conductance or IK channels and have only been identified in a few nonneuronal cell types, in particular epithelial and red blood cells (Gardos 1958; Ishii, Silvia, and others 1997). IK channels have been poorly studied largely owing to their sparse distribution. As for SK channels, IK channels are voltage insensitive but gated by rises in cytosolic calcium. Their pharmacological profile has been examined, and they have been shown to be sensitive to charybdotoxin, clotrimazole, and EBIO but insensitive to apamin and iberiotoxin (Ishii, Silvia, and others 1997; Joiner and others 1997; Logsdon and others 1997). Thus, these channels are clearly separable from both BK and SK channels. Macroscopic currents due to activation of IK channels have not been identified in central neurons; however, they have been suggested to underlie a slow afterhyperpolarization in myenteric neurons (Vogalis and others 2002).

### Physiological Roles

In many neurons, action potentials are followed by an afterhyperpolarization (AHP) that may last up to several seconds. Immediately following the action potential, there is a fast hyperpolarizing potential, the fast AHP, which typically lasts 1–10 ms and is due to the activation of voltage-gated potassium currents and/or a fast calcium-activated potassium current (Fig. 2). Following the fast AHP, there may be a prolonged hyperpolarization lasting between several hundred milliseconds and several seconds. In all cases, the slow component of the AHP results from activation of calcium-activated potassium conductances secondary to calcium entry during the action potential (Lancaster and Adams 1986; Storm 1987, 1990), typically through voltage-gated calcium channels (Lancaster and Adams 1986; Sah 1996; Marrion and Tavalin 1998). In some cell types, calcium release from intracellular stores also contributes to activation of the AHP (Sah and McLachlan 1991; Osmanovic and Schefner 1993; Yoshizaki and others 1995; Tanabe and others 1998). The slower calcium-activated potassium currents can be separated into two distinct components that are distinguishable on kinetic and pharmacological grounds, the medium AHP and the slow AHP (Sah 1996) (Fig. 2). The slower component of the AHP has two functions: it limits the firing frequency of the neuron and is responsible for generating the phenomenon of spike frequency adaptation. In some neurons, both the medium and the slow AHP are apparent, whereas in others only one or the other is present. It is now





**Fig. 1.** Proposed structure of large conductance (BK) channels and small conductance (SK) calcium-activated potassium channels. BK channels are formed from alpha and beta subunits in a heteromeric complex. The alpha subunit, a single gene product, has six transmembrane domains (S1-S6). S4 contains positively charged residues that form the proposed voltage sensor. An additional four hydrophobic domains have been identified in the extended C-terminal of the alpha subunit. Beta subunits have two transmembrane domains. SK channels are encoded by four genes SK1 to SK3 that assemble as tetramers to form a calcium-activated potassium channel. SK channels have six transmembrane domains (S1-S6) and a P domain between S5 and S6, which forms the pore region. Both the N terminus and the C terminus are cytoplasmic. The C-terminus contains a site for covalent linkage to calmodulin that binds calcium ions and is the calcium sensor for activation of these channels.


clear that different calcium-activated potassium channels contribute to the three phases of the AHP.

#### Fast AHP

In many neurons, the fast AHP is mediated by the macroscopic current generated by activation of a calcium-activated potassium current,  $I_c$  (Adams and others 1982; Pennefather and others 1985).  $I_c$  is blocked by low concentrations of TEA, iberiotoxin, and paxilline, indicating that the underlying channels are BK-type channels (Adams and others 1982; Lancaster and Nicoll 1987; Shao and others 1999). Due to their voltage dependence (Adams and others 1982), BK channels activate rapidly during the upstroke of the action potential and then close rapidly following return of the membrane potential to

negative values (Cui and others 1997), accounting for the rapid time course of the fast AHP. BK-type channels are modulated by protein kinase A (PKA)—both the probability of channel opening and the calcium and voltage sensitivity of the channel are modulated by phosphorylation. Levitan and collaborators have shown that the channel itself is the final target of phosphorylation by a closely associated protein kinase (Levitan 1994). Surprisingly, some types of BK-channel are up-regulated by PKA whereas other types are down-regulated. However, modulation of  $I_c$  has not been reported, and the functional effects of BK channel modulation in mammalian neurons, if any, are not known.

Consistent with its role in spike repolarization, blockade of  $I_c$  by TEA or paxilline slows action potential repolarization and reduces the fast AHP (Adams and others



**Fig. 2.** Action potentials are followed by fast and slow afterhyperpolarizations (AHP). *a*, A prolonged (600 ms) depolarizing current injection evokes a train of action potentials that cease firing after three to four spikes, a phenomenon known as spike frequency adaptation. The fast AHP (fAHP) immediately follows single-action potentials (*b*) and lasts up to tens of milliseconds. The fAHP results from the activity of several potassium channels that contribute to action potential repolarization. BK channels are activated during action potentials and contribute to the repolarization phase of the spike. Blockade of BK channels by paxilline broadens the action potential. The medium AHP (mAHP) is mediated by activation of the  $I_{AHP}$  current (*d*) and lasts tens to hundreds of milliseconds. The slow AHP (sAHP) is mediated by activation of  $sI_{AHP}$  and lasts several seconds. The slow AHP has a slow time to peak as compared to the medium AHP (*d*).


1982; Lancaster and Nicoll 1987; Storm 1987; Sah 1992) (Fig. 2). However, in other cell types, such as amygdala pyramidal neurons,  $I_C$  is active during spike repolarization but does not contribute to the fast AHP (Faber and Sah 2002). As discussed above, some BK channels show marked inactivation. This inactivation of  $I_C$  has been shown to contribute to spike broadening during repetitive firing in hippocampal pyramidal neurons (Shao and others 1999). Such broadening is likely to have consequences on transmitter release at axon terminals. Accordingly, BK channels have been shown to be located presynaptically on glutamatergic terminals in the hippocampus where they can control transmitter release under some conditions (Hu and others 2001; Gu and Storm 2002). Moreover, in olfactory bulb granule neurons, a coupling of BK channels with NMDA receptors has recently been demonstrated (Isaacson and Murphy 2001). This co-localization is thought to occur extrasynaptically because the BK channels can only be activated by tetanic stimulation. These latter studies suggest a role for BK channels in acting as a brake in times of excessive calcium influx.

#### Medium AHP

SK channels mediate the medium AHP and the underlying current,  $I_{AHP}$  (Sah 1996). The medium AHP is voltage

insensitive, is unaffected by BK channel blockers, but is blocked by apamin, indicating that it is due to the activation of SK-type channels (Pennefather and others 1985; Schwindt and others 1988; Sah and McLachlan 1991, 1992) (Figs. 2 and 3).  $I_{AHP}$  has a fast time to peak and decays with a time course dependent on the amount of calcium influx; as calcium load increases, peak amplitude increases and the current decay is slowed owing to saturation of local calcium buffers (Cassell and McLachlan 1987; Goh and Pennefather 1987; Sah 1992). Loading cells with low concentrations of the calcium chelator EGTA accelerates its decay (Sah 1992). These properties are due to the close anatomical proximity of N- or L-type voltage-gated calcium channels to SK channels (Viana and others 1993; Sah 1995a; Davies and others 1996; Marrion and Tavalin 1998; Bowden and others 2001).  $I_{AHP}$  was originally recorded from bullfrog sympathetic neurons (Pennefather and others 1985). A similar current has been termed  $gK_{Ca}$  in mammalian autonomic neurons (Cassell and McLachlan 1987; Sah and McLachlan 1991) and  $m-I_{AHP}$  in cortical neurons (Schwindt and others 1988).

Expression of SK1, SK2, and SK3 subunits as homo- or heteromultimers forms channels with properties similar to those of  $I_{AHP}$ . Thus, they are activated by rises in cytosolic calcium, are voltage independent and insensitive to low concentrations of TEA, and are blocked by



**Fig. 3.** Role of the apamin-sensitive medium AHP. *a*, Under voltage clamp, the currents underlying the medium and slow AHP are activated by a 100 ms depolarizing voltage step to 0 mV from a holding potential of  $-50$  mV. The fast-activating  $I_{\text{AHP}}$  current is blocked by apamin (100 nM), whereas the  $sI_{\text{AHP}}$  is not. Digital subtraction of the traces before and after apamin reveal  $I_{\text{AHP}}$ , which has rapid inactivation kinetics (trace on right). *b*, In cells that express only  $I_{\text{AHP}}$ , activation of this current is a major contributor to the firing frequency of these cells. Recording from a neuron in the dorsal motor nucleus of the vagus. *c*, In other neurons (pyramidal neuron in the lateral amygdala), the  $sI_{\text{AHP}}$  dominates spike-frequency adaptation and the apamin-sensitive current (shown in *a*) has only minor effects.

apamin (see below). The distribution of SK channel subunits closely mirrors the distribution of  $I_{\text{AHP}}$  type currents (Stocker and Pedarzani 2000; Sailer and others 2002) (see below). Within the central nervous system, these channels appear to largely assemble as homomultimers (Sailer and others 2002). Recently it has been determined that SK3 channels are the major contributors to the  $I_{\text{AHP}}$  in neurons of the rat dorsal motor nucleus of the

vagus (Pedarzani and others 2000), midbrain dopaminergic neurons (Wolfart and others 2001), and superior cervical ganglion neurons (Hosseini and others 2001).

In contrast to the fast AHP, the medium AHP does not contribute to action potential repolarization (Lancaster and Nicoll 1987; Storm 1990; Sah 1996) (Fig. 2). Cells that only express  $I_{\text{AHP}}$  fire tonically, and activation of  $I_{\text{AHP}}$  simply slows the maximal firing frequency (Engel and

others 1999) (Fig. 3). In contrast, in cells that express both the medium and slow AHP (see below), the medium AHP only controls the early interspike firing frequency (Stocker and others 1999; Pedarzani and others 2001; Stackman and others 2002). However, in amygdalar pyramidal neurons, although this current is also present, it does not appear to have any functional role in repetitive firing (Faber and Sah 2002) (Fig. 3). Recently, SK channels have been demonstrated to play a role in synaptic plasticity because blockade of SK channels with apamin shifted the threshold for induction of long-term potentiation and long-term depression in hippocampal neurons (see below) (Stackman and others 2002). Thus, the physiological role of this current in different cells needs to be independently evaluated.

### Slow AHP

The current that underlies the slow AHP was first described in neurons of the myenteric plexus (Hirst and others 1985). Although in some neurons the slow AHP has been described following a single action potential (Hirst and others 1985; Sah and McLachlan 1991), it is more commonly seen following a train (4–10) of spikes (Lancaster and Nicoll 1987; Schwindt and others 1988; Faber and Sah 2002). Following calcium influx, this current has a time to peak of hundreds of milliseconds then decays to baseline with a time constant of 1–2 seconds at 30°C (Figs. 2 and 4). To distinguish it from  $I_{\text{AHP}}$ , this current has been designated  $sI_{\text{AHP}}$  (Sah 1996). As with  $I_{\text{AHP}}$ ,  $sI_{\text{AHP}}$  requires a rise in cytosolic calcium for activation and is voltage insensitive.  $sI_{\text{AHP}}$  is not blocked by apamin or TEA. However,  $sI_{\text{AHP}}$  is modulated by a range of neurotransmitters, all of which block the current (Nicoll 1988) (Fig. 4). For the monoamines, the mechanism of this modulation is either by activation of protein kinase A (Pedarzani and Storm 1993; Torres and others 1996) or by inhibition of calcium-induced calcium release (Torres and others 1996). The second messenger pathways utilized by the other transmitters are as yet unknown, but acetylcholine has been suggested to act through activation of calcium calmodulin kinase II (Muller and others 1992; Pedarzani and Storm 1996). As with many other channels, the target for modulation appears to be the channel-opening probability, with both noradrenaline and acetylcholine lowering the probability for channel opening (Sah and Isaacson 1995).

The identity of the channels that underlie the  $sI_{\text{AHP}}$  remains a mystery. Direct recordings of these channels have not been successful; however, noise analysis indicates that their unitary conductance is in the range compatible with SK channels (Sah 1995b; Sah and Isaacson 1995; Valiante and others 1997). As SK2 and SK3 are both apamin sensitive, it seems unlikely that these subunits expressed alone could form  $sI_{\text{AHP}}$ . SK1 was initially reported to be apamin insensitive, and it has been suggested that these channels could underlie  $sI_{\text{AHP}}$  (Marrion and Tavalin 1998; Bowden and others 2001). However, this proposal has been challenged by others (Sah and Faber 2002).


Physiologically, the slow AHP is responsible for spike frequency adaptation in a number of neuronal cell types (Fig. 3). The presence of the slow AHP leads to a progressive slowing of the discharge frequency and eventual cessation of action potentials (Madison and Nicoll 1982; Sah 1996; Faber and Sah 2002). Modulation of the slow AHP by neurotransmitters (Nicoll 1988) dramatically changes the repetitive discharge properties of neurons that express  $sI_{\text{AHP}}$  (Fig. 4). Thus, the presence of  $sI_{\text{AHP}}$  allows for a greater level of control over the firing properties of neurons.

In the hippocampus, the slow AHP is also activated in response to synaptic activation of AMPA or NMDA receptors, and is capable of reducing the postsynaptic response to tetanic input (Sah and Bekkers 1996; Lancaster and others 2001). Furthermore, in lateral amygdala neurons, a slow calcium-activated potassium conductance has been shown to contribute to the inhibitory postsynaptic potential (IPSP) (Lang and Paré 1997; Danover and Pape 1998). In these studies, the AHP could be synaptically evoked by subthreshold stimuli. This suggests that the calcium influx through AMPA and NMDA receptors during the excitatory postsynaptic potential (EPSP) is sufficient to activate the channels and implies that the channels may be coupled to these receptors and located in dendritic spines in order to be activated by such elevations in intracellular calcium. Furthermore, in midbrain dopaminergic neurons, activation of metabotropic glutamate receptors has also been shown to activate apamin-sensitive SK channels by releasing calcium from intracellular stores (Fiorillo and Williams 1998). Thus, synaptic stimulation can activate SK channels by a variety of mechanisms.

### Distribution of SK Channels

The distribution of SK channels has been mapped using both in situ hybridization (Kohler and others 1996; Stocker and others 1999; Stocker and Pedarzani 2000) and immunohistochemistry using antibodies against each of the three subunits (Bowden and others 2001; Sailer and others 2002). These studies show that the three SK channel subunits have restricted distributions. SK1 and SK2 subunits are expressed at highest density in the hippocampus and cortex, whereas SK3 subunits are expressed at highest levels in regions such as the hypothalamus, thalamus, and midbrain (Stocker and Pedarzani 2000). Within the hippocampal formation, there are also differences in expression levels between area CA1, CA3, and dentate gyrus (Sailer and others 2002). These distinct distributions largely match the presence of the apamin-sensitive  $I_{\text{AHP}}$  in these regions (Sailer and others 2002). Interestingly, all three subunits are also found expressed in the neuropil throughout the cortex and hippocampus (Sailer and others 2002), indicating that as for BK channels (Hu and others 2001), SK channels may also play a role in the control of synaptic transmission. In CA1 hippocampal pyramidal neurons,  $sI_{\text{AHP}}$  channels have been suggested to be located in the proximal apical (Sah and Bekkers 1996; Lancaster and





**Fig. 4.** Role of the slow apamin-insensitive AHP. *a*, Under voltage clamp, the currents underlying the medium and slow AHP are activated by a 100 ms depolarizing voltage step to 0 mV from a holding potential of  $-50$  mV. Application of noradrenaline ( $10 \mu\text{M}$ ) blocks  $sI_{\text{AHP}}$ , leaving  $I_{\text{AHP}}$ . Digital subtraction of traces before and after application of noradrenaline reveals the kinetics of  $sI_{\text{AHP}}$ . This current has a slow rising phase and a slow decay. Physiologically, activation of the slow AHP contributes to spike-frequency adaptation (*b*).

others 2001) or basal (Bekkers 2000) dendrites where they play a role in the integration of synaptic inputs (see below).

## Functional Roles

### Learning and Memory

Due to its role in regulating both subthreshold and suprathreshold neuronal excitability, the AHP, and in particular the slow AHP, have been hypothesized to play a role in controlling the level of excitability of neurons and thus synaptic plasticity. As neuronal activity plays a key role in the processing of information within the central nervous system, it is perhaps not surprising that a number of recent studies have suggested that changes in both the medium and the slow AHP may be involved in aspects of learning and memory.

### Medium AHP

Blockade of SK channels with apamin has been shown to facilitate learning in a number of behavioral paradigms (van der Staay and others 1999). Almost all of the studies have focused on hippocampus-dependent learn-

ing paradigms, have administered apamin interperitoneally, and have found that the effects of apamin are on the acquisition of the learning task but not consolidation. Only two studies found an effect of apamin on the consolidation of learning (Messier and others 1991; Belcadi-Abbassi and Destrade 1995). Blockade of SK channels by systemic administration of apamin in rats enhanced learning in an object recognition task (Deschaux and others 1997). Furthermore, apamin reversed a spatial navigation deficit induced by medial septum and hippocampus lesions in mice in the Morris water maze spatial memory task (Ikonen and others 1998; Ikonen and Riekkinen 1999) and improved the performance of intact mice in this task (van der Staay and others 1999). In accordance with these studies, apamin also induced the expression of immediate early genes *c-fos* and *c-jun* in the hippocampus, which are thought to be the initial markers for memory formation (Heurteaux and others 1993). In addition to these hippocampal-dependent tasks, apamin also enhanced learning in an appetitively motivated bar-pressing response in mice (Messier and others 1991; Belcadi-Abbassi and Destrade 1995). Furthermore, Fournier and others (2001) applied apamin intracerebroventricularly in an olfactory discrimination learning task and found that



apamin enhanced learning but not consolidation of this task. Finally, an enhancement of the medium AHP recorded in hippocampal neurons was observed following eye blink conditioning in rabbits (de Jonge and others 1990).

In keeping with the view that the cellular substrate for learning and memory is synaptic plasticity (Moser and others 1998), SK channels have been demonstrated to facilitate the induction of plasticity in hippocampal neurons (Behnisch and Reymann 1998; Norris and others 1998; Stackman and others 2002). This effect of apamin was correlated with an increased rate of learning in the presence in hippocampal-dependent spatial (Morris water maze) and nonspatial (object recognition) memory tasks (Stackman and others 2002). As with previous findings, in these experiments, apamin accelerated the rate of learning but had no effect on long-term retention of the memory.

### *Slow AHP*

A number of studies have shown a reduction in the slow AHP in CA1 and CA3 hippocampal neurons in rabbits and rats in response to eye blink conditioning and Morris water maze tasks (Coulter and others 1989; de Jonge and others 1990; Moyer and others 1996; Thompson and others 1996; Oh and others 1999; Oh and others 2001). Both the duration and the amplitude of the slow AHP in hippocampal neurons is reduced following both hippocampus-dependent and -independent conditioning (Disterhoft and others 1986; Disterhoft and others 1988; Coulter and others 1989; Disterhoft and others 1996; Thompson and others 1996). This depression of the slow AHP is concurrent with a reduction in spike frequency adaptation (Coulter and others 1989; Moyer and others 1996; Thompson and others 1996). Similar reductions in the AHP were seen in the rat piriform cortex following conditioning using an olfactory discrimination task (Saar and others 1998). These studies suggest that reductions in the slow AHP are a general mechanism used to increase excitability during learning. As with the proposed role of the medium AHP in the induction but not consolidation of learning, changes in the slow AHP were transient and associated with the onset of learning but not the long-term retrieval (Moyer and others 1996; Thompson and others 1996; Saar and others 1998). This lends further support to the hypothesis that memories are learned in the hippocampus but stored elsewhere, possibly in the cortex (Moyer and others 1996; Thompson and others 1996).

In addition to the above studies, *in vitro* studies support the notion that the amplitude of the slow AHP is inversely related to the ability to learn, or for synapses to undergo plasticity. Thus, the dendritically located slow AHP in hippocampal CA1 pyramidal neurons is able to shunt synaptic potentials, thereby raising the threshold for synaptic plasticity (Sah and Bekkers 1996). Furthermore, there is a direct correlation between the degree of slow AHP suppression and the extent of induc-

tion of plasticity recorded in hippocampal neurons (Cohen and others 1999).

### *Aging*

During aging, the threshold for synaptic plasticity is increased (Deupree and others 1993; Moore and others 1993; Disterhoft and others 1996; Rosenzweig and others 1997) and the performance of aged animals and humans in learning is impaired (Thompson and others 1996; Knuttinen and others 2001; Moyer and others 2000). Impairments in trace eye blink conditioning and the Morris water maze, both of which are hippocampus dependent, have been observed in aged rabbits, rats, and humans (Deyo and others 1989; Gallagher and others 1993; Rapp and Gallagher 1996; Thompson and others 1996; Knuttinen and others 2001; Wu and others 2002). Conversely, an enhancement of the slow AHP recorded in hippocampal neurons with age has been demonstrated (for review, see Pitler and Landfield 1990; Moyer and Disterhoft 1994; Campbell and others 1996; Disterhoft and others 1996; Thibault and Landfield 1996; Power and others 2002; Wu and others 2002). Furthermore, the slow AHP is reduced during learning in aged animals (Landfield and Pitler 1984; Moyer and others 2000). The enlarged slow AHP has been attributed, at least in part, to an increased influx of calcium through L-type voltage-gated calcium channels, allowing a greater activation of the underlying calcium-activated potassium channels (Landfield and Pitler 1984; Moyer and others 1992; Campbell and others 1996; Thibault and Landfield 1996; Norris and others 1998; Power and others 2002). Because the slow AHP is modulated by a number of neurotransmitters (see above), it has been proposed that alterations in these neurotransmitter systems may also underlie the changes in the slow AHP (Wu and others 2002). In accordance with this, drugs that depress the slow AHP, such as calcium channel blockers or cholinergic agonists, have been shown to be effective in improving learning in aged animals (Deyo and others 1989; Moyer and others 1992; Kronforst-Collins and others 1997; Oh and others 1999; Weiss and others 2000).

The consequence of the enhanced slow AHP is a decrease in excitability, with neurons from aged animals showing greater spike frequency adaptation compared with those from young animals (Moyer and others 1992; Oh and others 1999; Moyer and others 2000). These findings, correlated with the role of the slow AHP in learning (see above), suggest that the reduction in excitability of hippocampal neurons caused by the enhancement of the slow AHP is partly responsible for the reduction in age-related learning deficits. Enhancement of the medium AHP also appears to contribute to a reduced ability of hippocampal neurons from aged animals to undergo synaptic plasticity (Norris and others 1998), albeit to a lesser extent than the slow AHP (Power and others 2002; Wu and others 2002). Such deficiencies in learning and memory with aging may contribute to the symptoms of neurodegenerative diseases such as Alzheimer's disease (see below).

## Disease

Recent reports have suggested that the medium and slow AHP may be important in Alzheimer's disease. An autoradiographic study showed that apamin binding sites are reduced in the hippocampus of brains from patients with Alzheimer's disease (Ikeda and others 1991). In addition, galantamine, an acetylcholinesterase inhibitor, has been shown to alleviate cognitive deficits in Alzheimer's disease patients, and a proposed mechanism of action is at least partly through inhibition of the slow AHP by an enhanced activation of muscarinic acetylcholine receptors (Wilkinson and Murray 2001; Wu and others 2002).

The slow AHP has been proposed to reduce excitability to prevent the onset of epilepsy in CA1 hippocampal neurons (Martin and others 2001). In this model, during the onset of the ictal state, enhancement of glutamate release by spontaneous EPSPs activates metabotropic glutamate receptors to depress the slow AHP. As the slow AHP plays a key role in neuronal excitability, its depression leads to increased excitability. Similarly, a synaptically evoked calcium-activated potassium conductance has been suggested to play a role in preventing the onset of epileptiform discharges in the amygdala (Danover and Pape 1998), a site of generation of temporal lobe epilepsy. Thus, calcium influx that occurs during strong excitation activates the slow AHP, which may act as a brake on the system to prevent overexcitation and the generation of epileptic-like activity (Danover and Pape 1998). This is in accordance with an *in vivo* study showing a reduced AHP during an epileptic state (Matsumoto and Ajmone-Marsan 1964). However, when measured directly in the rat piriform cortex, kindling did not cause any change in the AHP (Saar and others 1998). Thus, the precise contribution of the AHP to the prevention of epilepsy remains to be clarified.

A variety of mendelian genetic diseases are now known to be due to mutations in ion channels (Ashcroft 2000). The presence of polyglutamine repeat expansions is a newly recognized mechanism of disease that has been linked to a number of rare, late onset neurological diseases such as Huntington's disease and fragile X syndrome (Ashley and Warren 1995). Recently, a small conductance calcium-activated potassium channel gene (hSKCa3 or KCNN3), which codes for a channel with properties identical to SK channels, has been identified that has CAG repeats in its coding region (Chandy and others 1998). In genetic screens, the presence of trinucleotide repeats has been shown to be associated with schizophrenia and bipolar illness, suggestive of a possible link between SK channel function and these disorders (Gargus and others 1998; Dror and others 1999; Ritsner and others 2002).

## Conclusions

Calcium-activated potassium channels form a diverse family of potassium channels that play important roles in the control of neuronal excitability and transmitter release. Some of these channels are the target for modu-

lation of a number of transmitters. Of these channels, much is understood of the molecular structure and biophysical properties of one family of these channels, the BK channels. In contrast, the study of small conductance, SK channels is just beginning to reach a similar stage. In particular, little is known of the nature of the channels that underlie the macroscopic calcium-activated potassium currents. Mutations of these channels may underlie some neurological and psychiatric diseases. With the mapping of the human genome nearing completion, our understanding of these channels and their involvement in disease will no doubt improve in the future.

## References

- Adams PR, Constanti A, Brown DA, Clark RB. 1982. Intracellular  $Ca^{2+}$  activates a fast voltage-sensitive  $K^{+}$  current in vertebrate sensory neurones. *Nature* 296:746–9.
- Adelman JP, Shen K-Z, Kavanaugh MP, Warren RA, Wu Y-N, Lagrutta A, and others. 1992. Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron* 9:209–16.
- Artalejo AR, Garcia AG, Neher E. 1993. Small-conductance  $Ca^{2+}$ -activated  $K^{+}$  channels in bovine chromaffin cells. *Pflugers Arch* 423:97–103.
- Ashcroft FM. 2000. Ion channels and disease: channelopathies. New York: Academic Press.
- Ashley CT Jr, Warren ST. 1995. Trinucleotide repeat expansion and human disease. *Annu Rev Genet* 29:703–28.
- Atkinson NS, Robertson GA, Ganetzky B. 1992. A component of calcium-activated potassium channels encoded by the *drosophila slo* locus. *Science* 253:551–5.
- Behnisch T, Reymann KG. 1998. Inhibition of apamin-sensitive calcium dependent potassium channels facilitate the induction of long-term potentiation in the CA1 region of rat hippocampus *in vitro*. *Neurosci Lett* 253:91–4.
- Bekkers JM. 2000. Distribution of slow AHP channels on hippocampal CA1 pyramidal neurons. *J Neurophysiol* 83:1756–9.
- Belcadi-Abbassi W, Destradre C. 1995. Post-test apamin injection suppresses a Kamin-like effect following a learning session in mice. *Neuroreport* 6:1293–6.
- Blatz AL, Magleby KL. 1986. Single apamin-blocked Ca-activated  $K^{+}$  channels of small conductance in cultured rat skeletal muscle. *Nature* 323:718–20.
- Blatz AL, Magleby KL. 1987. Calcium-activated potassium channels. *Trends Neurosci* 10:463–7.
- Bowden SE, Fletcher S, Loane DJ, Marrion NV. 2001. Somatic colocalization of rat SK1 and D class ( $Ca(v)1.2$ ) L-type calcium channels in rat CA1 hippocampal pyramidal neurons. *J Neurosci* 21:RC175.
- Brenner R, Jegla TJ, Wickenden A, Liu Y, Aldrich RW. 2000. Cloning and functional characterisation of novel large conductance calcium-activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. *J Biol Chem* 275:6453–61.
- Campbell LW, Hao SY, Thibault O, Blalock EM, Landfield PW. 1996. Aging changes in voltage-gated calcium currents in hippocampal CA1 neurons. *J Neurosci* 16:6286–95.
- Campos Rosa J, Galanakis D, Piergentili A, Bhandari K, Ganellin CR, Dunn PM, and others. 2000. Synthesis, molecular modeling, and pharmacological testing of bis-quinolinium cyclophanes: potent, non-peptidic blockers of the apamin-sensitive  $Ca^{2+}$ -activated  $K^{+}$  channel. *J Med Chem* 43:420–31.
- Cassell JF, McLachlan EM. 1987. Two calcium-activated potassium conductances in a subpopulation of coeliac neurones of guinea-pig and rabbit. *J Physiol* 394:331–49.
- Chandy KG, Fantino E, Wittekindt O, Kalman K, Tong LL, Ho TH, and others. 1998. Isolation of a novel potassium channel gene hSKCa3 containing a polymorphic CAG repeat: a candidate for schizophrenia and bipolar disorder? *Mol Psychiatry* 3:32–7.

- Chen J-Q, Galanakis D, Ganellin CR, Dunn PM, Jenkinson DH. 2000. bis-Quinolium Cyclophanes: 8,14-diaza-1,7(1,4)-diquinoliny-clotradecaphane (UCL 1848), a highly potent and selective, non-peptidic blocker of the apamin-sensitive  $Ca^{2+}$ -activated  $K^{+}$  channel. *J Med Chem* 43:3478–81.
- Cohen AS, Coussens CM, Raymond CR, Abraham WC. 1999. Long-lasting increase in cellular excitability associated with the priming of LTP induction in rat hippocampus. *J Neurophysiol* 82:3139–48.
- Coulter DA, Lo Turco JJ, Kubota M, Disterhoft JF, Moore JW, Alkon DL. 1989. Classical conditioning reduces amplitude and duration of calcium-dependent afterhyperpolarization in rabbit hippocampal pyramidal cells. *J Neurophysiol* 61:971–81.
- Cui J, Cox DH, Aldrich RW. 1997. Intrinsic voltage dependence and  $Ca^{2+}$  regulation of mslo large conductance  $Ca$ -activated  $K^{+}$  channels. *J Gen Physiol* 109:647–73.
- Danober L, Pape H-C. 1998. Mechanisms and functional significance of a slow inhibitory potential in neurons of the lateral amygdala. *Eur J Neurosci* 10:853–67.
- Davies PJ, Ireland DR, McLachlan EM. 1996. Sources of  $Ca^{2+}$  for different  $Ca^{2+}$ -activated  $K^{+}$  conductances in neurones of the rat superior cervical ganglion. *J Physiol* 495:353–66.
- de Jonge MC, Black J, Deyo RA, Disterhoft JF. 1990. Learning-induced afterhyperpolarization reductions in hippocampus are specific for cell type and potassium conductance. *Exp Brain Res* 80:456–62.
- Deschaux O, Bizot JC, Goyffon M. 1997. Apamin improves learning in an object recognition task in rats. *Neurosci Lett* 222:159–62.
- Deupree DL, Bradley J, Turner DA. 1993. Age-related alterations in potentiation in the CA1 region in F344 rats. *Neurobiol Aging* 14:249–58.
- Deyo RA, Straube KT, Disterhoft JF. 1989. Nimodipine facilitates associative learning in aging rabbits. *Science* 243:809–11.
- Ding JP, Li ZW, Lingle CJ. 1998. Inactivating BK channels in rat chromaffin cells may arise from heteromultimeric assembly of distinct inactivation-competent and noninactivating subunits. *Biophys J* 74:268–89.
- Disterhoft JF, Coulter DA, Alkon DL. 1986. Conditioning-specific membrane changes of rabbit hippocampal neurons measured in vitro. *Proc Natl Acad Sci U S A* 83:2733–7.
- Disterhoft JF, Golden DT, Read HL, Coulter DA, Alkon DL. 1988. AHP reductions in rabbit hippocampal neurons during conditioning correlate with acquisition of the learned response. *Brain Res* 462:118–25.
- Disterhoft JF, Thompson LT, Moyer JR Jr, Mogul DJ. 1996. Calcium-dependent afterhyperpolarization and learning in young and aging hippocampus. *Life Sci* 59:413–20.
- Dror V, Shamir E, Ghanshani S, Kimhi R, Swartz M, Barak Y, and others. 1999. hKCa3/KCNN3 potassium channel gene: association of longer CAG repeats with schizophrenia in Israeli Ashkenazi Jews, expression in human tissues and localization to chromosome 1q21. *Mol Psychiatry* 4:254–60.
- Dunn PM. 1994. Dequalinium, a selective blocker of the slow afterhyperpolarization in rat sympathetic neurones in culture. *Eur J Pharmacol* 252:189–94.
- Dworetzky SI, Boissard CG, Lum-Ragan M, McKay C, Post-Munson DJ, Tronjnick JT, and others. 1996. Phenotypic alteration of a human BK (hsl) channels by hsl $\beta$  subunit coexpression: changes in blocker sensitivity, activation/relaxation and inactivation kinetics, and protein kinase A modulation. *J Neurosci* 16:4543–50.
- Dworetzky SI, Tronjnick JT, Gribkoff VK. 1994. Cloning and expression of a human large-conductance calcium-activated potassium channel. *Brain Res Mol Brain Res* 27:189–93.
- Engel J, Schultens HA, Schild D. 1999. Small conductance potassium channels cause an activity dependent spike frequency adaptation and make the transfer function of neurons logarithmic. *Biophys J* 76:1310–19.
- Faber ESL, Sah P. 2002. Physiological role of calcium-activated potassium currents in the rat lateral amygdala. *J Neurosci* 22:1618–28.
- Fiorillo CD, Williams JT. 1998. Glutamate mediates an inhibitory postsynaptic potential in dopamine neurons. *Nature* 394:78–82.
- Fournier C, Kourrich S, Soumireu-Mourat B, Mourre C. 2001. Apamin improves reference memory but not procedural memory in rats by blocking small conductance  $Ca^{2+}$ -activated  $K^{+}$  channels in an olfactory discrimination task. *Behav Brain Res* 121:81–93.
- Gallagher M, Burwell R, Burchinal M. 1993. Severity of spatial learning impairment in aging: development of a learning index for performance in the Morris water maze. *Behav Neurosci* 107:618–26.
- Galvez A, Gimenez-Gallego G, Reuben J, Roy-Contancin L, Feigenbaum P, Kaczorowski G, and others. 1990. Purification and characterization of a unique, potent, peptidyl probe for the high conductance calcium-activated potassium channel from venom of the scorpion *Buthus tamulus*. *J Biol Chem* 265:11083–90.
- Garcia-Calvo M, Knaus HG, McManus OB, Giangiacomo KM, Kaczorowski GJ, Garcia ML, and others. 1994. Purification and reconstruction of the high-conductance, calcium-activated potassium channel from tracheal smooth muscle. *J Biol Chem* 269:676–82.
- Gardos G. 1958. The function of calcium in the potassium permeability of human erythrocytes. *Biochem Biophys Acta* 30:653–4.
- Gargus JJ, Fantino E, Gutman GA. 1998. A piece in the puzzle: an ion channel candidate gene for schizophrenia. *Mol Med Today* 4:518–24.
- Goh JW, Pennefather PS. 1987. Pharmacological and physiological properties of the after-hyperpolarization current of bullfrog ganglion neurons. *J Physiol* 394:315–30.
- Grissmer S, Nguyen AN, Cahalan MD. 1993. Calcium-activated potassium channels in resting and activated human T lymphocytes. Expression levels, calcium dependence, ion selectivity, and pharmacology. *J Gen Physiol* 102:601–30.
- Gu N, Storm JF. 2002. Regulation of glutamate release by presynaptic calcium-activated potassium channels (BK channels) is enhanced under hypoxic conditions in rat hippocampus (CA1). *Soc Neurosci Abs* 340.15.
- Heurteaux C, Messier C, Destrade C, Lazdunski M. 1993. Memory processing and apamin induce immediate early gene expression in mouse brain. *Brain Res Mol Brain Res* 18:17–22.
- Hirschberg B, Maylie J, Adelman JP, Marrion NV. 1998. Gating of recombinant small-conductance  $Ca$ -activated  $K^{+}$  channels by calcium. *J Gen Physiol* 111:565–81.
- Hirst GDS, Johnson SM, van Helden DF. 1985. The slow calcium-dependent potassium current in a myenteric neurone of the guinea-pig ileum. *J Physiol* 361:315–37.
- Horrigan FT, Aldrich RW. 2002. Coupling between voltage sensor activation,  $Ca^{2+}$  binding and channel opening in large conductance (BK) potassium channels. *J Gen Physiol* 120:267–305.
- Hosseini R, Benton DCH, Dunn PM, Jenkinson DH, Moss GWJ. 2001. SK3 is an important component of  $K^{+}$  channels mediating the after-hyperpolarization in cultured rat SCG neurones. *J Physiol* 535:323–34.
- Hu H, Shao L-R, Chavoshy S, Gu N, Trieb M, Behrens R, and others. 2001. Presynaptic  $Ca^{2+}$ -activated  $K^{+}$  channels in glutamatergic hippocampal terminals and their role in spike repolarization and regulation of transmitter release. *J Neurosci* 21:9585–97.
- Ikeda M, Dewar D, McCulloch J. 1991. Selective reduction of [ $^{125}I$ ]apamin binding sites in Alzheimer hippocampus: a quantitative autoradiographic study. *Brain Res* 567:51–6.
- Ikonen S, Riekkinen P Jr. 1999. Effects of apamin on memory processing of hippocampal-lesioned mice. *Eur J Pharmacol* 382:151–6.
- Ikonen S, Schmidt B, Riekkinen P Jr. 1998. Apamin improves spatial navigation in medial septal-lesioned mice. *Eur J Pharmacol* 347:13–21.
- Isaacson JS, Murphy GJ. 2001. Glutamate-mediated extrasynaptic inhibition: direct coupling of NMDA receptors to  $Ca^{2+}$ -activated  $K^{+}$  channels. *Neuron* 31:1027–34.
- Ishii TM, Maylie J, Adelman JP. 1997. Determinants of apamin and d-tubocurarine block in SK potassium channels. *J Biol Chem* 272:23195–200.
- Ishii TM, Silvia C, Hirschberg B, Bond CT, Adelman JP, Maylie J. 1997. A human intermediate conductance calcium-activated potassium channel. *Proc Natl Acad Sci U S A* 94:11651–6.
- Jan LY, Jan YN. 1997. Cloned potassium channels from eukaryotes and prokaryotes. *Annu Rev Neurosci* 20:91–123.
- Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R. 2002. Crystal structure and mechanism of a calcium-gated potassium channel. *Nature* 417:515–22.



- Johnson SW, Seutin V. 1997. Bicuculline methiodide potentiates NMDA-dependent burst firing in rat dopamine neurons by blocking apamin-sensitive  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  currents. *Neurosci Lett* 231:13–16.
- Joiner WJ, Wang L-Y, Tang MD, Kaczmarek LK. 1997. hSK4, a member of a novel subfamily of calcium-activated potassium channels. *Proc Natl Acad Sci U S A* 94:11013–18.
- Keen JE, Khawaled R, Farrens DL, Neelands T, Rivard A, Bond CT, and others. 1999. Domains responsible for constitutive and  $\text{Ca}^{2+}$ -dependent interactions between calmodulin and small conductance  $\text{Ca}^{2+}$ -activated potassium channels. *J Neurosci* 19:8830–8.
- Knaus HG, Folander M, Garcia-Calvo ML, Garcia GJ, Kaczorowski GJ, Smith M, and others. 1994. Primary sequence and immunological characterisation of beta-subunit of high conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel from smooth muscle. *J Biol Chem* 269:17274–8.
- Knaus HG, McManus OB, Lee SH, Schmalhofer WA, Garcia-Calvo M, Helms LM, and others. 1994. Tremorgenic indole alkaloids potentially inhibit smooth muscle high-conductance calcium-activated potassium channels. *Biochemistry* 33:5819–28.
- Knutninen MG, Power JM, Preston AR, Disterhoft JF. 2001. Awareness in classical differential eyeblink conditioning in young and aging humans. *Behav Neurosci* 115:747–57.
- Kohler M, Hirschberg B, Bond CT, Kinzie JM, Marrion NV, Maylie J, and others. 1996. Small-conductance, calcium-activated potassium channels from mammalian brain. *Science* 273:1709–14.
- Kronforst-Collins MA, Moriearty PL, Schmidt B, Disterhoft JF. 1997. Metrifonate improves associative learning and retention in aging rabbits. *Behav Neurosci* 111:1031–40.
- Lancaster B, Adams PR. 1986. Calcium-dependent current generating the afterhyperpolarization of hippocampal neurons. *J Neurophysiol* 55:1268–82.
- Lancaster B, Hu H, Ramakers GM, Storm JF. 2001. Interaction between synaptic excitation and slow afterhyperpolarization current in rat hippocampal pyramidal cells. *J Physiol* 536:809–23.
- Lancaster B, Nicoll RA. 1987. Properties of two calcium-activated hyperpolarizations in rat hippocampal neurones. *J Physiol* 389:187–204.
- Landfield PW, Pitler TA. 1984. Prolonged  $\text{Ca}^{2+}$ -dependent afterhyperpolarizations in hippocampal neurons of aged rats. *Science* 226:1089–92.
- Lang EJ, Paré D. 1997. Synaptic and synaptically activated intrinsic conductances underlie inhibitory potentials in cat lateral amygdaloid projection neurons in vivo. *J Neurophysiol* 77:353–63.
- Latorre R, Oberhauser A, Labarca P, Alvarez O. 1989. Varieties of calcium-activated potassium channels. *Ann Rev Physiol* 51:385–99.
- Levitan IB. 1994. Modulation of ion channels by protein phosphorylation and dephosphorylation. *Annu Rev Physiol* 56:193–212.
- Logsdon NJ, Kang J, Togo JA, Christian EP, Aiyar J. 1997. A novel gene, hKCa4, encodes the calcium-activated potassium channel in human T lymphocytes. *J Biol Chem* 272:32723–6.
- Lovell PV, McCobb DP. 2001. Pituitary control of BK potassium channel function and intrinsic firing properties of adrenal chromaffin cells. *J Neurosci* 21:3429–42.
- Madison DV, Nicoll RA. 1982. Noradrenaline blocks accommodation of pyramidal cell discharge in the hippocampus. *Nature* 299:636–8.
- Marrion NV, Tavalin SJ. 1998. Selective activation of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels by co-localised  $\text{Ca}^{2+}$  channels in hippocampal neurons. *Nature* 395:900–5.
- Martin ED, Araque A, Buno W. 2001. Synaptic regulation of the slow  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  current in hippocampal CA1 pyramidal neurons: implication in epileptogenesis. *J Neurophysiol* 86:2878–86.
- Marty A. 1981. Ca-dependent K channels with large unitary conductance in chromaffin cell membranes. *Nature* 291:497–500.
- Matsumoto H, Ajmone-Marsan C. 1964. Cortical cellular phenomena in experimental epilepsy: ictal manifestations. *Exp Neurol* 9:305–26.
- McManus OB. 1991. Calcium-activated potassium channels: regulation by calcium. *J Bioenerg Biomembr* 23:537–60.
- McManus OB, Harris GH, Giangiacoio KM, Feigenbaum P, Reuben JP, Addy ME, and others. 1993. An activator of calcium-dependent potassium channels isolated from a medicinal herb. *Biochemistry* 32:128–33.
- McManus OB, Helms LM, Pallanck L, Ganetzky B, Swanson R, Leonard RJ. 1995. Functional role of the beta subunit of high conductance calcium-activated potassium channels. *Neuron* 14:645–50.
- Meech RW, Strumwasser F. 1970. Intracellular calcium injection activates potassium conductance in aplysia nerve cells. *Fed Proc* 29:834.
- Meera P, Wallner M, Song M, Toro L. 1997. Large conductance voltage- and calcium-dependent  $\text{K}^+$  channel, a distinct member of voltage-dependent ion channels with seven N-terminus transmembrane segments (S0-S6), and extracellular N terminus, and an intracellular (S9-S10) C terminus. *Proc Natl Acad Sci U S A* 94:14066–71.
- Meera P, Wallner M, Toro L. 2000. A neuronal beta subunit (KCNCB4) makes the large conductance, voltage- and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel resistant to charybdotoxin and iberiotoxin. *Proc Natl Acad Sci U S A* 97:5562–7.
- Messier C, Mourre C, Bontempi B, Sif J, Lazdunski M, Destrade C. 1991. Effect of apamin, a toxin that inhibits  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels, on learning and memory processes. *Brain Res* 551:322–6.
- Moore CI, Browning MD, Rose GM. 1993. Hippocampal plasticity induced by primed burst, but not long-term potentiation, stimulation is impaired in area CA1 of aged Fischer 344 rats. *Hippocampus* 3:57–66.
- Moser EI, Krobot KA, Moser MB, Morris RG. 1998. Impaired spatial learning after saturation of long-term potentiation. *Science* 281:2038–42.
- Moyer JR Jr, Disterhoft JF. 1994. Nimodipine decreases calcium action potentials in rabbit hippocampal CA1 neurons in an age-dependent and concentration-dependent manner. *Hippocampus* 4:11–17.
- Moyer JR Jr, Power JM, Thompson LT, Disterhoft JF. 2000. Increased excitability of aged rabbit CA1 neurons after trace eyeblink conditioning. *J Neurosci* 20:5476–82.
- Moyer JR Jr, Thompson LT, Black JP, Disterhoft JF. 1992. Nimodipine increases excitability of rabbit CA1 pyramidal neurons in an age- and concentration-dependent manner. *J Neurophysiol* 68:2100–9.
- Moyer JR Jr, Thompson LT, Disterhoft JF. 1996. Trace eyeblink conditioning increases CA1 excitability in a transient and learning-specific manner. *J Neurosci* 16:5536–46.
- Muller W, Petrozzino JJ, Griffith LC, Danho W, Connor JA. 1992. Specific involvement of  $\text{Ca}^{2+}$ -calmodulin kinase II in cholinergic modulation of neuronal responsiveness. *J Neurophysiol* 68:2264–8.
- Navaratnam DS, Bell TJ, Tu TD, Cohen EL, Oberholtzer JC. 1997. Differential distribution of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel splice variants among hair cells along the tonotopic axis of the chick cochlea. *Neuron* 19:1077–85.
- Nicoll RA. 1988. The coupling of neurotransmitter receptors to ion channels in the brain. *Science* 241:545–51.
- Nimigeon CM, Magleby KL. 1999. The beta subunit increases the  $\text{Ca}^{2+}$  sensitivity of large conductance  $\text{Ca}^{2+}$ -activated potassium channels by retaining the gating the bursting states. *J Gen Physiol* 113:425–39.
- Norris CM, Halpain S, Foster TC. 1998. Reversal of age-related alterations in synaptic plasticity by blockade of L-type  $\text{Ca}^{2+}$  channels. *J Neurosci* 18:3171–9.
- Oh MM, Gamelli AE, Wu WW, Sametsky E, Disterhoft JF. 2001. Morris water maze learning enhances neuronal excitability of CA1 hippocampal pyramidal neurons in rats. *Soc Neurosci Abs* 27:921.921.
- Oh MM, Power JM, Thompson LT, Moriearty PL, Disterhoft JF. 1999. Metrifonate increases neuronal excitability in CA1 pyramidal neurons from both young and aging rabbit hippocampus. *J Neurosci* 19:1814–23.
- Olesen SP, Munch E, Moldt P, Drejer J. 1994. Selective activation of  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels by novel benzimidazolone. *Eur J Pharmacol* 251:53–9.
- Osmanovic SS, Schefner SA. 1993. Calcium-activated hyperpolarizations in rat locus coeruleus neurons in vitro. *J Physiol* 469:89–109.
- Park Y-B. 1994. Ion selectivity and gating of small conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels in cultured rat adrenal chromaffin cells. *J Physiol* 481:555–70.



- Pedarzani P, Kulik A, Muller M, Ballanyi K, Stocker M. 2000. Molecular determinants of  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channel function in rat dorsal vagal neurones. *J Physiol* 527:283–90.
- Pedarzani P, Mosbacher J, Rivard A, Cingolani LA, Oliver D, Stocker M, and others. 2001. Control of electrical activity in central neurons by modulating the gating of small conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels. *J Biol Chem* 276:9762–9.
- Pedarzani P, Storm JF. 1993. PKA mediates the effects of monoamine transmitters on the  $\text{K}^+$  current underlying the slow spike frequency adaptation in hippocampal neurons. *Neuron* 11:1023–35.
- Pedarzani P, Storm JF. 1996. Evidence that the Ca/calmodulin-dependent protein kinase mediates the modulation of the  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  current IAHP by acetylcholine, but not by glutamate, in hippocampal neurons. *Pflügers Arch* 431:723–8.
- Pennefather P, Lancaster B, Adams PR, Nicoll RA. 1985. Two distinct Ca-dependent K currents in bullfrog sympathetic ganglionic cells. *Proc Natl Acad Sci U S A* 82:3040–4.
- Piskorowski R, Aldrich RW. 2002. Calcium activation of BK(Ca) potassium channels lacking the calcium bowl and RCK domains. *Nature* 420:499–502.
- Pitler TA, Landfield PW. 1990. Aging-related prolongation of calcium spike duration in rat hippocampal slice neurons. *Brain Res* 508:1–6.
- Power JM, Wu WW, Sametsky E, Oh MM, Disterhoft JF. 2002. Age-related enhancement of the slow outward calcium-activated potassium current in hippocampal CA1 pyramidal neurons in vitro. *J Neurosci* 22:7234–43.
- Rapp PR, Gallagher M. 1996. Preserved neuron number in the hippocampus of aged rats with spatial learning deficits. *Proc Natl Acad Sci U S A* 93:9926–30.
- Reinhart PH, Levitan IB. 1995. Kinase and phosphatase activities intimately associated with a reconstituted calcium-dependent potassium channel. *J Neurosci* 15:4572–9.
- Ritsner M, Modai I, Ziv H, Amir S, Halperin T, Weizman A, and others. 2002. An association of CAG repeats at the KCNN3 locus with symptom dimensions of schizophrenia. *Biol Psychiatry* 51:788–94.
- Romey G, Hugues M, Schmid-Antomarchi H, Lazdunski M. 1984. Apamin: a specific toxin to study a class of  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels. *J Physiol* 79:259–64.
- Romey G, Lazdunski M. 1984. The coexistence in rat muscle cells of two distinct classes of  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels with different pharmacological properties and different physiological functions. *Biochem Biophys Res Commun* 118:669–74.
- Rosenblatt KP, Sun Z-P, Heller S, Hudspeth AJ. 1997. Distribution of  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel isoforms along the tonotopic gradient of the chick cochlea. *Neuron* 19:1061–75.
- Rosenzweig ES, Rao G, McNaughton BL, Barnes CA. 1997. Role of temporal summation in age-related long-term potentiation-induction deficits. *Hippocampus* 7:549–58.
- Saar D, Grossman Y, Barkai E. 1998. Reduced after-hyperpolarization in rat piriform cortex pyramidal neurons is associated with increased learning capability during operant conditioning. *Eur J Neurosci* 10:1518–23.
- Sah P. 1992. The role of calcium influx and calcium buffering in the kinetics of a  $\text{Ca}^{2+}$  activated  $\text{K}^+$  current in rat vagal motoneurons. *J Neurophysiol* 68:2237–48.
- Sah P. 1995a. Different calcium channels are coupled to potassium channels with distinct physiological roles in vagal neurons. *Proc R Soc Lond B* 260:105–11.
- Sah P. 1995b. Properties of channels mediating the apamin-insensitive afterhyperpolarization in vagal motoneurons. *J Neurophysiol* 74:1772–6.
- Sah P. 1996.  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  currents in neurones: types, physiological roles and modulation. *Trends Neurosci* 19:150–4.
- Sah P, Bekkers JM. 1996. Apical dendritic location of slow-afterhyperpolarization current in hippocampal pyramidal neurons: implications for the integration of LTP. *J Neurosci* 16:4537–42.
- Sah P, Faber ESL. 2002. Channels underlying neuronal calcium-activated potassium currents. *Prog Neurobiol* 66:345–53.
- Sah P, Isaacson JS. 1995. Channels underlying the slow afterhyperpolarization in hippocampal pyramidal neurons: neurotransmitters modulate the open probability. *Neuron* 15:435–41.
- Sah P, McLachlan EM. 1991.  $\text{Ca}^{2+}$  activated  $\text{K}^+$  currents underlying the afterhyperpolarization in guinea pig vagal neurons: a role for  $\text{Ca}^{2+}$  activated  $\text{Ca}^{2+}$  release. *Neuron* 7:257–64.
- Sah P, McLachlan EM. 1992. Potassium currents contributing to action potential repolarization and the afterhyperpolarization in rat vagal motoneurons. *J Neurophysiol* 68:1834–41.
- Sailer CA, Hu H, Kaufmann WA, Trieb M, Schwarzer C, Storm JF, and others. 2002. Regional differences in distribution and functional expression of small-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels in rat brain. *J Neurosci* 22:9698–707.
- Sanchez M, McManus OB. 1996. Paxilline inhibition of the alpha-subunit of the high conductance calcium-activated potassium channel. *Neuropharmacology* 35:963–8.
- Sansom SC, Stockland JD, Hall D, Williams B. 1997. Regulation of calcium-activated potassium channels by protein phosphatase 2A. *J Biol Chem* 272:9902–6.
- Schreiber M, Salkoff L. 1997. A novel calcium-sensing domain in the BK channel. *Biophys J* 73:1355–63.
- Schumacher MA, Rivard AF, Bachinger HP, Adelman JP. 2001. Structure of the gating domain of a  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel complexed with  $\text{Ca}^{2+}$ /calmodulin. *Nature* 410:1120–4.
- Schwindt PC, Spain WJ, Foehring RC, Stafstrom CE, Chubb MC, Crill WE. 1988. Multiple potassium conductances and their functions in neurons from cat sensorimotor cortex in vitro. *J Neurophysiol* 59:424–49.
- Seutin V, Johnson SW. 1999. Recent advances in the pharmacology of quaternary salts of bicuculline. *Trends Pharmacol Sci* 20:268–70.
- Shah M, Haylett DG. 2000. The pharmacology of hSK1  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels expressed in mammalian cell lines. *Br J Pharmacol* 129:627–30.
- Shao LR, Halvorsrud R, Borg-Graham L, Storm JF. 1999. The role of BK-type  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  channels in spike broadening during repetitive firing in rat hippocampal pyramidal cells. *J Physiol* 521:135–46.
- Shen KZ, Lagrutta A, Davies NW, Standen NB, Adelman JP, North RA. 1994. Tetraethylammonium block of Slowpoke calcium-activated potassium channels expressed in *Xenopus* oocytes: evidence for tetrameric channel formation. *Pflügers Arch* 426:440–5.
- Shmukler BE, Bond CT, Wilhelm S, Bruening-Wright A, Maylie J, Adelman JP, and others. 2001. Structure and complex transcription pattern of the mouse SK1 K(Ca) channel gene, KCNN1. *Biochim Biophys Acta* 1518:36–46.
- Stackman RW, Hammond RS, Linardatos E, Gerlach A, Maylie J, Adelman JP, and others. 2002. Small conductance  $\text{Ca}^{2+}$ -activated  $\text{k}^+$  channels modulate synaptic plasticity and memory encoding. *J Neurosci* 22:10163–71.
- Stocker M, Krause M, Pedarzani P. 1999. An apamin-sensitive  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  current in hippocampal pyramidal neurons. *Proc Natl Acad Sci U S A* 96:4662–7.
- Stocker M, Pedarzani P. 2000. Differential distributions of three  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel subunits, SK1, SK2 and SK3 in the adult rat central nervous system. *Mol Cell Neurosci* 15:476–93.
- Storm JF. 1987. Action potential repolarization and a fast after-hyperpolarization in rat hippocampal pyramidal cells. *J Physiol* 385:733–59.
- Storm JF. 1990. Potassium currents in hippocampal pyramidal cells. *Prog Brain Res* 83:161–87.
- Strobaek D, Jorgensen TD, Christophersen P, Ahring PK, Olesen S-P. 2000. Pharmacological characterization of small-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels stably expressed in HEK 293 cells. *Br J Pharmacol* 129:991–9.
- Syme CA, Gerlach AC, Singh AK, Devor DC. 2000. Pharmacological activation of cloned intermediate- and small-conductance  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels. *Am J Physiol* 278:C570–81.
- Tanabe M, Gahwiler BH, Gerber U. 1998. L-type  $\text{Ca}^{2+}$  channels mediate the slow  $\text{Ca}^{2+}$ -dependent afterhyperpolarization current in rat CA3 pyramidal cells in vitro. *J Neurophysiol* 80:2268–73.
- Thibault O, Landfield PW. 1996. Increase in single L-type calcium channels in hippocampal neurons during aging. *Science* 272:1017–20.
- Thompson LT, Moyer JR Jr, Disterhoft JF. 1996. Transient changes in excitability of rabbit CA3 neurons with a time course appropriate to support memory consolidation. *J Neurophysiol* 76:1836–49.

- Toro L, Stefani E. 1991.  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels: metabolic regulation. *J Bioenerg Biomembr* 23:561–76.
- Torres GE, Arfken CL, Andrade R. 1996. 5-Hydroxytryptamine<sub>4</sub> receptors reduce afterhyperpolarization in hippocampus by inhibiting calcium-induced calcium release. *Mol Pharmacol* 50:1316–22.
- Tseng-Crank J, Godinot N, Johansen TE, Ahring PK, Strobaek D, Mertz R, and others. 1996. Cloning, expression, and distribution of a  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channel beta-subunit from human brain. *Proc Natl Acad Sci U S A* 93:9200–5.
- Valiante TA, Abdul-Ghani MA, Carlen PL, Pennefather P. 1997. Analysis of current fluctuations during after-hyperpolarization current in dentate granule neurones of the rat hippocampus. *J Physiol* 499:121–34.
- van der Staay FJ, Fanelli RJ, Blokland A, Schmidt BH. 1999. Behavioral effects of apamin, a selective inhibitor of the SK(Ca)-channel, in mice and rats. *Neurosci Biobehav Rev* 23:1087–110.
- Vegara C, Latorre R, Marrion NV, Adelman JP. 1998. Calcium-activated potassium channels. *Curr Opin Neurobiol* 8:321–9.
- Viana F, Bayliss DA, Berger AJ. 1993. Calcium conductances and their role in the firing behavior of neonatal rat hypoglossal motoneurons. *J Neurophysiol* 69:2137–49.
- Vogalis F, Harvey JR, Furness JB. 2002. TEA- and apamin-resistant K(Ca) channels in guinea-pig myenteric neurons: slow AHP channels. *J Physiol* 538:421–33.
- Wallner M, Meera P, Toro L. 1999. Molecular basis of fast inactivation in voltage and  $\text{Ca}^{2+}$ -activated  $\text{K}^+$  channels: a transmembrane beta-subunit homolog. *Proc Natl Acad Sci U S A* 96:4137–42.
- Wei A, Solaro C, Lingle C, Salkoff L. 1994. Calcium sensitivity of BK-type  $\text{K}_{\text{Ca}}$  channels determined by a separable domain. *Neuron* 13:671–81.
- Weiss C, Preston AR, Oh MM, Schwarz RD, Welty D, Disterhoft JF. 2000. The M1 muscarinic agonist CI-1017 facilitates trace eyeblink conditioning in aging rabbits and increases the excitability of CA1 pyramidal neurons. *J Neurosci* 20:783–90.
- Wilkinson D, Murray J. 2001. Galantamine: a randomized, double-blind, dose comparison in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 16:852–7.
- Wolfart J, Neuhoff H, Franz O, Roepe J. 2001. Differential expression of the small-conductance, calcium-activated potassium channel SK3 is critical for pacemaker control in dopaminergic midbrain neurons. *J Neurosci* 21:3443–56.
- Wu WW, Oh MM, Disterhoft JF. 2002. Age-related biophysical alterations of hippocampal pyramidal neurons: implications for learning and memory. *Ageing Res Rev* 1:181–207.
- Xia X-M, Ding JP, Lingle CJ. 1999. Molecular basis for the inactivation of  $\text{Ca}^{2+}$ - and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *J Neurosci* 19:5255–64.
- Xia X-M, Falker B, Rivard A, Wayman G, Johnson-Pais T, Keen JE, and others. 1998. Mechanism of calcium gating in small-conductance calcium-activated potassium channels. *Nature* 395:503–7.
- Xie J, McCobb DP. 1998. Control of alternative splicing of potassium channels by stress hormones. *Science* 280:443–6.
- Yoshizaki K, Hoshino T, Sato M, Koyano H, Nohmi M, Hua S-Y, and others. 1995.  $\text{Ca}^{2+}$ -induced  $\text{Ca}^{2+}$  release and its activation in response to a single action potential in rabbit otic ganglion cells. *J Physiol* 486.1:177–87.