

Functional significance of channels and transporters expressed in the inner ear and kidney

Florian Lang,¹ Volker Vallon,^{2,3} Marlies Knipper,⁴ and Philine Wangemann⁵

¹Department of Physiology, Eberhard-Karls-University of Tübingen, Tübingen, Germany; ²Departments of Medicine and Pharmacology, University of California, and ³Veterans Affairs San Diego Healthcare System, San Diego, California;

⁴Department of Otorhinolaryngology, Tübingen Hearing Research Centre, Molecular Neurobiology, Tübingen, Germany; and ⁵Department of Anatomy and Physiology, Kansas State University, Manhattan, Kansas

Lang F, Vallon V, Knipper M, Wangemann P. Functional significance of channels and transporters expressed in the inner ear and kidney. *Am J Physiol Cell Physiol* 293: C1187–C1208, 2007. First published August 1, 2007; doi:10.1152/ajpcell.00024.2007.—A number of ion channels and transporters are expressed in both the inner ear and kidney. In the inner ear, K⁺ cycling and endolymphatic K⁺, Na⁺, Ca²⁺, and pH homeostasis are critical for normal organ function. Ion channels and transporters involved in K⁺ cycling include K⁺ channels, Na⁺-2Cl⁻-K⁺ cotransporter, Na⁺/K⁺-ATPase, Cl⁻ channels, connexins, and K⁺/Cl⁻ cotransporters. Furthermore, endolymphatic Na⁺ and Ca²⁺ homeostasis depends on Ca²⁺-ATPase, Ca²⁺ channels, Na⁺ channels, and a purinergic receptor channel. Endolymphatic pH homeostasis involves H⁺-ATPase and Cl⁻/HCO₃⁻ exchangers including pendrin. Defective connexins (GJB2 and GJB6), pendrin (SLC26A4), K⁺ channels (KCNJ10, KCNQ1, KCNE1, and KCNMA1), Na⁺-2Cl⁻-K⁺ cotransporter (SLC12A2), K⁺/Cl⁻ cotransporters (KCC3 and KCC4), Cl⁻ channels (BSND and CLCNKA + CLCNKB), and H⁺-ATPase (ATP6V1B1 and ATPV0A4) cause hearing loss. All these channels and transporters are also expressed in the kidney and support renal tubular transport or signaling. The hearing loss may thus be paralleled by various renal phenotypes including a subtle decrease of proximal Na⁺-coupled transport (KCNE1/KCNQ1), impaired K⁺ secretion (KCNMA1), limited HCO₃⁻ elimination (SLC26A4), NaCl wasting (BSND and CLCNKB), renal tubular acidosis (ATP6V1B1, ATPV0A4, and KCC4), or impaired urinary concentration (CLCNKA). Thus, defects of channels and transporters expressed in the kidney and inner ear result in simultaneous dysfunctions of these seemingly unrelated organs.

cochlea; vestibular labyrinth; stria vascularis; deafness; renal tubule

THE INNER EAR is the sensory system for sound, motion, and gravity. It is housed within the temporal bone and consists of the cochlea, vestibular labyrinth, and endolymphatic sac (Fig. 1). The inner ear comprises an array of interconnected fluid compartments that are enclosed by a multitude of highly specialized epithelial cells. The luminal fluid, endolymph, differs in composition between different parts of the inner ear (Table 1). The epithelial cells enclosing the endolymph are highly diverse, as shown in more detail for the cochlea in Fig. 2. Among them are the sensory inner and outer hair cells (Fig. 2, B and C) and stria vascularis (Fig. 2D). Sensory hair cells transduce mechanical stimuli into electrical signals and release neurotransmitters to activate sensory neurons. The stria vascularis is a multilayered epithelium in the cochlea (Fig. 2, A and D), which secretes K⁺ into the endolymph and generates the endocochlear potential, which contributes significantly to the driving force of sensory transduction. In addition, the inner ear contains and depends on a multitude of highly specialized epithelial cells that control the ionic

composition of the endolymph and the magnitude of the transepithelial potential.

Different compartments of the inner ear serve the transduction of specific stimuli. The cochlea transduces mechanical stimuli associated with sound and provides the basis for hearing. The utricle, saccule, and ampullae of the semicircular canals belong to the vestibular labyrinth (Fig. 1), which transduces mechanical stimuli associated with head position and head motion. Vestibular sensory transduction provides input to the vestibular system that controls balance, posture, and eye movements. Sensory transduction in the cochlea and vestibular labyrinth has different electrochemical requirements, although all depend on the cycling of K⁺ between the endolymph and perilymph. In addition to the fluid compartments that house sensory hair cells, the vestibular labyrinth contains another fluid compartment, the endolymphatic sac, which is devoid of sensory hair cells. The function of the endolymphatic sac is poorly understood, although evidence suggests that it controls endolymph fluid volume (219).

Several epithelia in the cochlea appear to have functional equivalents in the vestibular labyrinth. Among such homology pairs are cochlear (Fig. 2A) and vestibular hair cells, stria marginal cells (Fig. 2D) and vestibular dark cells, outer sulcus

Address for reprint requests and other correspondence: F. Lang, Dept. of Physiology, Eberhard-Karls-Univ. of Tübingen, Gmelinstrasse 5, Tübingen D-72076, Germany (e-mail: florian.lang@uni-tuebingen.de).




Fig. 1. The most important compartments of the inner ear, including the cochlea, vestibular system, and endolymphatic sac [Modified from Ref. 83a.].

cells (Fig. 2A) and vestibular transitional cells, and Reissner’s membrane (Fig. 2A) and semicircular canal epithelial cells (163, 303). Several recent reviews have focused on ion transport in different inner ear epithelia (163, 304), hereditary hearing loss (197), cochlear fluid volume regulation (219), and ototoxicity (217).

The similarity between epithelial transport in the inner ear and kidney was first suggested by the observation more than 30 years ago that treatment with high doses of the loop diuretic furosemide causes reversible hearing loss (81, 248). Obviously, the inner ear and kidney have very different functions. Nevertheless, most of the genes encoding the epithelial transporters or channels in the inner ear are similarly expressed and/or similarly sensitive to pharmacological intervention in renal tubular epithelia. More importantly, defects of those genes can lead simultaneously to hearing loss and deranged renal tubular transport. Thus, even though several of the channels and transporters expressed in inner ear epithelia are similarly found in other epithelia or even in excitable tissues such as the heart, the pathophysiologically significant overlap is particularly striking between the inner ear and kidney. In several channelopathies, the renal defect is subtle and clinically overlooked in the face of striking hearing loss or life-threatening cardiac arrhythmia. Closer functional analysis reveals, however, the respective defect in renal function. Thus, much can be learned from a comparison of the transport organization in these two organs. The comparison could further serve as a paradigm that channels and transporters could serve different functions in different organs and that genetic defects or pharmacological inhibition of those channels and transporters could lead to seemingly unrelated functional consequences.

The present review first describes the channels and transporters required for inner ear function. The second part of the review is dedicated to the function of those transporters in renal epithelia. The function and pathophysiological significance of the channels and transporters expressed in both the inner ear and kidney is shown in Table 2.

K⁺ Cycling in the Inner Ear

Sensory transduction in the cochlea and vestibular labyrinth depends on the cycling of K⁺ between the endolymph and perilymph (Fig. 2A). K⁺ cycling in the cochlea consists of K⁺ flux from the endolymph through sensory hair cells into the

perilymph, uptake of K⁺ from the perilymph into fibrocytes of the spiral ligament, funneling of K⁺ via gap junctions into basal and intermediate cells of the stria vascularis, efflux of K⁺ from intermediate cells into the intrastrial fluid, and secretion of K⁺ by marginal cells of the stria vascularis into the endolymph (163, 304) (Fig. 2A). Similarly, K⁺ cycling in the vestibular labyrinth consists of K⁺ flux from the endolymph through hair cells into the perilymph and uptake of K⁺ from the perilymph and secretion into the endolymph by vestibular dark cells (163, 304). K⁺ cycling in the cochlea and vestibular labyrinth, however, is not limited to K⁺ efflux through hair cells and K⁺ secretion by the stria vascularis and vestibular dark cells. Additional pathways accomplish K⁺ and Na⁺ reabsorption from the endolymph. In the cochlea, these additional pathways are provided by Reissner’s membrane and by outer sulcus epithelial cells (Fig. 2A) and, in the vestibular system, by semicircular canal epithelial cells and transitional epithelial cells (163, 304).

K⁺ flux through hair cells. Sensory transduction in the cochlea and vestibular labyrinth depends on mechanically gated ion channels of hitherto elusive molecular identity that are located in the hair bundles of hair cells (32). Candidates that have been considered and rejected include acid-sensing ion channels (194) and members of the epithelial Na⁺ channel (ENaC)/degenerin (ENaC/DEG) superfamily, which are widely distributed in the central and peripheral nervous system and are also found in the cochlea (34, 70, 174, 335) and vestibular labyrinth (138, 202, 334). The current view, however, appears to favor the involvement of transient receptor potential (TRP) channels such as TRPN1, TRPV4, TRPML3, and TRPA1 for mechanotransduction (32).

Opening of the so far elusive transduction channels supports influx of K⁺ from the endolymph into hair cells, which depolarizes the basolateral membrane of the hair cell (Fig. 2, B and C). Influx of K⁺ from the endolymph into hair cells is balanced by efflux of K⁺ from hair cells via K⁺ channels into interstitial spaces that are continuous with the perilymph. The molecular entities of K⁺ efflux channels depend on the type of hair cell.


K⁺ efflux from cochlear inner hair cells involves the voltage-gated K⁺ channel KCNQ4 and the large-conductance (BK) Ca²⁺-activated K⁺ channel KCNMA1 (a BK channel) (48, 121, 122, 240, 278) (Fig. 2B). In addition, KCNQ4 contributes to the resting membrane potential of inner hair cells and thereby ensures the maintenance of the resting cytosolic

Table 1. Fluid composition of the cochlear endolymph and perilymph as well as the endolymph of the endolymphatic sac and cerebrospinal fluid


	Cochlear Perilymph	Cochlear Endolymph	Endolymph in the Endolymphatic Sac	Cerebrospinal Fluid
Na ⁺ , mM	148	1.3	129	149
K ⁺ , mM	4.2	157	8–13	3.1
Cl ⁻ , mM	119	132	124	129
HCO ₃ ⁻ , mM	21	31		19
Ca ²⁺ , mM	1.3	0.023		
Protein, mg/d	178	38		24
pH	7.3	7.5	6.7–7.1	7.3

Values were taken from a recent review (316) and were amended by additional data (35, 93–95, 178, 281, 315).


A Cochlea



B IHC



C OHC



D Stria vascularis




Fig. 2. Compartments of the cochlea. *A*: cross section through the cochlear duct. *B*: inner hair cell (IHC). *C*: outer hair cell (OHC). *D*: stria vascularis. Gene names of expressed ion channels and transporters are illustrated within the approximate position. DC, Deiter's cells; CC, Claudius' cells; HC, Hensen's cells; OS, outer sulcus cells; SC, supporting cells; I–V, specialized fibrocyte types.

Ca²⁺ concentration (156, 157, 189). Several splice variants of KCNQ4 are expressed in the inner ear, of which one variant, KCNQ4v3, is preferentially expressed in the high-frequency region base of the cochlea (12, 142, 143). Mice lacking KCNQ4 or expressing dominant negative mutations develop normal hearing but later in life suffer progressive hearing loss, which indicates that KCNQ4 is nonessential for basic inner hair cell function but required for the maintenance of hearing

(107). Consistently, mutations of KCNQ4 cause progressive high-frequency hearing loss in humans (33, 123, 269). Similarly, mice lacking KCNMA1 develop normal hearing but then suffer progressive hearing loss, indicating that also KCNMA1 is nonessential for basic inner hair cell function (190, 216). Loss of the KCNMB1 β-subunit, which associates with the KCNMA1 α-subunit, appears to have no effect on hearing (216).

Table 2. Channels, transporters, and pumps expressed in both the inner ear and kidney

Gene Symbol	Aliases	Function			Consequences of Loss of Function		
		Inner Ear	Kidney	Inner Ear	Inner Ear	Kidney	
Channels and associated proteins							
<i>KCNJ10</i>	Kir4.1 or Kir1.2	Formation of endocochlear potential	Unknown	Deafness ^b	Unknown	Unknown	
<i>KCNQ1/KCNE1</i>	KvLQT1/minK or ISK	K ⁺ secretion into endolymph; endolymph formation	Maintain PT brush border membrane potential ^b	Deafness ^{a,b}	No (KCNQ1) or modest (KCNE1) urinary loss of Na ⁺ and glucose ^b	No (KCNQ1) or modest (KCNE1) urinary loss of Na ⁺ and glucose ^b	
<i>KCNMA1/KCNMB1</i>	BK channel, α_1 and β_1	K ⁺ efflux from cochlear inner hair cells	(Flow-dependent) K ⁺ secretion in CNT and CD ^b	Progressive hearing loss (only KCNMA1) ^b	Impaired (flow-dependent) renal K ⁺ excretion ^b	Impaired (flow-dependent) renal K ⁺ excretion ^b	
<i>CLCNKA</i>	Cl ⁻ channel Ka	Recycling of Cl ⁻ in strial marginal cells and vestibular dark cells; endolymph formation	Basolateral Cl ⁻ reabsorption in thin ascending limb ^b	Deafness when both CLCNKB and CLCNKA are defective ^a	Nephrogenic diabetes insipidus ^b	Nephrogenic diabetes insipidus ^b	
<i>CLCNKB</i>	Cl ⁻ channel Kb	Recycling of Cl ⁻ in strial marginal cells and vestibular dark cells; endolymph formation	Basolateral Cl ⁻ reabsorption in thick ascending limb ^b	Deafness when both CLCNKB and CLCNKA are defective ^a	Barter syndrome ^{a,b}	Barter syndrome ^{a,b}	
<i>BSND</i>	Barttin	Recycling of Cl ⁻ in strial marginal cells and vestibular dark cells; endolymph formation	Basolateral Cl ⁻ reabsorption in thin and thick ascending limbs ^b	Deafness ^a	Barter syndrome ^{a,b}	Barter syndrome ^{a,b}	
<i>SCNN1A, SCNN1B, and SCNN1G</i>	ENaC, α , β , and γ	Uptake of Na ⁺ into epithelial cells of Reissner's membrane and semicircular canal	Na ⁺ reabsorption and secondary K ⁺ excretion in CNT and CD ^{a,b}	No known inner ear phenotype	Pseudohypoaldosteronism type I ^{a,b}	Pseudohypoaldosteronism type I ^{a,b}	
<i>GJA1</i>	Gap junction protein; connexin43	Metabolic coupling and glutamate buffering in the lateral and medial network; buffering of K ⁺ released from outer hair cells	Regulation of renin secretion ^b	Deafness ^a	Unknown	Unknown	
<i>GJB6</i>	Gap junction protein; connexin30	Metabolic coupling and glutamate buffering in the lateral and medial network; buffering of K ⁺ released from outer hair cells; endocochlear potential	Pathway for ATP release from intercalated cells; ^b	Deafness ^{a,b}	Unknown	Unknown	
<i>TRPV4</i>	Transient receptor potential V4	Ca ²⁺ homeostasis of endolymph	Cellular response to alterations of extracellular osmolarity ^b	Delayed-onset hearing loss and vulnerability to acoustic injury ^b	Unknown	Unknown	
<i>TRPV5</i>	Transient receptor potential V5; Ca _v T2	Ca ²⁺ homeostasis of endolymph	Ca ²⁺ reabsorption in late DCT and CNT ^b	Unknown	Urinary Ca ²⁺ loss ^b	Urinary Ca ²⁺ loss ^b	
<i>TRPV6</i>	Transient receptor potential V6; Ca _v T1	Ca ²⁺ homeostasis of endolymph	Ca ²⁺ reabsorption in CD ^b	Unknown	Urinary Ca ²⁺ loss ^b	Urinary Ca ²⁺ loss ^b	
<i>AQP1</i>	Aquaporin 1; CHIP28	Unknown	Water transport in PT, TDL, and OMDVR ^b	No overt phenotype ^{a,b}	Impaired near-isotonic reabsorption in PT and urine concentration ^b	Impaired near-isotonic reabsorption in PT and urine concentration ^b	
<i>AQP2</i>	Aquaporin 2	Unknown	Water reabsorption in CNT and CD (apical) ^b	Unknown	Urinary concentrating defect ^{a,b}	Urinary concentrating defect ^{a,b}	
<i>AQP3</i>	Aquaporin 3	Unknown	Water reabsorption in CNT and CD (basolateral) ^b	No overt phenotype ^b	Urinary concentrating defect ^b	Urinary concentrating defect ^b	
<i>AQP4</i>	Aquaporin 4	Unknown	Water reabsorption in CNT and CD (basolateral) ^b	Minor hearing loss ^b	Urinary concentrating defect ^b	Urinary concentrating defect ^b	
Transporters							
<i>SLC12A2</i>	Na ⁺ -2Cl ⁻ -K ⁺ cotransporter; NKCC1	K ⁺ uptake into strial marginal cells and vestibular dark cells; endolymph formation	Regulation of renin release ^b	Deafness ^b	Hypotension ^b	Hypotension ^b	

Continued

Table 2.—Continued

Gene Symbol	Aliases	Function			Consequences of Loss of Function		
		Inner Ear	Kidney	Inner Ear	Kidney	Kidney	
<i>SLC12A7</i>	KCl transporter; KCC4	K ⁺ exit (or uptake) in Deiter's cells?	Cl ⁻ recycling in type A intercalated cells ^b	Deafness ^b	Distal renal tubular acidosis ^b		
<i>SLC26A4</i>	Cl ⁻ /HCO ₃ ⁻ exchanger; pendrin	HCO ₃ ⁻ secretion into endolymph	HCO ₃ ⁻ secretion and Cl ⁻ reabsorption in type B and non-A, non-B intercalated cells ^b	Acidic endolymphatic pH; deafness ^{a,b}	No basal phenotype ^{a,b} ; elevated arterial pH and serum HCO ₃ ⁻ and lower blood pressure with NaCl restriction ^b		
Pumps							
<i>PMCA1B</i>	Plasma membrane Ca ²⁺ -ATPase	Ca ²⁺ homeostasis of endolymph	Ca ²⁺ reabsorption in late DCT and CNT	Unknown	Unknown		
<i>ATP6V1B1</i> and <i>ATP6V0A4</i>	H ⁺ -ATPase, B1 and A4	Acidification of endolymph	Acidification of urine ^a	Deafness	Distal renal tubular acidosis ^a		

CD, collecting duct; CNT, connecting tubule; DCT, distal convoluted tubule; OMDVR, outer medullary descending vasa recta; PT, proximal tubule; TDL, thin descending limb. ^aObserved in humans; ^bobserved in mouse models.

The K⁺ channels KCNQ4 and KCNMA1 also mediate K⁺ efflux from outer hair cells (Fig. 2C). In contrast to inner hair cells, both channels play a critical role for basic cell function in outer hair cells (24, 87, 155, 158). Mice lacking KCNMA1 or KCNQ4 lose outer hair cells, but not inner hair cells, during the progressive loss of hearing (52, 107, 216). The K⁺ channels KCNMA1 and KCNQ4 are apparently essential for the survival of outer hair cells.

KCNQ4 associates with the KCNE1 β-subunit and possibly with other KCNE subunits that are expressed in hair cells (255). This interaction may be critical for KCNQ4 function given that the KCNE1 mutation KCNE1(D76N) impairs KCNQ4 function and causes Jervell and Lange-Nielsen syndrome, consisting of life-threatening cardiac arrhythmias and deafness. Other mutations, such as KCNE1(S74L), which do not impair KCNQ4 function, cause Romano-Ward syndrome, consisting of arrhythmias without deafness (255).

K⁺ buffering near hair cells. Hair cells and neurons in the cochlea and vestibular labyrinth maintain their resting membrane potential via K⁺ channels in conjunction with high cytosolic and low extracellular K⁺ concentrations. Uncontrolled increases of the K⁺ concentration in the extracellular fluid are expected to affect the membrane potential and responsiveness of hair cells and neurons. Stimulation of cochlear and vestibular hair cells leads to measurable increases in the extracellular K⁺ concentration in the surrounding perilymph (104, 282). It is conceivable that K⁺-buffering mechanisms limit the magnitude of these increases. In general, multiple mechanisms have been recognized to limit the amplitude of K⁺ concentration changes in the extracellular environment near neurons. The predominant mechanism is diffusion into unobstructed open fluid spaces. Current measurements in scala tympani perilymph support the concept that the perilymph serves as an unobstructed open fluid space in the buffering of K⁺ (336). Furthermore, a strategic localization of K⁺ channels that differ in their rectification can provide buffering of localized K⁺ increases. Inward-rectifying K⁺ channels are well suited as uptake mechanisms in K⁺ buffering since they conduct K⁺ influx more efficiently than K⁺ efflux. A local increase in the extracellular K⁺ concentration may set the local K⁺ equilibrium potential below the membrane potential, which promotes K⁺ influx into the buffering cell. The ensuing elevation of the cytosolic K⁺ concentration sets the K⁺ equilibrium potential above the membrane potential and promotes K⁺ efflux preferentially through less inward-rectifying K⁺ channels or outward-rectifying K⁺ channels. Such a mechanism has been described in Müller glia of the retina (116, 208). It is conceivable that a similar mechanism is present in the organ of Corti. Deiter's cells have a membrane potential of -76 mV (186), which is near the K⁺ equilibrium potential. They express the inward-rectifying K⁺ channel KCNJ10 (Kir4.1), which is particularly abundant in the membrane area that faces KCNQ4 expressed in outer hair cells (83, 216) (Fig. 2C). K⁺ exit mechanisms in Deiter's cells may include outward-rectifying K⁺ channels (181) and the K⁺/Cl⁻ cotransporters SLC12A6 (KCC3) and SLC12A7 (KCC4) (15, 16) (Fig. 2C). Moreover, Deiter's cells are connected to neighboring supporting cells via gap junctions. K⁺ could thus be dispersed via gap junctions among epithelial cells that include Deiter's cells (Fig. 2A, DC), Claudius' cells (Fig. 2A, CC), Hensen's cells (Fig. 2A, HC), and outer sulcus cells (Fig. 2A, OC).

SLC12A6 and SLC12A7 may serve as a release mechanism for KCl not only in K⁺ buffering but also in cell volume regulation (131). The recent claim that SLC12A6 and SLC12A7 serve as a K⁺ uptake mechanism in Deiter's cells, however, would require an unusually low cytosolic Cl⁻, which has not been shown thus far (15, 16). Consistent with a role of SLC12A6 and SLC12A7 in cell volume regulation is the finding that mice lacking either transporter hear normal at the onset of hearing but suffer from a more or less early onset of hearing loss (15, 16).

K⁺ uptake from the perilymph. K⁺ released from sensory hair cells may travel along multiple pathways toward the spiral ligament in the lateral wall (246) (Fig. 2A). Pathways that avoid supporting cells or that involve buffering by Deiter's cells or neighboring cells and lead through the open perilymph space of scala tympani are supported by current measurements (336). An additional pathway may involve uptake of K⁺ into Deiter's cells, dispersion of K⁺ among Deiter's cells, Hensen's cells, and outer sulcus cells via gap junctions (Fig. 2A), and release of K⁺ into the interstitial space of the spiral ligament that is continuous with perilymph (110).

Uptake of K⁺ from the interstitial space of the spiral ligament occurs via specialized fibrocytes (named I–V within the spiral ligament in Fig. 2A). Fibrocyte types II, IV, and V express Na⁺/K⁺-ATPase, the Na⁺-2Cl⁻-K⁺ cotransporter SLC12A2 (NKCC1), and the Cl⁻ channels CLCNKA and CLCNKB (36, 154, 177, 206, 230) (Fig. 2D). Although functional data from fibrocytes are lacking, the resemblance of this array of transporters with the basolateral membrane of stria marginal cells and vestibular dark cells suggests that fibrocytes take up K⁺ from the perilymph. Gap junctions, in particular GJB2 (CX26) and GJB6 (CX30), connect fibrocyte types II, IV, and V among each other as well as to fibrocyte type I cells and basal and intermediate cells of the stria vascularis. Gap junctions form a network in the lateral wall that is thought to provide a pathway for K⁺ from the sites of uptake into fibrocyte types II, IV, and V to the sites of release from stria intermediate cells into the intrastrial fluid space.

K⁺ secretion into the endolymph. Strial marginal cells and vestibular dark cells take up K⁺ from the intrastrial fluid space and secrete it into the endolymph (Fig. 2D). K⁺ secretion by stria marginal cells and vestibular dark cells occurs via equivalent mechanisms (311). Both epithelial cells take up K⁺ across the basolateral cell membrane via the Na⁺-2Cl⁻-K⁺ cotransporter SLC12A2 (NKCC1) and Na⁺/K⁺-ATPase and secrete K⁺ across the apical membrane via the K⁺ channel KCNQ1/KCNE1 (Fig. 2D). Na⁺ and Cl⁻ taken up via the Na⁺-2Cl⁻-K⁺ cotransporter is recycled in the basolateral membrane via Na⁺/K⁺-ATPase and the Cl⁻ channels ClC-Ka/barttin (CLCNKA/BSND) and ClC-Kb (CLCNKB/BSND) (Fig. 2D). The following paragraphs focus on the Na⁺-2Cl⁻-K⁺ cotransporter, K⁺ channels, and Cl⁻ channels that are essential for K⁺ secretion in stria marginal cells and vestibular dark cells.

Na⁺/K⁺-ATPASE AND K⁺/H⁺-ATPASE. Strial marginal cells and vestibular dark cells absorb K⁺ from the intrastrial space and perilymph via Na⁺/K⁺ ATPase and the Na⁺-2Cl⁻-K⁺ cotransporter (170, 306). Na⁺/K⁺ ATPase takes up K⁺ and establishes a Na⁺ gradient that energizes further uptake of K⁺ via SLC12A2. The Na⁺/K⁺-ATPase in stria marginal cells and vestibular dark cells as well as in fibrocytes of the spiral

ligament consists of the subunits ATP1A1, ATP1B1, and ATP1B2 (173, 230) (Fig. 2D).

Inhibition of Na⁺/K⁺-ATPase with ouabain inhibits K⁺ secretion and consequently abolishes the endocochlear potential (126, 129). Strial marginal cells appear to express gastric K⁺/H⁺-ATPase in addition to Na⁺/K⁺-ATPase (135, 235). The functional significance of K⁺/H⁺-ATPase for the generation of the endocochlear potential is currently unclear since very high concentrations of K⁺/H⁺-ATPase inhibitors were necessary to affect the endocochlear potential.

Na⁺-2Cl⁻-K⁺ COTRANSPORTER. Strial marginal cells and vestibular dark cells absorb K⁺ from the intrastrial space and perilymph via the Na⁺-2Cl⁻-K⁺ cotransporter SLC12A2 (NKCC1) (162, 306, 313) (Fig. 2D). SLC12A2 is sensitive to the loop diuretics furosemide and bumetanide and to their analog piretanide (162, 313). SLC12A2 is an essential transporter for K⁺ secretion and endolymph production. Mice that lack SLC12A2 fail to produce endolymph, which leads to the collapse of Reissner's membrane onto the stria vascularis and organ of Corti (41, 43, 57).

K⁺ CHANNELS. Strial marginal cells and vestibular dark cells secrete K⁺ into the endolymph via the K⁺ channel KCNQ1/KCNE1 (165, 306) (Fig. 2D). KCNQ1/KCNE1 is a slowly activating delayed rectifier that carries the slow delayed rectifier K⁺ current and requires the assembly of the pore-forming KCNQ1 α -subunit with the KCNE1 β -subunit (11, 223). Mice lacking functional KCNE1 or KCNQ1 fail to produce endolymph, which leads to a collapse of Reissner's membrane onto the stria vascularis and organ of Corti due to loss of K⁺ secretion in the presence of ongoing reabsorptive processes (22, 139, 140, 294). Similar observations have been made in human patients (61). Homozygous or heterozygous compound mutations of KCNE1 or KCNQ1 lead to Jervell and Lange-Nielsen syndrome, which is characterized by deafness, prolonged cardiac action potentials, and potentially fatal cardiac arrhythmias (103, 182, 231, 233). Consistently, pharmacological inhibition of the KCNQ1/KCNE1 channel leads to hearing loss (79).

The rate of K⁺ secretion is controlled by the K⁺ concentration on the apical and basolateral membrane and by cell volume, pH, and a variety of receptors and signaling mechanisms. Transepithelial currents and currents through the apical KCNQ1/KCNE1 K⁺ channel are enhanced by lowering the apical K⁺ concentration or increasing the basolateral K⁺ concentration or by lowering the osmolarity on the basolateral side (161, 306, 308, 317). Furthermore, the rate of K⁺ secretion is increased by β_1 -adrenergic receptors via cAMP-dependent stimulation of the KCNQ1/KCNE1 K⁺ channel (258, 259, 309, 310). Conversely, muscarinic and purinergic receptors suppress K⁺ secretion (307). Purinergic P2Y₄ receptors decrease currents through the KCNQ1/KCNE1 K⁺ channel via protein kinase C (160, 166). In addition, KCNE1/KCNQ1 K⁺ channel activity is stimulated by the serum- and glucocorticoid-inducible kinase SGK1 (20, 50), which may contribute to the stimulation of cochlear ion transport and hearing improvement by glucocorticoids and mineralocorticoids (137, 280). The channel is also inhibited by estrogens (298), which may contribute to the inhibitory effect of those hormones on cochlear transport (136).

The rate of K⁺ secretion may further be regulated by trafficking of KCNE1/KCNQ1 K⁺ channels to the apical

membrane of strial marginal cells, which requires the participation of lysosomal integral membrane protein II (LIMP2) (115). LIMP2 is a transmembrane glycoprotein that is mainly located in lysosomal and endosomal membranes (127). Mice lacking LIMP2 suffer from progressive hearing loss correlated with a loss of surface expression of KCNQ1/KCNE1 in the apical membrane of marginal cells (115). In addition, mice lacking LIMP2 suffer from uni- or bilateral hydronephrosis due to hypertrophy of the smooth muscle layer at the ureteropelvic junction (65).

CL⁻ CHANNELS. K⁺ secretion by strial marginal cells and vestibular dark cells require Cl⁻ to recycle in the basolateral membrane via a major Cl⁻ conductance (306, 314) (Fig. 2D). This Cl⁻ conductance is composed of the Cl⁻ channels CLCNKA/BSND and CLCNKB/BSND (6, 53, 154, 167, 206, 218, 264, 265). Cl⁻ channels CLCNKA/BSND and CLCNKB/BSND consist of the pore-forming CLCNKA α -subunits and CLCNKB and the BSND β -subunit (53, 228). Mutations of BSND reduce channel conductivity and surface expression and thereby cause Bartter's syndrome type 4, which is characterized by deafness and renal salt wasting (14, 53). Similarly, simultaneous mutations of CLCNKA and CLCNKB also lead to Bartter's syndrome type 4, whereas mutations of CLCNKB lead to Bartter's syndrome without deafness (171, 226, 237). This observation is consistent with the finding that CLCNKA and CLCNKB are coexpressed in cells of the inner ear but not the kidney and with the notion that the two channels can substitute for each other in the inner ear (206) but not in the kidney (see below).

K⁺ and Na⁺ Reabsorption

Homeostasis of high K⁺ and low Na⁺ concentrations in the endolymph is maintained by K⁺ secretion and Na⁺ and K⁺ reabsorption. Reabsorption of K⁺ is not limited to the pathways through inner, outer, and vestibular hair cells. Indeed, currents generated by the stria vascularis in the cochlea flow not only through hair cells but also through the outer sulcus and through Reissner's membrane (222, 336). Consistently, outer sulcus and Reissner's membrane epithelial cells reabsorb Na⁺ and K⁺ from the endolymph (138, 159). Outer sulcus cells take up Na⁺ and K⁺ via apical nonselective cation channels, BK channels, and small-conductance K⁺ channels as well as P2X₂ receptor-gated nonselective cation channels. They release Na⁺ and K⁺ across the basolateral membrane via Na⁺/K⁺-ATPase and K⁺ channels, respectively (25, 26, 136). Subunits of ENaC (SCNN1) may contribute to the apical nonselective cation channels (Fig. 2A), although the channel involved is not the typical Na⁺-selective and amiloride-sensitive ENaC channel (25, 70).

Reissner's membrane epithelial cells take up Na⁺ via the amiloride-sensitive Na⁺ channel ENaC and extrude Na⁺ across the basolateral membrane via Na⁺/K⁺-ATPase (138) (Fig. 2A). The endocochlear potential contributes to the driving force of cation reabsorption in outer sulcus and Reissner's membrane epithelial cells much like it contributes to the transduction current through inner and outer hair cells.

Na⁺ and K⁺ reabsorption has also been found in the vestibular labyrinth. Cation reabsorption in vestibular transitional cells and semicircular canal epithelial cells bear some resemblance to cation reabsorption in outer sulcus and Reissner's

membrane epithelial cells, respectively. Vestibular transitional cells reabsorb Na⁺ and K⁺ via apical P2X₂ receptor-gated nonselective cation channels and extrude Na⁺ and K⁺ across the basolateral membrane via Na⁺/K⁺-ATPase and K⁺ channels, respectively (136, 312, 318).

Semicircular canal epithelial cells in the vestibular labyrinth reabsorb Na⁺ via ENaC, release Na⁺ across the basolateral membrane via Na⁺/K⁺-ATPase, and recycle K⁺ in the basolateral membrane via K⁺ channels (201, 202). Na⁺ reabsorption in semicircular canal epithelial cells is under the control of glucocorticoids but not mineralocorticoids (201, 202). ENaC is activated by the transmembrane serine protease TMPRSS3 (72). A defect of TMPRSS3 leads to deafness (72), which may, however, involve dysregulation of further transporters or channels besides ENaC activity. Loss of function mutations of ENaC do not lead to an inner ear phenotype, which is consistent with the presence of alternative Na⁺ reabsorption pathways in outer sulcus epithelial cells and transitional cells (136, 159).

Generation of the Endocochlear Potential

Mechanical stimuli associated with sound, head position, or gravity are transduced into electrical signals by sensory hair cells in the cochlea and vestibular labyrinth. Mechanically induced channel openings permit an influx of K⁺ from the endolymph into the hair cell. The driving force of this current is roughly the sum of the basolateral membrane potential of the hair cell and the transepithelial potential. The transepithelial potential in the cochlea, called the endocochlear potential, is enormous: +80 mV (Fig. 2A). For cochlear inner and outer hair cells, the driving force for sensory transduction is 120 mV (-40 mV + 80 mV) and 150 mV (-70 mV + 80 mV), respectively (38, 189). Driving forces for sensory transduction in the vestibular labyrinth are smaller due to the smaller endovestibular potential of 3–7 mV (130, 185, 220).

The endocochlear potential is a transepithelial potential that is generated by the stria vascularis (275, 295, 316). The stria vascularis is functionally a two-layered epithelium composed of a layer of marginal cells and a layer of basal cells that is penetrated by a capillary network (99). Marginal cell junctions contain a multitude of different claudins, whereas tight junctions between basal cells contain only claudin 11 (CLDN11) (58, 114). Tight junctions among basal cells define an inner membrane facing the intrastrial space and an outer membrane facing the spiral ligament (Fig. 2). The inner membrane is connected via gap junctions to strial intermediate cells, and strial intermediate cells are connected via gap junctions to strial pericytes and endothelial cells (108, 262). Gap junctions ensure that intermediate cells are electrically a part of the basal cell barrier. The outer membrane of basal cells is connected by gap junctions to type I fibrocytes of the spiral ligament.

The endocochlear potential is essentially a K⁺ equilibrium potential that is generated by the K⁺ channel KCNJ10 (Kir4.1) in intermediate cells of the stria vascularis (Fig. 2D) in conjunction with a very low K⁺ concentration of intrastrial fluid and a high cytosolic K⁺ concentration in intermediate cells (169, 263). A number of key findings provide support for this model. First, the endocochlear potential and the KCNJ10 K⁺ channel in intermediate cells share the same sensitivities to a panel of K⁺ channel blockers (164, 262, 266). Second, the

endocochlear potential can be measured across the basal cell barrier (221). Third, the expression of KCNJ10 correlates with the presence of the endocochlear potential in KCNJ10 knockout and pendrin knockout mouse models and in normal development (82, 169, 215, 305, 315). Fourth, loss of claudin 11, which is the only known claudin in basal cell tight junctions, renders the basal cell barrier leaky and leads to a loss of the endocochlear potential (67, 114). Fifth, increases of the K^+ concentration in the intrastrial fluid space suppress the endocochlear potential. Such increases can be achieved by vascular perfusion of solutions containing elevated K^+ concentrations, inhibitors of Na^+/K^+ -ATPase (ouabain), or inhibitors of the $Na^+-2Cl^- -K^+$ cotransporter (furosemide or bumetanide) (118, 126, 128, 129, 164). Finally, loss of GJB6 [connexin (Cx)30], which renders capillaries in the stria vascularis leaky to the intrastrial space, leads to a loss of the endocochlear potential (31). Collectively, these findings support the model that the endocochlear potential is a K^+ equilibrium potential that is generated by the K^+ channel KCNJ10.

Marginal cells of the stria vascularis and fibrocytes of the spiral ligament play important supporting roles in the generation of the endocochlear potential. Fibrocytes of the spiral ligament, which are connected via basal cells to intermediate cells, ensure a high cytosolic K^+ concentration in strial intermediate cells. Strial marginal cells reabsorb K^+ from the intrastrial fluid spaces and keep the K^+ concentration in the intrastrial fluid spaces as low as 2 mM (164, 263, 306).

Gap Junctional Networks

Several major networks of cells that are connected by gap junctions have been recognized in the cochlea (109). Notably excluded from these networks are marginal cells of the stria vascularis, inner hair cells and outer hair cells that are neither connected among each other nor to any of their neighbors. The importance of gap junctions for cochlear function is underscored by the fact that mutations of GJB2 (Cx26) and GJB6 (Cx30) are the most prevalent causes of hereditary childhood deafness consistent with the contribution of GJB2 and GJB6 to all major gap junctional networks in the cochlea (39, 42, 68, 73, 191, 193, 331).

One major network of gap junction interconnected cells links different types of fibrocytes in the spiral ligament as well as basal and intermediate cells, pericytes, and endothelial cells of the stria vascularis (Fig. 2A, spiral ligament). A major purpose of this network is to connect sites of K^+ uptake in fibrocyte types II, IV, and V to the site of K^+ release in intermediate cells of the stria vascularis. Most gap junctions in this network are formed by heteromeric complexes of GJB2 and GJB6 (2, 59, 109, 134, 325). In addition, endothelial cells of the stria vascularis express GJA1 (Cx43) and GJA7 (Cx45) (29, 134, 260) and fibrocytes of spiral ligament express GJB3 (Cx31) and GJB1 (Cx32) (150, 326). Mutations of GJA1, GJB1, and GJB3 are also associated with deafness (147, 148, 251, 326).

Mice that lack GJB6 are profoundly deaf despite the continued presence of GJB2 (277). The assumed limited gap junction coupling mediated by the remaining GJB2 and other connexins appears to be sufficient for the cycling of K^+ but insufficient to prevent leakiness of strial capillaries and breakdown of the endocochlear potential (31). Conse-

quently, mice lacking GJB6 failed to develop an endocochlear potential but had normal endolymphatic K^+ concentrations at least at a young age (277). Interestingly, the insufficiency of gap junction coupling, which is associated with leaky capillaries in mice lacking GJB6, can be restored by overexpression of GJB2. Mice lacking GJB6 and overexpressing GJB2 develop a normal endocochlear potential and have normal hearing (3).

Two further networks are formed by epithelial cells in and adjacent to the organ of Corti: the medial and lateral networks (98, 109, 247). Most gap junctions in these networks are formed by GJB2 and GJB6 (2, 59, 109, 134, 325). In addition, some cells express GJA1 (147, 260). The lateral network of gap junction interconnected cells in the organ of Corti includes outer pillar cells, Deiter's cells, Hensen's cells, Claudius' cells, outer sulcus cells, and root cells (98, 109) (Fig. 2A). The major purpose of this network may be metabolic coupling in addition to buffering of K^+ that is released from outer hair cells in response to sound stimulation (104).

The medial network includes inner pillar cells, supporting cells of the inner hair cells and interdental cells (Fig. 2, A and B). A major purpose of this network is to buffer glutamate, which is the neurotransmitter released from the inner hair cell. Expression of the glutamate uptake transporter SLC1A3 (GLAST) is limited to the immediate neighbor of the inner hair cell, whereas glutamine synthase, a key enzyme in the detoxification of glutamate, is mainly expressed in adjacent cells but not in SLC1A3-expressing cells (55). Gap junctions between SLC1A3-expressing cells and their glutamine synthase-expressing neighbors may be required for the transcellular metabolism of glutamate. Support for the concept of transcellular glutamate buffering comes from the finding that mice lacking SLC1A3 fail to buffer glutamate, which leads to an accumulation of glutamate in scala tympani perilymph during sound stimulation (76). SLC1A3-expressing supporting cells are also the first cells in the organ of Corti to undergo apoptosis in mice that lack GJB2 in this region of the cochlea (30). Further, several deafness-causing mutations of GJB2 and GJB6 impair the transfer of organic molecules but do not impede ionic coupling, which implies that these mutations do not affect K^+ cycling but could impair metabolic coupling and glutamate buffering (31, 332).

Several reasons may account for the intriguing observation that the loss of function of either GJB2 or GJB6 leads to deafness rather than simply being compensated by the remaining connexin-forming homomeric gap junctions (30, 124, 277). First, loss of GJB6 has been shown to suppress the protein expression of GJB2, which reduces intercellular coupling more than predicted by the simple omission of GJB6 (3). Mutations may also exert dominant negative effects on the function of wild-type isoforms (68, 210). Second, heteromeric gap junctions formed from GJB2 and GJB6 have slightly different biophysical properties than homomeric gap junctions (257). The finding that overexpression of GJB2 can rescue hearing of mice lacking GJB6 suggests that the biophysical differences between hetero- and homomeric GJB2 gap junctions are less important than the fact that loss of GJB6 leads to a loss of GJB2 expression and a reduction in intercellular coupling (3).

Ca²⁺ Homeostasis

The transduction channel in hair cells is a Ca²⁺-permeable nonselective cation channel. Although K⁺ is the major charge carrier, the transduction current is in part carried by Ca²⁺ and the reliability of the transduction process itself depends on the constancy of Ca²⁺ concentrations in the endolymph (Table 1). Both elevated and reduced concentrations of Ca²⁺ have been shown to suppress transduction currents and microphonic potentials (187, 271). Furthermore, Ca²⁺ homeostasis of vestibular endolymph during development affects the formation of otoconia, which are necessary for the detection of gravity and linear acceleration (106, 152). Consistent with the importance of Ca²⁺ homeostasis in the endolymph are the observations that mice and guinea pigs with reduced or elevated endolymphatic Ca²⁺ concentrations are deaf and have vestibular deficits (120, 185, 315, 324).

The endolymphatic Ca²⁺ concentration appears to be controlled by secretory and reabsorptive mechanisms. Ca²⁺ reabsorption may occur through paracellular and transcellular pathways and may at least in part be driven by the endocochlear potential (91). In general, transepithelial Ca²⁺ transport may employ Ca²⁺-permeable channels as Ca²⁺ uptake mechanisms, Ca²⁺ binding proteins as Ca²⁺ buffers in the cytosol, and Ca²⁺-ATPases or Na⁺/Ca²⁺ exchangers as Ca²⁺ extrusion mechanisms. Ca²⁺-ATPases appear to be most suitable for Ca²⁺ extrusion into the endolymph due to the low Na⁺ concentration in the endolymph (Table 1), which does not provide a driving force for Ca²⁺ extrusion via Na⁺/Ca²⁺ exchangers. Consistently, Ca²⁺ secretion into the endolymph has been shown to depend on Ca²⁺-ATPases rather than on Na⁺/Ca²⁺ exchangers (92, 324) and loss of function of Ca²⁺-ATPase ATP2B2 (PMCA2) leads to deafness and to a reduction in the endolymphatic Ca²⁺ concentration (253, 324).

Among the many different epithelial cells lining cochlear and vestibular endolymph, the cells best understood to be involved in the homeostasis of endolymph Ca²⁺ include outer hair cells in the cochlea and semicircular canal duct epithelial cells in the vestibular labyrinth, although it is currently unclear whether outer hair cells contribute to the homeostasis of bulk endolymph or only to the homeostasis of the endolymph in the nearest vicinity of the hair bundle. Nevertheless, outer hair cells have been shown to secrete Ca²⁺ into the endolymph (328). This Ca²⁺ secretion is required to remove Ca²⁺ from the cytosol of the hair bundle and to maintain an appropriate Ca²⁺ concentration in the endolymph surrounding the bundle (7, 84). Outer hair cells express Ca²⁺-ATPase ATP2B2 in the stereocilia (49, 64) (Fig. 2C), a high concentration of Ca²⁺ binding proteins in the cytosol (74), and Ca²⁺-permeable channels in the basolateral membrane including TRPC, TRPV1, TRPV4, and L-type and non-L-type Ca²⁺ channels (45, 144, 157, 207, 234, 333). Mice lacking TRPV4 develop normal hearing consistent with a redundancy of Ca²⁺-permeable channels. However, they suffer from a delayed-onset hearing loss and vulnerability to acoustic injury (261). It is currently unclear whether the delayed-onset hearing loss and vulnerability to acoustic injury are due to the loss of TRPV4 in outer hair cells or due to the loss of the channel from other cells including inner hair cells and spiral ganglion neurons (234).

Semicircular canal duct epithelial cells in the vestibular labyrinth have been shown to reabsorb Ca²⁺ from the en-

dolymph (180). Ductal epithelial cells express Ca²⁺-permeable TRPV5 and TRPV6 channels, Ca²⁺ binding proteins, Na⁺/Ca²⁺ exchangers, and Ca²⁺-ATPases (327). Consistent with an apical membrane expression of pH-sensitive TRPV5 and TRPV6 Ca²⁺ channels is the finding that the transepithelial Ca²⁺ flux was pH sensitive and that endolymph Ca²⁺ concentrations were elevated in mice that have acidic endolymph due to loss of pendrin (180).

Other epithelial cells in the cochlea and vestibular labyrinth may be involved in endolymph Ca²⁺ homeostasis since they express Ca²⁺-ATPases and Ca²⁺-permeable channels. Whether these cells secrete or reabsorb Ca²⁺ is currently not clear. Inner hair cells, in contrast to outer hair cells, may be involved in Ca²⁺ reabsorption (84). The transduction channel may serve as an uptake channel, and Ca²⁺-ATPase ATP2B1 (PMCA1) in the basolateral membrane may serve as a release mechanism (49). Similarly, vestibular hair cells express in their basolateral membrane ATP2B1 and ATP2B3 (PMCA3). However, inner and vestibular hair cells also express ATP2B2 in hair bundles (49) (Fig. 2B). It is thus unclear whether inner and vestibular hair cells support a transcellular Ca²⁺ flux. The stria vascularis expresses ATP2B1 and Ca²⁺-permeable TRPV4, TRPV5, and TRPV6 channels (1, 37, 144, 268, 315, 324). Reissner's membrane and interdental cells express Ca²⁺-ATPases (37, 64, 324). Outer sulcus cells express ATP2B2 (64), and inner and outer sulcus epithelial cells as well as ductal epithelial cells of the semicircular canals express TRPV5 and TRPV6 Ca²⁺ channels (180, 315). TRPV5 and TRPV6 Ca²⁺ channels may be located in the apical membrane of at least some cochlear epithelial cells (315).

pH Homeostasis

The pH of the endolymph varies greatly between different regions in the inner ear. In the cochlea and utricle, the endolymphatic pH is slightly alkaline (pH 7.5) (93, 180, 315). In the endolymphatic sac, on the other hand, the pH is more acidic (pH 6.6–7.1) (35, 281). The functional significance of these differences is largely elusive. The presence of these differences, however, underscores that fluid homeostasis in different compartments of the inner ear is controlled by local ion transport in adjacent epithelia rather than via a fluid flow between different compartments of the inner ear (219).

The homeostasis of endolymphatic pH depends on the secretion of H⁺ and HCO₃⁻. Epithelial cells that express H⁺-ATPase in their apical membrane include interdental cells of the spiral limbus (Fig. 2A) and stria marginal cells (Fig. 2D) as well as endolymphatic duct and sac epithelial cells (46, 105, 250). Furthermore, epithelial cells that express in their apical membrane the HCO₃⁻-permeable anion exchanger SLC26A4 (pendrin) include spiral prominence and outer sulcus epithelial cells and spindle cells of the stria vascularis as well as endolymphatic duct and sac epithelial cells (46, 54, 305, 330).

The main buffers, at least in the cochlear endolymph, appear to be CO₂ and HCO₃⁻. Glycosaminoglycans, which are found in high concentrations in the endolymph of the endolymphatic sac, may contribute to pH buffering (89, 203). Proteins, however, that contribute to the buffering capacity of blood plasma appear to play a lesser role in the buffering of cochlear endolymph due to their low concentration (Table 1). Marginal cells of the stria vascularis are a significant local source of CO₂

due to their high metabolic rate and their use of the hexose monophosphate pathway (168). Carbonic anhydrases in the stria vascularis, spiral ligament, and spiral limbus capture metabolically derived CO_2 and convert it to HCO_3^- (146, 188, 245). HCO_3^- generated within the fibrocyte gap junction network may be secreted into the endolymph via the HCO_3^- -permeable anion exchanger pendrin (SLC26A4). Consistent with HCO_3^- secretion into the endolymph is the observation that mice lacking pendrin have an acidic endolymphatic pH (180, 315). Furthermore, increased metabolic rates during acoustic stimulation cause an alkalization of the endolymph, which is consistent with an increased rate of HCO_3^- secretion (92).

Endolymphatic pH homeostasis is necessary for hearing and the prevention of hearing loss, although effects of pH may be indirect. For example, acidification of the endolymph inhibits Ca^{2+} reabsorption via pH-sensitive TRPV5 and TRPV6 Ca^{2+} channels and elevates the endolymphatic Ca^{2+} concentration, which impairs cochlear function (180, 315). Furthermore, acidification enhances free radical stress and promotes hearing loss (270).

Whether mutations of the B1 subunit (ATP6V1B1) or A4 subunit (ATPV0A4) of H^+ -ATPase cause an alkalization of endolymph pH is currently unknown. Nevertheless, mutations of either subunit may cause, in humans, a progressive sensorineural hearing loss in addition to renal tubular acidosis (105, 252, 288). The etiologies of these hearing losses, however, are unclear, in particular since mice lacking the B1 subunit (ATP6V1B1) develop normal hearing and show no overt morphological abnormalities in the inner ear (44).

Water Transport

Water transport follows osmotic gradients that are established by the transport, metabolism, or catabolism of solutes (131). Water can permeate most membranes freely with the notable exception of the apical membrane of thick ascending limb and of the cortical collecting duct of the kidney in the absence of the antidiuretic hormone vasopressin. The water permeability of cell membranes depends to a significant extent on the presence of aquaporins, which are water-permeable channels. According to a recent review (184), 13 different aquaporins (AQP0–AQP12) have so far been identified.

A multitude of aquaporins are expressed in the inner ear, including AQP1, AQP2, AQP3, AQP4, AQP5, AQP7, and AQP9 (88, 149, 175, 176, 224, 249, 267, 334). The functional significance of inner ear water channels is largely unclear. Loss of function of AQP1 associated with the Colton blood group does not cause an overt clinical phenotype, although it is associated with a reduction of the urinary concentration capacity (113, 205). Hearing loss or balance difficulties have not been reported in association with the Colton blood group. Neither have hearing loss or balance disorders been reported to be associated with diabetes insipidus due to loss of AQP2. Mice lacking AQP1, AQP3, or AQP5 have normal hearing; however, mice lacking AQP4 have a minor hearing loss of 10 dB at 4–5 wk of age (141) (Fig. 2A). Whether hearing is already impaired earlier is currently unknown.

Functions of Inner Ear Channels and Transporters in Renal Epithelia

Many of the channels, carriers, and pumps accomplishing transport in the inner ear are similarly expressed in the kidney and participate in renal tubular transport. Accordingly, the hearing loss in patients carrying genetic defects of defined transport molecules may be paralleled by deranged renal acid or electrolyte excretion that affects acid-base or electrolyte homeostasis of the body. Moreover, and possibly related to the different organization and function of the two organs, some of the transport proteins are used for quite different cellular functions in the inner ear and kidney, as discussed below.

The proximal tubule of the kidney (Fig. 3A) reabsorbs ~60% of filtered NaCl and fluid and most of the filtered amino acids and glucose. K^+ channels in the apical cell membrane of proximal tubules (Fig. 3A) contribute to the maintenance of the cell membrane potential during depolarizing Na^+ -coupled transport (e.g., cotransport of Na^+ with amino acids or glucose), thereby stabilizing the electrical driving force for electrogenic Na^+ reabsorption.

Henle's loop contributes to the generation of a hypertonic kidney medulla, a prerequisite for urinary concentration. Most importantly, the thick ascending limb of Henle's loop reabsorbs ~25% of NaCl filtered by the glomeruli without accompanying water reabsorption, thus enhancing interstitial osmolarity (Fig. 3B). The medullary collecting ducts (Fig. 3D) pass the hypertonic kidney medulla. During water retention, water channels allow water to leave the lumen of the collecting ducts, thus leading to urinary concentration.

The distal convoluted tubule (Fig. 3C), connecting tubule (not explicitly shown), and collecting duct (Fig. 3, D and E) allow the fine tuning of renal acid, fluid, and electrolyte excretion. In all nephron segments, the proximal tubule, Henle's loop, distal tubule, and collecting duct K^+ channels maintain the cell membrane potential and thus the driving force for electrogenic transport.

K^+ channels. K^+ channels expressed in both the inner ear and kidney include KCNE1/KCNQ1, KCNJ10, and BK channels (KCNMA1/KCNMB1) (Table 2).

KCNE1 and KCNQ1 have been localized to the brush border of the mid to late proximal tubule (256, 283) (Fig. 3A). Besides their potential role in net K^+ secretion into the early proximal tubule (284), they may polarize the brush border membrane and thus maintain the electrical driving force for Na^+ -coupled transport (132, 133). Studies (283, 284) in knockout mice indeed revealed that lack of functional KCNE1/KCNQ1 K^+ channels leads to moderate impairment of electrogenic Na^+ -glucose cotransport in proximal tubules (see Fig. 3A). KCNE1 may interact with additional K^+ channels, especially in the early proximal tubule, where most of the glucose, amino acids, and phosphate are reabsorbed by electrogenic cotransport with Na^+ and where KCNE1 but not KCNQ1 was detected. In the early proximal tubule, KCNE1 is likely to coassemble with another KCNQ isoform (Fig. 3A), similar to what has been recently shown for outer hair cells (255). KCNQ1-independent function of KCNE1 may explain the more severe phenotype (e.g., renal Na^+ and glucose loss) in mice lacking KCNE1 compared with KCNQ1 (283, 284). Thus, whereas KCNQ1/KCNE1 K^+ channels serve to establish high K^+ concentrations in the endolymph of the inner ear, they serve the very different




Fig. 3. Individual segments of the tubular and collecting duct system of the kidney. Positions within the nephron are indicated. Gene names of expressed ion channels and transporters are illustrated within the approximate position. S, substrate for Na⁺-coupled electrogenic transport.

function of stabilizing the membrane potential and thus electrogenic reabsorption of Na⁺ in the proximal tubule of the kidney, with the secreted K⁺ being subsequently reabsorbed by paracellular routes.

KCNJ10 is expressed in the basolateral cell membrane of renal distal tubules, including the thick ascending limb (97, 151, 272, 273) (see Fig. 3B). These channels are highly sensitive to cytosolic pH and are thus thought to link K⁺ metabolism with acid-base balance (21). To our knowledge, however, no data are available on K⁺ or acid-base balance in mice lacking KCNJ10. Whereas KCNJ10 is considered to be of primary importance for the endocochlear potential of the inner ear, its precise role in the kidney remains to be defined.

BK channels (KCNMA1) are expressed in the renal vasculature and tubular system (71, 198) (Fig. 3D). In the latter, they contribute to K⁺ secretion into the luminal fluid. BK channels in the luminal membrane of the distal nephron (Fig. 3D) are involved in K⁺ homeostasis in response to a high-K⁺ diet (9, 179, 212) and mediate renal K⁺ excretion in response to enhanced tubular flow rates (71, 198–200, 212, 274, 323). In the mouse, the β_1 -subunit of KCNMB1 was found exclusively in the connecting tubule (200). Notably, this β_1 -subunit confers protein kinase G activation of BK channels, dramatically increases the Ca²⁺ sensitivity of the channel, and leads to the activation of the channel at more negative potentials, thereby presumably enhancing the ability of the pore-forming α -subunit to induce significant K⁺ excretion in the distal nephron under physiological conditions (for a review, see Ref. 198). Moreover, mice lacking the α -subunit (KCNMA1) (212) but also mice deficient in the β_1 -subunit (KCNMB1) (199) exhibit blunted flow-induced renal K⁺ excretion. These studies implied a role for BK channels (KCNMA1/KCNMB1) in flow-induced renal K⁺ excretion and K⁺ homeostasis. BK channels are also expressed in other tubular segments, where their function is less clear. Whether circulation or flow of the

endolymph similarly affects BK channel activity in the inner ear remains to be determined.

Na⁺-2Cl⁻-K⁺ cotransporter. The Na⁺-2Cl⁻-K⁺ cotransporter NKCC1 (SLC12A2) is highly expressed in glomeruli of more mature nephrons (286) and may participate in the macula densa-dependent regulation of renin release (23, 66, 302). A closer look at SLC12A2 knockout mice more recently revealed that they suffer from hypotension, which was proposed to relate in part to an impaired responsiveness of the kidney to aldosterone and vasopressin (302).

The Na⁺-2Cl⁻-K⁺ cotransporter NKCC2 (SLC12A1), which is strongly expressed in the luminal membrane of the thick ascending limb (40) (Fig. 3B), is responsible for most of the NaCl reabsorption in that segment and is a prerequisite for the ability of the kidney to dilute and concentrate urine. Accordingly, genetic defects of SLC12A1 lead to isosthenuria and severe renal salt loss (80, 204, 238, 287). SLC12A1 is not expressed in the inner ear, and lack of functional SLC12A1 does not lead to hearing loss. Conversely, SLC12A2 deficiency, as discussed above, leads to deafness without leading to overt renal salt wasting (43, 57). Importantly, both SLC12A1 and SLC12A2 are inhibited by loop diuretics such as furosemide, and thus inhibition of Na⁺-2Cl⁻-K⁺ cotransport in the inner ear during excessive doses of loop diuretics leads to an accumulation of K⁺ in the intrastrial space, which abolishes the endocochlear potential (129) and leads to hearing loss (81, 96, 322). Much lower doses are sufficient to inhibit luminal SLC12A1 in the thick ascending limb, since the drug accumulates in the tubular fluid as a consequence of efficient secretion into proximal tubular fluid and fluid reabsorption along the tubule. Thus, natriuretic and diuretic actions can be achieved without hearing loss.

Cl⁻ channels. The Cl⁻ channel CLCNKA/BSND is expressed in the basolateral membrane of thin ascending limbs (not shown), whereas the Cl⁻ channel CLCNKB/BSND is

expressed in the basolateral membrane of thick ascending limbs of Henle's loop (53, 297) (see Fig. 3B). In the mouse, CIC-K1 (the rodent ortholog of CLCNKA) is also expressed in the thin ascending limb. Knockout of CIC-K1 in mice results in nephrogenic diabetes insipidus, establishing that CIC-K1 has a role in urine concentration and that the countercurrent system in the inner medulla is involved in the generation and maintenance of a hypertonic medullary interstitium (171). In the thick ascending limb, basolateral CLCNKB/BSND contributes to transcellular NaCl reabsorption. Defects of CLCNKB lead to the renal salt wasting of classical Bartter syndrome without hearing impairment (119, 237). The phenotype of patients suffering from defective CLCNKB (119, 237), however, is less severe than the phenotype of patients suffering from antenatal Bartter syndrome due to defective SLC12A1 (238) or apical K^+ channel ROMK (239). Genetic defects of BSND lead to renal salt wasting together with deafness (14) (Table 2).

Voltage-clamp experiments disclosed that a common (prevalence 20% in Caucasians and 40% in Africans) variant of the CLCNKB gene leading to the replacement of threonine by serine at the amino acid position 481 of the CIC-Kb protein (CIC-KbT481S) dramatically increases CIC-Kb Cl^- channel activity (100). Expression of the mutated channels should decrease the cytosolic Cl^- concentration and thus enhance the driving force and transport rate of $Na^+-2Cl^- -K^+$ cotransport. As a result, the gene variant may lead to enhanced transport in the inner ear and kidney. The gene variant was associated with increased blood pressure in one study (101) on a population of largely young, healthy individuals but not in two other studies (117, 244) on more elderly populations. The same gain of function mutation was associated with a slight but significant delay of hearing loss in female humans, whereas no significant differences were observed between male carriers and noncarriers of the mutation (60).

Thus, whereas CLCNKA/BSND and CLCNKB/BSND serve the recycling of Cl^- across the basolateral membrane of marginal cells of the inner ear to maintain the uptake of K^+ via SLC12A2, in the kidney the two channels serve the very different function of basolateral net transport of Cl^- along the ascending thin and thick limb.

K^+-Cl^- cotransport. The K^+-Cl^- cotransporter KCC4 (SLC12A7) is found along the basolateral cell membrane in several nephron segments (289) (see Fig. 3, A and E). KCC4 is colocalized with KCC3 (SLC12A6) in basolateral cell membranes of the proximal tubule (15, 47) (Fig. 3A), where it may contribute to proximal tubular cell volume regulation. This cotransporter is also important for Cl^- recycling in type A intercalated cells (Fig. 3E). In accordance with a crucial role of SLC12A7 for KCl release in K^+ buffering and volume regulation in both the kidney and inner ear, mice lacking KCC4 suffer from renal tubular acidosis (15, 102, 196) and deafness (15, 16) (Table 2). Whether KCC4 is of similar pathophysiological significance in humans remains elusive.

ENaC. ENaC (Fig. 3D, SCNN1) is mainly expressed in the luminal membrane of the aldosterone-sensitive distal nephron, where ~1–3% of filtered Na^+ is reabsorbed. ENaC is of critical importance for renal Na^+ reabsorption and secondary K^+ excretion and, thus, for salt and K^+ homeostasis and blood pressure regulation. Patients carrying loss of function mutations of ENaC ("dominant" pseudohypoaldosteronism type 1) as well as knockout mice for ENaC subunits (SCNN1A,

SCNN1B, and SCNN1C) suffer from renal salt wasting (69, 90). As mutant mice die soon after birth, it is still elusive whether the mutated gene would induce hearing loss. Patients with gain of function mutations of ENaC (Little's syndrome) suffer from hypertension (19, 77, 145, 225, 319, 320) but are not known to suffer from deafness.

Gap junctional channels. The classic gap junction channels have been shown by a freeze-fracture study (125) in the proximal tubule, and some of the ubiquitous connexin isoforms [GJA4 (Cx37), GJA5 (Cx40), GJA1 (Cx43), and GJA7 (Cx45)] have subsequently been identified in the kidney and localized to mainly vascular and glomerular components (8, 10). In the so-called juxtaglomerular apparatus, Cx40 and Cx43 have been implicated in the regulation of renin secretion (75, 296).

Moreover, GJB6 protein (Cx30), probably in the form of luminal hemichannels, was found to be expressed in renal tubular epithelial cells (Fig. 3E) and inserted into the apical cell membrane, particularly of intercalated cells (172). GJB6 proteins were upregulated by a high-salt diet in the distal nephron (172) (Fig. 3E). It has thus been speculated that GJB6 may function as an apical hemichannel allowing the passage of ATP and having a potential inhibitory role in the regulation of salt reabsorption in the distal nephron (172). Along those lines, a recent study (211) has showed that mice deficient for the ATP receptor P2Y₂ present a salt-resistant form of arterial hypertension that is associated with facilitated renal Na^+ and fluid reabsorption. It would be interesting to learn whether renal electrolyte excretion and/or blood pressure are altered in patients with defective GJB6 or in GJB6 knockout mice.

Ca^{2+} homeostasis. Together with the intestine and bone, the kidney is of primary importance for body Ca^{2+} homeostasis. Hormone-regulated renal Ca^{2+} reabsorption is mainly localized to the late distal convoluted tubule and connecting tubule, where TRPV5 channels, expressed in the luminal membrane (Fig. 3C), accomplish Ca^{2+} uptake and Ca^{2+} -ATPases (PMCA1B; Fig. 3C) as well as Na^+/Ca^{2+} exchangers (NCX1) basolateral exit (85). Accordingly, TRPV5-deficient mice suffer from impaired Ca^{2+} reabsorption leading to renal Ca^{2+} loss (86, 209). Notably, TRPV6, which mediates Ca^{2+} reabsorption in the intestine, is also expressed in the luminal membrane of the collecting duct (i.e., downstream of the segments primarily expressing TRPV5; Fig. 3D) and may contribute to renal tubular Ca^{2+} reabsorption. Accordingly, Ca^{2+} reabsorption in the collecting duct limits renal Ca^{2+} loss in mice lacking TRPV5 (86), and mice lacking TRPV6 also lose some Ca^{2+} into urine (13).

Renal TRPV4 is expressed mainly in the basolateral cell membrane of thin and thick ascending limbs and the distal convoluted tubule (28, 279) (Fig. 3B). TRPV4 channels were presumed to participate in the cellular response to alterations of extracellular osmolarity (195, 254). TRPV4 knockout mice tend to be hypercalcemic, which would indicate that TRPV4 rather decreases net renal Ca^{2+} reabsorption (63).

Together, these findings suggest that Ca^{2+} transport pathways that serve to stabilize the Ca^{2+} concentration of the endolymph in the inner ear are involved in kidney function to regulate the Ca^{2+} homeostasis of the whole body.

pH homeostasis. The kidney is of pivotal importance for the regulation of the acid-base balance of the body. Renal regulation of the acid-base balance primarily involves the reab-

sorption, generation, or excretion of HCO_3^- as well as the generation and excretion of NH_4^+ . Vacuolar H^+ -ATPase (ATP6V), H^+/K^+ -ATPase (ATP4A), and the $\text{Cl}^-/\text{HCO}_3^-$ exchangers pendrin (SLC26A4) and AE1 (SLC4A1) are all expressed in the kidney, where they contribute to acid-base balance. Gastric (and colonic) H^+/K^+ -ATPase is expressed in the collecting duct (Fig. 3E). It is responsible for H^+ secretion and K^+ reabsorption under normal conditions and may be stimulated by acid-base perturbations and/or K^+ depletion. The regulation may be species specific (for a review, see Ref. 236).

Vacuolar H^+ -ATPase, H^+/K^+ -ATPase, and $\text{Cl}^-/\text{HCO}_3^-$ exchangers SLC26A4 and SLC4A1 are all expressed in the intercalated cells of the kidney (Fig. 3E), which are critically involved in acid-base balance (for reviews, see Refs. 4, 232, and 299). The localization of H^+ -ATPase in the apical or basolateral membrane can vary between cortical intercalated cells, indicating that subpopulations of these cells have opposite polarities of an H^+ -ATPase, consistent with the presence of both H^+ - and HCO_3^- -secreting cells (17, 18). Along those lines, type A, type B, or non-A, non-B intercalated cells are defined according to the presence or absence of the $\text{Cl}^-/\text{HCO}_3^-$ exchanger SLC4A1 and the subcellular distribution of H^+ -ATPase (5, 111). Type A intercalated cells mediate net secretion of H^+ through apical H^+ -ATPase (ATP6V), which functions in series with basolateral SLC4A1 (51, 276, 321). Particularly during metabolic alkalosis, type B intercalated cells mediate the secretion of HCO_3^- by employing the apical $\text{Cl}^-/\text{HCO}_3^-$ exchanger pendrin (SLC26A4), which functions in series with basolateral H^+ -ATPase (5, 51, 62, 111, 112, 214, 243, 276, 300, 321). Non-A, non-B intercalated cells may be HCO_3^- - or H^+ -secreting cells or may interconvert between the two functions (111, 276). They express both pendrin as well as H^+ -ATPase in the apical membrane (300).

Under basal unstimulated conditions, persons with genetic disruption of pendrin (SLC26A4; Pendred syndrome) and mice lacking SLC26A4 exhibit no change in arterial pH, renal function, or fluid balance (213). Under conditions of dietary NaCl restriction or administration of mineralocorticoids, however, pendrin expression is increased in type B intercalated cells in rodents, and, under these conditions, mice lacking SLC26A4 show evidence for impaired renal HCO_3^- excretion as well as impaired Cl^- retention, which results in elevated arterial pH and serum HCO_3^- and lower blood pressure compared with wild-type mice (213, 293, 301). In contrast to the inner ear, where SLC26A4 mutations lead to hearing loss in humans and mice (54), in the kidney pendrin is essential for a normal response to low-salt conditions (Table 2), indicating that pendrin may be a new target for antihypertensive therapy. In humans, loss of pendrin leads, in addition, to defective iodide uptake into thyroid glands and thus in later life to goiter (54). The latter is not found in mice and is in humans of much later onset than the hearing loss.

Autosomal dominant and recessive forms of distal tubular acidosis are caused by mutations in ion transporters of acid-secreting type A intercalated cells (for a review, see Ref. 4). These include at least two subunits of apical H^+ -ATPase (Fig. 3E and Table 2). Loss of function mutations of the genes encoding for the B1 subunit of H^+ -ATPase lead to recessive distal tubular acidosis with sensorineural hearing loss (105, 241). HCO_3^- therapy successfully treats systemic symptoms of distal renal tubular acidosis but fails to correct deafness,

suggesting that transepithelial acid secretion is required for normal cochlear development and hair cell survival. Mice lacking the B1 subunit (ATP6V1B1) have preserved hearing but exhibit impaired maximal urinary acidification (56). Although patients with distal renal tubular acidosis due to homozygous B1-subunit mutations typically present as infants with spontaneous metabolic acidosis and failure to thrive (105), mice lacking ATP6V1B1 raised on a standard rodent diet were healthy, grew normally, and did not develop metabolic acidosis (56). The phenotypic discrepancy may be related to dietary differences, since a standard rodent diet provides a large net dietary alkali load, whereas the typical Western human diet, which has higher protein content, imposes a net acid load (56). On the other hand, apical expression of the alternative B-subunit isoform B2 is increased in the medulla of mice lacking ATP6V1B1 and may partially, although not completely, compensate for the loss of ATP6V1B1 (56).

Mutations in ATP6N1B, encoding a new kidney vacuolar H^+ -ATPase subunit, which was also localized to the apical membrane of type A intercalated cells, cause recessive distal renal tubular acidosis with preserved hearing (241). Similarly, multiple mutations have been described for the $\text{Cl}^-/\text{HCO}_3^-$ exchanger SLC4A1, which are associated with distal renal tubular acidosis in the absence of deafness (for a review, see Ref. 4).

In conclusion, many transport proteins that stabilize the pH of the endolymph of the inner ear are also involved in the renal transport of HCO_3^- and H^+ and, thus, contribute to the acid-base homeostasis of the body.

Water transport. Aquaporins AQP1–AQP4 play a central role in water reabsorption of the kidney (for a review, see Ref. 183). AQP1 is particularly expressed along the proximal tubule (Fig. 3A), and near-isosmolar fluid reabsorption, a hallmark of proximal tubular function, is dramatically impaired in mice lacking AQP1, indicating that proximal tubular fluid reabsorption is largely due to transcellular water movement through AQP1 (227, 285). AQP1 is the principal water channel in the thin descending limbs of Henle's loop and is also expressed in the outer medullary descending vasa recta, where it facilitates water transport and is thus an important component of the urinary concentrating mechanism (27, 153, 192, 291). A very recent study (78) has provided evidence for the involvement of AQP1 in the migration of proximal tubule cells and possibly in the response of the proximal tubule to injury. In comparison, AQP2 is exclusively expressed in the principal cells of the connecting tubule and collecting duct and is the predominant vasopressin-regulated water channel (Fig. 3D). AQP3 and AQP4 are both present in the basolateral plasma membrane of collecting duct principal cells (Fig. 3D) and represent exit pathways for water reabsorbed apically via AQP2. Studies in patients have demonstrated that AQP2 is essential for urinary concentration (183, 229, 290). Loss of function mutations of AQP2 cause nephrogenic diabetes insipidus (329). The inheritance is usually autosomal recessive but may, in some patients, be autosomal dominant (autosomal dominant nephrogenic diabetes insipidus) (329) and could be related to a dominant negative monomer that leads to a missorting of AQP2 to the basolateral instead of apical plasma membrane of collecting duct cells (242) (Table 2).

Moreover, mice lacking AQP2, AQP3, or AQP4 suffer from various degrees of nephrogenic diabetes insipidus (292).

Conclusions

Epithelial transport in the inner ear and kidney is critical for the function of both organs. Many of the proteins accomplishing ion transport within the inner ear and kidney are encoded by the same genes, as shown in Table 2. Most of the transporters are involved in K^+ cycling within the inner ear and simultaneously participate in the renal tubular transport of Na^+ and K^+ . Other transport systems are involved in regulating and stabilizing the Ca^{2+} concentration or pH of the endolymph and in the regulation of renal tubular transport of Ca^{2+} , HCO_3^- , and H^+ and thus participate in Ca^{2+} homeostasis and acid-base balance of the whole body. The different arrangement of channel proteins may lead to completely different transport functions. For example, KCNQ1/KCNE1 K^+ channels serve to establish high K^+ concentrations in the endolymph of the inner ear, whereas in proximal renal tubules, they stabilize the membrane potential across the apical cell membrane and thus contribute to the maintenance of the electrical driving force for Na^+ -coupled electrogenic transport. The comparison of the transport processes in the inner ear and kidney thus illustrates the amazing versatility of biology in the use of individual molecules. Moreover, the comparison leads to pathophysiological insight into syndromal genetic disease as well as into side effects of drugs targeting those channels and transporters, and it may provide clues to new therapeutic approaches. Our knowledge, though, is still far from complete, and many mechanisms are a matter of speculation. It is an aim of this brief synopsis to stimulate future interdisciplinary research in this exciting and clinically important area of physiology.

ACKNOWLEDGMENTS

The authors acknowledge the meticulous preparation of the manuscript by Jasmin Bühringer.

GRANTS

Work in the laboratory of the authors was supported by the Deutsche Forschungsgemeinschaft (to F. Lang and M. Knipper), American Heart Association Grant-In-Aid 655232Y, the Department of Veterans Affairs, and National Institutes of Health (NIH) Grants DK-56248, DK-28602, DK-70667, and GM-66232 (to V. Vallon). The support by NIH Research Grant R01-DC-01098 (to P. Wangemann) NIH Grant P20-RR-017686 is gratefully acknowledged.

REFERENCES

- Agrup C, Bagger-Sjoberg D, Fryckstedt J. Presence of plasma membrane-bound Ca^{2+} -ATPase in the secretory epithelia of the inner ear. *Acta Otolaryngol (Stockh)* 119: 437–445, 1999.
- Ahmad S, Chen S, Sun J, Lin X. Connexins 26 and 30 are co-assembled to form gap junctions in the cochlea of mice. *Biochem Biophys Res Commun* 307: 362–368, 2003.
- Ahmad S, Tang W, Chang Q, Qu Y, Hibshman J, Li Y, Sohl G, Willecke K, Chen P, Lin X. Restoration of connexin26 protein level in the cochlea completely rescues hearing in a mouse model of human connexin30-linked deafness. *Proc Natl Acad Sci USA* 104: 1337–1341, 2007.
- Alper SL. Genetic diseases of acid-base transporters. *Annu Rev Physiol* 64: 899–923, 2002.
- Alper SL, Natale J, Gluck S, Lodish HF, Brown D. Subtypes of intercalated cells in rat kidney collecting duct defined by antibodies against erythroid band 3 and renal vacuolar H^+ -ATPase. *Proc Natl Acad Sci USA* 86: 5429–5433, 1989.
- Ando M, Takeuchi S. mRNA encoding “Cl⁻-K⁺ channel” is expressed in marginal cells of the stria vascularis of rat cochlea: its possible contribution to Cl⁻ currents. *Neurosci Lett* 284: 171–174, 2000.
- Apicella S, Chen S, Bing R, Penniston JT, Llinas R, Hillman DE. Plasmalemmal ATPase calcium pump localizes to inner and outer hair bundles. *Neuroscience* 79: 1145–1151, 1997.
- Arensbaek B, Mikkelsen HB, Gustafsson F, Christensen T, Holstein-Rathlou NH. Expression of connexin 37, 40, and 43 mRNA and protein in renal preglomerular arterioles. *Histochem Cell Biol* 115: 479–487, 2001.
- Bailey MA, Cantone A, Yan Q, MacGregor GG, Leng Q, Amorim JB, Wang T, Hebert SC, Giebisch G, Malnic G. Maxi-K channels contribute to urinary potassium excretion in the ROMK-deficient mouse model of type II Bartter’s syndrome and in adaptation to a high-K diet. *Kidney Int* 70: 51–59, 2006.
- Barajas L, Liu L, Tucker M. Localization of connexin43 in rat kidney. *Kidney Int* 46: 621–626, 1994.
- Barhanin J, Lesage F, Guillemare E, Fink M, Lazdunski M, Romey G. K_v LQT1 and IsK (minK) proteins associate to form the I_{Ks} cardiac potassium current. *Nature* 384: 78–80, 1996.
- Beisel KW, Rocha-Sanchez SM, Morris KA, Nie L, Feng F, Kachar B, Yamoah EN, Fritsch B. Differential expression of KCNQ4 in inner hair cells and sensory neurons is the basis of progressive high-frequency hearing loss. *J Neurosci* 25: 9285–9293, 2005.
- Bianco SD, Peng JB, Takanaga H, Suzuki Y, Crescenzi A, Kos CH, Zhuang L, Freeman MR, Gouveia CH, Wu J, Luo H, Mauro T, Brown EM, Hediger MA. Marked disturbance of calcium homeostasis in mice with targeted disruption of the Trpv6 calcium channel gene. *J Bone Miner Res* 22: 274–285, 2007.
- Birkenhager R, Otto E, Schurmann MJ, Vollmer M, Ruf EM, Maier-Lutz I, Beekmann F, Fekete A, Omran H, Feldmann D, Milford DV, Jeck N, Konrad M, Landau D, Knoers NV, Antignac C, Sudbrak R, Kispert A, Hildebrandt F. Mutation of BSND causes Bartter syndrome with sensorineural deafness and kidney failure. *Nat Genet* 29: 310–314, 2001.
- Boettger T, Hubner CA, Maier H, Rust MB, Beck FX, Jentsch TJ. Deafness and renal tubular acidosis in mice lacking the K-Cl co-transporter Kcc4. *Nature* 416: 874–878, 2002.
- Boettger T, Rust MB, Maier H, Seidenbecher T, Schweizer M, Keating DJ, Faulhaber J, Ehmke H, Pfeiffer C, Scheel O, Lemcke B, Horst J, Leuwer R, Pape HC, Volkl H, Hubner CA, Jentsch TJ. Loss of K-Cl co-transporter KCC3 causes deafness, neurodegeneration and reduced seizure threshold. *EMBO J* 22: 5422–5434, 2003.
- Brown D, Hirsch S, Gluck S. An H^+ -ATPase in opposite plasma membrane domains in kidney epithelial cell subpopulations. *Nature* 331: 622–624, 1988.
- Brown D, Hirsch S, Gluck S. Localization of a proton-pumping ATPase in rat kidney. *J Clin Invest* 82: 2114–2126, 1988.
- Bubien JK, Ismailov II, Berdiev BK, Cornwell T, Lifton RP, Fuller CM, Achard JM, Benos DJ, Warnock DG. Liddle’s disease: abnormal regulation of amiloride-sensitive Na^+ channels by β -subunit mutation. *Am J Physiol Cell Physiol* 270: C208–C213, 1996.
- Busjahn A, Seebahn G, Maier G, Toliat MR, Nurnberg P, Aydin A, Luft FC, Lang F. Association of the serum and glucocorticoid regulated kinase (sgk1) gene with QT interval. *Cell Physiol Biochem* 14: 135–142, 2004.
- Casamassima M, D’Adamo MC, Pessia M, Tucker SJ. Identification of a heteromeric interaction that influences the rectification, gating, and pH sensitivity of Kir4.1/Kir5.1 potassium channels. *J Biol Chem* 278: 43533–43540, 2003.
- Casimiro MC, Knollmann BC, Ebert SN, Vary JC Jr, Greene AE, Franz MR, Grinberg A, Huang SP, Pfeifer K. Targeted disruption of the Kcnq1 gene produces a mouse model of Jervell and Lange-Nielsen syndrome. *Proc Natl Acad Sci USA* 98: 2526–2531, 2001.
- Castrop H, Lorenz JN, Hansen PB, Friis U, Mizel D, Oppermann M, Jensen BL, Briggs J, Skott O, Schnermann J. Contribution of the basolateral isoform of the Na-K-2Cl⁻ cotransporter (NKCC1/BSC2) to renin secretion. *Am J Physiol Renal Physiol* 289: F1185–F1192, 2005.
- Chambard JM, Ashmore JF. Regulation of the voltage-gated potassium channel KCNQ4 in the auditory pathway. *Pflügers Arch* 450: 34–44, 2005.

25. Chiba T, Marcus DC. Nonselective cation and BK channels in apical membrane of outer sulcus epithelial cells. *J Membr Biol* 174: 167–179, 2000.
26. Chiba T, Marcus DC. Basolateral K⁺ conductance establishes driving force for cation absorption by outer sulcus epithelial cells. *J Membr Biol* 184: 101–112, 2001.
27. Chou CL, Knepper MA, Hoek AN, Brown D, Yang B, Ma T, Verkman AS. Reduced water permeability and altered ultrastructure in thin descending limb of Henle in aquaporin-1 null mice. *J Clin Invest* 103: 491–496, 1999.
28. Cohen DM. TRPV4 and the mammalian kidney. *Pflügers Arch* 451: 168–175, 2005.
29. Cohen-Salmon M, Maxeiner S, Kruger O, Theis M, Willecke K, Petit C. Expression of the connexin43- and connexin45-encoding genes in the developing and mature mouse inner ear. *Cell Tissue Res* 316: 15–22, 2004.
30. Cohen-Salmon M, Ott T, Michel V, Hardelin JP, Perfettini I, Eybalin M, Wu T, Marcus DC, Wangemann P, Willecke K, Petit C. Targeted ablation of connexin26 in the inner ear epithelial gap junction network causes hearing impairment and cell death. *Curr Biol* 12: 1106–1111, 2002.
31. Cohen-Salmon M, Regnault B, Cayet N, Caille D, Demuth K, Hardelin JP, Janel N, Meda P, Petit C. Connexin30 deficiency causes intrastrial fluid-blood barrier disruption within the cochlear stria vascularis. *Proc Natl Acad Sci USA* 104: 6229–6234, 2007.
32. Corey DP. What is the hair cell transduction channel? *J Physiol* 576: 23–28, 2006.
33. Coucke PJ, Van Hauwe P, Kelley PM, Kunst H, Schatteman I, Van Velzen D, Meyers J, Ensink RJ, Verstreken M, Declau F, Marres H, Kastury K, Bhasin S, McGuirt WT, Smith RJ, Cremers CW, Van de HP, Willems PJ, Smith SD, Van Camp G. Mutations in the KCNQ4 gene are responsible for autosomal dominant deafness in four DFNA2 families. *Hum Mol Genet* 8: 1321–1328, 1999.
34. Couloigner V, Fay M, Djelidi S, Farman N, Escoubet B, Runembert I, Sterkers O, Friedlander G, Ferrary E. Location and function of the epithelial Na channel in the cochlea. *Am J Physiol Renal Physiol* 280: F214–F222, 2001.
35. Couloigner V, Teixeira M, Hulin P, Sterkers O, Bichara M, Escoubet B, Planelles G, Ferrary E. Effect of locally applied drugs on the pH of luminal fluid in the endolymphatic sac of guinea pig. *Am J Physiol Regul Integr Comp Physiol* 279: R1695–R1700, 2000.
36. Crouch JJ, Sakaguchi N, Lytle C, Schulte BA. Immunohistochemical localization of the Na-K-Cl co-transporter (NKCC1) in the gerbil inner ear. *J Histochem Cytochem* 45: 773–778, 1997.
37. Crouch JJ, Schulte BA. Expression of plasma membrane Ca-ATPase in the adult and developing gerbil cochlea. *Hear Res* 92: 112–119, 1995.
38. Dallos P. Response characteristics of mammalian cochlear hair cells. *J Neurosci* 5: 1591–1608, 1985.
39. del Castillo I, Moreno-Pelayo MA, del Castillo FJ, Brownstein Z, Marlin S, Adina Q, Cockburn DJ, Pandya A, Siemering KR, Chamberlin GP, Ballana E, Wuyts W, Maciel-Guerra AT, Alvarez A, Villamar M, Shohat M, Abeliovich D, Dahl HH, Estivill X, Gasparini P, Hutchin T, Nance WE, Sartorato EL, Smith RJ, Van Camp G, Avraham KB, Petit C, Moreno F. Prevalence and evolutionary origins of the del(GJB6-D13S1830) mutation in the DFNB1 locus in hearing-impaired subjects: a multicenter study. *Am J Hum Genet* 73: 1452–1458, 2003.
40. Delpire E, Kaplan MR, Plotkin MD, Hebert SC. The Na-(K)-Cl cotransporter family in the mammalian kidney: molecular identification and function(s). *Nephrol Dial Transplant* 11: 1967–1973, 1996.
41. Delpire E, Lu J, England R, Dull C, Thorne T. Deafness and imbalance associated with inactivation of the secretory Na-K-2Cl co-transporter. *Nat Genet* 22: 192–195, 1999.
42. Denoyelle F, Lina-Granada G, Plauchu H, Bruzzone R, Chaib H, Levi-Acobas F, Weil D, Petit C. Connexin 26 gene linked to a dominant deafness. *Nature* 393: 319–320, 1998.
43. Dixon MJ, Gazzard J, Chaudhry SS, Sampson N, Schulte BA, Steel KP. Mutation of the Na-K-Cl co-transporter gene Slc12a2 results in deafness in mice. *Hum Mol Genet* 8: 1579–1584, 1999.
44. Dou H, Finberg K, Cardell EL, Lifton R, Choo D. Mice lacking the B1 subunit of H⁺-ATPase have normal hearing. *Hear Res* 180: 76–84, 2003.
45. Dou H, Vazquez AE, Namkung Y, Chu H, Cardell EL, Nie L, Parson S, Shin HS, Yamoah EN. Null mutation of alpha1D Ca²⁺ channel gene results in deafness but no vestibular defect in mice. *J Assoc Res Otolaryngol* 5: 215–226, 2004.
46. Dou H, Xu J, Wang Z, Smith AN, Soleimani M, Karet FE, Greinwald JH Jr, Choo D. Co-expression of pendrin, vacuolar H⁺-ATPase alpha4-subunit and carbonic anhydrase II in epithelial cells of the murine endolymphatic sac. *J Histochem Cytochem* 52: 1377–1384, 2004.
47. Duan D, Winter C, Cowley S, Hume JR, Horowitz B. Molecular identification of a volume-regulated chloride channel. *Nature* 390: 417–421, 1997.
48. Dulon D, Sugawara M, Blanchet C, Erostequi C. Direct measurements of Ca²⁺-activated K⁺ currents in inner hair cells of the guinea-pig cochlea using photolabile Ca²⁺ chelators. *Pflügers Arch* 430: 365–373, 1995.
49. Dumont RA, Lins U, Filoteo AG, Penniston JT, Kachar B, Gillespie PG. Plasma membrane Ca²⁺-ATPase isoform 2a is the PMCA of hair bundles. *J Neurosci* 21: 5066–5078, 2001.
50. Embark HM, Bohmer C, Vallon V, Luft F, Lang F. Regulation of KCNE1-dependent K⁺ current by the serum and glucocorticoid-inducible kinase (SGK) isoforms. *Pflügers Arch* 445: 601–606, 2003.
51. Emmons C, Kurtz I. Functional characterization of three intercalated cell subtypes in the rabbit outer cortical collecting duct. *J Clin Invest* 93: 417–423, 1994.
52. Engel J, Braig C, Ruttiger L, Kuhn S, Zimmermann U, Blin N, Saubier M, Kalbacher H, Munkner S, Rohbock K, Ruth P, Winter H, Knipper M. Two classes of outer hair cells along the tonotopic axis of the cochlea. *Neuroscience* 143: 837–849, 2006.
53. Estevez R, Boettger T, Stein V, Birkenhager R, Otto E, Hildebrandt F, Jentsch TJ. Barttin is a Cl⁻ channel beta-subunit crucial for renal Cl⁻ reabsorption and inner ear K⁺ secretion. *Nature* 414: 558–561, 2001.
54. Everett LA, Morsli H, Wu DK, Green ED. Expression pattern of the mouse ortholog of the Pendred's syndrome gene (Pds) suggests a key role for pendrin in the inner ear. *Proc Natl Acad Sci USA* 96: 9727–9732, 1999.
55. Eybalin M, Norenberg MD, Renard N. Glutamine synthetase and glutamate metabolism in the guinea pig cochlea. *Hear Res* 101: 93–101, 1996.
56. Finberg KE, Wagner CA, Bailey MA, Paunescu TG, Breton S, Brown D, Giebisch G, Geibel JP, Lifton RP. The B1-subunit of the H⁺-ATPase is required for maximal urinary acidification. *Proc Natl Acad Sci USA* 102: 13616–13621, 2005.
57. Flagella M, Clarke LL, Miller ML, Erway LC, Giannella RA, Andringa A, Gawenis LR, Kramer J, Duffy JJ, Doetschman T, Lorenz JN, Yamoah EN, Cardell EL, Shull GE. Mice lacking the basolateral Na-K-2Cl cotransporter have impaired epithelial chloride secretion and are profoundly deaf. *J Biol Chem* 274: 26946–26955, 1999.
58. Florian P, Amasheh S, Lessidrensky M, Todt I, Bloedow A, Ernst A, Fromm M, Gitter AH. Claudins in the tight junctions of stria vascularis marginal cells. *Biochem Biophys Res Commun* 304: 5–10, 2003.
59. Forge A, Becker D, Casalotti S, Edwards J, Marziano N, Nevill G. Gap junctions in the inner ear: comparison of distribution patterns in different vertebrates and assessment of connexin composition in mammals. *J Comp Neurol* 467: 207–231, 2003.
60. Frey A, Lampert A, Waldegger S, Jeck N, Waldegger P, Artunc F, Seeböhm G, Lang UE, Kupka S, Pfister M, Hoppe J, Gerloff C, Schaeffeler E, Schwab M, Lang F. Influence of gain of function epithelial chloride channel ClC-Kb mutation on hearing thresholds. *Hear Res* 214: 68–75, 2006.
61. Friedmann I, Fraser GR, Froggatt P. Pathology of the ear in the cardioauditory syndrome of Jervell and Lange-Nielsen (recessive deafness with electrocardiographic abnormalities). *J Laryngol Otol* 80: 451–470, 1966.
62. Frische S, Kwon TH, Frokiaer J, Madsen KM, Nielsen S. Regulated expression of pendrin in rat kidney in response to chronic NH₄Cl or NaHCO₃ loading. *Am J Physiol Renal Physiol* 284: F584–F593, 2003.
63. Fu Y, Subramanya A, Rozansky D, Cohen DM. WNK kinases influence TRPV4 channel function and localization. *Am J Physiol Renal Physiol* 290: F1305–F1314, 2006.
64. Furuta H, Luo L, Hepler K, Ryan AF. Evidence for differential regulation of calcium by outer versus inner hair cells: plasma membrane Ca-ATPase gene expression. *Hear Res* 123: 10–26, 1998.
65. Gamp AC, Tanaka Y, Lullmann-Rauch R, Wittke D, D'Hooge R, De Deyn PP, Moser T, Maier H, Hartmann D, Reiss K, Illert AL, von Figura K, Saftig P. LIMP-2/LGP85 deficiency causes ureteric pelvic

- junction obstruction, deafness and peripheral neuropathy in mice. *Hum Mol Genet* 12: 631–646, 2003.
66. **Gimenez I.** Molecular mechanisms and regulation of furosemide-sensitive Na-K-Cl cotransporters. *Curr Opin Nephrol Hypertens* 15: 517–523, 2006.
 67. **Gow A, Davies C, Southwood CM, Frolenkov G, Chrustowski M, Ng L, Yamauchi D, Marcus DC, Kachar B.** Deafness in claudin 11-null mice reveals the critical contribution of basal cell tight junctions to stria vascularis function. *J Neurosci* 24: 7051–7062, 2004.
 68. **Grifa A, Wagner CA, D'Ambrosio L, Melchionda S, Bernardi F, Lopez-Bigas N, Rabionet R, Arbones M, Monica MD, Estivill X, Zelante L, Lang F, Gasparini P.** Mutations in GJB6 cause nonsyndromic autosomal dominant deafness at DFNA3 locus. *Nat Genet* 23: 16–18, 1999.
 69. **Grunder S, Firsov D, Chang SS, Jaeger NF, Gautschi I, Schild L, Lifton RP, Rossier BC.** A mutation causing pseudohypoaldosteronism type 1 identifies a conserved glycine that is involved in the gating of the epithelial sodium channel. *EMBO J* 16: 899–907, 1997.
 70. **Grunder S, Muller A, Ruppertsberg JP.** Developmental and cellular expression pattern of epithelial sodium channel alpha, beta and gamma subunits in the inner ear of the rat. *Eur J Neurosci* 13: 641–648, 2001.
 71. **Grunnet M, Hay-Schmidt A, Klaerke DA.** Quantification and distribution of big conductance Ca²⁺-activated K⁺ channels in kidney epithelia. *Biochim Biophys Acta* 1714: 114–124, 2005.
 72. **Guipponi M, Vuagniaux G, Wattenhofer M, Shibuya K, Vazquez M, Dougherty L, Scamuffa N, Guida E, Okui M, Rossier C, Hancock M, Buchet K, Reymond A, Hummler E, Marzella PL, Kudoh J, Shimizu N, Scott HS, Antonarakis SE, Rossier BC.** The transmembrane serine protease (TMPRSS3) mutated in deafness DFNB8/10 activates the epithelial sodium channel (ENaC) in vitro. *Hum Mol Genet* 11: 2829–2836, 2002.
 73. **Haack B, Schmalisch K, Palmada M, Bohmer C, Kohlschmidt N, Keilmann A, Zechner U, Limberger A, Beckert S, Zenner HP, Lang F, Kupka S.** Deficient membrane integration of the novel p.N14D-GJB2 mutant associated with non-syndromic hearing impairment. *Hum Mutat* 27: 1158–1159, 2006.
 74. **Hackney CM, Mahendrasingam S, Penn A, Fettiplace R.** The concentrations of calcium buffering proteins in mammalian cochlear hair cells. *J Neurosci* 25: 7867–7875, 2005.
 75. **Haefliger JA, Krattinger N, Martin D, Pedrazzini T, Capponi A, Doring B, Plum A, Charollais A, Willecke K, Meda P.** Connexin43-dependent mechanism modulates renin secretion and hypertension. *J Clin Invest* 116: 405–413, 2006.
 76. **Hakuba N, Koga K, Gyo K, Usami SI, Tanaka K.** Exacerbation of noise-induced hearing loss in mice lacking the glutamate transporter GLAST. *J Neurosci* 20: 8750–8753, 2000.
 77. **Hansson JH, Nelson-Williams C, Suzuki H, Schild L, Shimkets R, Lu Y, Canessa C, Iwasaki T, Rossier B, Lifton RP.** Hypertension caused by a truncated epithelial sodium channel gamma subunit: genetic heterogeneity of Liddle syndrome. *Nat Genet* 11: 76–82, 1995.
 78. **Hara-Chikuma M, Verkman AS.** Aquaporin-1 facilitates epithelial cell migration in kidney proximal tubule. *J Am Soc Nephrol* 17: 39–45, 2006.
 79. **Hartmann R, Gerlach U, Klinke R.** Ototoxic side-effects of the I_{Ks}-channel blocker HMR1556. *Hear Res* 172: 145–150, 2002.
 80. **Hebert SC.** Bartter syndrome. *Curr Opin Nephrol Hypertens* 12: 527–532, 2003.
 81. **Heidland A, Wigand ME.** [Hearing loss induced by the use of high doses of furosemide in the treatment of uremia]. *Klin Wochenschr* 48: 1052–1056, 1970.
 82. **Hibino H, Higashi-Shingai K, Fujita A, Iwai K, Ishii M, Kurachi Y.** Expression of an inwardly rectifying K⁺ channel, Kir5.1, in specific types of fibrocytes in the cochlear lateral wall suggests its functional importance in the establishment of endocochlear potential. *Eur J Neurosci* 19: 76–84, 2004.
 83. **Hibino H, Horio Y, Inanobe A, Doi K, Ito M, Yamada M, Gotow T, Uchiyama Y, Kawamura M, Kubo T, Kurachi Y.** An ATP-dependent inwardly rectifying potassium channel, KAB-2 (Kir4.1), in cochlear stria vascularis of inner ear: its specific subcellular localization and correlation with the formation of endocochlear potential. *J Neurosci* 17: 4711–4721, 1997.
 - 83a. **Highstein SM, Fay RR, Popper AN** (editors). *The Vestibular System*. Berlin: Springer, 2004.
 84. **Hill JK, Williams DE, LeMasurier M, Dumont RA, Strehler EE, Gillespie PG.** Splice-site A choice targets plasma-membrane Ca²⁺-ATPase isoform 2 to hair bundles. *J Neurosci* 26: 6172–6180, 2006.
 85. **Hoenderop JG, Nilius B, Bindels RJ.** Molecular mechanism of active Ca²⁺ reabsorption in the distal nephron. *Annu Rev Physiol* 64: 529–549, 2002.
 86. **Hoenderop JG, van Leeuwen JP, van der Eerden BC, Kersten FF, van der Kemp AW, Merillat AM, Waarsing JH, Rossier BC, Vallon V, Hummler E, Bindels RJ.** Renal Ca²⁺ wasting, hyperabsorption, and reduced bone thickness in mice lacking TRPV5. *J Clin Invest* 112: 1906–1914, 2003.
 87. **Housley GD, Ashmore JF.** Ionic currents of outer hair cells isolated from the guinea-pig cochlea. *J Physiol* 448: 73–98, 1992.
 88. **Huang D, Chen P, Chen S, Nagura M, Lim DJ, Lin X.** Expression patterns of aquaporins in the inner ear: evidence for concerted actions of multiple types of aquaporins to facilitate water transport in the cochlea. *Hear Res* 165: 85–95, 2002.
 89. **Hulterantz M, Bagger-Sjoberg D, Barbara M.** Presence of glycosaminoglycans in the endolymphatic sac. *Acta Otolaryngol (Stockh)* 117: 518–522, 1997.
 90. **Hummler E, Vallon V.** Lessons from mouse mutants of epithelial sodium channel and its regulatory proteins. *J Am Soc Nephrol* 16: 3160–3166, 2005.
 91. **Ikeda K, Kusakari J, Takasaka T, Saito Y.** The Ca²⁺ activity of cochlear endolymph of the guinea pig and the effect of inhibitors. *Hear Res* 26: 117–125, 1987.
 92. **Ikeda K, Morizono T.** Calcium transport mechanism in the endolymph of the chinchilla. *Hear Res* 34: 307–311, 1988.
 93. **Ikeda K, Morizono T.** Effects of carbon dioxide in the middle ear cavity upon the cochlear potentials and cochlear pH. *Acta Otolaryngol (Stockh)* 108: 88–93, 1989.
 94. **Ikeda K, Morizono T.** The preparation of acetic acid for use in otic drops and its effect on endocochlear potential and pH in inner ear fluid. *Am J Otolaryngol* 10: 382–385, 1989.
 95. **Ikeda K, Morizono T.** The ionic and electric environment in the endolymphatic sac of the chinchilla: relevance to the longitudinal flow. *Hear Res* 54: 118–122, 1991.
 96. **Ikeda M, Watanabe I.** Evaluation of hyperactive caloric responses in patients with inner ear diseases. *ORL J Otorhinolaryngol Relat Spec* 59: 326–331, 1997.
 97. **Ito M, Inanobe A, Horio Y, Hibino H, Isomoto S, Ito H, Mori K, Tonosaki A, Tomoike H, Kurachi Y.** Immunolocalization of an inwardly rectifying K⁺ channel, K(AB)-2 (Kir4.1), in the basolateral membrane of renal distal tubular epithelia. *FEBS Lett* 388: 11–15, 1996.
 98. **Jagger DJ, Forge A.** Compartmentalized and signal-selective gap junctional coupling in the hearing cochlea. *J Neurosci* 26: 1260–1268, 2006.
 99. **Jahnke K.** The fine structure of freeze-fractured intercellular junctions in the guinea pig inner ear. *Acta Otolaryngol Suppl (Stockh)* 336: 1–40, 1975.
 100. **Jeck N, Waldegger P, Doroszewicz J, Seyberth H, Waldegger S.** A common sequence variation of the CLCNKB gene strongly activates CIC-Kb chloride channel activity. *Kidney Int* 65: 190–197, 2004.
 101. **Jeck N, Waldegger S, Lampert A, Boehmer C, Waldegger P, Lang PA, Wissinger B, Friedrich B, Rislter T, Moehle R, Lang UE, Zill P, Bondy B, Schaeffeler E, Asante-Poku S, Seyberth H, Schwab M, Lang F.** Activating mutation of the renal epithelial chloride channel CIC-Kb predisposing to hypertension. *Hypertension* 43: 1175–1181, 2004.
 102. **Jentsch TJ.** Chloride transport in the kidney: lessons from human disease and knockout mice. *J Am Soc Nephrol* 16: 1549–1561, 2005.
 103. **Jervell A Lange-Nielsen F.** Congenital deaf-mutism, functional heart disease with prolongation of the Q-T interval and sudden death. *Am Heart J* 54: 59–68, 1957.
 104. **Johnstone BM, Patuzzi R, Syka J, Sykova E.** Stimulus-related potassium changes in the organ of Corti of guinea-pig. *J Physiol* 408: 77–92, 1989.
 105. **Karet FE, Finberg KE, Nelson RD, Nayir A, Mocan H, Sanjad SA, Rodriguez-Soriano J, Santos F, Cremers CW, Di Pietro A, Hoffbrand BI, Winiarski J, Bakkaloglu A, Ozen S, Dusunsel R, Goodyer P, Hulton SA, Wu DK, Skvorak AB, Morton CC, Cunningham MJ, Jha V, Lifton RP.** Mutations in the gene encoding B1 subunit of H⁺-ATPase cause renal tubular acidosis with sensorineural deafness. *Nat Genet* 21: 84–90, 1999.

106. Kawamata S, Igarashi Y. Growth and turnover of rat otoconia as revealed by labeling with tetracycline. *Anat Rec* 242: 259–266, 1995.
107. Kharkovets T, Dedek K, Maier H, Schweizer M, Khimich D, Nouvian R, Vardanyan V, Leuwer R, Moser T, Jentsch TJ. Mice with altered KCNQ4 K⁺ channels implicate sensory outer hair cells in human progressive deafness. *EMBO J* 25: 642–652, 2006.
108. Kikuchi T, Adams JC, Paul DL, Kimura RS. Gap junction systems in the rat vestibular labyrinth: immunohistochemical and ultrastructural analysis. *Acta Otolaryngol (Stockh)* 114: 520–528, 1994.
109. Kikuchi T, Kimura RS, Paul DL, Adams JC. Gap junctions in the rat cochlea: immunohistochemical and ultrastructural analysis. *Anat Embryol (Berl)* 191: 101–118, 1995.
110. Kikuchi T, Kimura RS, Paul DL, Takasaka T, Adams JC. Gap junction systems in the mammalian cochlea. *Brain Res Brain Res Rev* 32: 163–166, 2000.
111. Kim J, Kim YH, Cha JH, Tisher CC, Madsen KM. Intercalated cell subtypes in connecting tubule and cortical collecting duct of rat and mouse. *J Am Soc Nephrol* 10: 1–12, 1999.
112. Kim YH, Kwon TH, Frische S, Kim J, Tisher CC, Madsen KM, Nielsen S. Immunocytochemical localization of pendrin in intercalated cell subtypes in rat and mouse kidney. *Am J Physiol Renal Physiol* 283: F744–F754, 2002.
113. King LS, Choi M, Fernandez PC, Cartron JP, Agre P. Defective urinary-concentrating ability due to a complete deficiency of aquaporin-1. *N Engl J Med* 345: 175–179, 2001.
114. Kitajiri SI, Furuse M, Morita K, Saishin-Kiuchi Y, Kido H, Ito J, Tsukita S. Expression patterns of claudins, tight junction adhesion molecules, in the inner ear. *Hear Res* 187: 25–34, 2004.
115. Knipper M, Claussen C, Ruttiger L, Zimmermann U, Lullmann-Rauch R, Eskelinen EL, Schroder J, Schwake M, Saftig P. Deafness in LIMP2-deficient mice due to early loss of the potassium channel KCNQ1/KCNE1 in marginal cells of the stria vascularis. *J Physiol* 576: 73–86, 2006.
116. Kofuji P, Biedermann B, Siddharthan V, Raap M, Iandiev I, Milenkovic I, Thomzig A, Veh RW, Bringmann A, Reichenbach A. Kir potassium channel subunit expression in retinal glial cells: implications for spatial potassium buffering. *Glia* 39: 292–303, 2002.
117. Kokubo Y, Iwai N, Tago N, Inamoto N, Okayama A, Yamawaki H, Naraba H, Tomoike H. Association analysis between hypertension and CYBA, CLCNKB, and KCNMB1 functional polymorphisms in the Japanese population—the Suita study. *Circ J* 69: 138–142, 2005.
118. Konishi T, Mendelsohn M. Effect of ouabain on cochlear potentials and endolymph composition in guinea pigs. *Acta Otolaryngol (Stockh)* 69: 192–199, 1970.
119. Konrad M, Vollmer M, Lemmink HH, van den Heuvel LP, Jeck N, Vargas-Poussou R, Lakings A, Ruf R, Deschenes G, Antignac C, Guay-Woodford L, Knoers NV, Seyberth HW, Feldmann D, Hildebrandt F. Mutations in the chloride channel gene CLCNKB as a cause of classic Bartter syndrome. *J Am Soc Nephrol* 11: 1449–1459, 2000.
120. Kozel PJ, Friedman RA, Erway LC, Yamoah EN, Liu LH, Riddle T, Duffy JJ, Doetschman T, Miller ML, Cardell EL, Shull GE. Balance and hearing deficits in mice with a null mutation in the gene encoding plasma membrane Ca²⁺-ATPase isoform 2. *J Biol Chem* 273: 18693–18696, 1998.
121. Kros CJ, Crawford AC. Potassium currents in inner hair cells isolated from the guinea-pig cochlea. *J Physiol* 421: 263–291, 1990.
122. Kros CJ, Ruppertsberg JP, Rusch A. Expression of a potassium current in inner hair cells during development of hearing in mice. *Nature* 394: 281–284, 1998.
123. Kubisch C, Schroeder BC, Friedrich T, Lutjohann B, El Amraoui A, Marlin S, Petit C, Jentsch TJ. KCNQ4, a novel potassium channel expressed in sensory outer hair cells, is mutated in dominant deafness. *Cell* 96: 437–446, 1999.
124. Kudo T, Kure S, Ikeda K, Xia AP, Katori Y, Suzuki M, Kojima K, Ichinohe A, Suzuki Y, Aoki Y, Kobayashi T, Matsubara Y. Transgenic expression of a dominant-negative connexin26 causes degeneration of the organ of Corti and non-syndromic deafness. *Hum Mol Genet* 12: 995–1004, 2003.
125. Kuhn K, Reale E. Junctional complexes of the tubular cells in the human kidney as revealed with freeze-fracture. *Cell Tissue Res* 160: 193–205, 1975.
126. Kuijpers W, Bonting SL. The cochlear potentials. I. The effect of ouabain on the cochlear potentials of the guinea pig. *Pflügers Arch* 320: 348–358, 1970.
127. Kuronita T, Eskelinen EL, Fujita H, Saftig P, Himeno M, Tanaka Y. A role for the lysosomal membrane protein LGP85 in the biogenesis and maintenance of endosomal and lysosomal morphology. *J Cell Sci* 115: 4117–4131, 2002.
128. Kusakari J, Ise I, Comegys TH, Thalmann I, Thalmann R. Effect of ethacrynic acid, furosemide, and ouabain upon the endolymphatic potential and upon high energy phosphates of the stria vascularis. *Laryngoscope* 88: 12–37, 1978.
129. Kusakari J, Kambayashi J, Ise I, Kawamoto K. Reduction of the endocochlear potential by the new “loop” diuretic, bumetanide. *Acta Otolaryngol (Stockh)* 86: 336–341, 1978.
130. Kusakari J, Thalmann R. Effects of anoxia and ethacrynic acid upon ampullar endolymphatic potential and upon high energy phosphates in ampullar wall. *Laryngoscope* 86: 132–147, 1976.
131. Lang F, Busch GL, Ritter M, Volkl H, Waldegger S, Gulbins E, Haussinger D. Functional significance of cell volume regulatory mechanisms. *Physiol Rev* 78: 247–306, 1998.
132. Lang F, Oberleithner H, Giebisch G. Electrophysiological heterogeneity of proximal convoluted tubules in *Amphiuma* kidney. *Am J Physiol Renal Fluid Electrolyte Physiol* 251: F1063–F1072, 1986.
133. Lang F, Rehwald W. Potassium channels in renal epithelial transport regulation. *Physiol Rev* 72: 1–32, 1992.
134. Lautermann J, ten Cate WJ, Altenhoff P, Grummer R, Traub O, Frank H, Jahnke K, Winterhager E. Expression of the gap-junction connexins 26 and 30 in the rat cochlea. *Cell Tissue Res* 294: 415–420, 1998.
135. Lecain E, Robert JC, Thomas A, Tran Ba HP. Gastric proton pump is expressed in the inner ear and choroid plexus of the rat. *Hear Res* 149: 147–154, 2000.
136. Lee JH, Chiba T, Marcus DC. P2X2 receptor mediates stimulation of parasympathetic cation absorption by cochlear outer sulcus cells and vestibular transitional cells. *J Neurosci* 21: 9168–9174, 2001.
137. Lee JH, Marcus DC. Nongenomic effects of corticosteroids on ion transport by stria vascularis. *Audiol Neurootol* 7: 100–106, 2002.
138. Lee JH, Marcus DC. Endolymphatic sodium homeostasis by Reissner’s membrane. *Neuroscience* 119: 3–8, 2003.
139. Lee MP, Ravenel JD, Hu RJ, Lustig LR, Tomaselli G, Berger RD, Brandenburg SA, Litz T, Bunton TE, Limb C, Francis H, Gorelikow M, Gu H, Washington K, Argani P, Goldenring JR, Coffey RJ, Feinberg AP. Targeted disruption of the *Kvlqt1* gene causes deafness and gastric hyperplasia in mice. *J Clin Invest* 106: 1447–1455, 2000.
140. Letts VA, Valenzuela A, Dunbar C, Zheng QY, Johnson KR, Frankel WN. A new spontaneous mouse mutation in the *Kcne1* gene. *Mamm Genome* 11: 831–835, 2000.
141. Li J, Verkman AS. Impaired hearing in mice lacking aquaporin-4 water channels. *J Biol Chem* 276: 31233–31237, 2001.
142. Liang G, Moore EJ, Ulfendahl M, Rydqvist B, Jarlebark L. An M-like potassium current in the guinea pig cochlea. *ORL J Otorhinolaryngol Relat Spec* 67: 75–82, 2005.
143. Liang GH, Jin Z, Ulfendahl M, Jarlebark L. Molecular analyses of KCNQ1–5 potassium channel mRNAs in rat and guinea pig inner ears: expression, cloning, and alternative splicing. *Acta Otolaryngol (Stockh)* 126: 346–352, 2006.
144. Liedtke W, Choe Y, Marti-Renom MA, Bell AM, Denis CS, Sali A, Hudspeth AJ, Friedman JM, Heller S. Vanilloid receptor-related osmotically activated channel (VR-OAC), a candidate vertebrate osmoreceptor. *Cell* 103: 525–535, 2000.
145. Lifton RP. Molecular genetics of human blood pressure variation. *Science* 272: 676–680, 1996.
146. Lim DJ, Karabinas C, Trune DR. Histochemical localization of carbonic anhydrase in the inner ear. *Am J Otolaryngol* 4: 33–42, 1983.
147. Liu XZ, Xia XJ, Adams J, Chen ZY, Welch KO, Tekin M, Ouyang XM, Kristiansen A, Pandya A, Balkany T, Arnos KS, Nance WE. Mutations in *GJA1* (connexin 43) are associated with non-syndromic autosomal recessive deafness. *Hum Mol Genet* 10: 2945–2951, 2001.
148. Liu XZ, Xia XJ, Xu LR, Pandya A, Liang CY, Blanton SH, Brown SD, Steel KP, Nance WE. Mutations in connexin31 underlie recessive as well as dominant non-syndromic hearing loss. *Hum Mol Genet* 9: 63–67, 2000.
149. Lopez IA, Ishiyama G, Lee M, Baloh RW, Ishiyama A. Immunohistochemical localization of aquaporins in the human inner ear. *Cell Tissue Res* 328: 453–460, 2007.

150. Lopez-Bigas N, Arbones ML, Estivill X, Simonneau L. Expression profiles of the connexin genes, Gjb1 and Gjb3, in the developing mouse cochlea. *Mech Dev* 119, Suppl 1: S111–S115, 2002.
151. Lourdel S, Paulais M, Cluzeaud F, Bens M, Tanemoto M, Kurachi Y, Vandewalle A, Teulon J. An inward rectifier K⁺ channel at the basolateral membrane of the mouse distal convoluted tubule: similarities with Kir4-Kir5.1 heteromeric channels. *J Physiol* 538: 391–404, 2002.
152. Lundberg YW, Zhao X, Yamoah EN. Assembly of the otoconia complex to the macular sensory epithelium of the vestibule. *Brain Res* 1091: 47–57, 2006.
153. Ma T, Yang B, Gillespie A, Carlson EJ, Epstein CJ, Verkman AS. Severely impaired urinary concentrating ability in transgenic mice lacking aquaporin-1 water channels. *J Biol Chem* 273: 4296–4299, 1998.
154. Maehara H, Okamura HO, Kobayashi K, Uchida S, Sasaki S, Kitamura K. Expression of CLC-KB gene promoter in the mouse cochlea. *Neuroreport* 14: 1571–1573, 2003.
155. Mammamo F, Ashmore JF. Differential expression of outer hair cell potassium currents in the isolated cochlea of the guinea-pig. *J Physiol* 496: 639–646, 1996.
156. Marcotti W, Johnson SL, Holley MC, Kros CJ. Developmental changes in the expression of potassium currents of embryonic, neonatal and mature mouse inner hair cells. *J Physiol* 548: 383–400, 2003.
157. Marcotti W, Johnson SL, Rusch A, Kros CJ. Sodium and calcium currents shape action potentials in immature mouse inner hair cells. *J Physiol* 552: 743–761, 2003.
158. Marcotti W, Kros CJ. Developmental expression of the potassium current $I_{K,n}$ contributes to maturation of mouse outer hair cells. *J Physiol* 520: 653–660, 1999.
159. Marcus DC, Chiba T. K⁺ and Na⁺ absorption by outer sulcus epithelial cells. *Hear Res* 134: 48–56, 1999.
160. Marcus DC, Liu J, Lee JH, Scherer EQ, Scofield MA, Wangemann P. Apical membrane P2Y4 purinergic receptor controls K⁺ secretion by stria marginal cell epithelium. *Cell Commun Signal* 3: 13, 2005.
161. Marcus DC, Liu J, Wangemann P. Transepithelial voltage and resistance of vestibular dark cell epithelium from the gerbil ampulla. *Hear Res* 73: 101–108, 1994.
162. Marcus DC, Marcus NY, Greger R. Sidedness of action of loop diuretics and ouabain on nonsensory cells of utricle: a micro-Ussing chamber for inner ear tissues. *Hear Res* 30: 55–64, 1987.
163. Marcus DC, Pondugula SR, Lee JH, Chiba T, Yamauchi D, Sanne-man J, Harbidge DG, Kampalli SB, Wu T, Wangemann P. Ion transport pathways in the cochlea and vestibular labyrinth. *Proc Meniere's Meeting* 31–35, 2005.
164. Marcus DC, Rokugo M, Thalmann R. Effects of barium and ion substitutions in artificial blood on endocochlear potential. *Hear Res* 17: 79–86, 1985.
165. Marcus DC, Shen Z. Slowly activating voltage-dependent K⁺ conductance is apical pathway for K⁺ secretion in vestibular dark cells. *Am J Physiol Cell Physiol* 267: C857–C864, 1994.
166. Marcus DC, Sunose H, Liu J, Shen Z, Scofield MA. P2U purinergic receptor inhibits apical IsK/KvLQT1 channel via protein kinase C in vestibular dark cells. *Am J Physiol Cell Physiol* 273: C2022–C2029, 1997.
167. Marcus DC, Takeuchi S, Wangemann P. Two types of chloride channel in the basolateral membrane of vestibular dark cells. *Hear Res* 69: 124–132, 1993.
168. Marcus DC, Thalmann R, Marcus NY. Respiratory rate and ATP content of stria vascularis of guinea pig in vitro. *Laryngoscope* 88: 1825–1835, 1978.
169. Marcus DC, Wu T, Wangemann P, Kofuji P. KCNJ10 (Kir4.1) potassium channel knockout abolishes endocochlear potential. *Am J Physiol Cell Physiol* 282: C403–C407, 2002.
170. Marcus NY, Marcus DC. Potassium secretion by nonsensory region of gerbil utricle in vitro. *Am J Physiol Renal Physiol* 253: F613–F621, 1987.
171. Matsumura Y, Uchida S, Kondo Y, Miyazaki H, Ko SB, Hayama A, Morimoto T, Liu W, Arisawa M, Sasaki S, Marumo F. Overt nephrogenic diabetes insipidus in mice lacking the CLC-K1 chloride channel. *Nat Genet* 21: 95–98, 1999.
172. McCulloch F, Chambrey R, Eladari D, Peti-Peterdi J. Localization of connexin 30 in the luminal membrane of cells in the distal nephron. *Am J Physiol Renal Physiol* 289: F1304–F1312, 2005.
173. McGuirt JP, Schulte BA. Distribution of immunoreactive alpha- and beta-subunit isoforms of Na,K-ATPase in the gerbil inner ear. *J Histochem Cytochem* 42: 843–853, 1994.
174. Mercado F, Lopez IA, Acuna D, Vega R, Soto E. Acid-sensing ionic channels in the rat vestibular endorgans and ganglia. *J Neurophysiol* 96: 1615–1624, 2006.
175. Mhatre AN, Steinbach S, Hribar K, Hoque AT, Lalwani AK. Identification of aquaporin 5 (AQP5) within the cochlea: cDNA cloning and in situ localization. *Biochem Biophys Res Commun* 264: 157–162, 1999.
176. Mhatre AN, Stern RE, Li J, Lalwani AK. Aquaporin 4 expression in the mammalian inner ear and its role in hearing. *Biochem Biophys Res Commun* 297: 987–996, 2002.
177. Mizuta K, Adachi M, Iwasa KH. Ultrastructural localization of the Na-K-Cl cotransporter in the lateral wall of the rabbit cochlear duct. *Hear Res* 106: 154–162, 1997.
178. Mori N, Ninoyu O, Morgenstern C. Cation transport in the ampulla of the semicircular canal and in the endolymphatic sac. *Arch Otorhinolaryngol* 244: 61–65, 1987.
179. Najjar F, Zhou H, Morimoto T, Bruns JB, Li HS, Liu W, Kleyman TR, Satlin LM. Dietary K⁺ regulates apical membrane expression of maxi-K channels in rabbit cortical collecting duct. *Am J Physiol Renal Physiol* 289: F922–F932, 2005.
180. Nakaya K, Harbidge DG, Wangemann P, Schultz BD, Green E, Wall SM, Marcus DC. Lack of pendrin HCO₃⁻ transport elevates vestibular endolymphatic [Ca²⁺] by inhibition of acid-sensitive TRPV5 and TRPV6 channels. *Am J Physiol Renal Physiol* 292: F1314–F1321, 2007.
181. Nenov AP, Chen C, Bobbin RP. Outward rectifying potassium currents are the dominant voltage activated currents present in Deiters' cells. *Hear Res* 123: 168–182, 1998.
182. Neyroud N, Tesson F, Denjoy I, Leibovici M, Donger C, Barhanin J, Faure S, Gary F, Coumel P, Petit C, Schwartz K, Guicheney P. A novel mutation in the potassium channel gene KVLQT1 causes the Jervell and Lange-Nielsen cardioauditory syndrome. *Nat Genet* 15: 186–189, 1997.
183. Nielsen S, Frokiaer J, Marples D, Kwon TH, Agre P, Knepper MA. Aquaporins in the kidney: from molecules to medicine. *Physiol Rev* 82: 205–244, 2002.
184. Nielsen S, Kwon TH, Frokiaer J, Agre P. Regulation and dysregulation of aquaporins in water balance disorders. *J Intern Med* 261: 53–64, 2007.
185. Ninoyu O, Meyer zum Gotesberge AM. Calcium transport in the endolymphatic space of cochlea and vestibular organ. *Acta Otolaryngol (Stockh)* 102: 222–227, 1986.
186. Oesterle EC, Dallos P. Intracellular recordings from supporting cells in the guinea pig cochlea: DC potentials. *J Neurophysiol* 64: 617–636, 1990.
187. Ohmori H. Mechano-electrical transduction currents in isolated vestibular hair cells of the chick. *J Physiol* 359: 189–217, 1985.
188. Okamura HO, Sugai N, Suzuki K, Ohtani I. Enzyme-histochemical localization of carbonic anhydrase in the inner ear of the guinea pig and several improvements of the technique. *Histochem Cell Biol* 106: 425–430, 1996.
189. Oliver D, Knipper M, Derst C, Fakler B. Resting potential and submembrane calcium concentration of inner hair cells in the isolated mouse cochlea are set by KCNQ-type potassium channels. *J Neurosci* 23: 2141–2149, 2003.
190. Oliver D, Taberner AM, Thurm H, Saubier M, Arntz C, Ruth P, Fakler B, Liberman MC. The role of BKCa channels in electrical signal encoding in the mammalian auditory periphery. *J Neurosci* 26: 6181–6189, 2006.
191. Orzan E, Murgia A. Connexin 26 deafness is not always congenital. *Int J Pediatr Otorhinolaryngol* 71: 501–507, 2007.
192. Pallone TL, Edwards A, Ma T, Silldorff EP, Verkman AS. Requirement of aquaporin-1 for NaCl-driven water transport across descending vasa recta. *J Clin Invest* 105: 215–222, 2000.
193. Palmada M, Schmalisch K, Bohmer C, Schug N, Pfister M, Lang F, Blin N. Loss of function mutations of the GJB2 gene detected in patients with DFNB1-associated hearing impairment. *Neurobiol Dis* 22: 112–118, 2006.
194. Peng BG, Ahmad S, Chen S, Chen P, Price MP, Lin X. Acid-sensing ion channel 2 contributes a major component to acid-evoked excitatory responses in spiral ganglion neurons and plays a role in noise susceptibility of mice. *J Neurosci* 24: 10167–10175, 2004.

195. Peng JB, Hediger MA. A family of calcium-permeable channels in the kidney: distinct roles in renal calcium handling. *Curr Opin Nephrol Hypertens* 11: 555–561, 2002.
196. Peters TA, Monnens LA, Cremers CW, Curfs JH. Genetic disorders of transporters/channels in the inner ear and their relation to the kidney. *Pediatr Nephrol* 19: 1194–1201, 2004.
197. Petit C, Levilliers J, Hardelin JP. Molecular genetics of hearing loss. *Annu Rev Genet* 35: 589–646, 2001.
198. Pluznick JL, Sansom SC. BK channels in the kidney: role in K⁺ secretion and localization of molecular components. *Am J Physiol Renal Physiol* 291: F517–F529, 2006.
199. Pluznick JL, Wei P, Carmines PK, Sansom SC. Renal fluid and electrolyte handling in BKCa- β 1-/- mice. *Am J Physiol Renal Physiol* 284: F1274–F1279, 2003.
200. Pluznick JL, Wei P, Grimm PR, Sansom SC. BK- β 1 subunit: immunolocalization in the mammalian connecting tubule and its role in the kaliuretic response to volume expansion. *Am J Physiol Renal Physiol* 288: F846–F854, 2005.
201. Pondugula SR, Raveendran NN, Ergonul Z, Deng Y, Chen J, Sanneman JD, Palmer LG, Marcus DC. Glucocorticoid regulation of genes in the amiloride-sensitive sodium transport pathway by semicircular canal duct epithelium of neonatal rat. *Physiol Genomics* 24: 114–123, 2006.
202. Pondugula SR, Sanneman JD, Wangemann P, Milhaud PG, Marcus DC. Glucocorticoids stimulate cation absorption by semicircular canal duct epithelium via epithelial sodium channel. *Am J Physiol Renal Physiol* 286: F1127–F1135, 2004.
203. Porubsky ES, Marovitz WF, Arenberg IK. Presence of acidic protein-bound carbohydrates in the endolymphatic sac and duct of fetal, neonatal and adult rats, and adult humans. *Ann Otol Rhinol Laryngol* 81: 76–81, 1972.
204. Pressler CA, Heinzinger J, Jeck N, Waldegger P, Pechmann U, Reinalter S, Konrad M, Beetz R, Seyberth HW, Waldegger S. Late-onset manifestation of antenatal Bartter syndrome as a result of residual function of the mutated renal Na⁺-K⁺-2Cl⁻ cotransporter. *J Am Soc Nephrol* 17: 2136–2142, 2006.
205. Preston GM, Smith BL, Zeidel ML, Moulds JJ, Agre P. Mutations in aquaporin-1 in phenotypically normal humans without functional CHIP water channels. *Science* 265: 1585–1587, 1994.
206. Qu C, Liang F, Hu W, Shen Z, Spicer SS, Schulte BA. Expression of CLC-K chloride channels in the rat cochlea. *Hear Res* 213: 79–87, 2006.
207. Raybould NP, Jagger DJ, Kanjhan R, Greenwood D, Laslo P, Hoya N, Soeller C, Cannell MB, Housley GD. TRPC-like conductance mediates restoration of intracellular Ca²⁺ in cochlear outer hair cells in the guinea pig and rat. *J Physiol* 579: 101–113, 2007.
208. Reichenbach A, Henke A, Eberhardt W, Reichelt W, Dettmer D. K⁺ ion regulation in retina. *Can J Physiol Pharmacol* 70, Suppl: S239–S247, 1992.
209. Renkema KY, Nijenhuis T, van der Eerden BC, van der Kemp AW, Weinans H, van Leeuwen JP, Bindels RJ, Hoenderop JG. Hypervitaminosis D mediates compensatory Ca²⁺ hyperabsorption in TRPV5 knockout mice. *J Am Soc Nephrol* 16: 3188–3195, 2005.
210. Richard G, White TW, Smith LE, Bailey RA, Compton JG, Paul DL, Bale SJ. Functional defects of Cx26 resulting from a heterozygous missense mutation in a family with dominant deaf-mutism and palmoplantar keratoderma. *Hum Genet* 103: 393–399, 1998.
211. Rieg T, Bunday R, Chen Y, Deschenes G, Junger W, Insel PA, Vallon V. Mice lacking P2Y₂ receptors have salt-resistant arterial hypertension and facilitated renal Na⁺ and fluid reabsorption. *FASEB J*. In press.
212. Rieg T, Vallon V, Sausbier M, Sausbier U, Kaissling B, Ruth P, Osswald H. A role of BK channel in K⁺ homeostasis and flow-induced renal K⁺ excretion. *Kidney Int*. 72: 566–573, 2007.
213. Royaux IE, Belyantseva IA, Wu T, Kachar B, Everett LA, Marcus DC, Green ED. Localization and functional studies of pendrin in the mouse inner ear provide insight about the etiology of deafness in pendred syndrome. *J Assoc Res Otolaryngol* 4: 394–404, 2003.
214. Royaux IE, Wall SM, Karniski LP, Everett LA, Suzuki K, Knepper MA, Green ED. Pendrin, encoded by the Pendred syndrome gene, resides in the apical region of renal intercalated cells and mediates bicarbonate secretion. *Proc Natl Acad Sci USA* 98: 4221–4226, 2001.
215. Rozenfurt N, Lopez I, Chiu CS, Kofuji P, Lester HA, Neusch C. Time course of inner ear degeneration and deafness in mice lacking the Kir4.1 potassium channel subunit. *Hear Res* 177: 71–80, 2003.
216. Ruttiger L, Sausbier M, Zimmermann U, Winter H, Braig C, Engel J, Knirsch M, Arntz C, Langer P, Hirt B, Muller M, Kopschall I, Pfister M, Munkner S, Rohbock K, Pfaff I, Rusch A, Ruth P, Knipper M. Deletion of the Ca²⁺-activated potassium (BK) alpha-subunit but not the BKbeta1-subunit leads to progressive hearing loss. *Proc Natl Acad Sci USA* 101: 12922–12927, 2004.
217. Rybak LP, Whitworth CA. Ototoxicity: therapeutic opportunities. *Drug Discov Today* 10: 1313–1321, 2005.
218. Sage CL, Marcus DC. Immunolocalization of CIC-K chloride channel in strial marginal cells and vestibular dark cells. *Hear Res* 160: 1–9, 2001.
219. Salt AN. Regulation of endolymphatic fluid volume. *Ann NY Acad Sci* 942: 306–312, 2001.
220. Salt AN, Inamura N, Thalmann R, Vora A. Calcium gradients in inner ear endolymph. *Am J Otolaryngol* 10: 371–375, 1989.
221. Salt AN, Melichar I, Thalmann R. Mechanisms of endocochlear potential generation by stria vascularis. *Laryngoscope* 97: 984–991, 1987.
222. Salt AN, Ohyama K. Accumulation of potassium in scala vestibuli perilymph of the mammalian cochlea. *Ann Otol Rhinol Laryngol* 102: 64–70, 1993.
223. Sanguinetti MC, Curran ME, Zou A, Shen J, Spector PS, Atkinson DL, Keating MT. Coassembly of KvLQT1 and minK (IsK) proteins to form cardiac I_{Ks} potassium channel. *Nature* 384: 80–83, 1996.
224. Sawada S, Takeda T, Kitano H, Takeuchi S, Okada T, Ando M, Suzuki M, Kakigi A. Aquaporin-1 (AQP1) is expressed in the stria vascularis of rat cochlea. *Hear Res* 181: 15–19, 2003.
225. Schild L, Lu Y, Gautschi I, Schneeberger E, Lifton RP, Rossier BC. Identification of a PY motif in the epithelial Na channel subunits as a target sequence for mutations causing channel activation found in Liddle syndrome. *EMBO J* 15: 2381–2387, 1996.
226. Schlingmann KP, Konrad M, Jeck N, Waldegger P, Reinalter SC, Holder M, Seyberth HW, Waldegger S. Salt wasting and deafness resulting from mutations in two chloride channels. *N Engl J Med* 350: 1314–1319, 2004.
227. Schnermann J, Chou CL, Ma T, Traynor T, Knepper MA, Verkman AS. Defective proximal tubular fluid reabsorption in transgenic aquaporin-1 null mice. *Proc Natl Acad Sci USA* 95: 9660–9664, 1998.
228. Scholl U, Hebeisen S, Janssen AG, Muller-Newen G, Alekov A, Fahlke C. Barttin modulates trafficking and function of CIC-K channels. *Proc Natl Acad Sci USA* 103: 11411–11416, 2006.
229. Schrier RW. Body water homeostasis: clinical disorders of urinary dilution and concentration. *J Am Soc Nephrol* 17: 1820–1832, 2006.
230. Schulte BA, Adams JC. Distribution of immunoreactive Na⁺,K⁺-ATPase in gerbil cochlea. *J Histochem Cytochem* 37: 127–134, 1989.
231. Schulze-Bahr E, Wang Q, Wedekind H, Haverkamp W, Chen Q, Sun Y, Rubie C, Hordt M, Towbin JA, Borggrefe M, Assmann G, Qu X, Somberg JC, Breithardt G, Oberti C, Funke H. KCNE1 mutations cause jervell and Lange-Nielsen syndrome. *Nat Genet* 17: 267–268, 1997.
232. Schwartz GJ, Al Awqati Q. Role of hensen in mediating the adaptation of the cortical collecting duct to metabolic acidosis. *Curr Opin Nephrol Hypertens* 14: 383–388, 2005.
233. Schwartz PJ, Spazzolini C, Crotti L, Bathen J, Amlie JP, Timothy K, Shkolnikova M, Berul CI, Bitner-Glindzicz M, Toivonen L, Horie M, Schulze-Bahr E, Denjoy I. The Jervell and Lange-Nielsen syndrome: natural history, molecular basis, and clinical outcome. *Circulation* 113: 783–790, 2006.
234. Shen J, Harada N, Kubo N, Liu B, Mizuno A, Suzuki M, Yamashita T. Functional expression of transient receptor potential vanilloid 4 in the mouse cochlea. *Neuroreport* 17: 135–139, 2006.
235. Shibata T, Hibino H, Doi K, Suzuki T, Hisa Y, Kurachi Y. Gastric type H⁺,K⁺-ATPase in the cochlear lateral wall is critically involved in formation of the endocochlear potential. *Am J Physiol Cell Physiol* 291: C1038–C1048, 2006.
236. Silver RB, Soleimani M. H⁺-K⁺-ATPases: regulation and role in pathophysiological states. *Am J Physiol Renal Physiol* 276: F799–F811, 1999.
237. Simon DB, Bindra RS, Mansfield TA, Nelson-Williams C, Mendonca E, Stone R, Schurman S, Nayir A, Alpay H, Bakaloglu A, Rodriguez-Soriano J, Morales JM, Sanjad SA, Taylor CM, Pilz D, Brem A, Trachtman H, Griswold W, Richard GA, John E, Lifton RP. Mutations in the chloride channel gene, CLCNKB, cause Bartter's syndrome type III. *Nat Genet* 17: 171–178, 1997.

238. Simon DB, Karet FE, Hamdan JM, DiPietro A, Sanjad SA, Lifton RP. Bartter's syndrome, hypokalaemic alkalosis with hypercalciuria, is caused by mutations in the Na-K-2Cl cotransporter NKCC2. *Nat Genet* 13: 183–188, 1996.
239. Simon DB, Karet FE, Rodriguez-Soriano J, Hamdan JH, DiPietro A, Trachtman H, Sanjad SA, Lifton RP. Genetic heterogeneity of Bartter's syndrome revealed by mutations in the K⁺ channel, ROMK. *Nat Genet* 14: 152–156, 1996.
240. Skinner LJ, Enee V, Beurg M, Jung HH, Ryan AF, Hafidi A, Aran JM, Dulon D. Contribution of BK Ca²⁺-activated K⁺ channels to auditory neurotransmission in the Guinea pig cochlea. *J Neurophysiol* 90: 320–332, 2003.
241. Smith AN, Skaug J, Choate KA, Nayir A, Bakkaloglu A, Ozen S, Hulton SA, Sanjad SA, Al Sabban EA, Lifton RP, Scherer SW, Karet FE. Mutations in ATP6N1B, encoding a new kidney vacuolar proton pump 116-kD subunit, cause recessive distal renal tubular acidosis with preserved hearing. *Nat Genet* 26: 71–75, 2000.
242. Sahara E, Rai T, Yang SS, Uchida K, Nitta K, Horita S, Ohno M, Harada A, Sasaki S, Uchida S. Pathogenesis and treatment of autosomal-dominant nephrogenic diabetes insipidus caused by an aquaporin 2 mutation. *Proc Natl Acad Sci USA* 103: 14217–14222, 2006.
243. Soleimani M, Greeley T, Petrovic S, Wang Z, Amlal H, Kopp P, Burnham CE. Pendrin: an apical Cl⁻/OH⁻/HCO₃⁻ exchanger in the kidney cortex. *Am J Physiol Renal Physiol* 280: F356–F364, 2001.
244. Speirs HJ, Wang WY, Benjafeld AV, Morris BJ. No association with hypertension of CLCNKB and TNFRSF1B polymorphisms at a hypertension locus on chromosome 1p36. *J Hypertens* 23: 1491–1496, 2005.
245. Spicer SS, Schulte BA. Differentiation of inner ear fibrocytes according to their ion transport related activity. *Hear Res* 56: 53–64, 1991.
246. Spicer SS, Schulte BA. The fine structure of spiral ligament cells relates to ion return to the stria and varies with place-frequency. *Hear Res* 100: 80–100, 1996.
247. Spicer SS, Schulte BA. Evidence for a medial K⁺ recycling pathway from inner hair cells. *Hear Res* 118: 1–12, 1998.
248. Sprenger F. [Experiences with Terbolan in otorhinolaryngological practice]. *Med Welt* 9: 489–490, 1969.
249. Stankovic KM, Adams JC, Brown D. Immunolocalization of aquaporin CHIP in the guinea pig inner ear. *Am J Physiol Cell Physiol* 269: C1450–C1456, 1995.
250. Stankovic KM, Brown D, Alper SL, Adams JC. Localization of pH regulating proteins H⁺ATPase and Cl⁻/HCO₃⁻ exchanger in the guinea pig inner ear. *Hear Res* 114: 21–34, 1997. [Erratum. *Hear Res* 124(1–2): 191–192, 1998.]
251. Stojkovic T, Latour P, Vandenberghe A, Hurtevent JF, Vermersch P. Sensorineural deafness in X-linked Charcot-Marie-Tooth disease with connexin 32 mutation (R142Q). *Neurology* 52: 1010–1014, 1999.
252. Stover EH, Borthwick KJ, Bavalía C, Eady N, Fritz DM, Rungroj N, Giersch AB, Morton CC, Axon PR, Akil I, Al Sabban EA, Baguley DM, Bianca S, Bakkaloglu A, Bircan Z, Chauveau D, Clermont MJ, Guala A, Hulton SA, Kroes H, Li VG, Mir S, Mocan H, Nayir A, Ozen S, Rodriguez SJ, Sanjad SA, Tasic V, Taylor CM, Topaloglu R, Smith AN, Karet FE. Novel ATP6V1B1 and ATP6V0A4 mutations in autosomal recessive distal renal tubular acidosis with new evidence for hearing loss. *J Med Genet* 39: 796–803, 2002.
253. Street VA, McKee-Johnson JW, Fonseca RC, Tempel BL, Noben-Trauth K. Mutations in a plasma membrane Ca²⁺-ATPase gene cause deafness in deafwaddler mice. *Nat Genet* 19: 390–394, 1998.
254. Strotmann R, Harteneck C, Nunnenmacher K, Schultz G, Plant TD. OTRPC4, a nonselective cation channel that confers sensitivity to extracellular osmolarity. *Nat Cell Biol* 2: 695–702, 2000.
255. Strutz-Seeböhm N, Seeböhm G, Fedorenko O, Baltaev R, Engel J, Knirsch M, Lang F. Functional coassembly of KCNQ4 with KCNE-beta-subunits in *Xenopus* oocytes. *Cell Physiol Biochem* 18: 57–66, 2006.
256. Sugimoto T, Tanabe Y, Shigemoto R, Iwai M, Takumi T, Ohkubo H, Nakanishi S. Immunohistochemical study of a rat membrane protein which induces a selective potassium permeation: its localization in the apical membrane portion of epithelial cells. *J Membr Biol* 113: 39–47, 1990.
257. Sun J, Ahmad S, Chen S, Tang W, Zhang Y, Chen P, Lin X. Cochlear gap junctions coassembled from Cx26 and 30 show faster intercellular Ca²⁺ signaling than homomeric counterparts. *Am J Physiol Cell Physiol* 288: C613–C623, 2005.
258. Sunose H, Liu J, Marcus DC. cAMP increases K⁺ secretion via activation of apical IsK/KvLQT1 channels in strial marginal cells. *Hear Res* 114: 107–116, 1997.
259. Sunose H, Liu J, Shen Z, Marcus DC. cAMP increases apical IsK channel current and K⁺ secretion in vestibular dark cells. *J Membr Biol* 156: 25–35, 1997.
260. Suzuki M, Mizuno A, Kodaira K, Imai M. Impaired pressure sensation in mice lacking TRPV4. *J Biol Chem* 278: 22664–22668, 2003.
261. Tabuchi K, Suzuki M, Mizuno A, Hara A. Hearing impairment in TRPV4 knockout mice. *Neurosci Lett* 382: 304–308, 2005.
262. Takeuchi S, Ando M. Dye-coupling of melanocytes with endothelial cells and pericytes in the cochlea of gerbils. *Cell Tissue Res* 293: 271–275, 1998.
263. Takeuchi S, Ando M, Kakigi A. Mechanism generating endocochlear potential: role played by intermediate cells in stria vascularis. *Biophys J* 79: 2572–2582, 2000.
264. Takeuchi S, Ando M, Kozakura K, Saito H, Irimajiri A. Ion channels in basolateral membrane of marginal cells dissociated from gerbil stria vascularis. *Hear Res* 83: 89–100, 1995.
265. Takeuchi S, Irimajiri A. Cl⁻ and nonselective cation channels in the basolateral membrane of strial marginal cells in the cochlea. *Jpn J Physiol* 44, Suppl 2: S317–S319, 1994.
266. Takeuchi S, Kakigi A, Takeda T, Saito H, Irimajiri A. Intravascularly applied K⁺-channel blockers suppress differently the positive endocochlear potential maintained by vascular perfusion. *Hear Res* 101: 181–185, 1996.
267. Takumi Y, Nagelhus EA, Eidet J, Matsubara A, Usami S, Shinkawa H, Nielsen S, Ottersen OP. Select types of supporting cell in the inner ear express aquaporin-4 water channel protein. *Eur J Neurosci* 10: 3584–3595, 1998.
268. Takumida M, Kubo N, Ohtani M, Suzuka Y, Anniko M. Transient receptor potential channels in the inner ear: presence of transient receptor potential channel subfamily 1 and 4 in the guinea pig inner ear. *Acta Otolaryngol (Stockh)* 125: 929–934, 2005.
269. Talebizadeh Z, Kelley PM, Askew JW, Beisel KW, Smith SD. Novel mutation in the KCNQ4 gene in a large kindred with dominant progressive hearing loss. *Hum Mutat* 14: 493–501, 1999.
270. Tanaka F, Whitworth CA, Rybak LP. Round window pH manipulation alters the ototoxicity of systemic cisplatin. *Hear Res* 187: 44–50, 2004.
271. Tanaka Y, Asanuma A, Yanagisawa K. Potentials of outer hair cells and their membrane properties in cationic environments. *Hear Res* 2: 431–438, 1980.
272. Tanemoto M, Abe T, Ito S. PDZ-binding and di-hydrophobic motifs regulate distribution of Kir4.1 channels in renal cells. *J Am Soc Nephrol* 16: 2608–2614, 2005.
273. Tanemoto M, Abe T, Onogawa T, Ito S. PDZ binding motif-dependent localization of K⁺ channel on the basolateral side in distal tubules. *Am J Physiol Renal Physiol* 287: F1148–F1153, 2004.
274. Taniguchi J, Imai M. Flow-dependent activation of maxi K⁺ channels in apical membrane of rabbit connecting tubule. *J Membr Biol* 164: 35–45, 1998.
275. Tasaki I, Spyropoulos CS. Stria vascularis as source of endocochlear potential. *J Neurophysiol* 22: 149–155, 1959.
276. Teng-umnuay P, Verlander JW, Yuan W, Tisher CC, Madsen KM. Identification of distinct subpopulations of intercalated cells in the mouse collecting duct. *J Am Soc Nephrol* 7: 260–274, 1996.
277. Teubner B, Michel V, Pesch J, Lautermann J, Cohen-Salmon M, Sohl G, Jahnke K, Winterhager E, Herberhold C, Hardelin JP, Petit C, Willecke K. Connexin30 (Gjb6)-deficiency causes severe hearing impairment and lack of endocochlear potential. *Hum Mol Genet* 12: 13–21, 2003.
278. Thurm H, Fakler B, Oliver D. Ca²⁺-independent activation of BKCa channels at negative potentials in mammalian inner hair cells. *J Physiol* 569: 137–151, 2005.
279. Tian L, Hammond MS, Florance H, Antoni FA, Shipston MJ. Alternative splicing determines sensitivity of murine calcium-activated potassium channels to glucocorticoids. *J Physiol* 537: 57–68, 2001.
280. Trune DR, Kempton JB, Gross ND. Mineralocorticoid receptor mediates glucocorticoid treatment effects in the autoimmune mouse ear. *Hear Res* 212: 22–32, 2006.
281. Tsujikawa S, Yamashita T, Amano H, Kumazawa T, Vosteen KH. Acidity in the endolymphatic sac fluid of guinea pigs. *ORL J Otorhinolaryngol Relat Spec* 54: 198–200, 1992.

282. Valli P, Zucca G, Botta L. Perilymphatic potassium changes and potassium homeostasis in isolated semicircular canals of the frog. *J Physiol* 430: 585–594, 1990.
283. Vallon V, Grahmmer F, Richter K, Bleich M, Lang F, Barhanin J, Volkl H, Warth R. Role of KCNE1-dependent K⁺ fluxes in mouse proximal tubule. *J Am Soc Nephrol* 12: 2003–2011, 2001.
284. Vallon V, Grahmmer F, Volkl H, Sandu CD, Richter K, Rexhepaj R, Gerlach U, Rong Q, Pfeifer K, Lang F. KCNQ1-dependent transport in renal and gastrointestinal epithelia. *Proc Natl Acad Sci USA* 102: 17864–17869, 2005.
285. Vallon V, Verkman AS, Schnermann J. Luminal hypotonicity in proximal tubules of aquaporin-1-knockout mice. *Am J Physiol Renal Physiol* 278: F1030–F1033, 2000.
286. Vanden Heuvel GB, Payne JA, Igarashi P, Forbush B, III. Expression of the basolateral Na-K-Cl cotransporter during mouse nephrogenesis and embryonic development. *Gene Expr Patterns* 6: 1000–1006, 2006.
287. Vargas-Poussou R, Feldmann D, Vollmer M, Konrad M, Kelly L, van den Heuvel LP, Tebourbi L, Brandis M, Karolyi L, Hebert SC, Lemmink HH, Deschenes G, Hildebrandt F, Seyberth HW, Guay-Woodford LM, Knoers NV, Antignac C. Novel molecular variants of the Na-K-2Cl cotransporter gene are responsible for antenatal Bartter syndrome. *Am J Hum Genet* 62: 1332–1340, 1998.
288. Vargas-Poussou R, Houillier P, Le Pottier N, Stropff L, Loirat C, Baudouin V, Macher MA, Dechaux M, Ulinski T, Nobili F, Eckart P, Novo R, Cailliez M, Salomon R, Nivet H, Cochat P, Tack I, Fargeot A, Bouissou F, Kesler GR, Lorotte S, Godefroid N, Layet V, Morin G, Jeunemaitre X, Blanchard A. Genetic investigation of autosomal recessive distal renal tubular acidosis: evidence for early sensorineural hearing loss associated with mutations in the ATP6V0A4 gene. *J Am Soc Nephrol* 17: 1437–1443, 2006.
289. Velazquez H, Silva T. Cloning and localization of KCC4 in rabbit kidney: expression in distal convoluted tubule. *Am J Physiol Renal Physiol* 285: F49–F58, 2003.
290. Verbalis JG. Whole-body volume regulation and escape from antidiuresis. *Am J Med* 119: S21–S29, 2006.
291. Verkman AS. Aquaporins in endothelia. *Kidney Int* 69: 1120–1123, 2006.
292. Verkman AS. Roles of aquaporins in kidney revealed by transgenic mice. *Semin Nephrol* 26: 200–208, 2006.
293. Verlander JW, Hassell KA, Royaux IE, Glapion DM, Wang ME, Everett LA, Green ED, Wall SM. Deoxycorticosterone upregulates PDS (Slc26a4) in mouse kidney: role of pendrin in mineralocorticoid-induced hypertension. *Hypertension* 42: 356–362, 2003.
294. Vetter DE, Mann JR, Wangemann P, Liu J, McLaughlin KJ, Lesage F, Marcus DC, Lazdunski M, Heinemann SF, Barhanin J. Inner ear defects induced by null mutation of the isk gene. *Neuron* 17: 1251–1264, 1996.
295. von Békésy G. DC potentials and energy balance of the cochlear partition. *J Acoust Soc Am* 576–582, 1950.
296. Wagner C, de Wit C, Kurtz L, Grunberger C, Kurtz A, Schweda F. Connexin40 is essential for the pressure control of renin synthesis and secretion. *Circ Res* 100: 556–563, 2007.
297. Waldegger S, Jeck N, Barth P, Peters M, Vitzthum H, Wolf K, Kurtz A, Konrad M, Seyberth HW. Barttin increases surface expression and changes current properties of ClC-K channels. *Pflügers Arch* 444: 411–418, 2002.
298. Waldegger S, Lang U, Herzer T, Suessbrich H, Binder K, Lepple-Wienhues A, Nagl U, Paulmichl M, Franz HB, Kiesel L, Lang F, Busch AE. Inhibition of minK protein induced K⁺ channels in *Xenopus* oocytes by estrogens. *Naunyn Schmiedebergs Arch Pharmacol* 354: 698–702, 1996.
299. Wall SM. Recent advances in our understanding of intercalated cells. *Curr Opin Nephrol Hypertens* 14: 480–484, 2005.
300. Wall SM, Hassell KA, Royaux IE, Green ED, Chang JY, Shipley GL, Verlander JW. Localization of pendrin in mouse kidney. *Am J Physiol Renal Physiol* 284: F229–F241, 2003.
301. Wall SM, Kim YH, Stanley L, Glapion DM, Everett LA, Green ED, Verlander JW. NaCl restriction upregulates renal Slc26a4 through subcellular redistribution: role in Cl⁻ conservation. *Hypertension* 44: 982–987, 2004.
302. Wall SM, Knepper MA, Hassell KA, Fischer MP, Shodeinde A, Shin W, Pham TD, Meyer JW, Lorenz JN, Beierwaltes WH, Dietz JR, Shull GE, Kim YH. Hypotension in NKCC1 null mice: role of the kidneys. *Am J Physiol Renal Physiol* 290: F409–F416, 2006.
303. Wangemann P. Comparison of ion transport mechanisms between vestibular dark cells and stria marginal cells. *Hear Res* 90: 149–157, 1995.
304. Wangemann P. Supporting sensory transduction: cochlear fluid homeostasis and the endocochlear potential. *J Physiol* 576: 11–21, 2006.
305. Wangemann P, Itza EM, Albrecht B, Wu T, Jabba SV, Maganti RJ, Lee JH, Everett LA, Wall SM, Royaux IE, Green ED, Marcus DC. Loss of KCNJ10 protein expression abolishes endocochlear potential and causes deafness in Pendred syndrome mouse model. *BMC Med* 2: 30, 2004.
306. Wangemann P, Liu J, Marcus DC. Ion transport mechanisms responsible for K⁺ secretion and the transepithelial voltage across marginal cells of stria vascularis in vitro. *Hear Res* 84: 19–29, 1995.
307. Wangemann P, Liu J, Scherer EQ, Herzog M, Shimozone M, Scofield MA. Muscarinic receptors control K⁺ secretion in inner ear stria marginal cells. *J Membr Biol* 182: 171–181, 2001.
308. Wangemann P, Liu J, Shen Z, Shipley A, Marcus DC. Hypo-osmotic challenge stimulates transepithelial K⁺ secretion and activates apical IsK channel in vestibular dark cells. *J Membr Biol* 147: 263–273, 1995.
309. Wangemann P, Liu J, Shimozone M, Schimanski S, Scofield MA. K⁺ secretion in stria marginal cells is stimulated via beta 1-adrenergic receptors but not via beta 2-adrenergic or vasopressin receptors. *J Membr Biol* 175: 191–202, 2000.
310. Wangemann P, Liu J, Shimozone M, Scofield MA. Beta1-adrenergic receptors but not beta2-adrenergic or vasopressin receptors regulate K⁺ secretion in vestibular dark cells of the inner ear. *J Membr Biol* 170: 67–77, 1999.
311. Wangemann P, Liu P, Shen Z, Marcus DC. Similarity of ion transport properties between vestibular dark cells and stria marginal cells. *Proc Sendai Symp* 5: 145–150, 2007.
312. Wangemann P, Marcus DC. Membrane potential measurements of transitional cells from the crista ampullaris of the gerbil. Effects of barium, quinidine, quinine, tetraethylammonium, cesium, ammonium, thallium and ouabain. *Pflügers Arch* 414: 656–662, 1989.
313. Wangemann P, Marcus DC. K⁺-induced swelling of vestibular dark cells is dependent on Na⁺ and Cl⁻ and inhibited by piretanide. *Pflügers Arch* 416: 262–269, 1990.
314. Wangemann P, Marcus DC. The membrane potential of vestibular dark cells is controlled by a large Cl⁻ conductance. *Hear Res* 62: 149–156, 1992.
315. Wangemann P, Nakaya K, Wu T, Maganti RJ, Itza EM, Sanneman JD, Harbidge DG, Billings S, Marcus DC. Loss of cochlear HCO₃⁻ secretion causes deafness via endolymphatic acidification and inhibition of Ca²⁺ reabsorption in a Pendred syndrome mouse model. *Am J Physiol Renal Physiol* 292: F1345–F1353, 2007.
316. Wangemann P, Schacht J. Homeostatic mechanisms in the cochlea. In: *Springer Handbook of Auditory Research: the Cochlea*, edited by Dallos P, Popper AN, Fay R. New York: Springer, 1996, p. 130–185.
317. Wangemann P, Shen Z, Liu J. K⁺-induced stimulation of K⁺ secretion involves activation of the IsK channel in vestibular dark cells. *Hear Res* 100: 201–210, 1996.
318. Wangemann P, Shiga N. Ba²⁺ and amiloride uncover or induce a pH-sensitive and a Na⁺ or non-selective cation conductance in transitional cells of the inner ear. *Pflügers Arch* 426: 258–266, 1994.
319. Warnock DG. The epithelial sodium channel in hypertension. *Curr Hypertens Rep* 1: 158–163, 1999.
320. Warnock DG. Liddle syndrome: genetics and mechanisms of Na⁺ channel defects. *Am J Med Sci* 322: 302–307, 2001.
321. Weiner ID, Hamm LL. Regulation of intracellular pH in the rabbit cortical collecting tubule. *J Clin Invest* 85: 274–281, 1990.
322. Wigand ME, Heidland A. Ototoxic side-effects of high doses of frusemide in patients with uraemia. *Postgrad Med J* 47, Suppl: 54–56, 1971.
323. Woda CB, Bragin A, Kleyman TR, Satlin LM. Flow-dependent K⁺ secretion in the cortical collecting duct is mediated by a maxi-K channel. *Am J Physiol Renal Physiol* 280: F786–F793, 2001.
324. Wood JD, Muchinsky SJ, Filoteo AG, Penniston JT, Tempel BL. Low endolymph calcium concentrations in deafwaddler2J mice suggest that PMCA2 contributes to endolymph calcium maintenance. *J Assoc Res Otolaryngol* 5: 99–110, 2004.
325. Xia A, Katori Y, Oshima T, Watanabe K, Kikuchi T, Ikeda K. Expression of connexin 30 in the developing mouse cochlea. *Brain Res* 898: 364–367, 2001.

326. Xia AP, Ikeda K, Katori Y, Oshima T, Kikuchi T, Takasaka T. Expression of connexin 31 in the developing mouse cochlea. *Neuroreport* 11: 2449–2453, 2000.
327. Yamauchi D, Raveendran NN, Pondugula SR, Kampalli SB, Sanne-man JD, Harbidge DG, Marcus DC. Vitamin D upregulates expression of ECaC1 mRNA in semicircular canal. *Biochem Biophys Res Commun* 331: 1353–1357, 2005.
328. Yamoah EN, Lumpkin EA, Dumont RA, Smith PJ, Hudspeth AJ, Gillespie PG. Plasma membrane Ca^{2+} -ATPase extrudes Ca^{2+} from hair cell stereocilia. *J Neurosci* 18: 610–624, 1998.
329. Yang B, Gillespie A, Carlson EJ, Epstein CJ, Verkman AS. Neonatal mortality in an aquaporin-2 knock-in mouse model of recessive nephrogenic diabetes insipidus. *J Biol Chem* 276: 2775–2779, 2001.
330. Yoshino T, Sato E, Nakashima T, Nagashima W, Teranishi MA, Nakayama A, Mori N, Murakami H, Funahashi H, Imai T. The immunohistochemical analysis of pendrin in the mouse inner ear. *Hear Res* 195: 9–16, 2004.
331. Zelante L, Gasparini P, Estivill X, Melchionda S, D'Agruma L, Govea N, Mila M, Monica MD, Lutfi J, Shohat M, Mansfield E, Delgrosso K, Rappaport E, Surrey S, Fortina P. Connexin26 mutations associated with the most common form of non-syndromic neurosensory autosomal recessive deafness (DFNB1) in Mediterraneans. *Hum Mol Genet* 6: 1605–1609, 1997.
332. Zhang Y, Tang W, Ahmad S, Sipp JA, Chen P, Lin X. Gap junction-mediated intercellular biochemical coupling in cochlear supporting cells is required for normal cochlear functions. *Proc Natl Acad Sci USA* 102: 15201–15206, 2005.
333. Zheng J, Dai C, Steyger PS, Kim Y, Vass Z, Ren T, Nuttall AL. Vanilloid receptors in hearing: altered cochlear sensitivity by vanilloids and expression of TRPV1 in the organ of corti. *J Neurophysiol* 90: 444–455, 2003.
334. Zhong SX, Liu ZH. Expression of aquaporins in the cochlea and endolymphatic sac of guinea pig. *ORL J Otorhinolaryngol Relat Spec* 65: 284–289, 2003.
335. Zhong SX, Liu ZH. Immunohistochemical localization of the epithelial sodium channel in the rat inner ear. *Hear Res* 193: 1–8, 2004.
336. Zidanic M, Brownell WE. Fine structure of the intracochlear potential field. I. The silent current. *Biophys J* 57: 1253–1268, 1990.

