

# EXTENSION FUNCTORS OF LOCAL COHOMOLOGY MODULES

M. AGHAPOURNAHR<sup>1</sup>, A. J. TAHERIZADEH<sup>2</sup>, AND ALIREZA VAHIDI<sup>3</sup>

ABSTRACT. Let  $R$  be a commutative Noetherian ring with non-zero identity,  $\mathfrak{a}$  an ideal of  $R$ , and  $X$  an  $R$ -module. In this paper, for fixed integers  $s, t$  and a finite  $\mathfrak{a}$ -torsion  $R$ -module  $N$ , we first study the membership of  $\text{Ext}_R^{s+t}(N, X)$  and  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X))$  in Serre subcategories of the category of  $R$ -modules. Then we present some conditions which ensure the existence of an isomorphism between them. Finally, we introduce the concept of Serre cofiniteness as a generalization of cofiniteness and study this property for certain local cohomology modules.

## 1. INTRODUCTION

Throughout  $R$  will denote a commutative Noetherian ring with non-zero identity and  $\mathfrak{a}$  an ideal of  $R$ . Also  $N$  will be a finite  $\mathfrak{a}$ -torsion module and  $X$  an  $R$ -module. For unexplained terminology from homological and commutative algebra we refer the reader to [10] and [11].

The following conjecture was made by Grothendieck in [19].

**Conjecture 1.1.** *For any ideal  $\mathfrak{a}$  and finite  $R$ -module  $X$ , the module  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is finite for all  $n \geq 0$ .*

This conjecture is false in general as shown by Hartshorne in [21]. However, he defined an  $R$ -module  $X$  to be  $\mathfrak{a}$ -cofinite if  $\text{Supp}_R(X) \subseteq V(\mathfrak{a})$  and  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is finite for each  $i$ , and he asked the following question.

**Question 1.2.** *If  $\mathfrak{a}$  is an ideal of  $R$  and  $X$  is a finite  $R$ -module when is  $\text{Ext}_R^i(R/\mathfrak{a}, H_{\mathfrak{a}}^j(X))$  finite for every  $i$  and  $j$ ?*

There are some attempts to show that under some conditions, for fixed integers  $s$  and  $t$ , the  $R$ -module  $\text{Ext}_R^s(R/\mathfrak{a}, H_{\mathfrak{a}}^t(X))$  is finite, for example see [3, Theorem 3.3], [16, Theorems A and B], [17, Theorem 6.3.9] and [24, Theorem 3.3].

Recently, the first author and Melkersson in [1] and [2], and Asgharzadeh and Tousi in [5] approached the study of local cohomology modules by means of Serre subcategories and it is noteworthy that their approach enables us to deal with several important problems on local cohomology modules comprehensively. For more information, we refer the reader to [23] to see a survey of some important problems on finiteness, vanishing, Artinianness, and finiteness of associated primes of local cohomology modules.

In this paper, we study some properties of extension functors of local cohomology modules by using Serre classes. Recall that a class of  $R$ -modules is a *Serre subcategory* of the category of  $R$ -modules when it is closed under taking submodules, quotients and extensions. Always,  $\mathcal{S}$  stands for a Serre subcategory of the category of  $R$ -modules.

---

2000 *Mathematics Subject Classification.* 13D45, 13D07.

*Key words and phrases.* Local cohomology modules, Serre subcategories, Cofinite modules.

The crucial points of Section 2 are Theorems 2.1 and 2.3 which show that when  $R$ -modules  $\text{Ext}_R^{s+t}(N, X)$  and  $\text{Ext}_R^s(N, H_a^t(X))$  belong to  $\mathcal{S}$ . These two theorems, which are frequently used through the paper, enable us to demonstrate some new facts and improve some older facts about the extension functors of local cohomology modules. We find the weakest possible conditions for finiteness of associated primes of local cohomology modules and, improve and give a new proof for [24, Theorem 3.3] in Corollaries 2.5 and 2.7. The relation between  $R$ -modules  $\text{Ext}_R^{s+t}(N, X)$  and  $\text{Ext}_R^s(N, H_a^t(X))$  to be in a Serre subcategory of the category of  $R$ -modules is shown in Corollary 2.8.

In Section 3, we first introduce the class of Melkersson subcategory as a special case of Serre classes and next investigate the extension functors of local cohomology modules in these subcategories. In Propositions 3.2, 3.3 and 3.4, we give new proofs for [1, Theorems 2.9 and 2.13] and study the membership of the local cohomology modules of an  $R$ -module  $X$  with respect to different ideals in Melkersson subcategories. Our main result in this section is Theorem 3.5 which provides an isomorphism between the  $R$ -modules  $\text{Ext}_R^{s+t}(N, X)$  and  $\text{Ext}_R^s(N, H_a^t(X))$ . Corollaries 3.6 through 3.9 are some applications of this theorem.

In Section 4, we present a generalization of the concept of cofiniteness with respect to an ideal to Serre subcategories of the category of  $R$ -modules. Theorems 4.2, 4.4 and 4.6 generalize [26, Proposition 2.5], [27, Proposition 3.11], [14, Theorem 3.1], [16, Theorems A and B] and [13, Corollary 2.7]. The Change of ring principle for Serre cofiniteness is presented in Theorem 4.8. We also give a proposition about  $\mathfrak{a}$ -cofinite minimax local cohomology modules in Proposition 4.10. Corollaries 4.11 and 4.12 are immediate results of this proposition where Corollary 4.11 improves [6, Theorem 2.3].

## 2. LOCAL COHOMOLOGY MODULES AND SERRE SUBCATEGORIES

Let  $\mathfrak{a}$  be an ideal of  $R$ ,  $N$  a finite  $\mathfrak{a}$ -torsion module and  $s, t$  non-negative integers. In this section, we present sufficient conditions which convince us the  $R$ -modules  $\text{Ext}_R^t(N, X)$  and  $\text{Ext}_R^s(N, H_a^t(X))$  are in a Serre subcategory of the category of  $R$ -modules. Even though we can provide elementary proofs by using induction for our main theorems, for shortening the proofs we use spectral sequences argument.

**Theorem 2.1.** *Let  $X$  be an  $R$ -module and  $t$  be a non-negative integer such that  $\text{Ext}_R^{t-r}(N, H_a^r(X))$  is in  $\mathcal{S}$  for all  $r$ ,  $0 \leq r \leq t$ . Then  $\text{Ext}_R^t(N, X)$  is in  $\mathcal{S}$ .*

*Proof.* By [29, Theorem 11.38], there is a Grothendieck spectral sequence

$$E_2^{p,q} := \text{Ext}_R^p(N, H_a^q(X)) \Rightarrow \text{Ext}_R^{p+q}(N, X).$$

For all  $r$ ,  $0 \leq r \leq t$ , we have  $E_\infty^{t-r,r} = E_{t+2}^{t-r,r}$  since  $E_i^{t-r-i, r+i-1} = 0 = E_i^{t-r+i, r+1-i}$  for all  $i \geq t+2$ ; so that  $E_\infty^{t-r,r}$  is in  $\mathcal{S}$  from the fact that  $E_{t+2}^{t-r,r}$  is a subquotient of  $E_2^{t-r,r}$  which is in  $\mathcal{S}$  by assumption. There exists a finite filtration

$$0 = \phi^{t+1}H^t \subseteq \phi^tH^t \subseteq \cdots \subseteq \phi^1H^t \subseteq \phi^0H^t = \text{Ext}_R^t(N, X)$$

such that  $E_\infty^{t-r,r} = \phi^{t-r}H^t / \phi^{t-r+1}H^t$  for all  $r$ ,  $0 \leq r \leq t$ . Now the exact sequences

$$0 \longrightarrow \phi^{t-r+1}H^t \longrightarrow \phi^{t-r}H^t \longrightarrow E_\infty^{t-r,r} \longrightarrow 0,$$

for all  $r$ ,  $0 \leq r \leq t$ , yield the assertion. □

Recall that, an  $R$ -module  $X$  is said to be *weakly Laskerian* if the set of associated primes of any quotient module of  $X$  is finite (see [13, Definition 2.1]). Also, we say that  $X$  is  $\mathfrak{a}$ -*weakly cofinite* if  $\text{Supp}_R(X) \subseteq V(\mathfrak{a})$  and  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is weakly Laskerian for all  $i \geq 0$  (see [14, Definition 2.4]). We denote the category of  $R$ -modules (resp. the category of finite  $R$ -modules, the category of weakly Laskerian  $R$ -modules) by  $\mathcal{C}(R)$  (resp.  $\mathcal{C}_{f.g}(R)$ ,  $\mathcal{C}_{w.l}(R)$ ).

**Corollary 2.2.** (cf. [17, Theorem 6.3.9(i)]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that for all  $r$ ,  $0 \leq r \leq n$ ,  $\text{Ext}_R^{n-r}(N, H_{\mathfrak{a}}^r(X))$  is weakly Laskerian (resp. finite). Then  $\text{Ext}_R^n(N, X)$  is weakly Laskerian (resp. finite) and so  $\text{Ass}_R(\text{Ext}_R^n(N, X))$  is finite.*

The next theorem is related to the  $R$ -module  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X))$  to be in a Serre subcategory of the category of  $R$ -modules.

**Theorem 2.3.** *Let  $X$  be an  $R$ -module and  $s, t$  be non-negative integers such that*

- (i)  $\text{Ext}_R^{s+t}(N, X)$  is in  $\mathcal{S}$ ,
- (ii)  $\text{Ext}_R^{s+t+1-i}(N, H_{\mathfrak{a}}^i(X))$  is in  $\mathcal{S}$  for all  $i$ ,  $0 \leq i < t$ , and
- (iii)  $\text{Ext}_R^{s+t-1-i}(N, H_{\mathfrak{a}}^i(X))$  is in  $\mathcal{S}$  for all  $i$ ,  $t+1 \leq i < s+t$ .

Then  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X))$  is in  $\mathcal{S}$ .

*Proof.* Consider the Grothendieck spectral sequence

$$E_2^{p,q} := \text{Ext}_R^p(N, H_{\mathfrak{a}}^q(X)) \Rightarrow \text{Ext}_R^{p+q}(N, X).$$

For all  $r \geq 2$ , let  $Z_r^{s,t} = \ker(E_r^{s,t} \rightarrow E_r^{s+r, t+1-r})$  and  $B_r^{s,t} = \text{Im}(E_r^{s-r, t+r-1} \rightarrow E_r^{s,t})$ . We have the exact sequences:

$$0 \longrightarrow Z_r^{s,t} \longrightarrow E_r^{s,t} \longrightarrow E_r^{s,t}/Z_r^{s,t} \longrightarrow 0$$

and

$$0 \longrightarrow B_r^{s,t} \longrightarrow Z_r^{s,t} \longrightarrow E_{r+1}^{s,t} \longrightarrow 0.$$

Since, by assumptions (ii) and (iii),  $E_2^{s+r, t+1-r}$  and  $E_2^{s-r, t+r-1}$  are in  $\mathcal{S}$ ,  $E_r^{s+r, t+1-r}$  and  $E_r^{s-r, t+r-1}$  are also in  $\mathcal{S}$ , and so  $E_r^{s,t}/Z_r^{s,t}$  and  $B_r^{s,t}$  are in  $\mathcal{S}$ . It shows that  $E_r^{s,t}$  is in  $\mathcal{S}$  whenever  $E_{r+1}^{s,t}$  is in  $\mathcal{S}$ .

We have  $E_r^{s-r, t+r-1} = 0 = E_r^{s+r, t+1-r}$  for all  $r$ ,  $r \geq t+s+2$ . Therefore we obtain  $E_{t+s+2}^{s,t} = E_{\infty}^{s,t}$ . To complete the proof, it is enough to show that  $E_{\infty}^{s,t}$  is in  $\mathcal{S}$ . There exists a finite filtration

$$0 = \phi^{s+t+1}H^{s+t} \subseteq \phi^{s+t}H^{s+t} \subseteq \dots \subseteq \phi^1H^{s+t} \subseteq \phi^0H^{s+t} = \text{Ext}_R^{s+t}(N, X)$$

such that  $E_{\infty}^{s+t-j, j} = \phi^{s+t-j}H^{s+t}/\phi^{s+t-j+1}H^{s+t}$  for all  $j$ ,  $0 \leq j \leq s+t$ . Since  $\text{Ext}_R^{s+t}(N, X)$  is in  $\mathcal{S}$ ,  $\phi^sH^{s+t}$  is in  $\mathcal{S}$  and so  $E_{\infty}^{s,t} = \phi^sH^{s+t}/\phi^{s+1}H^{s+t}$  is in  $\mathcal{S}$  as we desired.  $\square$

**Corollary 2.4.** (cf. [5, Theorem 2.2]) *Suppose that  $X$  is an  $R$ -module and  $n$  is a non-negative integer such that*

- (i)  $\text{Ext}_R^n(N, X)$  is in  $\mathcal{S}$ , and
- (ii)  $\text{Ext}_R^{n+1-i}(N, H_{\mathfrak{a}}^i(X))$  is in  $\mathcal{S}$  for all  $i$ ,  $0 \leq i < n$ .

Then  $\text{Hom}_R(N, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{S}$ .

*Proof.* Apply Theorem 2.3 with  $s = 0$  and  $t = n$ .  $\square$

We can deduce from the above corollary the main results of [25, Theorem B], [9, Theorem 2.2], [28, Theorem 5.6], [13, Corollary 2.7], [17, Theorem 6.3.9(ii)], [7, Theorem 2.3], [15, Corollary 3.2], [8, Corollary 2.3] and [6, Lemma 2.2] concerning the finiteness of associated primes of local cohomology modules. We just state the weakest possible conditions which yield the finiteness of associated primes of local cohomology modules in the next corollary.

**Corollary 2.5.** *Suppose that  $X$  is an  $R$ -module and  $n$  is a non-negative integer such that*

- (i)  $\text{Ext}_R^n(R/\mathfrak{a}, X)$  is weakly Laskerian, and
- (ii)  $\text{Ext}_R^{n+1-i}(R/\mathfrak{a}, H_{\mathfrak{a}}^i(X))$  is weakly Laskerian for all  $i$ ,  $0 \leq i < n$ .

*Then  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is weakly Laskerian, and so  $\text{Ass}_R(H_{\mathfrak{a}}^n(X))$  is finite.*

*Proof.* Apply Corollary 2.4 with  $N = R/\mathfrak{a}$  and  $\mathcal{S} = \mathcal{C}_{w.l}(R)$ , and note that we have the equality  $\text{Ass}_R(\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))) = V(\mathfrak{a}) \cap \text{Ass}_R(H_{\mathfrak{a}}^n(X)) = \text{Ass}_R(H_{\mathfrak{a}}^n(X))$ .  $\square$

It is easy to see that if  $R$  is a local ring and  $\mathcal{S}$  is a non-zero Serre subcategory of the category of  $R$ -modules, then every  $R$ -module with finite length belongs to  $\mathcal{S}$ .

**Corollary 2.6.** (cf. [5, Theorem 2.12]) *Let  $R$  be a local ring with maximal ideal  $\mathfrak{m}$  and  $X$  be an  $R$ -module. Assume also that  $\mathcal{S}$  is a non-zero Serre subcategory of  $\mathcal{C}(R)$  and  $n$  is a non-negative integer such that*

- (i)  $\text{Ext}_R^n(R/\mathfrak{m}, X)$  is finite, and
- (ii)  $\text{Ext}_R^{n+1-i}(R/\mathfrak{m}, H_{\mathfrak{a}}^i(X))$  is in  $\mathcal{S}$  for all  $i$ ,  $0 \leq i < n$ .

*Then  $\text{Hom}_R(R/\mathfrak{m}, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{S}$ .*

*Proof.* Since  $\mathcal{S} \neq 0$ ,  $\text{Ext}_R^n(R/\mathfrak{m}, X)$  is in  $\mathcal{S}$ . Now, the assertion follows from Corollary 2.4.  $\square$

Khashayarmanesh, in [24, Theorem 3.3], by using the concept of  $\mathfrak{a}$ -filter regular sequence, proved the following corollary with stronger assumptions. His assumptions were  $X$  is a finite  $R$ -module with finite Krull dimension and  $N = R/\mathfrak{b}$ , where  $\mathfrak{b}$  is an ideal of  $R$  contains  $\mathfrak{a}$ , while it is a simple conclusion of Theorem 2.3 for an arbitrary  $R$ -module  $X$  and a finite  $\mathfrak{a}$ -torsion module  $N$ .

**Corollary 2.7.** (cf. [24, Theorem 3.3]) *Suppose that  $X$  is an  $R$ -module and  $s, t$  are non-negative integers such that*

- (i)  $\text{Ext}_R^{s+t}(N, X)$  is finite,
- (ii)  $\text{Ext}_R^{s+t+1-i}(N, H_{\mathfrak{a}}^i(X))$  is finite for all  $i$ ,  $0 \leq i < t$ , and
- (iii)  $\text{Ext}_R^{s+t-1-i}(N, H_{\mathfrak{a}}^i(X))$  is finite for all  $i$ ,  $t+1 \leq i < s+t$ .

*Then  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X))$  is finite.*

*Proof.* Apply Theorem 2.3 for  $\mathcal{S} = \mathcal{C}_{f.g}(R)$ .  $\square$

Theorem 2.1 in conjunction with Theorem 2.3 arise the following corollary.

**Corollary 2.8.** *Let  $X$  be an  $R$ -module and  $n, m$  be non-negative integers such that  $n \leq m$ . Assume also that  $H_{\mathfrak{a}}^i(X)$  is in  $\mathcal{S}$  for all  $i$ ,  $i \neq n$  (resp.  $0 \leq i \leq n-1$  or  $n+1 \leq i \leq m$ ). Then, for all  $i$ ,  $i \geq 0$  (resp.  $0 \leq i \leq m-n$ ),  $\text{Ext}_R^i(N, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{S}$  if and only if  $\text{Ext}_R^{i+n}(N, X)$  is in  $\mathcal{S}$ .*

In the course of the remaining parts of the paper by  $\text{cd}_{\mathcal{S}}(\mathfrak{a}, X)$  ( $\mathcal{S}$ -cohomological dimension of  $X$  with respect to  $\mathfrak{a}$ ) we mean the largest integer  $i$  in which  $H_{\mathfrak{a}}^i(X)$  is not in  $\mathcal{S}$  (see [5, Definition 3.4] or [1, Definition 3.5]). Note that when  $\mathcal{S} = 0$ , then  $\text{cd}_{\mathcal{S}}(\mathfrak{a}, X) = \text{cd}(\mathfrak{a}, X)$  as in [20].

**Corollary 2.9.** *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer. Then the following statements hold true.*

- (i) *If  $\text{cd}_{\mathcal{S}}(\mathfrak{a}, X) = 0$ , then  $\text{Ext}_R^n(N, \Gamma_{\mathfrak{a}}(X))$  is in  $\mathcal{S}$  if and only if  $\text{Ext}_R^n(N, X)$  is in  $\mathcal{S}$ .*
- (ii) *If  $\text{cd}_{\mathcal{S}}(\mathfrak{a}, X) = 1$ , then  $\text{Ext}_R^n(N, H_{\mathfrak{a}}^1(X))$  is in  $\mathcal{S}$  if and only if  $\text{Ext}_R^{n+1}(N, X/\Gamma_{\mathfrak{a}}(X))$  is in  $\mathcal{S}$ .*
- (iii) *If  $\text{cd}_{\mathcal{S}}(\mathfrak{a}, X) = 2$ , then  $\text{Ext}_R^n(N, H_{\mathfrak{a}}^2(X))$  is in  $\mathcal{S}$  if and only if  $\text{Ext}_R^{n+2}(N, D_{\mathfrak{a}}(X))$  is in  $\mathcal{S}$ .*

*Proof.* (i) This is clear from Corollary 2.8.

(ii) For all  $i \neq 1$ ,  $H_{\mathfrak{a}}^i(X/\Gamma_{\mathfrak{a}}(X))$  is in  $\mathcal{S}$  by assumption. Now, the assertion follows from Corollary 2.8.

(iii) By [10, Corollary 2.2.8],  $H_{\mathfrak{a}}^i(D_{\mathfrak{a}}(X))$  is in  $\mathcal{S}$  for all  $i \neq 2$ . Again, use Corollary 2.8.  $\square$

### 3. SPECIAL SERRE SUBCATEGORIES

In this section, we study the extension functors of local cohomology modules in some special Serre subcategories of the category of  $R$ -modules. We begin with a definition.

**Definition 3.1.** (see [1, Definition 2.1]) *Let  $\mathcal{M}$  be a Serre subcategory of the category of  $R$ -modules. We say that  $\mathcal{M}$  is a Melkersson subcategory with respect to the ideal  $\mathfrak{a}$  if for any  $\mathfrak{a}$ -torsion  $R$ -module  $X$ ,  $0 :_X \mathfrak{a}$  is in  $\mathcal{M}$  implies that  $X$  is in  $\mathcal{M}$ .  $\mathcal{M}$  is called Melkersson subcategory when it is a Melkersson subcategory with respect to all ideals of  $R$ .*

In honor of Melkersson who proved this property for Artinian category (see [10, Theorem 7.1.2]) and Artinian  $\mathfrak{a}$ -cofinite category (see [27, Proposition 4.1]), we named the above subcategory as Melkersson subcategory. To see some examples of Melkersson subcategories, we refer the reader to [1, Examples 2.4 and 2.5].

The next two propositions show that how properties of Melkersson subcategories behave similarly at the initial points of Ext and local cohomology modules. These propositions give new proofs for [1, Theorems 2.9 and 2.13] based on Theorems 2.1 and 2.3.

**Proposition 3.2.** (see [1, Theorem 2.13]) *Let  $X$  be an  $R$ -module,  $\mathcal{M}$  be a Melkersson subcategory with respect to the ideal  $\mathfrak{a}$ , and  $n$  be a non-negative integer such that  $\text{Ext}_R^{j-i}(R/\mathfrak{a}, H_{\mathfrak{a}}^i(X))$  is in  $\mathcal{M}$  for all  $i, j$  with  $0 \leq i \leq n-1$  and  $j = n, n+1$ . Then the following statements are equivalent.*

- (i)  $\text{Ext}_R^n(R/\mathfrak{a}, X)$  is in  $\mathcal{M}$ .
- (ii)  $H_{\mathfrak{a}}^n(X)$  is in  $\mathcal{M}$ .

*Proof.* (i)  $\Rightarrow$  (ii). Apply Theorem 2.3 with  $s = 0$  and  $t = n$ . It shows that  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{M}$ . Thus  $H_{\mathfrak{a}}^n(X)$  is in  $\mathcal{M}$ .

(ii)  $\Rightarrow$  (i). Apply Theorem 2.1 with  $t = n$ .  $\square$

**Proposition 3.3.** (see [1, Theorem 2.9]) *Let  $X$  be an  $R$ -module,  $\mathcal{M}$  be a Melkersson subcategory with respect to the ideal  $\mathfrak{a}$ , and  $n$  be a non-negative integer. Then the following statements are equivalent.*

- (i)  $H_{\mathfrak{a}}^i(X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n$ .
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n$ .

*Proof.* (i)  $\Rightarrow$  (ii). Let  $0 \leq t \leq n$ . Since  $H_{\mathfrak{a}}^r(X)$  is in  $\mathcal{M}$  for all  $r$ ,  $0 \leq r \leq t$ ,  $\text{Ext}_R^{t-r}(R/\mathfrak{a}, H_{\mathfrak{a}}^r(X))$  is in  $\mathcal{M}$  for all  $r$ ,  $0 \leq r \leq t$ . Hence  $\text{Ext}_R^t(R/\mathfrak{a}, X)$  is in  $\mathcal{M}$  by Theorem 2.1.

(ii)  $\Rightarrow$  (i). We prove by using induction on  $n$ . Let  $n = 0$  and consider the isomorphism  $\text{Hom}_R(R/\mathfrak{a}, X) \cong \text{Hom}_R(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(X))$ . Since  $\text{Hom}_R(R/\mathfrak{a}, X)$  is in  $\mathcal{M}$ ,  $\text{Hom}_R(R/\mathfrak{a}, \Gamma_{\mathfrak{a}}(X))$  is in  $\mathcal{M}$ . Thus  $\Gamma_{\mathfrak{a}}(X)$  is in  $\mathcal{M}$ .

Now, suppose that  $n > 0$  and that  $n - 1$  is settled. Since  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n - 1$ ,  $H_{\mathfrak{a}}^i(X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n - 1$  by the induction hypothesis. Now, by the above proposition,  $H_{\mathfrak{a}}^n(X)$  is in  $\mathcal{M}$ .  $\square$

In the next proposition, we study the membership of the local cohomology modules of an  $R$ -module  $X$  with respect to different ideals in Melkersson subcategories which, among other things, shows that  $\text{cd}_{\mathcal{M}}(\mathfrak{b}, X) \leq \text{cd}_{\mathcal{M}}(\mathfrak{a}, X) + \text{ara}(\mathfrak{b}/\mathfrak{a})$  where  $\mathcal{M}$  is a Melkersson subcategory of  $\mathcal{C}(R)$  and  $\mathfrak{b}$  is an ideal of  $R$  contains  $\mathfrak{a}$ .

**Proposition 3.4.** *Let  $X$  be an  $R$ -module and  $\mathfrak{b}$  be an ideal of  $R$  such that  $\mathfrak{a} \subseteq \mathfrak{b}$ . Assume also that  $\mathcal{M}$  is a Melkersson subcategory of  $\mathcal{C}(R)$  and  $n$  is a non-negative integer such that  $H_{\mathfrak{a}}^i(X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n$  (resp.  $i \geq n$ ). Then  $H_{\mathfrak{b}}^i(X)$  is in  $\mathcal{M}$  for all  $i$ ,  $0 \leq i \leq n$  (resp.  $i \geq n + \text{ara}(\mathfrak{b}/\mathfrak{a})$ ).*

*Proof.* Let  $r = \text{ara}(\mathfrak{b}/\mathfrak{a})$ . There exist  $x_1, \dots, x_r \in R$  such that  $\sqrt{\mathfrak{b}} = \sqrt{\mathfrak{a} + (x_1, \dots, x_r)}$ . We can, and do, assume that  $\mathfrak{b} = \mathfrak{a} + \mathfrak{c}$  where  $\mathfrak{c} = (x_1, \dots, x_r)$ . By [29, Theorem 11.38], there is a Grothendieck spectral sequence

$$E_2^{p,q} := H_{\mathfrak{c}}^p(H_{\mathfrak{a}}^q(X)) \underset{p}{\Rightarrow} H_{\mathfrak{b}}^{p+q}(X).$$

Assume that  $t$  is a non-negative integer such that  $0 \leq t \leq n$  (resp.  $t \geq n + r$ ). For all  $i$ ,  $0 \leq i \leq t$ ,  $E_{\infty}^{t-i,i} = E_{t+2}^{t-i,i}$  since  $E_j^{t-i-j,i+j-1} = 0 = E_j^{t-i+j,i-j+1}$  for all  $j \geq t + 2$ . Therefore  $E_{\infty}^{t-i,i}$  is in  $\mathcal{M}$  from the fact that  $E_{t+2}^{t-i,i}$  is a subquotient of  $E_2^{t-i,i} = H_{\mathfrak{c}}^{t-i}(H_{\mathfrak{a}}^i(X))$  which belongs to  $\mathcal{M}$  by assumption and Proposition 3.3. There exists a finite filtration

$$0 = \phi^{t+1}H^t \subseteq \phi^tH^t \subseteq \dots \subseteq \phi^1H^t \subseteq \phi^0H^t = H_{\mathfrak{b}}^t(X)$$

such that  $E_{\infty}^{t-i,i} = \phi^{t-i}H^t / \phi^{t-i+1}H^t$  for all  $i$ ,  $0 \leq i \leq t$ . Now the exact sequences

$$0 \longrightarrow \phi^{t-i+1}H^t \longrightarrow \phi^{t-i}H^t \longrightarrow E_{\infty}^{t-i,i} \longrightarrow 0,$$

for all  $i$ ,  $0 \leq i \leq t$ , show that  $H_{\mathfrak{b}}^t(X)$  is in  $\mathcal{M}$ .  $\square$

Let  $\mathfrak{a}$  be an ideal of  $R$ ,  $N$  a finite  $\mathfrak{a}$ -torsion module and  $s, t$  non-negative integers. In the following theorem, we find some sufficient conditions for validity of the isomorphism  $\text{Ext}_R^{s+t}(N, X) \cong \text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X))$  which concerns to the case  $\mathcal{S} = 0$ .

**Theorem 3.5.** *Let  $X$  be an  $R$ -module and  $s, t$  be non-negative integers such that*

- (i)  $\text{Ext}_R^{s+t-i}(N, H_{\mathfrak{a}}^i(X)) = 0$  for all  $i$ ,  $0 \leq i < t$  or  $t < i \leq s + t$ ,
- (ii)  $\text{Ext}_R^{s+t+1-i}(N, H_{\mathfrak{a}}^i(X)) = 0$  for all  $i$ ,  $0 \leq i < t$ , and
- (iii)  $\text{Ext}_R^{s+t-1-i}(N, H_{\mathfrak{a}}^i(X)) = 0$  for all  $i$ ,  $t + 1 \leq i < s + t$ .

*Then we have  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X)) \cong \text{Ext}_R^{s+t}(N, X)$ .*

*Proof.* Consider the Grothendieck spectral sequence

$$E_2^{p,q} := \text{Ext}_R^p(N, H_{\mathfrak{a}}^q(X)) \xrightarrow{\cong} \text{Ext}_R^{p+q}(N, X)$$

and, for all  $r \geq 2$ , the exact sequences

$$0 \rightarrow B_r^{s,t} \rightarrow Z_r^{s,t} \rightarrow E_{r+1}^{s,t} \rightarrow 0 \quad \text{and} \quad 0 \rightarrow Z_r^{s,t} \rightarrow E_r^{s,t} \rightarrow E_r^{s,t}/Z_r^{s,t} \rightarrow 0$$

as we used in Theorem 2.3. Since  $E_2^{s+r,t+1-r} = 0 = E_2^{s-r,t+r-1}$ ,  $E_r^{s+r,t+1-r} = 0 = E_r^{s-r,t+r-1}$ . Therefore  $E_r^{s,t}/Z_r^{s,t} = 0 = B_r^{s,t}$  which shows that  $E_r^{s,t} = E_{r+1}^{s,t}$ . Hence we have

$$\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X)) = E_2^{s,t} = E_3^{s,t} = \dots = E_{s+t+1}^{s,t} = E_{s+t+2}^{s,t} = E_{\infty}^{s,t}.$$

There is a finite filtration

$$0 = \phi^{s+t+1}H^{s+t} \subseteq \phi^{s+t}H^{s+t} \subseteq \dots \subseteq \phi^1H^{s+t} \subseteq \phi^0H^{s+t} = \text{Ext}_R^{s+t}(N, X)$$

such that  $E_{\infty}^{s+t-j,j} = \phi^{s+t-j}H^{s+t}/\phi^{s+t-j+1}H^{s+t}$  for all  $j$ ,  $0 \leq j \leq s+t$ . Note that for each  $j$ ,  $0 \leq j \leq t-1$  or  $t+1 \leq j \leq s+t$ , by assumption (i), we have  $E_{\infty}^{s+t-j,j} = 0$ . Therefore we get

$$0 = \phi^{s+t+1}H^{s+t} = \phi^{s+t}H^{s+t} = \dots = \phi^{s+2}H^{s+t} = \phi^{s+1}H^{s+t}$$

and

$$\phi^sH^{s+t} = \phi^{s-1}H^{s+t} = \dots = \phi^1H^{s+t} = \phi^0H^{s+t} = \text{Ext}_R^{s+t}(N, X).$$

Thus  $\text{Ext}_R^s(N, H_{\mathfrak{a}}^t(X)) = E_{\infty}^{s,t} = \phi^sH^{s+t}/\phi^{s+1}H^{s+t} = \text{Ext}_R^{s+t}(N, X)$  as desired.  $\square$

The following corollaries are immediate applications of the above theorem which give us some useful isomorphisms and equalities about the extension functors and Bass numbers of local cohomology modules, respectively.

**Corollary 3.6.** (cf. [2, Corollary 4.2.(c)]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer. Then the isomorphism  $\text{Hom}_R(N, H_{\mathfrak{a}}^n(X)) \cong \text{Ext}_R^n(N, X)$  holds in either of the following cases:*

- (i)  $\text{Ext}_R^{j-i}(N, H_{\mathfrak{a}}^i(X)) = 0$  for all  $i, j$  with  $0 \leq i \leq n-1$  and  $j = n, n+1$ ;
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X) = 0$  for all  $i$ ,  $0 \leq i \leq n-1$ .

*Proof.* (i) Apply Theorem 3.5 with  $s = 0$  and  $t = n$ .

(ii) By Proposition 3.3,  $H_{\mathfrak{a}}^i(X) = 0$  for all  $i$ ,  $0 \leq i \leq n-1$ . Now, use case (i).  $\square$

**Corollary 3.7.** *Let  $X$  be an  $R$ -module and  $n, m$  be non-negative integers such that  $n \leq m$ . Assume also that  $H_{\mathfrak{a}}^i(X) = 0$  for all  $i$ ,  $i \neq n$  (resp.  $0 \leq i \leq n-1$  or  $n+1 \leq i \leq m$ ). Then we have  $\text{Ext}_R^i(N, H_{\mathfrak{a}}^n(X)) \cong \text{Ext}_R^{i+n}(N, X)$  for all  $i$ ,  $i \geq 0$  (resp.  $0 \leq i \leq m-n$ ).*

*Proof.* For all  $i$ ,  $i \geq 0$  (resp.  $0 \leq i \leq m-n$ ), apply Theorem 3.5 with  $s = i$  and  $t = n$ .  $\square$

**Corollary 3.8.** (cf. [18, Proposition 3.1]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that  $H_{\mathfrak{a}}^i(X) = 0$  for all  $i$ ,  $i \neq n$ . Then we have  $\mu^i(\mathfrak{p}, H_{\mathfrak{a}}^n(X)) = \mu^{i+n}(\mathfrak{p}, X)$  for all  $i \geq 0$  and all  $\mathfrak{p} \in V(\mathfrak{a})$ .*

*Proof.* Let  $\mathfrak{p} \in V(\mathfrak{a})$ . By assumption,  $H_{\mathfrak{a}R_{\mathfrak{p}}}^i(X_{\mathfrak{p}}) = 0$  for all  $i$ ,  $i \neq n$ ; so that  $\text{Ext}_{R_{\mathfrak{p}}}^i(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}, H_{\mathfrak{a}R_{\mathfrak{p}}}^n(X_{\mathfrak{p}})) \cong \text{Ext}_{R_{\mathfrak{p}}}^{i+n}(R_{\mathfrak{p}}/\mathfrak{p}R_{\mathfrak{p}}, X_{\mathfrak{p}})$  for all  $i \geq 0$  by Corollary 3.7. Hence  $\mu^i(\mathfrak{p}, H_{\mathfrak{a}}^n(X)) = \mu^{i+n}(\mathfrak{p}, X)$  for all  $i \geq 0$ .  $\square$

**Corollary 3.9.** *For an arbitrary  $R$ -module  $X$ , the following statements hold true.*

- (i) If  $\text{cd}(\mathfrak{a}, X) = 0$ , then  $\text{Ext}_R^i(N, \Gamma_{\mathfrak{a}}(X)) \cong \text{Ext}_R^i(N, X)$  for all  $i \geq 0$ .
- (ii) If  $\text{cd}(\mathfrak{a}, X) = 1$ , then  $\text{Ext}_R^i(N, H_{\mathfrak{a}}^1(X)) \cong \text{Ext}_R^{i+1}(N, X/\Gamma_{\mathfrak{a}}(X))$  for all  $i \geq 0$ .
- (iii) If  $\text{cd}(\mathfrak{a}, X) = 2$ , then  $\text{Ext}_R^i(N, H_{\mathfrak{a}}^2(X)) \cong \text{Ext}_R^{i+2}(N, D_{\mathfrak{a}}(X))$  for all  $i \geq 0$ .

*Proof.* By considering Corollary 3.7, this is similar to that of Corollary 2.9.  $\square$

#### 4. COFINITE MODULES

We first introduce the class of cofinite modules with respect to an ideal and a Serre subcategory of the category of  $R$ -modules.

**Definition 4.1.** Let  $\mathfrak{a}$  be an ideal of  $R$ ,  $X$  be an  $R$ -module and  $\mathcal{S}$  be a Serre subcategory of  $\mathcal{C}(R)$ . We say that  $X$  is  $\mathcal{S}$ -cofinite with respect to the ideal  $\mathfrak{a}$  if  $\text{Supp}_R(X) \subseteq V(\mathfrak{a})$  and  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is in  $\mathcal{S}$  for all  $i \geq 0$ . We will denote this concept by  $(\mathcal{S}, \mathfrak{a})$ -cofinite.

Note that when  $\mathcal{S}$  is  $\mathcal{C}_{f,g}(R)$  (resp.  $\mathcal{C}_{w,l}(R)$ ),  $X$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite exactly when  $X$  is  $\mathfrak{a}$ -cofinite (resp.  $\mathfrak{a}$ -weakly cofinite).

**Theorem 4.2.** Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that  $H_{\mathfrak{a}}^i(X)$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite for all  $i$ ,  $i \neq n$ . Then the following statements are equivalent.

- (i)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is in  $\mathcal{S}$  for all  $i \geq 0$ .
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is in  $\mathcal{S}$  for all  $i \geq n$ .
- (iii)  $H_{\mathfrak{a}}^n(X)$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite.

*Proof.* (i)  $\Rightarrow$  (ii). This is clear.

(ii)  $\Rightarrow$  (iii). For all  $i \geq 0$ , apply Theorem 2.3 with  $N = R/\mathfrak{a}$ ,  $s = i$  and  $t = n$ .

(iii)  $\Rightarrow$  (i). Apply Theorem 2.1 with  $N = R/\mathfrak{a}$ .  $\square$

As an immediate result, the following corollary recovers and improves [26, Proposition 2.5], [27, Proposition 3.11] and [14, Theorem 3.1].

**Corollary 4.3.** (cf. [26, Proposition 2.5], [27, Proposition 3.11] and [14, Theorem 3.1]) Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite (resp.  $\mathfrak{a}$ -weakly cofinite) for all  $i$ ,  $i \neq n$ . Then the following statements are equivalent.

- (i)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is finite (resp. weakly Laskerian) for all  $i \geq 0$ .
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is finite (resp. weakly Laskerian) for all  $i \geq n$ .
- (iii)  $H_{\mathfrak{a}}^n(X)$  is  $\mathfrak{a}$ -cofinite (resp.  $\mathfrak{a}$ -weakly cofinite).

**Theorem 4.4.** Suppose that  $X$  is an  $R$ -module and  $n$  is a non-negative integer such that

- (i)  $H_{\mathfrak{a}}^i(X)$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite for all  $i$ ,  $0 \leq i \leq n-1$ , and
- (ii)  $\text{Ext}_R^{1+n}(N, X)$  is in  $\mathcal{S}$ .

Then  $\text{Ext}_R^1(N, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{S}$ .

*Proof.* Consider [22, Proposition 3.4] and apply Theorem 2.3 with  $s = 1$  and  $t = n$ .  $\square$

The following result is an application of the above theorem.



**Corollary 4.5.** (cf. [16, Theorem A] and [13, Corollary 2.7]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer. Assume also that*

- (i)  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite (resp.  $\mathfrak{a}$ -weakly cofinite) for all  $i$ ,  $0 \leq i \leq n-1$ , and
- (ii)  $\text{Ext}_R^{1+n}(N, X)$  is finite (resp. weakly Laskerian).

*Then  $\text{Ext}_R^1(N, H_{\mathfrak{a}}^n(X))$  is finite (resp. weakly Laskerian).*

**Theorem 4.6.** *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that  $\text{Ext}_R^{n+1}(N, X)$  and  $\text{Ext}_R^{n+2}(N, X)$  are in  $\mathcal{S}$ , and  $H_{\mathfrak{a}}^i(X)$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite for all  $i$ ,  $0 \leq i < n$ . Then the following statements are equivalent.*

- (i)  $\text{Hom}_R(N, H_{\mathfrak{a}}^{n+1}(X))$  is in  $\mathcal{S}$ .
- (ii)  $\text{Ext}_R^2(N, H_{\mathfrak{a}}^n(X))$  is in  $\mathcal{S}$ .

*Proof.* (i)  $\Rightarrow$  (ii). Consider [22, Proposition 3.4] and apply Theorem 2.3 with  $s = 2$  and  $t = n$ .

(ii)  $\Rightarrow$  (i). Again consider [22, Proposition 3.4] and apply Theorem 2.3 with  $s = 0$  and  $t = n + 1$ .  $\square$

Asadollahi and Schenzel proved that over local ring  $(R, \mathfrak{m})$ , if  $X$  is a Cohen-Macaulay  $R$ -module and  $t = \text{grade}(\mathfrak{a}, X)$  then  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^{t+1}(X))$  is finite if and only if  $\text{Ext}_R^2(R/\mathfrak{a}, H_{\mathfrak{a}}^t(X))$  is finite (see [4, Theorem 1.2]). Dibaei and Yassemi, in [16], generalized this result with weaker assumptions on  $R$  and  $X$ . As an immediate consequence of Theorem 4.6, the following is a generalization of [16, Theorem B].

**Corollary 4.7.** (cf. [16, Theorem B]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer. Assume also that  $\text{Ext}_R^{n+1}(N, X)$  and  $\text{Ext}_R^{n+2}(N, X)$  are finite (resp. weakly Laskerian), and  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite (resp.  $\mathfrak{a}$ -weakly cofinite) for all  $i$ ,  $0 \leq i < n$ . Then the following statements are equivalent.*

- (i)  $\text{Hom}_R(N, H_{\mathfrak{a}}^{n+1}(X))$  is finite (resp. weakly Laskerian).
- (ii)  $\text{Ext}_R^2(N, H_{\mathfrak{a}}^n(X))$  is finite (resp. weakly Laskerian).

In [12, Proposition 2], Delfino and Marley proved the Change of ring principle for cofiniteness. In the following theorem, we prove it for Serre cofiniteness. The proof is an adaption of the proof of [12, Proposition 2].

**Theorem 4.8.** *Let  $\phi : A \rightarrow B$  be a homomorphism between Noetherian rings such that  $B$  is a finite  $A$ -module,  $\mathfrak{a}$  be an ideal of  $A$  and  $X$  be a  $B$ -module. Let  $\mathcal{S}$  and  $\mathcal{T}$  be Serre subcategories of  $\mathcal{C}(A)$  and  $\mathcal{C}(B)$ , respectively. Assume also that for any  $B$ -module  $Y$ ,  $Y$  is in  $\mathcal{T}$  exactly when  $Y$  is in  $\mathcal{S}$  (as an  $A$ -module). Then  $X$  is  $(\mathcal{T}, \mathfrak{a}B)$ -cofinite if and only if  $X$  is  $(\mathcal{S}, \mathfrak{a})$ -cofinite (as an  $A$ -module).*

*Proof.* By [29, Theorem 11.65], there is a Grothendieck spectral sequence

$$E_2^{p,q} := \text{Ext}_B^p(\text{Tor}_q^A(B, A/\mathfrak{a}), X) \xrightarrow{\cong} \text{Ext}_A^{p+q}(A/\mathfrak{a}, X).$$

( $\Rightarrow$ ). For all  $p$  and  $q$ , by [22, Proposition 3.4],  $E_2^{p,q}$  is in  $\mathcal{S}$ . Therefore  $E_{\infty}^{p,q}$  belongs to  $\mathcal{S}$  since  $E_{\infty}^{p,q} = E_{p+q+2}^{p,q}$  and  $E_{p+q+2}^{p,q}$  is a subquotient of  $E_2^{p,q}$ . Let  $n$  be a non-negative integer. There exists a finite filtration

$$0 = \phi^{n+1}H^n \subseteq \phi^n H^n \subseteq \dots \subseteq \phi^1 H^n \subseteq \phi^0 H^n = \text{Ext}_A^n(A/\mathfrak{a}, X)$$

such that  $E_{\infty}^{n-i,i} = \phi^{n-i}H^n / \phi^{n-i+1}H^n$  for all  $i$ ,  $0 \leq i \leq n$ . Now, by the exact sequences

$$0 \longrightarrow \phi^{n-i+1}H^n \longrightarrow \phi^{n-i}H^n \longrightarrow E_{\infty}^{n-i,i} \longrightarrow 0,$$

for all  $i$ ,  $0 \leq i \leq n$ ,  $\text{Ext}_A^n(A/\mathfrak{a}, X)$  is in  $\mathcal{S}$ .

( $\Leftarrow$ ). By using induction on  $n$ , we show that  $E_2^{n,0} = \text{Ext}_B^n(B/\mathfrak{a}B, X)$  is in  $\mathcal{T}$  for all  $n \geq 0$ . The case  $n = 0$  is clear from the isomorphism  $\text{Hom}_B(B/\mathfrak{a}B, X) \cong \text{Hom}_A(A/\mathfrak{a}, X)$ . Assume that  $n > 0$  and that  $E_2^{p,0}$  is in  $\mathcal{T}$  for all  $p$ ,  $0 \leq p \leq n - 1$ . For all  $r \geq 2$ , we have  $E_{r+1}^{n,0} \cong E_r^{n,0}/\text{Im}(E_r^{n-r,r-1} \rightarrow E_r^{n,0})$ . Thus  $E_r^{n,0}$  is in  $\mathcal{T}$  whenever  $E_{r+1}^{n,0}$  is in  $\mathcal{T}$  because  $E_r^{n-r,r-1}$  is in  $\mathcal{T}$  by the induction hypotheses and [22, Proposition 3.4]. Since  $E_\infty^{n,0} = E_{n+2}^{n,0}$ , to complete the proof it is enough to show that  $E_\infty^{n,0}$  is in  $\mathcal{T}$ . By assumption,  $\text{Ext}_A^n(A/\mathfrak{a}, X)$  is in  $\mathcal{T}$  and hence  $\phi^n H^n$  is in  $\mathcal{T}$ . That is  $E_\infty^{n,0}$  belongs to  $\mathcal{T}$  as desired.  $\square$

**Definition 4.9.** (see [30]) *The  $R$ -module  $X$  is a minimax module if it has a finite submodule  $X'$  such that  $X/X'$  is Artinian.*

The class of minimax modules thus includes all finite and all Artinian modules. Note that the category of minimax modules and the category of  $\mathfrak{a}$ -cofinite minimax modules are two Serre subcategories of the category of  $R$ -modules (see [27, Corollary 4.4]).

**Proposition 4.10.** *Let  $X$  be an  $R$ -module and  $n, m$  be non-negative integers such that  $n \leq m$ . Assume also that*

- (i)  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite for all  $i$ ,  $0 \leq i \leq n - 1$ ,
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is finite for all  $i$ ,  $n \leq i \leq m$ , and
- (iii)  $H_{\mathfrak{a}}^i(X)$  is minimax for all  $i$ ,  $n \leq i \leq m$ .

Then  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite for all  $i$ ,  $0 \leq i \leq m$ .

*Proof.* Apply Theorem 2.3 with  $s = 0$  and  $t = n$  for  $N = R/\mathfrak{a}$  and  $\mathcal{S} = \mathcal{C}_{f.g.}(R)$ . It shows that  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is finite. Thus  $H_{\mathfrak{a}}^n(X)$  is  $\mathfrak{a}$ -cofinite from [27, Proposition 4.3].  $\square$

**Corollary 4.11.** (cf. [6, Theorem 2.3]) *Let  $X$  be an  $R$ -module and  $n$  be a non-negative integer such that*

- (i)  $H_{\mathfrak{a}}^i(X)$  is minimax for all  $i$ ,  $0 \leq i \leq n - 1$ , and
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  is finite for all  $i$ ,  $0 \leq i \leq n$ .

Then  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is finite.

*Proof.* By [27, Proposition 4.3],  $\Gamma_{\mathfrak{a}}(X)$  is  $\mathfrak{a}$ -cofinite. Hence  $H_{\mathfrak{a}}^i(X)$  is  $\mathfrak{a}$ -cofinite for all  $i$ ,  $0 \leq i \leq n - 1$ , from Proposition 4.10. Thus, by Theorem 2.3,  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^n(X))$  is finite.  $\square$

**Corollary 4.12.** *Suppose that  $X$  is an  $R$ -module and that  $n$  is a non-negative integer. Then the following statements are equivalent.*

- (i)  $H_{\mathfrak{a}}^i(X)$  is Artinian  $\mathfrak{a}$ -cofinite for all  $i$ ,  $0 \leq i \leq n$ .
- (ii)  $\text{Ext}_R^i(R/\mathfrak{a}, X)$  has finite length for all  $i$ ,  $0 \leq i \leq n$ .

*Proof.* (i)  $\Rightarrow$  (ii). Let  $0 \leq t \leq n$ . Since  $\text{Ext}_R^{t-i}(R/\mathfrak{a}, H_{\mathfrak{a}}^i(X))$  has finite length for all  $i$ ,  $0 \leq i \leq t$ ,  $\text{Ext}_R^t(R/\mathfrak{a}, X)$  has also finite length by Theorem 2.1.

(ii)  $\Rightarrow$  (i). By Proposition 3.3,  $H_{\mathfrak{a}}^i(X)$  is Artinian for all  $i$ ,  $0 \leq i \leq n$ . Let  $0 \leq t \leq n$  and consider Corollary 4.11. It shows that  $\text{Hom}_R(R/\mathfrak{a}, H_{\mathfrak{a}}^t(X))$  is finite and so has finite length. Now, the assertion follows from [27, Proposition 4.3].  $\square$

## REFERENCES

1. M. Aghapournahr, L. Melkersson, *Local cohomology and Serre subcategories*, J. Algebra, **320** (2008), 1275–1287.
2. M. Aghapournahr, L. Melkersson, *A natural map in local cohomology*, To appear in Ark. Mat.
3. J. Asadollahi, K. Khashyarmansh, Sh. Salarian, *A generalization of the cofiniteness problem in local cohomology*, J. Aust. Math. Soc., **75** (2003), 313–324.
4. J. Asadollahi and P. Schenzel, *Some results on associated primes of local cohomology modules*, Japan. J. Math., **29** (2003), 285–296.
5. M. Asgharzadeh, M. Tousi, *A unified approach to local cohomology modules using serre classes*, arXiv: 0712.3875v2 [math.AC].
6. K. Bahmanpour, R. Naghipour, *On the cofiniteness of local cohomology modules*, Proc. Amer. Math. Soc., **136** (2008), 2359–2363.
7. K. Borna Lorestani, P. Sahandi, T. Sharif, *A note on the associated primes of local cohomology modules*, Comm. Algebra, **34** (2006), 3409–3412.
8. K. Borna Lorestani, P. Sahandi, S. Yassemi, *Artinian local cohomology*, Can. Math. Bull., **50** (2007), 598–602.
9. M. P. Brodmann, A. Lashgari, *A finiteness result for associated primes of local cohomology modules*, Proc. Amer. Math. Soc., **128**(10) (2000), 2851–2853.
10. M. P. Brodmann, R. Y. Sharp, *Local cohomology : an algebraic introduction with geometric applications*, Cambridge University Press, 1998.
11. W. Bruns, J. Herzog, *Cohen-Macaulay rings*, Cambridge University Press, revised ed., 1998.
12. D. Delfino, T. Marley, *Cofinite modules of local cohomology*, J. Pure Appl. Algebra, **121**(1) (1997), 45–52.
13. K. Divaani-Aazar, A. Mafi, *Associated primes of local cohomology modules*, Proc. Amer. Math. Soc., **133** (2005), 655–660.
14. K. Divaani-Aazar, A. Mafi, *Associated primes of local cohomology modules of weakly Laskerian modules*, Comm. Algebra, **34** (2006), 681–690.
15. M. T. Dibaei, A. Nazari, *Graded local cohomology : attached and associated primes, asymptotic behaviors*, Comm. Algebra, **35** (2007), 1567–1576.
16. M. T. Dibaei, S. Yassemi, *Finiteness of extention functors of local cohomology modules*. Comm. Algebra, **34** (2006), 3097–3101.
17. M. T. Dibaei, S. Yassemi, *Associated primes of the local cohomology modules*, Abelian groups, rings, modules, and homological algebra, 49–56, Chapman and Hall/CRC, 2006.
18. M. T. Dibaei, S. Yassemi, *Bass numbers of local cohomology modules with respect to an ideal*, Algebr Represent Theor., **11** (2008), 299–306.
19. A. Grothendieck, *Cohomologie locale des faisceaux cohérents et théorèmes de Lefschetz locaux et globaux (SGA 2)*, North-Holland, Amsterdam, 1968.
20. R. Hartshorne, *Cohomological dimension of algebraic varieties*, Ann. of Math., **88** (1968), 403–450.
21. R. Hartshorne, *Affine duality and cofiniteness*, Invent. Math., **9** (1970), 145–164.
22. S. H. Hasanzadeh, A. Vahidi, *On vanishing and cofinitness of generalized local cohomology modules*, to appear in Comm. Algebra.
23. C. Huneke, *Problems on local cohomology : Free resolutions in commutative algebra and algebraic geometry*, (Sundance, UT, 1990), 93–108, Jones and Bartlett, 1992.
24. K. Khashyarmansh, *On the finiteness properties of extention and torsion functors of local cohomology modules*, Proc. Amer. Math. Soc., **135** (2007), 1319–1327.
25. K. Khashyarmansh, Sh. Salarian, *On the associated primes of local cohomology modules*, Comm. Algebra, **27** (1999), 6191–6198.
26. T. Marley and J. C. Vassilev, *Cofiniteness and associated primes of local cohomology modules*, J. Algebra, **256**(1) (2002), 180–193.
27. L. Melkersson, *Modules cofinite with respect to an ideal*, J. Algebra, **285** (2005), 649–668.
28. L. T. Nhan, *On generalized regular sequences and the finiteness for associated primes of local cohomology modules*, Comm. Algebra, **33** (2005), 793–806.

29. J. Rotman, *An introduction to homological algebra*, Academic Press, 1979.

30. H. Zöschinger, *Minimax Moduln*, J. Algebra, **102** (1986), 1–32.

<sup>1</sup> ARAK UNIVERSITY, BEHESHTI ST, P.O. BOX: 879, ARAK, IRAN.

*E-mail address:* m-aghapour@araku.ac.ir

<sup>2,3</sup> FACULTY OF MATHEMATICAL SCIENCES AND COMPUTER, TARBIAT MOALLEM UNIVERSITY, TEHRAN, IRAN.

*E-mail address:* taheri@saba.tmu.ac.ir

<sup>3</sup> PAYAME NOOR UNIVERSITY (PNU), IRAN.

*E-mail address:* vahidi.ar@gmail.com