

Evaluation Effects of Quercetin on Kidney Apoptosis in Streptozotocin-induced Diabetic Rat.¹Zahedi Afshin, ²Khaki Arash, ³Nori Mohammad, ⁴Fathiazad Fatemeh¹Department of Pathology, Rasht Branch, Islamic Azad University, Rasht-Iran.²Department of Pathology, Tabriz Branch, Islamic Azad University, Tabriz-Iran.³Department of Biochemistry, Tabriz University of medical sciences, Tabriz-Iran.⁴Department of Pharmacognosy, Tabriz University of medical sciences, Tabriz-Iran.⁵Department of Anatomical sciences, Tabriz University of medical sciences, Tabriz-Iran.

Zahedi Afshin, Khaki Arash, Nori Mohammad, Fathiazad Fatemeh; Evaluation Effects of Quercetin on Kidney Apoptosis in Streptozotocin-induced Diabetic Rat.

ABSTRACT

Aim: Quercetin is a strong antioxidant and long-term treatment of STZ-diabetic animals and it has been shown to reduce oxidative stress. Enhanced oxidative stress and changes in antioxidant capacity are considered to play an important role in the pathogenesis of chronic diabetes mellitus. **Methods:** Wistar male rat (n=40) were allocated into three groups, control group (n=10) and Quercetin (QR) group that received 15mg/kg (IP) QR, (n=10), and Diabetic group that received 55mg/kg (IP) streptozotocin (STZ) (n=20) which was subdivided to two groups of 10; STZ group and treatment group. Treatment group received 55mg/kg (IP) STZ plus 15mg/kg QR, daily for 4 weeks, respectively; however, the control group just received an equal volume of distilled water daily (IP). Diabetes was induced by a single (IP) injection of streptozotocin (55mg/kg). Animals were kept in standard condition. In 28 days after inducing diabetic 5cc blood were collected for TAC, MDA and Ox-LDL levels and kidney tissues of Rat in whole groups were removed then prepared for Apoptosis analysis by Tunel method. **Results:** Apoptotic cells significantly decreased in group that has received 15mg/kg (IP) Quercetin (P<0.05) in comparison to experimental groups (P<0.05). **Conclusion:** Since in our study 15mg/kg (IP) Quercetin have significantly preventive effect on kidney cells damages by reducing number of Apoptotic cells in kidney, so it seems that using it can be effective for treatment in Diabetic Rat.

Key words: Apoptosis, Diabetic, Quercetin, Streptozotocin, kidney, Rat.**Introduction**

Diabetes is a chronic disease that due to hyperglycemic. Hyperglycemic in long time have side effect in other tissues especially in liver. Due attention has been paid to the search of effective drugs in the field of traditional Chinese medicine (TCM). Diabetes mellitus is a group of syndromes characterized by hyperglycemia, altered metabolism of lipids, carbohydrates and proteins and an increased risk of complications from vascular diseases [3]. Enhanced oxidative stress and changes in antioxidant capacity are considered to play an important role in the pathogenesis of chronic diabetes mellitus [4,5]. Although the mechanisms underlying the alterations associated with diabetes mellitus are presently not well understood, hyperglycemia lead patients to increased oxidative stress because the production of several reducing sugars (through glycolysis and the polyol pathway) is enhanced [6]. These reducing sugars can easily react with lipids and proteins (nonenzymatic glycation reaction),

increasing the production of reactive oxygen species (ROS) [6]. Diabetes is the most common endocrine disease that leads to metabolic abnormalities involving regulation of carbohydrate metabolism. In addition to imbalanced carbohydrate metabolism, yet another major concern in diabetes is increased oxidative stress. Increased production of free radicals or ROS formation may induce oxidized LDL (Ox-LDL), which is key step in the sequence of events leading to atherosclerosis. Sustained hyperglycemia and increased oxidative stress, are the major players in the development of secondary complications in diabetes. These abnormalities produce pathologies including vasculopathies, neuropathies, ophthalmopathies and nephropathies, among many other medical derangements [7]. The balance of ROS and antioxidant is a major mechanism in preventing damage by oxidative stress. Therefore, the dietary supplement of antioxidants such as vitamins, flavonoids has been used to prevent the occurrence of many chronic diseases [8]. Many herbal such as: Barberry, Estragon, Rhus coriaria, Cinnamomum

Corresponding Author

Khaki Arash, Department of Pathology, Rasht Branch, Islamic Azad University, Rasht-Iran.

zelenicum, Hypericum perforatum and onion known anti diabetic effects and use to patient treatment. Quercetin is a well-known flavonoid and a strong antioxidant and long-term treatment of STZ-diabetic animals and it has been shown to reduce oxidative stress [9] we want to study the effect of Quercetin as a protective on kidney cells apoptosis.

Material and Methods

Animals:

Forty adult Wistar albino male rats were 8 weeks old and weighing 250 ± 10 g, they were obtained from animal facility of pasture institute of Iran. Male rats were housed in temperature controlled rooms (25°C) with constant humidity (40-70%) and 12h/12h light/dark cycle prior to use in experimental protocols. All animals were treated in accordance to the Principles of Laboratory Animal Care. The experimental protocol was approved by the Animal Ethical Committee in accordance with the guide for the care and use of laboratory animals prepared by Tabriz medical University. All Rats were fed a standard diet and water. The daily intake of animal water was monitored at least one week prior to start of treatments in order to determine the amount of water needed per experimental animal. Thereafter, the rats were randomly selected and divided into control ($n=10$) and Quercetin (QR) group that received 15mg/kg QR (IP), ($n=10$), and Diabetic group that received 55mg/kg (IP) streptozotocin (STZ) ($n=20$) which was subdivided to two groups of 10; STZ group and treatment group. Treatment group received 55mg/kg (IP) STZ plus 15mg/kg QR (IP). The control group just received an equal volume of 1cc distilled water daily (IP). Diabetes was induced by a single intra peritoneal (I.P) injection of streptozotocin (STZ, Sigma- U.S.A.) in 0.1 M citrate buffer (pH 4.0) at a dose of 55 mg/kg body weight. Quercetin (QR) injections were continued to the end of the study (for 4 weeks), [12].

Induction of experimental type 1, Diabetes:

Experimental type 1 diabetes was induced in rats by intra peritoneal (I.P) injection of 55 mg/kg streptozotocin (STZ) in distilled water. Control rats were received distilled water, only.

Blood glucose determination:

Blood samples were collected from the tail vein. Basal glucose levels were determined prior to STZ injection, using an automated blood glucose analyzer (Glucometer Elite XL). Sample collections were then made 48 h after STZ injection and blood glucose concentrations were determined and compared between groups. Rats with blood glucose concentrations above 300 mg/dl were declared

diabetic and were used in the experimental group. One week after the induction of experimental diabetes, protocol was started.

Quercetin preparation:

Quercetin powder was obtained from Sigma Chemical Company (St. Louis, MO, USA). It was dissolved and diluted with 20% glycerol in 0.9% normal saline, mixed vigorously and stored in a dark bottle at 4°C . The quercetin solution was freshly prepared each week.

Surgical Procedure:

In the 28th day, (at the end of the treatment period), the rats were killed with diethyl ether, and kidney tissues in control & experimental groups were immediately removed.

TUNEL analysis of apoptosis:

The *in-situ* DNA fragmentation was visualized by TUNEL method [16]. Briefly, dewaxed tissue sections were predigested with 20 mg/ml proteinase K for 20 min and incubated in phosphate buffered saline solution (PBS) containing 3 % H_2O_2 for 10 min to block the endogenous peroxidase activity. The sections were incubated with the TUNEL reaction mixture, fluorescein-dUTP (*in situ* Cell Death Detection, POD kit, Roche, Germany), for 60 min at 37°C . The slides were then rinsed three times with PBS and incubated with secondary anti fluorescein-POD-conjugate for 30 min. After washing three times in PBS, diaminobenzidine- H_2O_2 (DAB, Roche, Germany) chromogenic reaction was added on sections and counterstained with hematoxylin. As a control for method specificity, the step using the TUNEL reaction mixture was omitted in negative control serial sections, and nucleotide mixture in reaction buffer was used instead. Apoptotic cells were quantified by counting the number of TUNEL stained nuclei per liver tissues cross sections. Cross sections of 100 kidney tissues per specimen were assessed and the mean number of TUNEL positive apoptotic cells per cross-section was calculated.

Measurement of Serum Total Antioxidant capacity (TAS):

TAS was measured in serum by means of a commercial kit (Randox Co-England). The assay is based on the incubation of 2, 2'-azino-di-(3-ethylbenzthiazoline sulphonate) (ABTS) with a peroxidase (methmyoglobin) and hydrogen peroxide to produce the radical cation ABTS^+ , which has a relatively stable blue-green color, measured at 600 nm. The suppression of the color is compared with that of the Trolox, which is widely used as a traditional standard for TAS measurement assays,

and the assay results are expressed as Trolox equivalent (mmol/L), [14].

Measurement of Serum MDA:

Tissue MDA levels were determined by the thiobarbituric acid (TBA) method and expressed as nmol MDA formed/mL. Plasma MDA concentrations were determined with spectrophotometer. A calibration curve was prepared by using 1,1',3,3'-tetramethoxypropane as the standard [15].

Statistical analysis:

Statistical analysis was done using the ANOVA and test for comparison of data in the control group with the experimental groups. The results were expressed as mean \pm S.E.M (standard error of means). P-value less than 0.05 were considered significant and are written in the parentheses.

Amount of apoptotic cells among kidney cells

Number of apoptotic cells colored brown, in diabetic group was (12.25 \pm 1.14) and in Quercetin received diabetic group and control group was (6.15 \pm 8.17) and (1.05 \pm 0.41) respectively. These changes was significant as p value less than 0.05 (P<0.05).

Results of total blood anti oxidant capacity

Amount of total blood anti oxidant capacity in control group was (0.70 \pm 0.03 mmol/ml) and in experimental groups was 0.75 \pm 0.03, 0.32 \pm 0.04 and 0.61 \pm 0.05 mmol/ml respectively. Statistic analysis Dunnett(one side) shows significant differences between experimental groups in comparison to control group(P<0.05).(table 1).

Results of MDA (malondi aldehyde) level in blood

MDA level in control group was 0.25 \pm 0.04mmol/L and in experimental groups was 0.30 \pm 0.212, 4.1 \pm 0.06, 1.1 \pm 0.08 mmol/L respectively.

Statistic analysis Dunnett(one side) shows significant differences between experimental groups in comparison to control group(P<0.05).(Table 1).

Table 1:

Groups	control (n=10)	Quercetin (15mg/kg(IP)) (n=10)	STZ (55mg/kg (IP)) (n=10)	Quercetin +Stz 55mg/kg (IP) streptozotocin plus 15mg/kg Quercetin (n=10)
Apoptotic cells	1.05 \pm 0.41	1.01 \pm 8.17	12.25 \pm 1.14* (0.006)	6.15 \pm 8.17* (0.004)
(TAC) (nmol/ml)	0.70 \pm 0.03	0.75 \pm 0.03 * (0.006)	0.32 \pm 0.04 * (0.003)	0.61 \pm 0.05 * (0.004)
(MDA) (nmol/ml)	0.25 \pm 0.04	0.30 \pm 0.212 * (0.004)	4.1 \pm 0.06 * (0.008)	1.1 \pm 0.08 * (0.005)

Data are presented as mean \pm SE.

* P-value less than 0.05 were considered significant and are writing in the parentheses, (compared with the control group).

Discussion and conclusion:

Make use of onion and Quercetin in diabetic patient treatment has been experimented [17]. Investigations show onion and Quercetin decrease serum glucose level [18] but this reduction with onion has been significant [19]. Also Quercetin decreases oxidative stress and blood vessels damage in diabetic rats [20,21]. Other investigations show Quercetin increases the level of blood insulin and serum Ca²⁺ and Mg²⁺ [22]. Investigations show liver has an important role in carbohydrate metabolism since it is responsible for the balance of blood glucose level by means of glycogen sis and glycogenolysis therefore impaired hepatic function impairs metabolic homeostasis of glucose [2,3]. In the presence of impaired glucose metabolism and occurrence of hyperglycemia, genes involved in fatty acid storage were activated [4]. On the other hand, liver diseases can induce diabetes mellitus. This type of diabetes mellitus is clinically different from that of type II diabetes mellitus since it is less frequently associated with microangiopathy [22]. Insulin resistance occurs in muscular and adipose tissues combined with hyper insulinemia are pathophysiological bases of diabetes in liver disease

[23]. The etiology of liver disease is important in the incidence of diabetes mellitus since non alcoholic fatty liver disease (NALFD), alcohol, hepatitis C virus (HCV) and hemachromatosis are frequently associated with diabetes mellitus (22). Investigations show liver tissue damage and apoptosis induced by diabetes mellitus increase active O₂ specious. Flavonoids as an antioxidant factor found in nutrient such as fruit, vegetables, tea and black burgundy grape [24]. Flavonoids value in daily mail varies from 16 mg to 1000 mg. Quercetin as an important and main flavonoids found in human mail [25]. Investigations show Quercetin absorbs in small bowel. Useful effect of Quercetin in human health involves prevention of diabetes induced cataract, reduced blood vessels fragility, anti microbial, anti viral, anti allergy, and anti inflammatory effects and prevention of platelet aggregation [26,27,28]. One of the Quercetin anti oxidant mechanism is removal of free radical such as xanthine superoxide and xanthine oxidase [29]. Investigations show Quercetin in the chicken spermatogonial cell culture not only doesn't have harmful effects but also increases amount of spermatogonial cells and decreases oxidative effects. In this study like pervious investigations Quercetin decreases malondy aldeid and increases serum anti

oxidant capacity [30,31,32,33,34]. According our results it seems quercetin as an anti-oxidant can protection kidney cells from cell injury by modulating TAC and MDA in diabetic conditions, and it advise to using it in diabetes diseases.

References

- Jiang, G.Y., Practical Diabetes, 1996. 1st Edition. Beijing: People's Health Publishing House, 295.
- Holstein, A., S. Hinze, E. Thiessen, A. Plaschke, E.H. Egberts, 2002. Clinical implications of hepatogenous diabetes in liver irrhosis. *J Gastroenterol Hepatol.*, 17: 677-68.
- Tappy, L., K. Minehira, 2001. New data and new concepts on the role of the liver in glucose homeostasis. *Curr Opin Clin Nutr Metab Care*, 4: 273-277.
- Baynes, J.W., S.R. Thorpe, 1999. Role of oxidative stress in diabetic complications: a new perspective on an old paradigm. *Diabetes*, 48: 1-9.
- Wolff, S.P., Z.Y. Jiang, J.V. Hunt, 1991. Protein glycation and oxidative stress in diabetes mellitus and ageing. *Free Radic Biol Med.*, 10: 339-52.
- Palmeira, C.M., D.L. Santos, R. Seica, A.J. Moreno, M.S. Santos, 2001. Enhanced mitochondrial testicular antioxidant capacity in Goto-Kakizaki diabetic rats: role of coenzyme Q. *Am J Physiol Cell Physiol.*, 281: C1023-8.
- Sexton, W.J., J.P. Jarow, 1997. Effect of diabetes mellitus upon male reproductive function. *Urology*, 49: 508-13.
- Peluso, M.R., 2006. Flavonoids attenuate cardiovascular disease, inhibit phosphodiesterase, and modulate lipid homeostasis in adipose tissue and liver. *Exp Biol Med (Maywood)*, 231(8): 1287-99.
- Mahesh, T., V.P. Menon, 2004. Quercetin alleviates oxidative stress in streptozotocin-induced diabetic rats. *Phytother Res.*, 18: 123-127.
- Shrilatha, B., Muralidhara, 2007. Early oxidative stress in testis and epididymal sperm in streptozotocin-induced diabetic mice: its progression and genotoxic consequences. *Reprod Toxicol.*, 23(4): 578-87.
- Khaki, A., F. Fathi Azad, M. Nouri, A.A. Khaki, H. Jabbari-kh, M. Hammadeh, 2009. Evaluation of Androgenic Activity of *Allium cepa* on Spermatogenesis in Rat. *Folia Morphologica*, 68(1): 45-51.
- Coskun, O., M. Kanter, A. Korkmaz, S. Oter, 2005. Quercetin, a flavonoid antioxidant, prevents and protects streptozotocin-induced oxidative stress and beta-cell damage in rat pancreas. *Pharmacol Res.*, 51(2): 117-23.
- Khaki, A., M.G. Novin, A.A. Khaki, M. Nouri, E. Sanati, M. Nikmanesh, 2008. Comparative study of the effects of gentamicin, neomycin, streptomycin and ofloxacin antibiotics on sperm parameters and testis apoptosis in rats. *Pak J Biol Sci.*, 11(13): 1683-9.
- Feng, R., W. He, H. Ochi, 2001. A new murine oxidative stress model associated with senescence, *Mech Ageing Dev.*, 122: 547-559.
- Packer, A.T., J.M. Davies, T.L. Racanelli, K.J. Davies, 1982. Membrane effects of vitamin E deficiency: bioenergetics and surface charge density studies of skeletal muscle and liver mitochondria. *Ann NY Acad Sci.*, 393: 32-47.
- Huang, H.F.S., T.A. Linsenmeyer, M.T. Li, W. Giglio, R. Anesetti, J. von Hagen, J.E. Ottenweller, L. Pogach, 1995. Acute effects of spinal cord injury on the pituitary-testicular hormone axis and Sertoli cell functions: a time course study. *J Androl.*, 16: 148-157.
- Machha, Francis I. Achike, Ali Mohd Mustafa,nd Mohd Rais Mustafa, 2007. Quercetin, a flavonoid antioxidant, modulates endothelium-derived nitric oxide bioavailability in diabetic rat aortas. *Nitric Oxide*, 16: 442-447.
- Cunha, W.R., G.M. Arantes, D.S. Ferreira, R. Lucarini, M.L.A. Silva, N.A.J.C. Furtado, A.A. da Silva Filho, A.E.M. Crottiand A.R.B. Araújo, 2008. Hypoglycemic effect of *Leandra lacunosa* in normal and alloxan-induced diabetic rats. *Fitoterapia*, 79: 356-360.
- Mehmet, K., F. Mehmet, D. Senayi, Ayse, E. Murat, The effects of quercetin on bone minerals, biomechanical behavior and structure in streptozotocin-induced diabetic rats. *Cell Biochemistry and Function*, pp: 747-752.
- Kato, Yuka Minoshima, Jo Yamamoto, Isao Adachi, Alison A. Watson, and Robert J. Nash, 2008. Protective Effects of Dietary Chamomile Tea on Diabetic Complications *J. Agric. Food Chem.*, 56(17).
- Custro, N., A. Carroccio, A. Ganci, V. Scafidi, P. Campagna, L. Di Prima, G. Montalto, 2001. Glycemic homeostasis in chronic viral hepatitis and liver cirrhosis. *Diabetes Metab.*, 27: 476-481.
- Lecube, A., C. Hernandez, J. Genesca, J.I. Esteban, R. Jardi, R. Simo, 2004. High prevalence of glucose abnormalities in patients with hepatitis C virus infection: a multivariate analysis Considering the liver injury. *Diabetes Care*, 27: 1171-1175.
- Skibola, C.F., M.T. Smith, 2000. Potential health impacts of excessive flavonoid intake. *Free Radic. Biol. Med.*, 29: 375-383.
- Manach, C., C. Morand, V. Crespy, C. Demigne, O. Texier, F. Regeat, C. Remesy, 1998. Quercetin is recovered in human plasma as conjugated derivatives which retain antioxidant properties. *FEBS Lett.*, 426: 331-336.

25. Formica, J.V., W. Regelson, 1995. Review of the biology of Quercetin and related bioflavonoids. *Food Chem. Toxicol.*, 33: 1061-1080.
26. Hertog, M.G., P.C. Hollman, 1996. Potential health effects of the dietary flavonol quercetin. *Eur. J. Clin. Nutr.*, 50: 63-71.
27. Bors, W., C. Michel, K. Stettmaier, 1977. Antioxidant effects of flavonoids. *Biofactors*, 6: 399-402.
28. Dok-Go, H., K.H. Lee, H.J. Kim, E.H. Lee, J. Lee, Y.S. Song, Y.H. Lee, C. Jin, Y.S. Lee, J. Cho, 2003. Neuroprotective effects of antioxidative flavonoids, quercetin, (1)-dihydroquercetin and quercetin 3-methyl ether, isolated from *Opuntia ficus-indica* var. *saboten*. *Brain Res.*, 965: 130-136.
29. Mi, Y., C. Zhang, 2005. Protective Effect of Quercetin on Aroclor 1254-Induced Oxidative Damage in Cultured Chicken Spermatogonial Cells. *Toxicological Sciences*, 88(2): 545-550.
30. Mi, Y., C. Zhang, K. Taya, 2007. Quercetin protects spermatogonial cells from 2,4-d-induced Oxidative damage in embryonic chickens. *J Reprod Dev.*, 53(4): 749-54.
31. Chandel, A., S. Dhindsa, S. Topiwala, A. Chaudhuri, P. Dandona, 2008. Testosterone concentration in young patients with diabetes. *Diabetes Care*, 31(10): 2013-7.
32. Naziroğlu, M., 2003. Enhanced testicular antioxidant capacity in streptozotocin-induced diabetic rats: protective role of vitamins C and E and selenium. *Biol Trace Elem Res.*, 94(1): 61-72.
33. Sanders, R.A., F.M. Rauscher, J.B. Watkins, 2001. 3rd. Effects of quercetin on antioxidant defense in streptozotocin-induced diabetic rats. *J Biochem Mol Toxicol.*, 15(3): 143-9.
34. Drobiova, H., M. Thomson, K. Al-Qattan, R. Peltonen-Shalaby, Z. Al-Amin, M. Ali, 2009. Garlic Increases Antioxidant Levels in Diabetic and Hypertensive Rats Determined by a Modified Peroxidase Method. *Evid Based Complement Alternat Med.*, pp: 1-7.