

Socioeconomic Status in Health Research

One Size Does Not Fit All

Paula A. Braveman, MD, MPH

Catherine Cubbin, PhD

Susan Egerter, PhD

Sekai Chideya, MD, MPH

Kristen S. Marchi, MPH

Marilyn Metzler, RN

Samuel Posner, PhD

THE TERMS “SOCIOECONOMIC STATUS,” “socioeconomic position,” and “social class” (collectively, “SES”) are widely used in health research, reflecting widespread albeit often implicit recognition of the importance of socioeconomic factors for diverse health outcomes. Most health studies that consider SES treat socioeconomic characteristics as potential confounders of relationships between other variables and health. Others explicitly examine relationships between SES and health, seeking to better understand the associations that have been repeatedly observed.¹⁻²² The social science and social epidemiology literature consistently treats SES as a multidimensional construct comprising diverse socioeconomic factors (typically, economic resources, power, and/or prestige).^{6,7,23-30} Different socioeconomic factors could affect health at different times in the life course,³¹⁻³³ operating at different levels (eg, individual, household, neighborhood)^{30,34-37} and through different causal pathways^{3,6,7,9,38-45} (eg, by determining exposures, vulnerability, or even direct physiological effects).^{40,42,44,45} Socioeconomic factors can interact with other social characteristics, such as racial/ethnic group and sex, to produce different health effects across groups.^{9,10,15,27,46-52} Despite expert consensus that SES is

Problems with measuring socioeconomic status (SES)—frequently included in clinical and public health studies as a control variable and less frequently as the variable(s) of main interest—could affect research findings and conclusions, with implications for practice and policy. We critically examine standard SES measurement approaches, illustrating problems with examples from new analyses and the literature. For example, marked racial/ethnic differences in income at a given educational level and in wealth at a given income level raise questions about the socioeconomic comparability of individuals who are similar on education or income alone. Evidence also shows that conclusions about nonsocioeconomic causes of racial/ethnic differences in health may depend on the measure—eg, income, wealth, education, occupation, neighborhood socioeconomic characteristics, or past socioeconomic experiences—used to “control for SES,” suggesting that findings from studies that have measured limited aspects of SES should be reassessed. We recommend an outcome- and social group-specific approach to SES measurement that involves (1) considering plausible explanatory pathways and mechanisms, (2) measuring as much relevant socioeconomic information as possible, (3) specifying the particular socioeconomic factors measured (rather than SES overall), and (4) systematically considering how potentially important unmeasured socioeconomic factors may affect conclusions. Better SES measures are needed in data sources, but improvements could be made by using existing information more thoughtfully and acknowledging its limitations.

JAMA. 2005;294:2879-2888

www.jama.com

complex and multifactorial, most health studies that consider SES use a single socioeconomic variable measured at a single period and level. Often, very few categories are used (eg, poor/nonpoor, less than high school/high school graduation or more schooling), which could obscure important social gradients in health that apply across the entire socioeconomic spectrum.^{2,8,9} Occupation is frequently used as a measure of SES in Europe,^{11-14,16,17,52} while income or education is more commonly used in the United States. Regardless of the measure(s) used, most studies include SES

variables without justifying why a given measure was selected over others, without explaining its meaning for a given analysis, and without discussing how

Author Affiliations: Center on Social Disparities in Health and Department of Family and Community Medicine, University of California, San Francisco (Drs Braveman, Cubbin, and Egerter and Ms Marchi); and Epidemic Intelligence Service (Dr Chideya) and National Center for Chronic Disease Prevention and Health Promotion, Divisions of Adult and Community Health (Ms Metzler) and Reproductive Health (Dr Posner), Centers for Disease Control and Prevention, Atlanta, Ga.

Corresponding Author: Paula A. Braveman, MD, MPH, Center on Social Disparities in Health, University of California, San Francisco, 500 Parnassus Ave, MU-3E, Box 0900, San Francisco, CA 94143-0900 (braveman@fcm.ucsf.edu).

unmeasured socioeconomic differences might have affected findings.

This article critically examines several widespread standard practices in SES measurement, discussing key concepts and providing examples to illustrate problems with those practices. The examples were chosen to illustrate SES measurement issues, not to gain new knowledge about the role of socioeconomic or other social factors in the selected health indicators. Our goals were to increase awareness—not only among health researchers but also among policy makers and practitioners who use health research findings—about inherent problems with standard methods of measuring and interpreting socioeconomic characteristics in health research, as well as to encourage improved approaches. While excellent critiques have been available,^{10,25-28,53,54} the persistence of problematic ap-

proaches to SES measurement indicates that these documents have not yet influenced most researchers' practices. In many ways, this article parallels a recent *JAMA* Editorial critiquing the ways in which race/ethnicity is commonly measured in health research⁵⁵ and suggesting improved approaches based on more sound conceptualization.

Methods

To illustrate key points, we selected examples from existing literature and from new analyses of several national and statewide surveys, using the most recent data available that were suitable for the desired analyses. The new analyses were planned a priori to examine a range of (1) widely used, high-quality, population-based data sources that represent a range of data collection methods (eg, household vs tele-

phone survey); (2) indicators of different aspects of health (status, behaviors, care) in (3) different life stages; and (4) different commonly used and reasonable measures of income and education. Data sources, samples, and key SES and health-related variables for the new analyses are summarized in TABLE 1. We used 5 nationally or statewide-representative data sources with well-documented strengths and limitations: the multistate Behavioral Risk Factor Surveillance System (BRFSS), 2004⁵⁶; the National Health Interview Survey (NHIS) linked with the National Death Index, 1989-1994, with mortality follow-up through 1997 (mortality follow-up is not currently available for later years of the NHIS)⁵⁷; the Third National Health and Nutrition Examination Survey (NHANES III), 1988-1994 (both income and education were measured

Table 1. Summary of Data Sources, Samples, Measures of Socioeconomic Status (SES), and Dependent Variables Used in New Analyses of Health Surveys

Data Source	Age Group, y	Racial/Ethnic Groups	Measures of SES*	Dependent Variables (Health-Related Indicators)
Behavioral Risk Factor Surveillance System, 2004 ⁵⁶	18-74 (n = 255 896)	Black (non-Hispanic) Hispanic White (non-Hispanic)	Poverty level Income (categories) Educational level	Cigarette smoking Obesity Sedentary
National Health Interview Survey, 1989-1994 (mortality follow-up through 1997) ⁵⁷	18-64 (n = 380 552)	Black (non-Hispanic) Mexican American White (non-Hispanic)	Poverty level Income (continuous) Income/needs (continuous) Income (quartiles) Income/needs (quartiles) Educational level Education, y	Fair or poor health status All-cause mortality Heart disease mortality Homicide Motor vehicle fatalities
Third National Health and Nutrition Examination Survey, 1988-1994 ⁵⁸	1-5 (n = 5493) <17 (n = 12 388) 25-64 (n = 10 029)	Black (non-Hispanic) Mexican American White (non-Hispanic)	Poverty level Income (continuous) Educational level	Elevated lead levels (ages 1-5 y) Asthma (ages <17 y) Cigarette smoking (ages 25-64 y) Obesity (ages 25-64 y) Hypertension (ages 25-64 y) Diabetes (ages 25-64 y)
National Longitudinal Study of Adolescent Health, 1994-1995 ⁵⁹	11-21 (n = 14 282)	Black (non-Hispanic) Cuban Mexican American Puerto Rican White (non-Hispanic)	Poverty level Income (continuous) Income/needs (continuous) Income (quartiles) Income/needs (quartiles) Educational level	Cigarette smoking Initiation of sexual intercourse Violent behavior Binge drinking Marijuana use
California Maternal and Infant Health Assessment, 1999-2004 ⁶⁰	>24 (n = 13 952)	Asian/Pacific Islander (immigrant or US-born) Black Latina (immigrant or US-born) White	Poverty level Income (continuous) Income/needs (continuous) Income (quartiles) Income/needs (quartiles) Educational level Education, y	Unintended pregnancy Delayed or no prenatal care Low birth weight Never breastfed

*Poverty level indicates annual income estimated in categories defined by 100% increments of the federal poverty level according to family or household size (0%-100%, 101%-200%, 201%-300%, 301%-400%, >400%, unknown).

Income (categories) indicates annual income in categories corresponding to ranges in dollars; income (continuous), annual income in continuous dollars estimated as the midpoints of a given income range; income/needs (continuous), annual income in continuous dollars divided by family size; income (quartiles), annual income in continuous dollars grouped into empirical quartiles; and income/needs (quartiles), annual income in continuous dollars divided by family size, grouped into empirical quartiles.

Educational level indicates completed education in levels according to earned credentials (<9 years, some high school, high school graduate/General Equivalency Diploma, some college, college graduate or more); for California Maternal and Infant Health Assessment, <9 years and some high school are combined; the National Longitudinal Study of Adolescent Health also includes a category for unknown education. Education, y indicates completed education in years.

Table 2. Mean Family Income by Educational Level and Racial/Ethnic Group Among Adults Aged 18-64 Years—National Health Interview Survey, 1989-1994 (n = 380 552)

Educational Level, y	Mean Income (95% Confidence Interval), \$			
	Black	Mexican American	White	Overall
<9	15 503 (14 851-16 154)	19 104 (17 549-20 659)	22 707 (21 935-23 479)	20 551 (19 868-21 234)
9-11	17 743 (17 051-18 434)	22 377 (20 807-23 948)	28 573 (28 062-29 083)	25 945 (25 486-26 403)
12	25 337 (24 529-26 144)	30 945 (29 791-32 098)	37 853 (37 400-38 305)	36 022 (35 613-36 432)
13-15	33 026 (31 898-34 154)	37 642 (36 089-39 195)	43 197 (42 510-43 884)	41 857 (41 228-42 487)
≥16	46 815 (45 535-48 095)	48 055 (46 137-49 973)	55 277 (54 725-55 829)	54 606 (54 064-55 149)
Overall	27 937 (27 123-28 752)	27 927 (26 498-29 356)	42 113 (41 626-42 600)	39 716 (39 274-40 159)

more adequately in NHANES III than in the most recent NHANES data⁵⁸; the National Longitudinal Study of Adolescent Health (Add Health), 1994-1995 (the baseline survey, when parents or guardians of the adolescents responded to questions on income and education)⁵⁹; and the Maternal and Infant Health Assessment (MIHA), 1999-2004,⁶⁰ a statewide survey of postpartum women in California supported by the California Department of Health Services Maternal, Child and Adolescent Health Branch.⁶¹⁻⁶⁴ Using SAS software,⁶⁵ we examined correlations among multiple individual- or household-level measures of current education and income. To provide additional illustrative examples, we used SUDAAN software⁶⁶ and the specified data sources and samples to construct a series of weighted unadjusted and multivariate logistic regression models for each of 23 health-related indicators (listed as dependent variables in Table 1, counting obesity and smoking as separate indicators in different data sources or samples). These models examined how observed associations—expressed as odds ratios with 95% confidence intervals—between racial/ethnic groups and the health-related indicators varied by the income and/or education measures (see Table 1 footnotes) used as covariables to reflect SES; models also included age and sex (or parity, in MIHA), as well as family structure (in Add Health). Reference groups were those with the highest income or educational attainment. Log likelihood ratio tests were used to assess goodness of fit.⁶⁷ Further de-

tails about methods are included in a technical appendix available from the authors on request or at <http://www.ucsf.edu/csdh>.

Race/ethnicity, regarded as a social construct, was assessed in the study primarily to examine whether observed associations between racial/ethnic groups and health-related indicators varied by the income measures, education measures, or both used as covariables to reflect SES; we also examined how a range of socioeconomic factors varied by racial/ethnic group. While racial/ethnic groups were categorized differently depending on the data source (see Table 1 or text below), all categories were based on self-reported information about respondents' primary racial/ethnic identification. The MIHA included Latino/Hispanic as a separate racial/ethnic category; the BRFSS, NHIS, NHANES III, and Add Health surveys included separate questions about race and Hispanic ethnicity, which we used to create mutually exclusive racial/ethnic groups.

Critique of SES Measurement Approaches in Health Research

Education and Income: Not Interchangeable. Socioeconomic status is often implicitly or explicitly equated with income, especially in the United States. Despite wide recognition of its importance for health, income information is considered to be sensitive and is not measured in many studies. Information about education (typically measured as years completed or credentials of formal schooling) is more easily obtained and is frequently treated as a

proxy for income (or for SES overall). When both education and income are available, researchers may hesitate to include both in analytic models because of concerns about colinearity. Evidence from the literature and our new analyses indicates, however, that while standard measures of education and income are correlated, these correlations are generally not strong enough to justify using education as a proxy for income (or vice versa). Earnings can vary at similar educational levels, particularly across different social (eg, racial/ethnic, sex, age) groups.

For example, TABLE 2 shows, based on NHIS data, that black and Mexican American adults at every educational level had significantly lower mean incomes compared with white adults of similar educational attainment (eg, 33% and 18% lower for those with 12 years of schooling). This difference may reflect not only unequal employment opportunities and rewards but variations in educational quality.¹⁰ While measuring credentials is generally preferred to years of schooling for the purposes of reflecting SES,^{27,28,68,69} neither captures potentially dramatic differences across schools in prestige or resources, which may contribute to differences in future earnings. TABLE 3 displays correlations between education and income (see Table 3 footnotes) overall and within racial/ethnic groups for each data source or sample. Consistent with other studies,^{25,51,70,71} income-education correlations—most less than 0.50—were not strong enough to justify using income and education as proxies for each other.

Table 3. Spearman Correlation Coefficients Between Poverty Level and Educational Level Overall and by Racial/Ethnic Group, by Data Source and Age Group*

Race/Ethnicity	Data Source/Age Group						
	BRFSS, 2004, 18-74 y	NHIS, 1989-1994, 18-64 y†	NHANES III, 1988-1994			Add Health, 1994-1995, 11-21 y	MIHA, 1999-2004, >24y
			1-5 y	<17 y	25-64 y		
Black	0.45	0.42	0.44	0.42	0.42	0.45	0.59
Hispanic	0.50						
Mexican American		0.44	0.46	0.52	0.50	0.40	
Cuban						0.42	
Puerto Rican						0.37	
Asian/Pacific Islander							
Immigrant							0.48
US-born							0.47
Latina							
Immigrant							0.35
US-born							0.56
White	0.41	0.37	0.54	0.50	0.40	0.44	0.49
Overall	0.48	0.41	0.58	0.55	0.44	0.48	0.67

Abbreviations: Add Health, National Longitudinal Study of Adolescent Health; BRFSS, Behavioral Risk Factor Surveillance System; MIHA, California Maternal and Infant Health Assessment; NHANES III, Third National Health and Nutrition Examination Survey; NHIS, National Health Interview Survey.

*Poverty level indicates annual income estimated in categories defined by 100% increments of the federal poverty level according to family or household size (0%-100%, 101%-200%, 201%-300%, 301%-400%, >400%, unknown); educational level indicates completed education in levels according to earned credentials (<9 years, some high school, high school graduate/General Equivalency Diploma, some college, college graduate or more); for MIHA <9 years and some high school are combined; Add Health also includes a category for unknown education.

†With mortality follow-up through 1997.

Table 4. Odds Ratios for Racial/Ethnic Disparities in Fair or Poor Health Among Adults Aged 18-64 Years—National Health Interview Survey, 1989-1994 (n = 380 552)

SES Measure in Model*	OR (95% CI)†	
	Black Non-Hispanic	Mexican American
None (baseline)	2.47 (2.30-2.64)	2.16 (2.01-2.32)
Poverty level‡	1.65 (1.56-1.76)	1.30 (1.22-1.39)
Income, per \$1000§	1.67 (1.57-1.78)	1.44 (1.32-1.56)
Educational level	1.96 (1.85-2.08)	1.06 (0.99-1.14)
Education, per 1 y	2.02 (1.91-2.14)	0.89 (0.82-0.97)
Poverty level and educational level	1.53 (1.45-1.61)	0.86 (0.81-0.93)

Abbreviations: CI, confidence interval; OR, odds ratio; SES, socioeconomic status.

*All models included race/ethnicity, age, and sex.

†White non-Hispanic is the reference group.

‡Annual income estimated in categories defined by 100% increments of the federal poverty level according to family or household size (0%-100%, 101%-200%, 201%-300%, 301%-400%, >400%, unknown).

§Annual income in continuous dollars estimated as the midpoints of a given income range.

||Completed education in levels according to earned credentials (<9 years, some high school, high school graduate/General Equivalency Diploma, some college, college graduate or more).

To further illustrate practical implications of treating education as an income proxy, we examined how conclusions regarding racial/ethnic disparities in health might vary depending on whether income or education was used to measure SES. TABLE 4 displays results on racial/ethnic differences in self-rated health among adults aged 18 to 64 years from 6 multivariate models using NHIS data. The models differed by whether and how income and education were included as

covariables (see Table 4 footnotes); all models included race/ethnicity, age, and sex. When income (measured continuously or grouped by poverty level) was used to “adjust for SES,” Mexican Americans appeared to be at significantly increased risk of fair or poor health compared with non-Hispanic whites; however, risks appeared similar when educational level alone was included and significantly lower when either years of education or both poverty and educational levels were in-

cluded. Comparing blacks and whites, higher risks were seen for blacks regardless of the income or education measure used; however, the relevant odds ratios were significantly smaller after adjustment for income or poverty level. Similarly, TABLE 5 reveals disparities in receipt of delayed (after the first trimester) or no prenatal care among women who gave birth in California during 1999-2004; all models used MIHA data and included race/ethnicity, age, and parity. In separate comparisons of black and immigrant Latina women with white women, significant disparities in delayed or no prenatal care were seen when adjusting for either education measure alone but not when adjusting for either income measure alone or together with education. Although differences often were not statistically significant, one would have reached different conclusions about the significance, magnitude, or direction of racial/ethnic disparities in 10 of the 23 health indicators we examined, depending on whether (or less frequently, how) income and education were measured. For 20 of the 23 health indicators, model fit was significantly better

when both income and education were included, indicating that both factors should be considered (data available from authors on request or at <http://www.ucsf.edu/csdlh>).

Both income and education can influence the etiology of many health outcomes, in part through pathways involving material resources. Education also can reflect a range of noneconomic social characteristics (eg, general and health-related knowledge, literacy, and problem-solving skills; prestige; influence over others and one's own life)^{68,72-76} with important health effects; thus, for many health outcomes, education should be considered in addition to—not instead of— income, accumulated wealth, and other more directly economic factors.

Income: Not a Proxy for Wealth. Important links between economic resources and health, operating through diverse causal pathways, are widely recognized. While income is the most commonly used measure of economic resources in affluent countries, total accumulated economic resources or “wealth” could be at least as important for health; for example, wealth can buffer the effects of temporarily low income due to unemployment or illness and can reflect power or influence over others.⁷⁷ Furthermore, wealth can vary dramatically across different social groups with similar incomes. For example, TABLE 6, based on 2000 data from the Census Bureau, shows that, in the lowest income quintile, households headed by whites had on average more than 400 times as much wealth as those headed by blacks, and that, in higher income quintiles, whites had approximately 3 to 9 times the wealth of blacks; these differences were statistically significant (see Table 6 footnotes).⁷⁸

With notable exceptions,^{7,18,25,29,46,79-84} few health studies in affluent countries have measured wealth. Like income, wealth can be a sensitive topic, and standard methods of calculating net worth can be laborious. Some studies have observed wealth effects on health using simpler measures such as

home^{7,29,46,80} or car^{46,84} ownership or a single question on “liquid assets,”⁸³ even after controlling for income, another socioeconomic measure, or both.^{7,29,46,83,84} In summary, there are strong conceptual and empirical grounds for measuring wealth in health studies and for concluding that income is not an adequate measure of wealth.

Inadequacy of Standard US Occupational Categories. Occupational categories based on prestige, skills, social influence, and/or power have been the primary basis for socioeconomic classification in western European countries. Studies have repeatedly found strong relationships between occupational status using such classifications (eg, manual vs nonmanual labor, or graded hierarchies according to prestige or skills) and diverse health indicators,^{2,8,12,15-17,46,84-90} even after controlling for car ownership⁸⁴ or income and education.⁴⁸ Stepwise gradients in mortality and cardiovascular outcomes have

been observed in the British civil service hierarchy, with individuals in each occupational grade experiencing worse outcomes than those in the grade immediately above,⁵² even after adjusting for health-related behaviors^{2,17}; no group was poor, and financial access to medical care was unlikely to account for the differences. One explanation is variation in the psychosocial characteristics of one's occupation,¹⁷ including control over one's work.^{17,91-93}

Health studies in the United States rarely measure occupation, however. Occupational information is included in vital statistics and some national health surveys, but these data were not intended—and do not appear to be meaningful—as SES measures, because the categories include workers with diverse prestige, skills, power, and/or earnings. For example, the 2000 Standard Occupational Classification System⁹⁴ classifies chief executive officers, town clerks, and tenant farmers under “Man-

Table 5. Odds Ratios for Racial/Ethnic Disparities in Delayed or No Prenatal Care Among Childbearing Women Aged 25 Years and Older—California Maternal and Infant Health Assessment, 1999-2004 (n = 13 952)

SES Measure in Model*	OR (95% CI)†	
	Black	Latina Immigrant
None (baseline)	1.86 (1.53-2.26)	2.21 (1.90-2.57)
Poverty level‡	1.16 (0.95-1.43)	1.08 (0.91-1.28)
Income, per \$1000‡	1.12 (0.91-1.40)	1.05 (0.88-1.26)
Educational level§	1.57 (1.29-1.92)	1.47 (1.22-1.76)
Education, per 1 y	1.72 (1.41-2.10)	1.37 (1.12-1.67)
Poverty level and educational level	1.13 (0.91-1.39)	0.98 (0.82-1.18)

Abbreviations: CI, confidence interval; OR, odds ratio; SES, socioeconomic status.

*All models included race/ethnicity, age, and parity.

†White is the reference group.

‡See Table 4 footnotes for definition.

§Completed education in levels according to earned credentials (<9 years/some high school, high school graduate/General Equivalency Diploma, some college, college graduate or more).

Table 6. Median Net Worth in Dollars (Excluding Home Equity) by Quintile of Monthly Household Income and Householder's Racial/Ethnic Group, 2000*


Quintile	Black†	Hispanic‡	White Non-Hispanic
Lowest	57	500	24 000
Second	5275	5670	48 500
Third	11 500	11 200	59 500
Fourth	32 600	36 225	92 842
Highest	65 141	73 032	208 023

*From US Census Bureau Survey of Income and Program Participation.⁷⁸ Standard errors are not reported by the source; however, both the comparisons of blacks with whites and of Hispanics with whites are stated to be significantly different.

†Report does not indicate whether blacks are Non-Hispanic.

‡Persons of Hispanic origin may be of any race.

Figure 1. Percentage of College-Graduate Childbearing Women Aged 25 Years and Older With ≥ 1 College-Graduate Parent—California Maternal and Infant Health Assessment, 2003-2004 (n=1702)



Error bars indicate 95% confidence intervals.

agement” and head chefs, waitresses, and dishwashers under “Food Preparation and Serving Related.” Several authors^{25-28,53} have provided detailed critiques of occupational classifications. Some researchers have grouped the standard occupational classifications into manual and nonmanual work categories and found strong associations with health outcomes.⁴⁸ While this may be worthwhile when better alternatives are unavailable, more meaningful occupational classifications (eg, such as questions from Karasek^{92,93} measuring control and demands at work) are needed to capture information about important differences in occupation-related prestige or power that could affect health. Occupational measures, however, present challenges for classifying persons outside the paid labor force (thus disproportionately affecting women) or the chronically unemployed.

Importance of Past Socioeconomic Experiences. Childhood SES may influence adult health independently of adult SES.^{6,13,31-33,79,95-99} Acknowledging that the relative importance of SES early in life varies with health outcomes, Smith and Ben-Shlomo¹³ concluded that “studies with data on socioeconomic circumstances at only one stage of life are inadequate for fully elucidating the contribution of socioeconomic factors to health and mortality risk.” Past socioeconomic factors could

act independently or modify the effects of current factors on health.^{13,39,100} Changes in SES over time, including dramatic loss of income, can affect later health,^{90,101,102} and poverty can have cumulative health effects.^{99,103,104} Different socioeconomic factors could be more or less important at different stages of life.⁸³

Standard practice for measuring SES in most health studies, however, is to include only measures of current or recent socioeconomic characteristics; apart from some occupational health studies, socioeconomic experiences during earlier life stages are rarely measured. Socioeconomic status in childhood generally is related to SES in adulthood,¹⁰⁵⁻¹⁰⁷ but important SES characteristics earlier in life (eg, parents’ educational attainment) may not be reflected in measures of current SES, particularly for some population groups. For example, when their own parents’ education was examined among women older than 24 years who gave birth in California during 2003-2004, only 50% of college-graduate women overall had been raised by a college-graduate parent. In addition, this percentage varied significantly across racial/ethnic groups (for all comparisons except between blacks and Latinas); as shown in FIGURE 1, 50% of Asian/Pacific Islander, 34% of black, 20% of Latina, and 58% of white col-

lege-graduate women had been raised by a college-graduate parent. Lack of correspondence between educational attainment of women and their parents was seen for women at all educational levels (data available on request or at <http://www.ucsf.edu/csdlh>). Given the potentially important role that past socioeconomic experiences may play in health outcomes, practical measures of socioeconomic experiences earlier in life—eg, the highest educational attainment of either parent/guardian when an adult respondent was a child^{79,108}—should be developed and tested in diverse populations.

Importance of Neighborhood Socioeconomic Conditions. Accumulating evidence suggests that an individual’s health can be influenced by the socioeconomic characteristics of the neighborhood in which she or he lives, above and beyond her or his own individual-level SES.³⁴⁻³⁷ Socioeconomic characteristics of neighborhoods could affect health through features of the physical (“built”), social, or service environments¹⁰⁹⁻¹¹³ via multiple pathways.¹¹⁴ People with similar individual- or household-level socioeconomic characteristics can live in very different local environments. For example, as seen in FIGURE 2, Add Health data show that poor black and Puerto Rican adolescents (in families with incomes at or below the poverty line) lived in neighborhoods with higher poverty concentrations than their poor white counterparts; mean differences were statistically significant and similar patterns of racial/ethnic differences in neighborhood conditions were apparent among higher-income adolescents (data available on request or at <http://www.ucsf.edu/csdlh>) and among nationally representative samples of adults¹¹⁵ and women of reproductive age.¹¹⁶

Despite increasing recognition that both individual- and neighborhood-level SES can influence health, few health studies measure neighborhood features along with—rather than as proxies for—individual-level SES measures. In practice, however, many re-

searchers could consider characteristics of both individuals and their neighborhoods, despite theoretical and methodological challenges.¹¹⁷⁻¹¹⁹ By linking residential addresses with census geographic codes such as census tracts, census variables (eg, percentages of poor households or of unemployed adults) can be used to describe neighborhoods; similarly, geographic coordinates (longitude/latitude) can be linked with various data sources to describe physical environments in ways that are relevant for health research.

Conclusions and Recommendations

Based on the empirical evidence from new analyses and from the literature, we have reached 5 general conclusions, with corresponding recommendations for improving the measurement and interpretation of SES in health studies.


Different socioeconomic measures cannot be assumed to be interchangeable. The generally modest correlations between income and education indicate that measures of these 2 socioeconomic factors are not interchangeable; this is further supported by the examples showing that associations between racial/ethnic groups and health indicators can depend on which socioeconomic measures are used as covariables. Evidence of racial/ethnic differences in income at a given level of education, in wealth at a given level of income, in past SES at a given level of current SES, and in neighborhood SES at a given level of individual SES also supports the noninterchangeability of these variables across diverse populations. Multiple socioeconomic measures often can be included simultaneously in multivariable models without colinearity problems^{20-22,61-64,70,71}; stratified analyses also should be considered. Composite SES measures, or “indices,” also have been used to reflect multiple socioeconomic factors. However, few of the individual- or household-level (distinguished from community-level) indices have been validated. Most involve multiple questionable assumptions and, to

an even greater extent than simpler measures, may not apply over time and across populations.^{26,28} Furthermore, such composite measures, while potentially useful for classification in some studies, do not permit study of how particular SES factors influence health. Health researchers should justify the particular socioeconomic measures they have studied, avoiding claims to have measured SES overall.

Standard measures may not reflect important and relevant aspects of SES. Even studies that include multiple standard SES measures cannot examine all potentially important socioeconomic influences on health. Years of schooling received or earned credentials may not capture significant differences in educational quality that may also be relevant. Conventional US occupational groupings were not designed and are unlikely to adequately measure job-related socioeconomic characteristics that may affect health. Subjective SES¹²⁰ and experiences of economic hardship¹²¹ could plausibly influence health through psychophysiological pathways discussed earlier but are not explicitly reflected in standard SES measures. Rather than claiming to have “controlled for SES,” researchers should acknowledge the potentially relevant aspects of SES that could not be measured and explicitly consider the implications of unmeasured socioeconomic influences when interpreting findings.⁷⁰

Racial/ethnic differences are likely to reflect unmeasured socioeconomic differences. The concerns expressed above underscore the fact that—without measuring all relevant SES dimensions, life stages, and aggregation levels (eg, individual, household, neighborhood, city)—an observed racial/ethnic disparity in health cannot be considered “independent of SES.”¹²² However, racial/ethnic differences also cannot be assumed to be reducible to socioeconomic issues; for example, systematic socioeconomic differences between racial/ethnic groups can reflect racial discrimination at the institutional/structural level, personal experience, or both.^{10,54,123,124} Researchers who ob-

Figure 2. Distribution of Neighborhood Poverty Among Poor Adolescents by Racial/Ethnic Group—National Longitudinal Study of Adolescent Health, 1994-1995



Neighborhood poverty defined as percentage of families in the census tract who had 1989 income that was below the federal poverty level; poor adolescents defined as those aged 11-21 years and living in families with 1994 income that was at or below the federal poverty level.

serve racial/ethnic disparities in health outcomes should explicitly acknowledge the plausible role of unmeasured aspects of SES and other potentially relevant explanations,⁵⁵ including institutional or personal experiences of discrimination. We believe that conclusions from prior research about the nature of racial/ethnic (and other social) differences in health should be thoughtfully reassessed in light of findings and recommendations in this article as well as prior literature.*

A given SES measure may have different meanings in different social groups. Although our examples focused on racial/ethnic differences, similar issues may apply to other social groups categorized, for example, by age,⁸³ sex,^{50,52,127} or urban/rural location.¹²⁸ Whenever possible, researchers should examine how findings using a given SES measure vary across social groups.

Measures of SES should be selected and interpreted thoughtfully in the context of plausible explanatory pathways through which socioeconomic factors may influence health. Researchers should select socioeconomic factors systematically, considering whether eco-

*References 10, 27, 54, 55, 70, 114, 122, 123, 125, 126.

conomic resources, education, occupation, socioeconomic factors earlier in life, and neighborhood socioeconomic conditions at any life stage could plausibly be relevant to the particular health outcome and population of interest. They should assess the feasibility of adequately measuring each potentially relevant factor using available data; when a relevant factor cannot be measured, the implications of its exclusion should be considered thoughtfully and acknowledged. If a theoretically plausible argument or evidence from previous studies indicates that a particular socioeconomic factor is unlikely to be relevant for the given outcome or population of interest, it may not be necessary to include that factor. However, the approach we recommend would shift the burden to the researcher to justify why a study does not include measures of economic resources, education, occupation, past SES, and neighborhood socioeconomic conditions and to discuss fully the implications of unmeasured factors when stating conclusions.

Final Comments

We have presented conceptual arguments and empirical examples from diverse data sources, populations, and health-related indicators to illustrate several major and pervasive problems with the standard ways in which SES is measured in health research. The examples presented here highlight the potential consequences of inadequately measuring socioeconomic factors, particularly in studies of racial/ethnic disparities; similar caveats apply to studies of other social disparities, eg, by age, sex, or geographic residence. Our aim has been to provide convincing evidence for a wide audience of health researchers and users of health research findings. While previous authors have written about many of these issues, the messages apparently have not reached most clinical and public health researchers and policy makers. In particular, the evidence presented here, reinforced by the discussion in a recent *JAMA* Editorial on measuring racial/

ethnic group,⁵⁵ calls for a careful reassessment of conclusions about the etiologic basis of racial/ethnic differences in health based on studies with limited socioeconomic information.

These findings highlight the challenges in capturing the multidimensional nature of SES and the limitations of standard measures. Should we despair of measuring SES adequately? We think not. Better measures are needed, along with work to develop and validate measures and measurement approaches that are as comparable across studies and populations as possible. However, health research could be improved significantly with a more conceptually and empirically sound approach to measurement of SES. Building on our own and others' work, we call for a fundamentally different conceptual approach to measuring SES in health studies. This approach is outcome- and social group-specific and rests on considering explanatory pathways and mechanisms, measuring as much relevant socioeconomic information as possible, claiming to measure only what was measured, and systematically considering how important unmeasured socioeconomic factors may affect conclusions.

Author Contributions: Drs Braveman, Cubbin, and Egarter and Ms Marchi had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Study concept and design: Braveman, Cubbin, Egarter, Metzler.

Acquisition of data: Braveman, Cubbin, Chideya, Marchi.

Analysis and interpretation of data: Braveman, Cubbin, Egarter, Chideya, Marchi, Metzler, Posner.

Drafting of the manuscript: Braveman, Cubbin, Egarter.

Critical revision of the manuscript for important intellectual content: Braveman, Cubbin, Egarter, Chideya, Marchi, Metzler, Posner.

Statistical analysis: Cubbin, Marchi.

Obtained funding: Braveman, Cubbin, Egarter, Marchi.

Administrative, technical, or material support: Cubbin, Chideya, Marchi, Metzler, Posner.

Study supervision: Braveman, Cubbin.

Financial disclosures: None reported.

Funding/Support: The research for this article was supported by the Division of Reproductive Health, Coordinating Center for Health Promotion, Centers for Disease Control and Prevention (CDC-ATPM Cooperative Agreement TS-0842). One of the data sources (Maternal and Infant Health Assessment, MIHA) is a collaborative effort of the authors (P.A.B., S.E., K.S.M.) with the California Department of Health Services Maternal and Child Health Branch (MCAH), supported by MCAH (see Acknowledgment). Dr Chideya's participation in this effort was funded by the Department of Family and Community Medicine, Univer-

sity of California, San Francisco, which supported her as a Kellogg Health Disparities Scholar.

Role of the Sponsors: None of the funding sources for this study had any role in the design or conduct of the study; the collection, management, analysis, or interpretation of the data; or the preparation of the manuscript. Anonymous reviewers at the Centers for Disease Control and Prevention reviewed and approved the technical quality of the manuscript. The role of the California Department of Health Services in the collection of the data from the MIHA is noted above.

Acknowledgment: We acknowledge MCAH researchers Shabbir Ahmad, DVM, MPH, Eugene Takahashi, PhD, and Moreen Libet, PhD, for their contributions to the design and conduct of MIHA. We thank Nicole Wojtal and Pinal Shah, Department of Family and Community Medicine, University of California, San Francisco (UCSF), for their assistance with research and preparing the manuscript, and Roberto Vargas, MD, MPH, Department of Medicine, University of California, Los Angeles, Craig Pollack, MD, Department of Medicine, UCSF, and Thomas Bodenheimer, MD, Department of Family and Community Medicine, UCSF, for helpful comments on drafts.

REFERENCES

1. Isaacs SL, Schroeder SA. Class—the ignored determinant of the nation's health. *N Engl J Med*. 2004; 351:1137-1142.
2. Marmot MG, Rose G, Shipley M, Hamilton PJ. Employment grade and coronary heart disease in British civil servants. *J Epidemiol Community Health*. 1978; 32:244-249.
3. Adler NE, Boyce WT, Chesney MA, Folkman S, Syme SL. Socioeconomic inequalities in health: no easy solution. *JAMA*. 1993;269:3140-3145.
4. Kaplan GA. People and places: contrasting perspectives on the association between social class and health. *Int J Health Serv*. 1996;26:507-519.
5. Backlund E, Sorlie PD, Johnson NJ. The shape of the relationship between income and mortality in the United States: evidence from the National Longitudinal Mortality Study. *Ann Epidemiol*. 1996;6:12-20.
6. Lynch JW, Kaplan GA, Salonen JT. Why do poor people behave poorly? variations in adult health behaviour and psychosocial characteristics, by stage of the socioeconomic life course. *Soc Sci Med*. 1997;44: 809-820.
7. Macintyre S, Ellaway A, Der G, Ford G, Hunt K. Do housing tenure and car access predict health because they are simply markers of income or self esteem? A Scottish study. *J Epidemiol Community Health*. 1998;52:657-664.
8. Marmot MG, Smith GD, Stansfeld S, et al. Health inequalities among British civil servants: the Whitehall II study. *Lancet*. 1991;337:1387-1393.
9. Marmot M, Ryff CD, Bumpass LL, Shipley M, Marks NF. Social inequalities in health: next questions and converging evidence. *Soc Sci Med*. 1997;44:901-910.
10. Williams DR. Race, socioeconomic status, and health: the added effects of racism and discrimination. *Ann N Y Acad Sci*. 1999;896:173-188.
11. Avendano M, Kunst AE, van Lenthe F, et al. Trends in socioeconomic disparities in stroke mortality in six European countries between 1981-1985 and 1991-1995. *Am J Epidemiol*. 2005;161:52-61.
12. Black D, Morris JN, Smith C, Townsend P. The Black Report. In: Townsend P, Davidson N, Whitehead M, eds. *Inequalities in Health: The Black Report/The Health Divide*. 2nd ed. London, England: Penguin; 1999:29-213.
13. Smith GD, Ben-Shlomo Y. Geographical and social class differentials in stroke mortality—the influence of early-life factors: comment on papers by Maheswaran and colleagues. *J Epidemiol Community Health*. 1997;51:134-137.

14. Stansfeld SA, Head J, Marmot MG. Explaining social class differences in depression and well-being. *Soc Psychiatry Psychiatr Epidemiol*. 1998;33:1-9.
15. Mackenbach JP, Kunst AE, Groenof F, et al. Socioeconomic inequalities in mortality among women and among men: an international study. *Am J Public Health*. 1999;89:1800-1806.
16. Kunst AE, Groenof F, Mackenbach JP, Working EU; Group on Socioeconomic Inequalities in Health. Occupational class and cause specific mortality in middle aged men in 11 European countries: comparison of population-based studies. *BMJ*. 1998;316:1636-1642.
17. Marmot MG, Bosma H, Hemingway H, Brunner E, Stansfeld S. Contribution of job control and other risk factors to social variations in coronary heart disease incidence. *Lancet*. 1997;350:235-239.
18. Benzeval M, Judge K, Shouls S. Understanding the relationship between income and health: how much can be gleaned from cross-sectional data? *Soc Policy Adm*. 2001;35:376-396.
19. Mackenbach JP, Martikainen P, Looman CW, Dalstra JA, Kunst AE, Lahelma E. The shape of the relationship between income and self-assessed health: an international study. *Int J Epidemiol*. 2005;34:286-293.
20. Lantz PM, House JS, Lepkowski JM, Williams DR, Mero RP, Chen J. Socioeconomic factors, health behaviors, and mortality: results from a nationally representative prospective study of US adults. *JAMA*. 1998;279:1703-1708.
21. Lannin DR, Mathews HF, Mitchell J, Swanson MS, Swanson FH, Edwards MS. Influence of socioeconomic and cultural factors on racial differences in late-stage presentation of breast cancer. *JAMA*. 1998;279:1801-1807.
22. Lowry R, Kann L, Collins JL, Kolbe LJ. The effect of socioeconomic status on chronic disease risk behaviors among US adolescents. *JAMA*. 1996;276:792-797.
23. Weber M. Classes, status groups and parties. In: Runciman WG, ed. *Max Weber: Selections in Translation*. Cambridge, England: Cambridge University Press; 1978:43-61.
24. Wright EO. Review of Coleman RP, Rainwater L, McClelland KA. Social standing in America—new dimensions of class. *Am J Sociol*. 1980;85:1443-1449.
25. Abramson JH, Gofin R, Habib J, Pridan H, Gofin J. Indicators of social class: a comparative appraisal of measures for use in epidemiological studies. *Soc Sci Med*. 1982;16:1739-1746.
26. Berkman LF, Macintyre S. The measurement of social class in health studies: old measures and new formulations. In: Kogevinas M, Pearce N, Susser M, Boffetta P, eds. *Social Inequalities and Cancer*. Lyon, France: International Agency for Research on Cancer; 1997:51-64. IARC Scientific Publications No. 138.
27. Krieger N, Williams DR, Moss NE. Measuring social class in US public health research: concepts, methodologies, and guidelines. *Annu Rev Public Health*. 1997;18:341-378.
28. Liberatos P, Link BG, Kelsey JL. The measurement of social class in epidemiology. *Epidemiol Rev*. 1988;10:87-121.
29. Sundquist J, Johansson SE. Indicators of socioeconomic position and their relation to mortality in Sweden. *Soc Sci Med*. 1997;45:1757-1766.
30. Diez Roux AV, Merkin SS, Arnett D, et al. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med*. 2001;345:99-106.
31. Smith GD, Hart C, Blane D, Hole D. Adverse socioeconomic conditions in childhood and cause specific adult mortality: prospective observational study. *BMJ*. 1998;316:1631-1635.
32. Rahkonen O, Lahelma E, Huuhka M. Past or present? childhood living conditions and current socioeconomic status as determinants of adult health. *Soc Sci Med*. 1997;44:327-336.
33. Galobardes B, Lynch JW, Smith GD. Childhood socioeconomic circumstances and cause-specific mortality in adulthood: systematic review and interpretation. *Epidemiol Rev*. 2004;26:7-21.
34. Yen IH, Syme SL. The social environment and health: a discussion of the epidemiologic literature. *Annu Rev Public Health*. 1999;20:287-308.
35. Robert SA. Socioeconomic position and health: the independent contribution of community socioeconomic context. *Annu Rev Sociol*. 1999;25:489-516.
36. Pickett KE, Pearl M. Multilevel analyses of neighbourhood socioeconomic context and health outcomes: a critical review. *J Epidemiol Community Health*. 2001;55:111-122.
37. Sampson RJ, Morenoff JD, Gannon-Rowley T. Assessing "neighborhood effects": social processes and new directions in research. *Annu Rev Sociol*. 2002;28:443-478.
38. Marmot M. Inequalities in health. *N Engl J Med*. 2001;345:134-136.
39. Lynch JW, Everson SA, Kaplan GA, Salonen R, Salonen JT. Does low socioeconomic status potentiate the effects of heightened cardiovascular responses to stress on the progression of carotid atherosclerosis? *Am J Public Health*. 1998;88:389-394.
40. Taylor SE, Repetti RL, Seeman T. Health psychology: what is an unhealthy environment and how does it get under the skin? *Annu Rev Psychol*. 1997;48:411-447.
41. Krieger N. Theories for social epidemiology in the 21st century: an ecosocial perspective. *Int J Epidemiol*. 2001;30:668-677.
42. Singer B, Ryff C. Racial and ethnic inequalities in health: environmental, psychological, and physiological pathways. In: Devlin B, Fienberg SE, Resnick DP, Roeder K, eds. *Intelligence, Genes, and Success: Scientists Respond to the Bell Curve*. New York, NY: Copernicus; 1997:89-122.
43. Wilkinson RG. Socioeconomic determinants of health: health inequalities: relative or absolute material standards? *BMJ*. 1997;314:591-595.
44. Steptoe A, Marmot M. The role of psychobiological pathways in socio-economic inequalities in cardiovascular disease risk. *Eur Heart J*. 2002;23:13-25.
45. McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med*. 1998;338:171-179.
46. Arber S, Ginn J. Gender and inequalities in health in later life. *Soc Sci Med*. 1993;36:33-46.
47. Nicolaidis C, Ko CW, Saha S, Koepsell TD. Racial discrepancies in the association between paternal vs. maternal educational level and risk of low birthweight in Washington State. *BMC Pregnancy Childbirth*. 2004;4:10.
48. Barbeau EM, Krieger N, Soobader MJ. Working class matters: socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. *Am J Public Health*. 2004;94:269-278.
49. Pearl M, Braveman P, Abrams B. Relationship of neighborhood socioeconomic characteristics to birth weight among 5 ethnic groups in California. *Am J Public Health*. 2001;91:1808-1814.
50. Stafford M, Cummins S, Macintyre S, Ellaway A, Marmot M. Gender differences in the associations between health and neighbourhood environment. *Soc Sci Med*. 2005;60:1681-1692.
51. Gazmararian JA, Adams MM, Pamuk ER. Associations between measures of socioeconomic status and maternal health behavior. *Am J Prev Med*. 1996;12:108-115.
52. Griffin JM, Fuhrer R, Stansfeld SA, Marmot M. The importance of low control at work and home on depression and anxiety: do these effects vary by gender and social class? *Soc Sci Med*. 2002;54:783-798.
53. Krieger N, Fee E. Social class: the missing link in U.S. health data. *Int J Health Serv*. 1994;24:25-44.
54. Williams DR. Race/ethnicity and socioeconomic status: measurement and methodological issues. *Int J Health Serv*. 1996;26:483-505.
55. Winker MA. Measuring race and ethnicity: why and how? *JAMA*. 2004;292:1612-1614.
56. Centers for Disease Control and Prevention. Behavior Risk Factor Surveillance System (BRFSS). 2000. Available at: <http://www.cdc.gov/brfss>. Accessibility verified October 24, 2005.
57. Centers for Disease Control and Prevention. National Health Interview Survey (NHIS). Available at: <http://www.cdc.gov/nchs/nhis.htm>. Accessibility verified October 24, 2005.
58. Centers for Disease Control and Prevention. Third National Health and Nutrition Examination Survey (NHANES III), 1988-1994, Public Use Data Files. Available at: http://www.cdc.gov/nchs/products/elec_prods/subject/nhanes3.htm. Accessibility verified October 24, 2005.
59. National Institutes of Health. National Longitudinal Study of Adolescent Health (Add Health), 1994-1995. Available at: http://www.nichd.nih.gov/about/cpr/dbs/res_add.htm. Accessibility verified October 24, 2005.
60. Maternal and Infant Health Assessment (MIHA). Available at: http://www.ucsf.edu/csdp/research_projects.htm. Accessibility verified October 24, 2005.
61. Cubbin C, Braveman P, Marchi K, Chavez G, Santelli J, Gilbert BJ. Socioeconomic and racial/ethnic disparities in unintended pregnancy among postpartum women in California. *Matern Child Health J*. 2002;6:237-246.
62. Galbraith AA, Egerter SA, Marchi KS, Chavez G, Braveman PA. Newborn early discharge revisited: are California newborns receiving recommended postnatal services? *Pediatrics*. 2003;111:364-371.
63. Chung E, Hung Y, Marchi K, Chavez G, Braveman P. Infant sleep position: associated maternal and infant factors. *Ambul Pediatr*. 2003;3:234-239.
64. Heck KE, Schoendorf KC, Chavez GF, Braveman P. Does postpartum length of stay affect breastfeeding duration? a population-based study. *Birth*. 2003;30:153-159.
65. SAS. Version 9.1. Cary, NC: SAS Institute Inc; 2002.
66. SUDAAN. Version 9.0.1. Research Triangle Park, NC: Research Triangle Institute; 2005.
67. McCullagh P, Nelder JA. *Generalized Linear Models*. London, England: Chapman & Hall; 1989. *Monographs on Statistics and Applied Probability*. 2nd ed. Vol 37.
68. Ross CE, Mirowsky J. Refining the association between education and health: the effects of quantity, credential, and selectivity. *Demography*. 1999;36:445-460.
69. Hadden WC. Annotation: the use of educational attainment as an indicator of socioeconomic position. *Am J Public Health*. 1996;86:1525-1526.
70. Braveman P, Cubbin C, Marchi K, Egerter S, Chavez G. Measuring socioeconomic status/position in studies of racial/ethnic disparities: maternal and infant health. *Public Health Rep*. 2001;116:449-463.
71. Winkleby MA, Jatulis DE, Frank E, Fortmann SP. Socioeconomic status and health: how education, income, and occupation contribute to risk factors for cardiovascular disease. *Am J Public Health*. 1992;82:816-820.
72. Winch C. The economic aims of education. *J Philos Educ*. 2002;36:101-117.
73. Mirowsky J, Ross CE. Education, personal control, lifestyle and health: a human capital hypothesis. *Res Aging*. 1998;20:415-449.
74. Reynolds JR, Ross CE. Social stratification and health: education's benefit beyond economic status and social origins. *Soc Probl*. 1998;45:221-247.
75. Ross CE, Van Willigen M. Education and the subjective quality of life. *J Health Soc Behav*. 1997;38:275-297.
76. Ross CE, Wu C. The links between education and health. *Am Sociol Rev*. 1995;60:719-745.
77. Brenner MH. Health, productivity, and the eco-

- nomie environment: dynamic role of socioeconomic status. In: Green GM, Baker F, eds. *Work, Health and Productivity*. New York, NY: Oxford University Press; 1991:241-255.
78. US Census Bureau. *Net Worth and Asset Ownership of Households: 1998 and 2000*. Washington, DC: US Census Bureau; May 2003. Current Population Reports P70-88.
79. Cohen S, Doyle WJ, Turner RB, Alper CM, Skoner DP. Childhood socioeconomic status and host resistance to infectious illness in adulthood. *Psychosom Med*. 2004;66:553-558.
80. Macintyre S, McKay L, Der G, Hiscock R. Socioeconomic position and health: what you observe depends on how you measure it. *J Public Health Med*. 2003;25:288-294.
81. Smith GD, Shipley MJ, Rose G. Magnitude and causes of socioeconomic differentials in mortality: further evidence from the Whitehall study. *J Epidemiol Community Health*. 1990;44:265-270.
82. Von dem Knesebeck O, Luschen G, Cockerham WC, Siegrist J. Socioeconomic status and health among the aged in the United States and Germany: a comparative cross-sectional study. *Soc Sci Med*. 2003;57:1643-1652.
83. Robert S, House JS. SES differentials in health by age and alternative indicators of SES. *J Aging Health*. 1996;8:359-388.
84. Smith GD, Bartley M, Blane D. The Black Report on socioeconomic inequalities in health 10 years on. *BMJ*. 1990;301:373-377.
85. Chandola T. Social inequality in coronary heart disease: a comparison of occupational classifications. *Soc Sci Med*. 1998;47:525-533.
86. Diderichsen F, Hallqvist J. Trends in occupational mortality among middle-aged men in Sweden 1961-1990. *Int J Epidemiol*. 1997;26:782-787.
87. Vagero D, Erikson R. Socioeconomic inequalities in morbidity and mortality in western Europe. *Lancet*. 1997;350:516.
88. Macintyre S. The patterning of health by social position in contemporary Britain: directions for sociological research. *Soc Sci Med*. 1986;23:393-415.
89. Pamuk ER. Social-class inequality in infant mortality in England and Wales from 1921 to 1980. *Eur J Popul*. 1988;4:1-21.
90. Basso O, Olsen J, Johansen AMT, Christensen K. Change in social status and risk of low birth weight in Denmark: population based cohort study. *BMJ*. 1997;315:1498-1502.
91. Theorell T, Tsutsumi A, Hallqvist J. Decision latitude, job strain, and myocardial infarction: a study of working men in Stockholm. *Am J Public Health*. 1998;88:382-388.
92. Karasek R, Brisson C, Kawakami N, Houtman I, Bongers P, Amick B. The Job Content Questionnaire (JCQ): an instrument for internationally comparative assessments of psychosocial job characteristics. *J Occup Health Psychol*. 1998;3:322-355.
93. Karasek R. Lower health risk with increased job control among white collar workers. *J Organ Behav*. 1990;11:171-185.
94. US Census Bureau. 2000 Standard Occupational Classification System. Available at: <http://www.census.gov/hhes/www/oiindex/view.html>. Accessibility verified October 24, 2005.
95. Blane D, Hart CL, Smith GD, et al. Association of cardiovascular disease risk factors with socioeconomic position during childhood and during adulthood. *BMJ*. 1996;313:1434-1438.
96. Kuh D, Ben-Shlomo Y, eds. *A Life Course Approach to Chronic Disease Epidemiology*. New York, NY: Oxford University Press; 1997.
97. van de Mheen H, Stronks K, Looman CWN, Mackenbach JP. Role of childhood health in the explanation of socioeconomic inequalities in early adult health. *J Epidemiol Community Health*. 1998;52:15-19.
98. Bosma H, van de Mheen HD, Mackenbach JP. Social class in childhood and general health in adulthood: questionnaire study of contribution of psychological attributes. *BMJ*. 1999;318:18-22.
99. Power C, Matthews S, Manor O. Inequalities in self-rated health: explanations from different stages of life. *Lancet*. 1998;351:1009-1014.
100. Frankel S, Elwood P, Sweetnam P, Yamell J, Smith GD. Birthweight, body-mass index in middle age, and incident coronary heart disease. *Lancet*. 1996;348:1478-1480.
101. Duncan G. Income dynamics and health. *Int J Health Serv*. 1996;26:419-444.
102. McDonough P, Duncan G, Williams D, House J. Income dynamics and adult mortality in the United States, 1972 through 1989. *Am J Public Health*. 1997;87:1476-1483.
103. Korenman S, Miller JE, Sjaastad JE. Long-term poverty and child development in the United States: results from the NLSY. *Child Youth Serv Rev*. 1995;17:127-155.
104. Miller JE, Korenman S. Poverty and children's nutritional status in the United States. *Am J Epidemiol*. 1994;140:233-242.
105. Duncan GJ, Yeung WJ, Brooks-Gunn J, Smith JR. How much does childhood poverty affect the life chances of children? *Am Sociol Rev*. 1998;63:406-423.
106. Warren JR, Hauser RM. Social stratification across three generations: new evidence from the Wisconsin longitudinal study. *Am Sociol Rev*. 1997;62:561-572.
107. Rodgers JR. Intergenerational transmission of poverty in the United States. *Soc Sci Q*. 1995;76:178-194.
108. US Department of Labor. NLSY79: National Longitudinal Survey of Youth, 1979. Available at: <http://www.bls.gov/nls/nlsy79.htm>. Accessibility verified October 24, 2005.
109. Morland K, Wing S, Diez Roux A. The contextual effect of the local food environment on residents' diets: the atherosclerosis risk in communities study. *Am J Public Health*. 2002;92:1761-1767.
110. Macintyre S. The social patterning of exercise behaviours: the role of personal and local resources. *Br J Sports Med*. 2000;34:6.
111. Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and violent crime: a multilevel study of collective efficacy. *Science*. 1997;277:918-924.
112. Macintyre S, Ellaway A, Cummins S. Place effects on health: how can we conceptualise, operationalise and measure them? *Soc Sci Med*. 2002;55:125-139.
113. Morrison RS, Wallenstein S, Natale DK, Senzel RS, Huang L-L. "We don't carry that"—failure of pharmacies in predominantly nonwhite neighborhoods to stock opioid analgesics. *N Engl J Med*. 2000;342:1023-1026.
114. Massey DS. Segregation and stratification: a biosocial perspective. *The DuBois Review: Social Science Research on Race*. 2004;1:1-19.
115. Cubbin C, Hadden WC, Winkleby MA. Neighborhood context and cardiovascular disease risk factors: the contribution of material deprivation. *Ethn Dis*. 2001;11:687-700.
116. Brewster KL. Race differences in sexual activity among adolescent women: the role of neighborhood characteristics. *Am Sociol Rev*. 1994;59:408-424.
117. Diez Roux AV. Invited commentary: places, people, and health. *Am J Epidemiol*. 2002;155:516-519.
118. Diez-Roux AV. Bringing context back into epidemiology: variables and fallacies in multilevel analysis. *Am J Public Health*. 1998;88:216-222.
119. Tienda M. Poor people, poor places: deciphering neighborhood effects on poverty outcomes. In: Huber J, ed. *Macro-micro Linkages in Sociology*. Newbury Park, Calif: Sage; 1991:244-262.
120. Singh-Manoux A, Adler NE, Marmot MG. Subjective social status: its determinants and its association with measures of ill-health in the Whitehall II study. *Soc Sci Med*. 2003;56:1321-1333.
121. Mayer SE, Jencks C. Poverty and the distribution of material hardship. *J Hum Resour*. 1989;24:88-114.
122. Kaufman JS, Cooper RS, McGee DL. Socioeconomic status and health in blacks and whites: the problem of residual confounding and the resiliency of race. *Epidemiology*. 1997;8:621-628.
123. Williams DR, Collins C. Racial residential segregation: a fundamental cause of racial disparities in health. *Public Health Rep*. 2001;116:404-416.
124. Muntaner C, Nieto FJ, O'Campo P. Additional clarification re: "The Bell curve: on race, social class, and epidemiologic research." *Am J Epidemiol*. 1997;146:607-608.
125. Bamshad M. Genetic influences on health: does race matter [published correction appears in *JAMA*. 2005;294:1620]? *JAMA*. 2005;294:937-946.
126. Sankar P, Cho MK, Condit CM, et al. Genetic research and health disparities. *JAMA*. 2004;291:2985-2989.
127. American Academy of Pediatrics; Committee on Pediatric Research. Race/ethnicity, gender, socioeconomic status—research exploring their effects on child health: a subject review. *Pediatrics*. 2000;105:1349-1351.
128. Gilthorpe MS, Wilson RC. Rural/urban differences in the association between deprivation and healthcare utilisation. *Soc Sci Med*. 2003;57:2055-2063.