

Kabuki Syndrome and Anorectal Malformations: Implications for Diagnosis and Treatment

Sotirios Siminas¹ Colin Tennant Baillie² Richard Turnock²

¹Department of Paediatric Surgery and Urology, Manchester Children's Hospital, Manchester, United Kingdom

²Paediatric Surgery Department, Alder Hey Children's Hospital, Liverpool, United Kingdom

Address for correspondence Sotirios Siminas, MD, PhD, FEBPS, Department of Paediatric Surgery and Urology, Manchester Children's Hospital, Oxford Road, Manchester M13 9WL, United Kingdom (e-mail: siminas1@gmail.com).

Eur J Pediatr Surg Rep 2015;3:54–58.

Abstract

Kabuki syndrome (KS) is a rare genetic condition characterized by a distinctive facies, intellectual disability, growth delay, and a variety of skeletal, visceral, and other anomalies, including anorectal malformations (ARMs). We present two cases of female patients with KS, diagnosed and successfully managed at our institution, one with a perineal fistula and one with a rectovestibular fistula. Our report, along with a literature review, shows that the syndrome is usually associated with “low” anomalies, with a potential for a good prognosis. Management of the anorectal anomaly in patients with KS is not essentially different from that in other nonsyndromic patients, taking into account the frequent association of the syndrome with serious congenital heart disease, which might affect the decision-making and timing of the stages of anorectal reconstruction. The frequent occurrence of learning and feeding difficulties makes establishment of toilet training and bowel management rather more challenging, requiring the expertise of a multidisciplinary team. The finding of ARMs in female patients with other characteristics of KS, although inconstant, could support the clinical suspicion for the syndrome until genetic confirmation is available, and should alert the physician for the potential of severe cardiac defects.

Keywords

- ▶ Kabuki syndrome
- ▶ anorectal malformations
- ▶ chromosomal abnormalities

Introduction

Kabuki syndrome (KS) is a rare genetic condition characterized by clinical variability and genetic heterogeneity, whose molecular basis was only recently identified.¹ Niikawa and Kuroki in 1981 first described the constellation of a distinctive facies, intellectual disability, growth delay, and a variety of skeletal, visceral, and other malformations.² The syndrome derives its name from the resemblance of the patient's facies to the makeup of the actors of the Japanese dance-drama theater Kabuki. Although literature review as well as our own experience, shows that the syndrome is most frequently related with anorectal anomalies of favorable prognosis, the association with major congenital heart defects and other anomalies, makes short- and long-term management challenging. We

present two cases of patients with KS treated in our institution with emphasis on the anorectal aspects of the syndrome.

Case 1

The patient was referred to us at the 22nd day of life and was noted to have a perineal fistula and groove and an anal index of less than 0.35. She was born prematurely at 32 weeks of gestation and was also diagnosed with pulmonary artery stenosis and multiple other congenital anomalies (▶ **Table 1**). KS was clinically suspected initially by the characteristic facies and the constellation of other anomalies and eventually confirmed, when the tests became available by demonstrating the introduction of a premature stop mutation in the *MLL2* gene (*MLL2/KMT2D* MIM# 602113). Her stenotic anus

received
November 10, 2014
accepted after revision
February 2, 2015
published online
May 28, 2015

DOI <http://dx.doi.org/10.1055/s-0035-1547529>.
ISSN 2194-7619.

© 2015 Georg Thieme Verlag KG
Stuttgart · New York

License terms



Table 1 Constellation of anomalies in the reported patients

	Patient 1	Patient 2
Face	Kabuki facies (ear malformations, epicanthus) Diversion squint in right eye	Facies, natal teeth
Musculoskeletal	Congenital hip dysplasia: Surgery Bilateral polydactyly of feet	Congenital hip dysplasia
Cranium	Trigonocephaly: Surgery	Plagiocephaly
Cognitive	Mild learning difficulties	Learning difficulties
Anorectal	Anterior anus and perineal groove	Anovestibular fistula decompressing through fistula though—dilataions—washouts
Surgery	Sphincter mapping—dilataions—table-contrast enema: No megarectum	PSARP + colostomy and colostomy closure (8 wks)
Urogenital		Bilateral duplex kidney—vesicoureteral reflux
Cardiac	Pulmonary stenosis and patent ductus arteriosus Surgery at the age of 1 y	Fallot tetralogy—double outlet right ventricle—surgery
Other GI		Feeding difficulties Gastroesophageal reflux
Spinal		US: Normal
Others	Prematurity 32/40 w	Twin, premature, 35/40 w Hypotonia
Management	Dilataions Laxatives	Preoperative and postoperative dilataions Laxatives

was initially treated with dilataions (starting from a Hegar no. 9) that allowed for bowel decompression and was started on stool softeners (lactulose) from the 1st month of life, while being followed regularly in the anorectal clinic. Spinal ultrasound early in neonatal life revealed no structural abnormalities of the caudal spine. After dilataions reached a caliber of a Hegar no. 12 at the age of 1 year, they were discontinued. At the age of 1.5 years and after undergoing cardiac surgery, we performed anal sphincter mapping under anesthetic which showed that although the position of the fistula-anus, based on the anal index was rather anterior, approximately three quadrants of the fistula were passing through the external sphincter. An on-table contrast enema showed absence of megarectum, and a decision was made to manage the anorectal anomaly conservatively, given the lack of retentive symptoms. She has been regularly followed up, and at the last clinic visit (4-years-old) she was doing very well, being entirely toilet trained and requiring regular use of small doses of oral and rectal laxatives.

Case 2

The patient was first seen by us at the age of 4 months due to an imperforate anus with a rectovestibular fistula. She was born as one of the twins at 35 weeks of gestation and was soon diagnosed with a large atrioventricular septal defect and patent ductus arteriosus which caused intractable heart failure for which she eventually required cardiac surgery. She was also found to have other multiple congenital anomalies (→Table 1). Spinal ultrasound early in neonatal life revealed no structural abnormalities of the caudal spine. As

per the institutional practice from the referring center, she was initially managed with rectal washouts through the fistula, in view of a primary definite repair after stabilization of the cardiac condition. Following the cardiac surgery, she underwent a posterior sagittal anorectoplasty with covering sigmoidostomy at the age of 9 months. Following the genetic review, the constellation of her anomalies was attributed to KS, mainly based on the characteristic facies associated with the syndrome and was later confirmed after genetic testing demonstrating anomalies in the *MLL2* gene. After dilating her anus to a Hegar no. 13, the colostomy was closed and the patient was followed up in the anorectal clinic for bowel management. At the time of the last follow-up (3.5 years of age), she had a good functional result with frequent bowel motions and occasional only use of laxatives, having started toilet training.

Discussion

The two cases highlighted here, as well as a literature review²⁻⁷ (→Table 2) show that when KS is associated with anorectal malformations (ARMs), the affected patients are females and the ARM type is either a rectovestibular or a rectoperineal fistula, both considered to have a good functional prognosis. As some of the reviewed articles were published before the Krickbeck consensus, the terms used to describe the anomalies vary, and some of the older cases were described as anterior stenotic anus.

KS is a rare disease, with estimates of prevalence ranging from 1 to 32,000 for Japanese children and 1 to 86,000 for Australian and New Zealand children.¹ In 2010, mutations in

Table 2 Reported cases with Kabuki syndrome in the literature

S. no.	Author (Y)	Type of ARM	Surgery	Other anomalies
1	Matsumura et al (1992) ³	Vestibular fistula	Anal transposition ^a	Cleft palate, ASD, webbed neck, renal anomalies, long great toes
2	Matsumura et al (1992) ³	Vestibular fistula	Anal transposition ^a	Cleft palate, VSD, aortic aneurysm, horse-shoe kidney, bifid renal pelvis, pigmented retina, abnormal EEG
3	Matsumura et al (1992) ³	Vestibular fistula	“Cut-back” procedure ^b	Autism, retardation in bone age
4	Matsumura et al (1992) ³	Vestibular fistula	Anal transposition ^b	Renal agenesis, hip dysplasia, retardation in bone age, VSD
5	Philip et al (1992) ⁴	Vestibular fistula	PSARP	N/A
6	Kokitsu-Nakata et al (1999) ⁵	Vestibular fistula	Bowel management	Facies, growth retardation, cleft palate, ectopic right kidney, finger anomalies
7	Kawame et al (1999) ⁶	Perineal fistula	PSARP	N/A
8	Abdel-Salam et al (2008) ⁷	Vestibular fistula	PSARP	Diaphragmatic defect, congenital heart defects, cleft palate, lower lip pits, hypopigmentation, seizures, hyperlaxity of joints and premature breast development
9	Present study	Perineal fistula	Bowel management	Facies, hip dysplasia, trigonocephaly, finger anomalies, pulmonary artery stenosis, learning difficulties
10	Present study	Vestibular fistula	PSARP	Facies, hip dysplasia, plagiocephaly, Fallot tetralogy, duplex kidneys, hypotonia, learning difficulties

Abbreviation: PSARP, posterior sagittal anorectoplasty.

Note: All patients studied are females.

^aProcedure listed as described by reporting authors, although terms do not reflect current nomenclature. Most likely term: anterior sagittal approach.

^bProcedure listed as described by reporting authors, although terms do not reflect current nomenclature. Most likely term: perineal operation—anooplasty.

the gene *KMT2D* (*MLL2*) were identified as the major genetic cause for the disease in 56 to 75% of the patients affected.⁸ Other mutations have also been identified in the gene *KDM6A* (*UTX*),⁹ but almost 30% of the patients with the syndrome have no recognizable genetic anomaly identified.¹ The mainstay of diagnosis is the typical facial gestalt with long palpebral fissures, eversion of the lower eyelid, arched eyebrows, short columella, thin upper lip, slanted downward mouth corners, and sometimes prognathism. Other maxillofacial anomalies include cleft lip and palate (12–50%), dentition anomalies (53–100%), and craniosynostosis. Failure to thrive due to feeding difficulties has been documented in 65 to 74% of the patients, with short stature in approximately 58%. Learning difficulties are rather the rule than the exception, with more than 90% of the patients having some form of impairment. Other anomalies include a variety of skeletal malformations (more than 80%, vertebral anomalies or scoliosis, brachydactyly, or clinodactyly), ocular anomalies (38–61%), endocrine disorders (early breast development the most common), cardiac defects (40–50%, most common atrial septal defect—ventricular septal defect), urogenital anomalies in 30 to 40% (hydronephrosis and ectopia), hearing loss, and susceptibility to middle ear infections. Characteristic features are also the presence of persistent fetal finger pads (up to 92%) and abnormalities of the dermatoglyphic pattern. The syndrome has also been associated with a variety of autoimmune

disorders and malignancy (medulloblastoma, non-Hodgkin lymphoma), and a susceptibility to respiratory infections is well documented.^{1,2,6} The association with ARMs is estimated to be around 7%.³ The management of the anorectal anomaly in patients with KS is not essentially different from that in other nonsyndromic patients, but one must take into account the frequent association of the syndrome with serious congenital heart disease, which might affect the decision-making and timing of the stages in anorectal reconstruction. A major cardiac defect might warrant surgical treatment before definite anorectal reconstruction, and in these instances, dilatations of the perineal fistula or a decompressing colostomy might prove useful. The frequent occurrence of learning and feeding difficulties make the establishment of toilet training and bowel management rather more challenging. The importance of a multidisciplinary team in the management of these complex patients cannot be overemphasized. The fact though that these anomalies are usually in the mild end of the ARM spectrum offer a good anorectal functional prognosis, as was the case for our two patients.

The association of specific syndromes with specific types of ARMs is well documented, as in the case of trisomy 21 linked with anomalies without fistula,¹⁰ or Currarino syndrome presenting with a perineal fistula or funnel anus,^{10,11} while other syndromes or chromosomal anomalies do not

follow a set pattern. Due to the rarity of the disease and the rather infrequent association with ARMs, it is uncertain whether KS follows such a pattern, although all reported cases represent ARM at the mild end of the spectrum, which could be associated with good prognosis, provided there are no other risk factors of a poor outcome (dysraphic anomalies, severe developmental delay). The etiopathogenesis of this association, and of the occurrence in females in the reported cases, is yet unclear due to the limited knowledge regarding the gene transcription regulated by the *KMT2D* complex during embryogenesis.

In conclusion, our data, along with data from the literature, show that the finding of ARMs in female patients with other characteristics of KS, although inconstant, could support the clinical suspicion for the syndrome until genetic confirmation is available, although pediatric surgeons should be aware of the high probability of serious cardiac defects.

Conflict of Interest

None.

References

- 1 Bögershausen N, Wollnik B. Unmasking Kabuki syndrome. *Clin Genet* 2013;83(3):201–211
- 2 Niikawa N, Kuroki Y, Kajii T, et al. Kabuki make-up (Niikawa-Kuroki) syndrome: a study of 62 patients. *Am J Med Genet* 1988; 31(3):565–589
- 3 Matsumura M, Yamada R, Kitani Y, et al. Anorectal anomalies associated with Kabuki make-up syndrome. *J Pediatr Surg* 1992; 27(12):1600–1602
- 4 Philip N, Meinecke P, David A, et al. Kabuki make-up (Niikawa-Kuroki) syndrome: a study of 16 non-Japanese cases. *Clin Dysmorphol* 1992;1(2):63–77
- 5 Kokitsu-Nakata NM, Vendramini S, Guion-Almeida ML. Lower lip pits and anorectal anomalies in Kabuki syndrome. *Am J Genet* 1993;47:423–425
- 6 Kawame H, Hannibal MC, Hudgins L, Pagon RA. Phenotypic spectrum and management issues in Kabuki syndrome. *J Pediatr* 1999;134(4):480–485
- 7 Abdel-Salam GM, Afifi HH, Eid MM, el-Badry TH, Kholoussi NM. Anorectal anomalies, diaphragmatic defect, cleft palate, lower lip pits, hypopigmentation and hypogammaglobulinemia A in Kabuki syndrome: a rare combination. *Genet Couns* 2008;19(3):309–317
- 8 Ng SB, Bigham AW, Buckingham KJ, et al. Exome sequencing identifies *MLL2* mutations as a cause of Kabuki syndrome. *Nat Genet* 2010;42(9):790–793
- 9 Lederer D, Grisart B, Digilio MC, et al. Deletion of *KDM6A*, a histone demethylase interacting with *MLL2*, in three patients with Kabuki syndrome. *Am J Hum Genet* 2012;90(1):119–124
- 10 Martucciello G. Genetics of anorectal malformations. In: *Anorectal Malformations in Children*. Holschneider AM, Hutson JM, eds. Berlin, Heidelberg: Springer Verlag; 2006:17–30
- 11 Suomalainen A, Wester T, Koivusalo A, Rintala RJ, Pakarinen MP. Congenital funnel anus in children: associated anomalies, surgical management and outcome. *Pediatr Surg Int* 2007;23(12):1167–1170