

Determination of Gestational Age by Ultrasound

This clinical practice guideline has been prepared by the Diagnostic Imaging Committee, reviewed by the Family Physician Advisory Committee, and approved by the Executive and Council of the Society of Obstetricians and Gynaecologists of Canada.

PRINCIPAL AUTHORS

Kimberly Butt, MD, Fredericton NB

Ken Lim, MD, Vancouver BC

DIAGNOSTIC IMAGING COMMITTEE

Ken Lim, MD (Chair), Vancouver BC

Stephen Bly, MD, Ottawa ON

Kimberly Butt, MD, Fredericton NB

Yvonne Cargill, MD, Ottawa ON

Greg Davies, MD, Kingston ON

Nanette Denis, RDMS, Saskatoon SK

Gail Hazlitt, RN, RDMS, Winnipeg MB

Lucie Morin, MD, Montreal QC

Annie Ouellet, MD, Sherbrooke QC

Shia Salem, MD, Toronto, ON

Disclosure statements have been received from all contributors.

Evidence: Published literature was retrieved through searches of PubMed or MEDLINE and The Cochrane Library in 2013 using appropriate controlled vocabulary and key words (gestational age, ultrasound biometry, ultrasound dating). Results were restricted to systematic reviews, randomized control trials/controlled clinical trials, and observational studies written in English. There were no date restrictions. Searches were updated on a regular basis and incorporated in the guideline to July 31, 2013. Grey (unpublished) literature was identified through searching the websites of health technology assessment and health technology-related agencies, clinical practice guideline collections, clinical trial registries, and national and international medical specialty societies.

Values: The quality of evidence in this document was rated using the criteria described in the Report of the Canadian Task Force on Preventive Health Care (Table 1).

Benefits, harms, and costs: Accurate assignment of gestational age may reduce post-dates labour induction and may improve obstetric care through allowing the optimal timing of necessary interventions and the avoidance of unnecessary ones. More accurate dating allows for optimal performance of prenatal screening tests for aneuploidy. A national algorithm for the assignment of gestational age may reduce practice variations across Canada for clinicians and researchers. Potential harms include the possible reassignment of dates when significant fetal pathology (such as fetal growth restriction or macrosomia) result in a discrepancy between ultrasound biometric and clinical gestational age. Such reassignment may lead to the omission of appropriate—or the performance of inappropriate—fetal interventions.

Abstract

Objective: To assist clinicians in assigning gestational age based on ultrasound biometry.

Outcomes: To determine whether ultrasound dating provides more accurate gestational age assessment than menstrual dating with or without the use of ultrasound. To provide maternity health care providers and researchers with evidence-based guidelines for the assignment of gestational age. To determine which ultrasound biometric parameters are superior when gestational age is uncertain. To determine whether ultrasound gestational age assessment is cost effective.

Keywords: ultrasound, gestational age, dating

Summary Statements

1. When performed with quality and precision, ultrasound alone is more accurate than a “certain” menstrual date for determining gestational age in the first and second trimesters (≤ 23 weeks) in spontaneous conceptions, and it is the best method for estimating the delivery date. (II)
2. In the absence of better assessment of gestational age, routine ultrasound in the first or second trimester reduces inductions for post-term pregnancies. (I)

J Obstet Gynaecol Can 2014;36(2):171–181

This document reflects emerging clinical and scientific advances on the date issued and is subject to change. The information should not be construed as dictating an exclusive course of treatment or procedure to be followed. Local institutions can dictate amendments to these opinions. They should be well documented if modified at the local level. None of these contents may be reproduced in any form without prior written permission of the SOGC.

Table 1. Key to evidence statements and grading of recommendations, using the ranking of the Canadian Task Force on Preventive Health Care

Quality of evidence assessment*	Classification of recommendations†
I: Evidence obtained from at least one properly randomized controlled trial	A. There is good evidence to recommend the clinical preventive action
II-1: Evidence from well-designed controlled trials without randomization	B. There is fair evidence to recommend the clinical preventive action
II-2: Evidence from well-designed cohort (prospective or retrospective) or case-control studies, preferably from more than one centre or research group	C. The existing evidence is conflicting and does not allow to make a recommendation for or against use of the clinical preventive action; however, other factors may influence decision-making
II-3: Evidence obtained from comparisons between times or places with or without the intervention. Dramatic results in uncontrolled experiments (such as the results of treatment with penicillin in the 1940s) could also be included in this category	D. There is fair evidence to recommend against the clinical preventive action
III: Opinions of respected authorities, based on clinical experience, descriptive studies, or reports of expert committees	E. There is good evidence to recommend against the clinical preventive action
	L. There is insufficient evidence (in quantity or quality) to make a recommendation; however, other factors may influence decision-making

*The quality of evidence reported in these guidelines has been adapted from The Evaluation of Evidence criteria described in the Canadian Task Force on Preventive Health Care.¹¹⁸

†Recommendations included in these guidelines have been adapted from the Classification of Recommendations criteria described in the Canadian Task Force on Preventive Health Care.¹¹⁸

- Ideally, every pregnant woman should be offered a first-trimester dating ultrasound; however, if the availability of obstetrical ultrasound is limited, it is reasonable to use a second-trimester scan to assess gestational age. (I)
- Notwithstanding Summary Statements 1, 2, and 3, women vary greatly in their awareness of their internal functions, including ovulation, and this self-knowledge can sometimes be very accurate. (III)

Recommendations

- First-trimester crown-rump length is the best parameter for determining gestational age and should be used whenever appropriate. (I-A)
- If there is more than one first-trimester scan with a mean sac diameter or crown-rump length measurement, the earliest ultrasound with a crown-rump length equivalent to at least 7 weeks (or 10 mm) should be used to determine the gestational age. (III-B)
- Between the 12th and 14th weeks, crown-rump length and biparietal diameter are similar in accuracy. It is recommended that crown-rump length be used up to 84 mm, and the biparietal diameter be used for measurements > 84 mm. (II-1A)
- Although transvaginal ultrasound may better visualize early embryonic structures than a transabdominal approach, it is not more accurate in determining gestational age. Crown-rump length measurement from either transabdominal or transvaginal ultrasound may be used to determine gestational age. (II-1C)
- If a second- or third-trimester scan is used to determine gestational age, a combination of multiple biometric parameters (biparietal diameter, head circumference, abdominal circumference, and femur length) should be used to determine gestational age, rather than a single parameter. (II-1A)
- When the assignment of gestational age is based on a third-trimester ultrasound, it is difficult to confirm an accurate due date. Follow-up of interval growth is suggested 2 to 3 weeks following the ultrasound. (III-C)

INTRODUCTION

The accurate dating of pregnancy is critically important for pregnancy management from the first trimester to delivery, and is particularly necessary for determining viability in premature labour and in post-dates deliveries.¹ Prior to the widespread use of ultrasound, caregivers relied on a combination of history and physical examination to clinically determine gestational age. Ultrasound gave clinicians a method to measure the fetus and therefore to estimate gestational age. Much of our current clinical practice is based on studies from the 1980s and 1990s. As new information emerges in fields, such as reproductive biology, perinatal epidemiology, and medical imaging, our current clinical practice is being challenged. “Certain” menstrual dating, for example, is less certain than previously thought.

When ultrasound is performed with quality and precision, there is evidence to suggest that dating a pregnancy using ultrasound measurements is clinically superior to using menstrual dating with or without ultrasound, and this has been advocated and adopted in other jurisdictions.²⁻⁶

GESTATIONAL AGE ESTIMATES USING CLINICAL DATING

The clinical estimate of gestational age typically relies on clinical history (menstrual cycle length, regularity, and recall of the first day of the last menstrual period), followed by confirmation by physical examination or other signs and symptoms.⁷⁻⁹

Dating Based on Menstrual History

Dating by certain menstrual history is inexpensive and readily available. Typically, the EDD is based on a 280-day gestation from the first day of the LMP. Certain menstrual dating criteria assume regular cycles, ovulation at the midpoint of the cycle, fertilization on the middle day of the cycle, correct recall of the onset of the LMP, and the woman having been free of oral contraceptives for several months prior. Women vary greatly in their awareness of their internal functions, including ovulation. Their self-knowledge of ovulation can sometimes be very accurate; however, the only truly certain clinical history is one in which the dates of ovulation, fertilization, and implantation are precisely known, as in ART, in which records include the date of oocyte retrieval, and other methods of timed ovulation and fertilization. Unfortunately, without timed ovulation and fertilization as in ART and other timed methods, clinical history is often not reliable.¹⁰ Campbell et al. demonstrated that 45% of pregnant women are uncertain of menstrual dates as a result of poor recall, irregular cycles, bleeding in early pregnancy, or oral contraceptive use within 2 months of conception.¹¹

Even if menstrual history is correct, the exact time of ovulation, fertilization, and implantation cannot be known. Women may undergo several “waves” of follicular development during a normal menstrual cycle, which may mean ovulatory inconsistency during any given cycle.^{12,13} Sperm may survive for 5 to 7 days in the female reproductive tract, a “known” conception date is therefore not completely reliable.¹⁴ Recent studies suggest the ovulation-to-implantation duration can vary by as much as

11 days, and this may affect fetal size and growth.¹⁵ Even in women who are certain of menstrual dating, delayed ovulation is an important cause of perceived prolonged pregnancy and is more likely to occur than early ovulation.¹⁶ Some authors have suggested that 282 days should be used instead of 280 to improve dating accuracy, since it is more likely that women will ovulate later rather than earlier than predicted.⁹ All of these factors conspire to make it difficult to accurately predict gestational age based on menstrual history.

Dating Based on Clinical Examination

The size of the uterus, estimated through pelvic or abdominal examination, can be roughly correlated with gestational age; however, factors that affect uterine size (such as fibroids) and maternal body characteristics (such as obesity) will affect such an estimate. The uterus is approximately the size of a grapefruit at 10 to 12 weeks. At 20 weeks the fundus reaches the umbilicus. After 20 weeks the symphysis fundal height, in centimetres, should correlate with the week of gestation.^{7,17} Fetal heart tones are audible at 11 to 12 weeks with electronic Doppler devices, and this audibility can also assist in the clinical assignment of gestational age.¹⁰

Gestational Age Estimation Based on Ultrasound Findings

Ultrasound biometric measurements determine gestational age based on the assumption that the size of the embryo or fetus is consistent with its age. Biological variation in size is less during the first trimester than in the third trimester. Ultrasound estimation of gestational age in the first trimester is therefore more accurate than later in pregnancy.¹⁸ Full descriptions of each parameter and published ranges of accuracy are outlined in Table 2.

The determination of gestational age in the first trimester uses the mean gestational sac diameter and/or the crown–rump length. During the first 3 to 5 menstrual weeks an intrauterine pregnancy is first signaled by the presence of a gestational sac.¹⁹ The gestational sac represents the chorionic cavity, and its echogenic rim represents the implanting chorionic villi and associated decidual tissue.²⁰ The smallest gestational sac size that can be clearly distinguished by high frequency transvaginal transducers is 2 to 3 mm, which corresponds to a gestational age of about 32 to 33 days.²¹ The MSD is a commonly used, standardized, way to estimate gestational age during early pregnancy. It is less reliable when the MSD exceeds 14 mm or when the embryo can be identified.²² The growth of the MSD is approximately 1 mm per day.²³ CRL has lower interobserver variability than MSD, and may thus be better for dating a pregnancy.²⁴

ABBREVIATIONS

AC	abdominal circumference
ART	assisted reproductive technology
BPD	biparietal diameter
CRL	crown–rump length
EDD	estimated due date
FL	femur length
HC	head circumference
ISUOG	International Society of Ultrasound in Obstetrics and Gynecology
LMP	last menstrual period
MSD	mean sac diameter
PPV	positive predictive value
TA	transabdominal
TV	transvaginal
US	ultrasound

Yolk Sac

When the yolk sac appears in the gestational sac it provides confirmation of an intrauterine pregnancy and may be initially visible as early as the start of the 5th week or as late as the 6th week. It grows to a maximal size of 6 mm by 10 weeks and gradually migrates to the periphery of the chorionic cavity. At the end of the first trimester it becomes undetectable. Although the presence of the yolk sac is helpful in determining the presence of an intrauterine pregnancy, direct measurement of this structure is not useful in determining gestational age.²⁵

Crown-Rump Length

Direct measurement of the CRL provides the most accurate estimate of gestational age once the embryo is clearly seen. Ideally, either the best CRL or the average of several satisfactory measurements should be used.²⁶ The CRL measurement is reported to be accurate for dating to within 3 to 8 days.^{22,27–39} The MSD should not be used to estimate gestational age once the CRL can be measured.^{22,39,40}

The narrowest confidence interval appears to be between 7 and 60 mm for CRL.^{3,29,30,40} The slope of the embryonic growth curve is small before this time and it can be difficult to clearly identify a very early fetus; thus, it is this committee's expert opinion that reliability and measurability is best when the CRL is at least 10 mm. If more than one scan is performed in the first trimester, the earliest scan with a CRL of at least 10 mm should be used. To avoid performing extra ultrasounds, it is acceptable to time the dating scan to coincide with nuchal translucency screening (if available).

Several studies have evaluated CRL and BPD between 12 and 14 weeks, with conflicting results. The majority suggest there is no clinically important difference among confidence intervals, suggesting that either CRL or BPD should be used at this gestational age.^{41–44} The 84 mm threshold for CRL for estimating gestational age, as suggested by the ISUOG, seems reasonable.⁶

Recommendations

1. First-trimester crown-rump length is the best parameter for determining gestational age and should be used whenever appropriate. (I-A)
2. If there is more than one first-trimester scan with a mean sac diameter or crown-rump length measurement, the earliest ultrasound with a crown-rump length equivalent to at least 7 weeks (or 10 mm) should be used to determine the gestational age. (III-B)

3. Between the 12th and 14th weeks, crown-rump length and biparietal diameter are similar in accuracy. It is recommended that crown-rump length be used up to 84 mm, and the biparietal diameter be used for measurements > 84 mm. (II-1A)

Transabdominal Versus Transvaginal Ultrasonography

Transvaginal ultrasound is typically used to evaluate early first trimester pregnancy structures, such as the gestational sac, yolk sac, and embryo. Original studies comparing transabdominal and transvaginal techniques in early pregnancy demonstrated that TV was the superior method.^{45–47} Perhaps because of better technology, more recent studies have not found the same result. Grisolia et al. concluded that the accuracy of TV ultrasound has not been shown to be superior to TA ultrasound in dating pregnancies.³¹ Other authors have found that TA is comparable to TV in determining gestational age if CRL is measurable after 6 weeks.^{31,48–50}

Recommendation

4. Although transvaginal ultrasound may better visualize early embryonic structures than a transabdominal approach, it is not more accurate in determining gestational age. Crown-rump length measurement from either transabdominal or transvaginal ultrasound may be used to determine gestational age. (II-1C)

Crown-Rump Length in Pregnancies Conceived by Assisted Reproductive Technology

When the conception date is known absolutely, as with timed ovulation/fertilization during ART, the EDD should be calculated based on the fertilization date. Studies have demonstrated that the CRL measurements in IVF pregnancies are those to be expected in naturally conceived pregnancies, suggesting that study results can be extrapolated between the 2 populations.^{30,51,52}

Second and Third Trimester

In the second and third trimesters, estimation of gestational age is accomplished by measuring the biparietal diameter, head circumference, abdominal circumference, and femur length. These measurements are only as good as the quality of the images. Optimal imaging can be difficult in some clinical situations, such as in a late pregnancy abnormal lie when the head is deep in the maternal pelvis, maternal obesity, or multiple gestation. Normal biological variation appears to have more influence on measurements in the second and third trimester. Thus, in the second half of pregnancy these measurements are less reliable than first trimester CRL, and they become increasingly inaccurate as gestation progresses

Table 2. Common definitions of ultrasound biometry parameters and estimates of accuracy for predicting gestational age

Parameter	Description	Notes	Approximate accuracy of dates	References
Mean sac diameter	The mean of 3 orthogonal sac "inner to inner" diameter measurements (mm). Cursors should be placed on the gestational sac and not the surrounding echogenic region.	Should not be averaged with the CRL. Should not be used once CRL can be measured. GA = 30 days plus MSD measured in mm.	4 to 11 days	Grisolia (2003) ³¹ Daya (1993) ³⁰
Crown-rump length	The crown-rump length is the longest straight line length of the embryo from the outer margin of the cephalic end to the rump. The neck position should be neutral.	The best CRL or the average of several satisfactory measurements should be used.	3 to 8 days	Grisolia (2003) ³¹ Daya (1993) ³⁰ Sladevickus (2004) ⁴²
Biparietal diameter	Axial plane through a symmetrical calvarium that includes the third ventricle, thalami, falx cerebri, and cavum septi pellucidum anteriorly and the tentorial hiatus posteriorly. The calipers should be placed at the maximal diameter from the outer edge of the proximal skull wall to the inner edge of the distal skull.		1st T: 3 to 8 days 2nd T: 7 to 12 days	Grisolia (2003) ³¹ Daya (1993) ³⁰ Sladevickus (2004) ⁴² Bovicelli (1981) ²⁸ Hadlock (1984) ⁷² Hadlock (1987) ⁷³ Hadlock (1991) ⁵⁴ Chervenak (1998) ⁶¹
Head circumference	The head circumference is obtained in the identical plane to the BPD. The trace/ellipse should follow the outer perimeter of the bony skull, not the overlying skin, as that will falsely increase the head circumference	The cerebellum is not included in this image.	2nd T: 7 to 12 days	Hadlock (1984) ⁷² Hadlock (1987) ⁷³ Hadlock (1991) ⁵⁴ Chervenak (1998) ⁶¹
Abdominal circumference	True axial plane at the level of the bifurcation of the portal vein (into right and left branches) and the stomach. The measurement should be as tight to skin as possible.		2nd T: 7 to 15 days 3rd T: 18 to 35 days	Hadlock (1984) ⁷² Hadlock (1987) ⁷³ Hadlock (1991) ⁵⁴ Chervenak (1998) ⁶¹
Femur length	Both the femoral head or greater trochanter and the femoral condyle are simultaneously visualized. The cursor should be placed at the junction of bone and cartilage and only the bone measured	Ideally, the ultrasound transducer should be aligned perpendicular to the long axis of the femur. Varies with ethnicity.	2nd T: 7 to 17 days 3rd T: 21 days	Hadlock (1984) ⁷² Hadlock (1987) ⁷³ Hadlock (1991) ⁵⁴ Chervenak (1998) ⁶¹
T: trimester				

(Table 2).^{18,53} Maternal and fetal pathology may affect them, so their inclusion or exclusion in the determination of gestational age requires clinical judgement.

The BPD is less reliable in determining gestational age when there are variations in skull shape, such as dolichocephaly or brachycephaly; hence some authors feel that BPD is less reliable than HC.^{9,53–57} As a single parameter, HC correlates better to gestational age than the other 3 standard parameters in the second trimester, and as with all others, it becomes less accurate with increasing gestational age.^{58–61}

It is more challenging to measure the fetal AC than the other parameters. The abdomen has no bright echoes of bone, it is not always symmetrical, and its size will vary with fetal respiration and central body flexion/extension. Of all the fetal biometric parameters, this measurement has the most variability as it is somewhat dependant on fetal growth factors and body position.^{54,61–63}

Femur length varies somewhat with ethnicity. Short femurs are commonly a normal variant, however this finding may also indicate fetal growth restriction, aneuploidy, and—when severely shortened—skeletal dysplasias.^{53,64–69}

Composite Versus Single Biometry Measurement

Using multiple parameters is superior to using a single second trimester parameter.^{63,70,71} As more parameters are used, accuracy improves; however, there is no significant benefit beyond 3 commonly used parameters.^{3,61,70–73}

Multiple parameters are also useful if any one parameter is affected by a fetal condition/syndrome, such as achondroplasia on femur length. It is prudent to evaluate the etiology of an aberrant measurement to determine its clinical significance.

Commonly, clinicians use the unweighted mean of all 4 biometric parameters (BPD, HC, AC, and FL). However, it is clear that all 4 are not equally correlative, thus many authors have created regression equations using various combinations of biometric parameters to improve accuracy.^{54,61,74,75} It is not clear which method is superior in determining gestational age.^{61,72} Until more research is available, it is reasonable to use either when estimating second or third trimester gestational age. Up to 23 weeks, second trimester US has a 95% CI of less than a week for predicting gestational age, comparable to first trimester US.⁶¹ In the late second trimester, clinical judgement should be exercised.

Since the confidence intervals in the third trimester are quite large (2 to 2.4 weeks), it is not clear that US determined gestational age is superior to clinical history and the application of judicious clinical judgement may be

warranted. There is also a concern that when gestational age assessment is based on a third trimester scan only, the fetus may in fact be growth restricted,⁶¹ and gestational age therefore underestimated. Hence, a follow-up scan for growth in such circumstances should be considered to evaluate interval growth.

Recommendations

5. If a second- or third-trimester scan is used to determine gestational age, a combination of multiple biometric parameters (biparietal diameter, head circumference, abdominal circumference, and femur length) should be used to determine gestational age, rather than a single parameter. (II-1A)
6. When the assignment of gestational age is based on a third-trimester ultrasound, it is difficult to confirm an accurate due date. Follow-up of interval growth is suggested 2 to 3 weeks following the ultrasound. (III-C)

Other Biometry

Measurement of the transcerebellar diameter, foot length, clavicle length, intra/interorbital diameters, kidney length, sacral length, scapula length, as well as the length of other long bones of the extremity have also been evaluated to determine gestational age. Studies have not shown that these parameters improve the assessment of gestational age beyond that achieved with standard biometry, however they may be useful in clinical situations in which traditional biometry is difficult to attain (such as uteroplacental insufficiency) or when fetal abnormalities are present.^{76–86}

Signs of Fetal Maturity

Identification of certain US findings suggest that a fetus has reached the third trimester and may correlate with fetal lung maturity and gestational age. These parameters are the epiphyseal ossification centres of the distal femur, proximal tibia, and proximal humerus. The measurement of these ossification centres does not precisely correlate with gestational age; however, their presence may be helpful late in pregnancy when the gestational age is not known. The presence of distal femoral epiphysis has a PPV of 96% for indicating a pregnancy of at least 32 weeks, the proximal tibial epiphysis has a PPV of 83% for indicating a pregnancy of at least 37 weeks, and the proximal humeral epiphysis has a PPV of 100% for indicating a pregnancy of at least 38 weeks.^{87–91}

WHAT IS THE BEST METHOD FOR ASSIGNING GESTATIONAL AGE?

Currently, most centres in Canada use a combined approach in which US is used to confirm reliable

menstrual dating. If there is an unreliable menstrual history, the US prediction of EDD is used. Clinical judgement is used to resolve conflicting data. However, centres vary in terms of confidence intervals and biometry charts used.

Many studies evaluating menstrual dating, compared with US dating, in the first and second trimesters have found US dating superior for predicting the actual date of delivery.^{92–98} No study has shown that it is inferior to menstrual dating. Menstrual dating underestimates the US-based EDD by an average of 2 to 3 days.⁹⁵ Ultrasound dating alone was significantly better in predicting the actual date of delivery than any of the dating policies using menstrual dates alone or in combination with US.^{9,99}

Many studies document that the use of US dates reduces the rate of post-dates pregnancy by about 70% even in the face of certain menstrual history.^{92,96,97,99,100} The most recent Cochrane systematic review found reduced rates of induction for post-term pregnancy (OR 0.59; 95% CI 0.42 to 0.83) among women who underwent routine US in early pregnancy (< 24 weeks) and concluded that early pregnancy US enables better gestational age assessment, as well as conferring other benefits.¹⁰¹

Using US-based gestational age assignment would also result in improved performance of prenatal screening programs. Using US estimates exclusively would increase sensitivity for Down syndrome anywhere from 9% to 16%, and/or decrease false-positive rates (for a set sensitivity) by 2.6%.^{102,103} There might be a very slight increase in the screening positive rate for open neural tube defects, but this is more than offset by the decrease in false-positive rates for Down syndrome. The common practice of using certain menstrual dates confirmed by US is inferior to using US alone.¹⁰⁴ In the context of serum screening, first and early second-trimester US parameters perform similarly.^{100,102,104–106}

Some clinicians fear that the exclusive use of US-based estimates of gestational age would result in pregnancy complications because of the potential to miss early growth discordance. A large-for-gestational-age fetus might be mistakenly assigned a greater gestational age because of its larger size or an early growth-restricted fetus may be incorrectly assigned a later due date. This may potentially mask an underlying fetal or placental problem leading to pregnancy complications, such as preterm birth, preeclampsia, and fetuses small for gestational age, or it may cause a delay in the recognition of fetal abnormalities. There is disagreement in the literature as to whether a significant discordance

between menstrual and US estimates of gestational age is associated with an increased risk of obstetrical complications.^{107–114} Interestingly, an unreliable menstrual history itself confers an increased risk of adverse pregnancy outcomes.¹¹² Although there may be a risk in using US dating exclusively, some of this risk would remain whenever there is discordance between menstrual and US estimates, regardless of which method of gestational age assignment is used. Furthermore, the clinical management of such situations is unclear. More research is needed in this area.

Summary Statements

1. When performed with quality and precision, ultrasound alone is more accurate than a “certain” menstrual date for determining gestational age in the first and second trimesters (≤ 23 weeks) in spontaneous conceptions, and it is the best method for estimating the delivery date. (II)
2. In the absence of better assessment of gestational age, routine ultrasound in the first or second trimester reduces inductions for post-term pregnancies. (I)

SHOULD ROUTINE FIRST TRIMESTER DATING ULTRASOUNDS BE OFFERED TO ALL PREGNANT WOMEN?

A routine first trimester US has many advantages: early identification of gross anomalies and multiple gestations, more precise dating, improved performance of prenatal screening, and an opportunity to perform nuchal translucency as part of prenatal genetic screening. In many jurisdictions, a dating US is performed routinely in all women, regardless of menstrual history. The availability, quality, and health care cost of obstetrical US is a significant factor in local patterns of practice.² Crowther et al. found routine early US (< 17 weeks) provided more precise estimates of gestational age than later US (18 to 22 weeks), reduced the need to adjust the EDD in mid-gestation, and decreased maternal anxiety.¹¹⁵

The balance of the literature supports using first trimester US to reduce the incidence of induction for post-dates pregnancy.^{18,99,100,116–118} Although no comprehensive cost benefit analysis has been done on routine early US for dating, the current literature suggests significant benefits are present. Ideally, where it is readily available, a first trimester US for dating should be performed. Where nuchal translucency is available, this scan can serve both functions. When a first trimester US cannot be obtained, the available evidence suggests the second trimester US can be used for similar benefits.¹⁰¹

Summary Statements

3. Ideally, every pregnant woman should be offered a first trimester dating ultrasound; however, if the availability of obstetrical ultrasound is limited, it is reasonable to use a second trimester scan to assess gestational age. (I)
4. Notwithstanding Summary Statements 1, 2, and 3, women vary greatly in their awareness of their internal functions, including ovulation, and this self-knowledge can sometimes be very accurate. (III)

SUMMARY

The accurate determination of gestational age is required for many aspects of antenatal care. In the past, it was probably felt that a few days of inaccuracy was acceptable; however, emerging data suggests that a few days inaccuracy can affect things, such as the performance of maternal serum screening, the assessment of post-dates pregnancy, and the subsequent induction of labour. Based on the available research, the use of US-derived dates is the best method to determine gestational age for clinical use. It is not, however, intended to be used to determine the exact date of conception because of biological variability in reproduction, fetal size, and development. Clinical history may have value in determining gestational age, and on rare occasions may supersede US dating; however, in order to achieve the most clinical benefit, the use of US dating should predominate.

REFERENCES

1. Kalish RB, Chervenak FA. Sonographic determination of gestational age. *Ultrasound Rev Obstet Gynecol* 2005;5:254–8.
2. Hughes R, Aitken E, Anderson J, Barry C, Benton M, Elliot J; National Institute for Health and Clinical Excellence. Antenatal care. Routine care for the healthy pregnant woman. NICE clinical guideline 62. London: RCOG Press; 2008.
3. Bottomley C, Bourne T. Dating and growth in the first trimester. *Best Pract Res Clin Obstet Gynaecol* 2009;23:439–52.
4. Gardosi J. Dating of pregnancy: time to forget the last menstrual period. *Ultrasound Obstet Gynecol* 1997;9:367–8.
5. Gardosi J, Geirsson RT. Routine ultrasound is the method of choice for dating pregnancy. *Br J Obstet Gynaecol* 1998;105:933–6.
6. Salomon LJ, Alfirevic Z, Bilardo CM, Chalouhi GE, Ghi T, Kagan KO, et al. ISUOG practice guidelines: performance of first-trimester fetal ultrasound scan. *Ultrasound Obstet Gynecol* 2013;41(1):102–13.
7. Andersen HF, Johnson TR Jr, Flora JD Jr, Barclay ML. Gestational age assessment. II. Prediction from combined clinical observations. *Am J Obstet Gynecol* 1981;140(1):770–4.
8. Andersen HF, Johnson TR Jr, Barclay ML, Flora JD Jr. Gestational age assessment. I. analysis of individual clinical observations. *Am J Obstet Gynecol* 1981;139:173–7.
9. Nguyen TH, Larsen T, Engholm G, Moller H. Evaluation of ultrasound-estimated date of delivery in 17,450 spontaneous singleton births: do we need to modify Naegele's rule? *Ultrasound Obstet Gynecol*. 1999;14:23–8.
10. Johnson TR, Niebyl JR. Preconception and prenatal care: part of the continuum. In: obstetrics: normal and problem pregnancies. 4th ed. Philadelphia: Churchill Livingstone; 2002.
11. Campbell S, Warsof SL, Little D, Cooper DJ. Routine ultrasound screening for the prediction of gestational age. *Obstet Gynecol* 1985;65:613–20.
12. Baerwald AR, Adams GP, Pierson RA. A new model for ovarian follicular development during the human menstrual cycle. *Fertil Steril* 2003;80:116–22.
13. Baerwald AR, Adams GP, Pierson RA. Characterization of ovarian follicular wave dynamics in women. *Biol Reprod* 2003;69:1023–31.
14. Leppaluoto P. Vaginal flora and sperm survival. *J Reprod Med* 1974;12:99–107.
15. Mahendru AA, Daemen A, Everett TR, Wilkinson IB, McEniery CM, Abdallah Y, et al. Impact of ovulation and implantation timing on first trimester crown-rump length and gestational age. *Ultrasound Obstet Gynecol* 2012;40:630–5.
16. Saito M, Keijiro Y, Akinori H, Takahiro K, Nozumu N, Kohei K. Time of ovulation and prolonged pregnancy. *Am J Obstet Gynecol* 1976;112:31–8.
17. Beazley JM, Underhill RA. Fallacy of the fundal height. *BMJ* 1970;4:404–6.
18. Caughey AB, Nicholson JM, Washington AE. First- vs second-trimester ultrasound: the effect on pregnancy dating and perinatal outcomes. *Am J Obstet Gynecol* 2008;198:703–5.
19. Laing FC, Frates MC. Ultrasound evaluation during the first trimester of pregnancy. In: Callen PW, ed. *Ultrasonography in obstetrics and gynecology*, 4th ed. Philadelphia: WB Saunders, 2000.
20. Timor-Tritsh IE, Farine D, Rosen MG. A close look at early embryonic development with the high-frequency transvaginal transducer. *Am J Obstet Gynecol* 1988;159:676–81.
21. Rowling SE, Langer JE, Coleman BG, Nisenbaum HL, Horii SC, Arger PH, et al. Sonography during early pregnancy: dependence of threshold and discriminatory values on transducer frequency. *AJR Am J Roentgenol* 1999;172:983–8.
22. Nyberg DA, Mack LA, Laing FC, Patten R. Distinguishing normal from abnormal gestational sac growth in early pregnancy. *J Ultrasound Med* 1987;6(1):23–7.
23. MacKenzie AP, Stephenson CD, Funai EF. Prenatal assessment of gestational age. UpToDate. Version 16.3, October 1 2008. Available at: <http://www.uptodate.com/contents/prenatal-assessment-of-gestational-age?> Accessed on April 21, 2013.
24. Pexsters A. Clinical implications of intra- and interobserver reproducibility of transvaginal sonographic measurement of gestational sac and crown-rump length at 6–9 weeks' gestation. *Ultrasound Obstet Gynecol* 2011;38:510–5.
25. Sauerbrei E, Cooperbeg PL, Poland JB. Ultrasound demonstration of the normal fetal yolk sac. *J Clin Ultrasound* 1980;8:217–20.
26. Salomon LJ, Bernard JP, Duyme M, Dorion A, Ville Y. Revisiting first-trimester biometry. *Ultrasound Obstet Gynecol* 2003;22:63–6.
27. Tezuka N, Saito H, Hiroi M. Comparison of the accuracy of gestational age estimation from fetal heart rate and crown-rump length. *Prim Care Update Ob Gyns* 1998;5:193.
28. Bovicelli L, Orsini LF, Rizzo N, Calderoni P, Pazzaglia FL, Michelacci L. Estimation of gestational age during the first trimester by real-time measurement of fetal crown-rump length and biparietal diameter. *J Clin Ultrasound* 1981;9:71–5.
29. Hadlock FP, Shah YP, Kanon DJ, Lindsey JV. Fetal crown-rump length: reevaluation of relation to menstrual age (5–18 weeks) with high-resolution real-time US. *Radiology* 1992;182:501–5.

30. Daya S. Accuracy of gestational age estimation by means of fetal crown-rump length measurement. *Am J Obstet Gynecol* 1993;168:903–8.
31. Grisolia G, Milano K, Pilu G, Banzi C, David C, Gabrielli S, et al. Biometry of early pregnancy with transvaginal sonography. *Ultrasound Obstet Gynecol*. 1993;3:403–11.
32. Robinson HP, Fleming JE. A critical evaluation of sonar “crown-rump length” measurements. *Br J Obstet Gynaecol* 1975;82:702–10.
33. van de Velde EH, Broeders GH, Horbach JG, Esser-Rath VW. Estimation of pregnancy duration by means of ultrasonic measurements of the fetal crown-rump length. *Eur J Obstet Gynecol Reprod Biol* 1980;10:225–30.
34. Vollebbergh JH, Jongsma HW, van Dongen PW. The accuracy of ultrasonic measurement of fetal crown-rump length. *Eur J Obstet Gynecol Reprod Biol* 1989;30:253–6.
35. Silva PD, Mahairas G, Schaper AM, Schauburger CW. Early crown rump length. A good predictor of gestational age. *J Reprod Med* 1990;35:641–4.
36. Wisser J, Dirschedl P, Krone S. Estimation of gestational age by transvaginal sonographic measurement of greatest embryonic length in dated human embryos. *Ultrasound Obstet Gynecol* 1994;4:457–62.
37. Goldstein SR, Wolfson R. Endovaginal ultrasonographic measurement of early embryonic size as a means of assessing gestational age. *J Ultrasound Med* 1994;13:27–31.
38. Nyberg DA, Laing FC, Filly RA. Threatened abortion: sonographic distinction of normal and abnormal gestation sac. *Radiology* 1986;158:397–400.
39. Goldstein SR. Early detection of pathologic pregnancy by transvaginal ultrasonography. *J Clin Ultrasound* 1990;18:262–73.
40. Piantelli G, Sacchini C, Coltri A, Ludovici G, Paita Y, Gramellini D. Ultrasound dating-curve analysis in the assessment of gestational age. *Clin Exp Obstet Gynecol* 1994;2:108–18.
41. Saltved S, Almstrom H, Kublickas M. Ultrasound dating at 12-14 or 15-20 weeks of gestation? A prospective cross-validation of established dating formulae in a population of in-vitro fertilized pregnancies randomized to early or late dating scan. *Ultrasound Obstet Gynecol* 2004;24:42–50.
42. Sladkevicius P, Saltved S, Almstrom H, Kublickas M, Grunewald C, Valentin L, et al. Ultrasound dating at 12-14 weeks of gestation. A prospective cross-validation of established dating formulae in in-vitro fertilized pregnancies. *Ultrasound Obstet Gynecol* 2005;26:504–11.
43. Wu FS, Hwu YM, Lee RK, Li SH, Sun FJ, Lin MH, et al. First trimester ultrasound estimation of gestational age in pregnancies conceived after in vitro fertilization. *Eur J Obstet Gynecol Reprod Biol* 2012;160:151–5.
44. Chalouhi GE, Bernard JP, Benoist G, Nasr B, Ville Y, Salomon LJ. A comparison of first trimester measurements for prediction of delivery date. *J Matern Fetal Neonatal Med* 2011;24:51–7.
45. Coleman BG, Arger PH, Grumbach K, Menard MK, Mintz MC, Allen KS, et al. Transvaginal and transabdominal sonography: prospective comparison. *Radiology* 1988;168:639–43.
46. Jain KA, Hamper UM, Sanders RC. Comparison of transvaginal and transabdominal sonography in the detection of early pregnancy and its complications. *AJR Am J Roentgenol* 1988;151:1139–43.
47. Pennell RG, Needleman L, Pajak T, Baltarowich O, Vilaro M, Goldberg BB, et al. Prospective comparison of vaginal and abdominal sonography in normal early pregnancy. *J Ultrasound Med* 1991;10:63–7.
48. Lohr PA, Reeves MF, Creinin MD. A comparison of transabdominal and transvaginal ultrasonography for determination of gestational age and clinical outcomes in women undergoing early medical abortion. *Contraception* 2010;81:240–4.
49. Ferrazzi E, Garbo S, Sulpizio P, Ghisoni L, Levi SP, Buscaglia M. Miscarriage diagnosis and gestational age estimation in the early first trimester of pregnancy: transabdominal versus transvaginal sonography. *Ultrasound Obstet Gynecol* 1993;3:36–41.
50. Kaur A. Transvaginal ultrasonography in first trimester of pregnancy and its comparison with transabdominal ultrasonography. *J Pharm Bioallied Sci* 2011;3:329–38.
51. Gjerris AC, Loft A, Pinborg A, Tabor A, Christiansen M. First-trimester screening in pregnancies conceived by assisted reproductive technology: significance of gestational dating by oocyte retrieval or sonographic measurement of crown-rump length. *Ultrasound Obstet Gynecol* 2008;32:612–7.
52. Tunón K, Eik-Nes SH, Grøttum P, Von Düring V, Kahn JA. Gestational age in pregnancies conceived after in vitro fertilization: a comparison between age assessed from oocyte retrieval, crown-rump length and biparietal diameter. *Ultrasound Obstet Gynecol* 2000;15(1):41–6.
53. Nyberg DA. Diagnostic imaging of fetal anomalies, 2nd ed. Philadelphia: Lippincott Williams Wilkins; 2003: pp 32–4.
54. Hadlock FP, Harrist RB, Martinez-Poyer J. How accurate is second trimester fetal dating? *J Ultrasound Med* 1991;10:557–61.
55. Hadlock FP, Deter RL, Harrist RB, Park SK. Fetal biparietal diameter: rational choice of plane section for sonographic measurement. *AJR Am J Roentgenol* 1982;138:871–4.
56. Hadlock FP, Deter RL, Harrist RB, Park SK. Fetal biparietal diameter: a critical re-evaluation of the relation to menstrual age by means of real-time ultrasound. *J Ultrasound Med* 1982;1:97–104.
57. Hadlock FP, Deter RL, Carpenter RJ, Park SK. Estimating fetal age: effect of head shape on BPD. *AJR Am J Roentgenol* 1981;137:83–5.
58. Law RG, MacRae KD. Head circumference as an index of fetal age. *J Ultrasound Med* 1982;1:281–8.
59. Hadlock FP, Deter RL, Harrist RB, Park SK. Fetal head circumference: relation to menstrual age. *AJR Am J Roentgenol* 1982;138:649–53.
60. Ott WJ. The use of ultrasonic fetal head circumference for predicting expected date of confinement. *J Clin Ultrasound* 1984;12:411–5.
61. Chervenak FA, Skupski DW, Romero R, Myers MK, Smith-Levitin M, Rosenwaks Z, et al. How accurate is fetal biometry in the assessment of fetal age? *Am J Obstet Gynecol* 1998;178:678–87.
62. Hadlock FP, Deter RL, Harrist RB, Park SK. Fetal abdominal circumference as a predictor of menstrual age. *AJR Am J Roentgenol* 1982;139:367–70.
63. Benson C, Doubilet PM. Sonographic prediction of gestational age: accuracy of second and third trimester fetal measurements. *AJR Am J Roentgenol* 1991;157:1275–7.
64. Shalev E, Feldman E, Weiner E, Zuckerman H. Assessment of gestational age by ultrasonic measurement of the femur length. *Acta Obstet Gynecol Scand* 1985;64:71–4.
65. Hadlock FP, Harrist RB, Deter RL, Park SK. A prospective evaluation of femur length as a predictor of gestational age. *J Ultrasound Med* 1983;2:111–2.
66. Yeh MN, Bracero L, Reilly KB, Murtha L, Aboulafia M, Barron BA. Ultrasonic measurement of the femur length as an index of fetal gestational age. *Am J Obstet Gynecol* 1982;144:519–22.
67. Hadlock FP, Harrist RB, Deter RL, Park SK. Fetal femur length as a predictor of menstrual age: sonographically measured. *AJR Am J Roentgenol* 1982;138:875–8.
68. Weisz B, David AL, Chitty L, Peebles D, Pandya P, Patel P, et al. Association of isolated short femur in the mid-trimester fetus with perinatal outcome. *Ultrasound Obstet Gynecol* 2008;31:512–6.
69. Papageorgiou AT, Fratelli N, Leslie K, Bhide A, Thilaganathan B. Outcome of fetuses with antenatally diagnosed short femur. *Ultrasound Obstet Gynecol* 2008;31:507–11.

70. Hill LM, Guzick D, Hixson J, Peterson CS, Rivello DM. Composite assessment of gestational age: a comparison of institutionally derived and published regression equations. *Am J Obstet Gynecol* 1992;166:551–5.
71. Geirsson RT, Have G. Comparison of actual and ultrasound estimated second trimester gestational length in in-vitro fertilized pregnancies. *Acta Obstet Gynecol Scand* 1993;72:344–6.
72. Hadlock FP, Deter RL, Harrist RB, Park SK. Estimating fetal age: computer-assisted analysis of multiple fetal growth parameters. *Radiology* 1984;152:497–501.
73. Hadlock FP, Harrist RB, Shah YP, King DE, Park SK, Sharman RS. Estimating fetal age using multiple parameters: a prospective evaluation in a racially mixed population. *Am J Obstet Gynecol* 1987;156:955–57.
74. Mongelli M, Yuxin NG, Biswas A, Chew S. Accuracy of ultrasound dating formulae in the late second-trimester in pregnancies conceived with in-vitro fertilization. *Acta Radiol* 2003;44:452–5.
75. Mongelli M, Chew S, Yuxin NG, Biswas A. Third-trimester ultrasound dating algorithms derived from pregnancies conceived with artificial reproductive techniques. *Ultrasound Obstet Gynecol* 2005;26:129–31.
76. Gottlieb AG, Galen HL. Nontraditional sonographic pearls in estimating gestational age. *Semin Perinatol* 2008;32:154–60.
77. Jeanty P, Rodesch F, Delbeke D, Dumont JE. Estimation of gestational age from measurements of fetal long bones. *J Ultrasound Med* 1984;3:75–9.
78. Yarkoni S, Schmidt W, Jeanty P, Reece EA, Hobbins JC. Clavicular measurement: a new biometric parameter for fetal evaluation. *J Ultrasound Med* 1985;4:467–70.
79. Mercer BM, Sklar S, Shariatmadar A, Gillieson MS, D'Alton ME. Fetal foot length as a predictor of gestational age. *Am J Obstet Gynecol* 1987;156:350–5.
80. Platt LD, Medearis AL, DeVore GR, Horenstein JM, Carlson DE, Brar HS. Fetal foot length: relationship to menstrual age and fetal measurements in the second trimester. *Obstet Gynecol* 1988;71:526–31.
81. Murao F, Shibukawa T, Takamiya O, Yamamoto K, Hasegawa K. Antenatal measurement of the scapula length using ultrasound. *Gynecol Obstet Invest* 1989;28:195–7.
82. Reece EA, Gabrielli S, Degennaro N, Hobbins JC. Dating through pregnancy: a measure of growing up. *Obstet Gynecol Surv* 1989;44:544–55.
83. Lee W, Barton S, Comstock CH, Bajorek S, Batton D, Kirk JS. Transverse cerebellar diameter: a useful predictor of gestational age for fetuses with asymmetric growth retardation. *Am J Obstet Gynecol* 1991;165(4 Pt 1):1044–50.
84. Sherer DM, Abramowicz JS, Plissinger MA, Woods JR Jr. Fetal sacral length in the ultrasonographic assessment of gestational age. *Am J Obstet Gynecol* 1993;168:626–33.
85. Konje JC, Abrams KR, Bell SC, Taylor DJ. Determination of gestational age after the 24th week of gestation from fetal kidney length measurements. *Ultrasound Obstet Gynecol* 2002;19:592–7.
86. Drey EA, Kang MS, McFarland W, Taylor DJ. Improving the accuracy of fetal foot length to confirm gestation duration. *Obstet Gynecol* 2005;105:773–8.
87. Donne HD, Faúndes A, Tristão EG, de Sousa MH, Urbanetz AA. Sonographic identification and measurement of the epiphyseal ossification centers as markers of fetal gestational age. *J Clin Ultrasound* 2005;33:394–400.
88. Goldstein I, Lockwood CJ, Reece, Hobbins JC. Sonographic assessment of the distal femoral and proximal tibial ossification centers in the prediction of pulmonary maturity in normal women and women with diabetes. *Am J Obstet Gynecol* 1988;159:72–6.
89. Goldstein I, Lockwood C, Belanger K, Hobbins J. Ultrasonographic assessment of gestational age with the distal femoral and proximal tibial ossification centers in the third trimester. *Am J Obstet Gynecol* 1988;158:127–30.
90. Gentili P, Trasimeni A, Giorlando C. Fetal ossification centers as predictors of gestational age in normal and abnormal pregnancies. *J Ultrasound Med* 1984;3:193–7.
91. Mahoney BS, Callen PW, Filly RA. The distal femoral epiphyseal ossification center in the assessment of third-trimester menstrual age: sonographic identification and measurement. *Radiology* 1985;155:201–4.
92. Mongelli M, Wilcox M, Gardosi J. Estimating the date of confinement: ultrasonographic biometry versus certain menstrual dates. *Am J Obstet Gynecol* 1996;174:278–81.
93. Rossavik IK, Fishburne JI. Conception age, menstrual age and ultrasound age: a second trimester comparison of pregnancies of known conception date with pregnancies dates from last menstrual period. *Obstet Gynecol* 1987;73:243–9.
94. Kramer MS, McLean FH, Boyd ME, Usher RH. The validity of gestational age estimation by menstrual dating in term, preterm, and postterm gestations. *JAMA* 1988;260:3306–8.
95. Tunón K, Eik-Nes SH, Grøttum P. A comparison between ultrasound and a reliable last menstrual period as predictors of the day of delivery in 15,000 examinations. *Ultrasound Obstet Gynecol* 1996;8:178–85.
96. Waldenström U, Axelsson O, Nilsson S. A comparison of the ability of a sonographically measured biparietal diameter and last menstrual period to predict the spontaneous onset of labor. *Obstet Gynecol* 1990;76:336–8.
97. Kieler H, Axelsson O, Nilsson S, Waldenström U. Comparison of ultrasonic measurement of biparietal diameter and last menstrual period as a predictor of day of delivery in women with regular 28 day-cycles. *Acta Obstet Gynecol Scand* 1993;72:347–9.
98. Backe B, Nakling J. Term prediction in routine ultrasound practice. *Acta Obstet Gynecol Scand* 1994;73:113–8.
99. Taipale P, Hiilesmaa V. Predicting delivery date by ultrasound and last menstrual period in early gestation. *Obstet Gynecol* 2001;97:189–94.
100. Blondel B, Morin I, Platt RW, Kramer MS, Usher R, Breart G. Algorithms for combining menstrual and ultrasound estimates of gestational age: consequences for rates of preterm and postterm birth. *BJOG* 2002;109:718–20.
101. Whitworth M, Bricker L, Neilson JP, Dowswell T. Ultrasound for fetal assessment in early pregnancy. *Cochrane Database Syst Rev* 2010;4:CD007058.
102. Benn PA, Borgida A, Horne D, Briganti S, Collins R, Rodis JF. Down syndrome and neural tube defect screening: the value of using gestational age by ultrasonography. *Am J Obstet Gynecol* 1997;176:1056–61.
103. Wald NJ, Cuckle HS, Densem JW, Kennard A, Smith D. Maternal serum screening for Down's syndrome: the effect of routine ultrasound scan determination of gestational age and adjustment for maternal weight. *Br J Obstet Gynaecol* 1992;99:144–9.
104. Rahim RR, Cuckle HS, Sehmi IK, Jones RG. Compromise ultrasound dating policy in maternal serum screening for Down syndrome. *Prenat Diagn* 2002;22:1181–4.
105. Koster MP, Van Leeuwen-Spruijt M, Wortelboer EJ, Stoutenbeek P, Elvers LH, Loeber JG, et al. Lack of standardization in determining gestational age for prenatal screening. *Ultrasound Obstet Gynecol* 2008;32:607–11.
106. Wald NJ, Smith D, Kennard A, Palomaki GE, Salonen R, Holzgreve W, et al. Biparietal diameter and crown-rump length in fetuses with Down's syndrome: implications for antenatal serum screening for Down's syndrome. *Br J Obstet Gynaecol* 1993;100:430–5.

107. Gardosi J, Francis A. Early pregnancy predictors of preterm birth: the role of a prolonged menstruation-conception interval. *BJOG* 2000;107:228–37.
108. Kallen K. Increased risk of perinatal/neonatal death in infants who were smaller than expected at ultrasound fetometry in early pregnancy. *Ultrasound Obstet Gynecol* 2004;24:30–4.
109. Nakling J, Backe B. Adverse obstetric outcome in fetuses that are smaller than expected at second trimester routine ultrasound examination. *Acta Obstet Gynecol Scand* 2002;81:846–51.
110. Tunón K, Eik-Nes SH, Grøttum P. Fetal outcome when the ultrasound estimate of the day of delivery is more than 14 days later than the last menstrual period estimate. *Ultrasound Obstet Gynecol* 1999;14:17–22.
111. Dietz PM, England LJ, Callaghan WM, Pearl M, Wier ML, Kharrazi M. A comparison of LMP-based and ultrasound-based estimates of gestational age using linked California livebirth and prenatal screening records. *Paediatr Perinat Epidemiol* 2007;21(Suppl 2):62–71.
112. Nguyen TH, Larsen T, Engholm G, Moller H. Increased adverse pregnancy outcomes with unreliable last menstruation. *Obstet Gynecol*. 2000;95:867–73.
113. Smith GC, Smith MF, McNay MB, Fleming JE. First-trimester growth and the risk of low birth weight. *N Engl J Med* 1998;339:1817–22.
114. Lynch CD, Zhang J. The research implications of the selection of a gestational age estimation method. *Paediatr Perinat Epidemiol* 2007;21(Suppl 2):86–96.
115. Crowther CA, Kornman L, O'Callaghan S, George K, Furness M, Willson K. Is an ultrasound assessment of gestational age at the first antenatal visit of value? A randomised clinical trial. *Br J Obstet Gynaecol* 1999;106:1273–9.
116. Bennett KA, Crane JM, O'Shea P, Lacelle J, Hutchens D, Copel JA. First trimester ultrasound screening is effective in reducing postterm labor induction rates: a randomized controlled trial. *Am. J Obstet Gynecol* 2004;190:1077–81.
117. Harrington DJ, MacKenzie IZ, Thompson K, Fleminger M, Greenwood C. Does a first trimester dating scan using crown rump length measurement reduce the rate of induction of labour for prolonged pregnancy? An uncompleted randomised controlled trial of 463 women. *BJOG* 2006;113:171–6.
118. Woolf SH, Battista RN, Angerson GM, Logan AG, Eel W. Canadian Task Force on Preventive Health Care. New grades for recommendations from the Canadian Task Force on Preventive Health Care. *CMAJ* 2003;169:207–8.