

Arteriosclerosis, Thrombosis, and Vascular Biology



JOURNAL OF THE AMERICAN HEART ASSOCIATION

Contribution of Gene Sequence Variations of the Hepatic Cytochrome P450 3A4 Enzyme to Variability in Individual Responsiveness to Clopidogrel

Dominick J. Angiolillo, Antonio Fernandez-Ortiz, Esther Bernardo, Celia Ramírez, Ugo Cavallari, Elisabetta Trabetti, Manel Sabaté, Rosana Hernández, Raul Moreno, Javier Escaned, Fernando Alfonso, Camino Bañuelos, Marco A. Costa, Theodore A. Bass, Pier Franco Pignatti and Carlos Macaya

Arterioscler Thromb Vasc Biol. 2006;26:1895-1900; originally published online April 27, 2006;
doi: 10.1161/01.ATV.0000223867.25324.1a

Arteriosclerosis, Thrombosis, and Vascular Biology is published by the American Heart Association, 7272
Greenville Avenue, Dallas, TX 75231

Copyright © 2006 American Heart Association, Inc. All rights reserved.
Print ISSN: 1079-5642. Online ISSN: 1524-4636

The online version of this article, along with updated information and services, is located on the
World Wide Web at:

<http://atvb.ahajournals.org/content/26/8/1895>

Data Supplement (unedited) at:

<http://atvb.ahajournals.org/content/suppl/2006/04/27/01.ATV.0000223867.25324.1a.DC1.html>

Permissions: Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Arteriosclerosis, Thrombosis, and Vascular Biology* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

Reprints: Information about reprints can be found online at:
<http://www.lww.com/reprints>

Subscriptions: Information about subscribing to *Arteriosclerosis, Thrombosis, and Vascular Biology* is online at:
<http://atvb.ahajournals.org/subscriptions/>

Contribution of Gene Sequence Variations of the Hepatic Cytochrome P450 3A4 Enzyme to Variability in Individual Responsiveness to Clopidogrel

Dominick J. Angiolillo, Antonio Fernandez-Ortiz, Esther Bernardo, Celia Ramírez, Ugo Cavallari, Elisabetta Trabetti, Manel Sabaté, Rosana Hernández, Raul Moreno, Javier Escaned, Fernando Alfonso, Camino Bañuelos, Marco A. Costa, Theodore A. Bass, Pier Franco Pignatti, Carlos Macaya

Objectives—Metabolic activity of cytochrome P450 (CYP) 3A4 has been associated with clopidogrel response variability. Because metabolic activity of CYP3A4 is genetically regulated, we hypothesized that genetic variations of this enzyme may contribute to clopidogrel response variability.

Methods and Results—The CYP3A4*1B, CYP3A4*3, IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A polymorphisms of the CYP3A4 gene were assessed in 82 patients in a steady phase of clopidogrel therapy. Glycoprotein (platelet glycoprotein (GP) IIb/IIIa receptor activation and platelet aggregation were assessed. A cohort of 45 clopidogrel-naïve patients was studied to determine the modulating effects of these polymorphisms after loading dose (300 mg) administration. Only the IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A polymorphisms were sufficiently polymorphic. During the steady phase of clopidogrel treatment, IVS10+12A allele carriers had reduced GP IIb/IIIa activation ($P=0.025$) and better responsiveness ($P=0.02$); similarly, clopidogrel-naïve patients carriers of the IVS10+12A allele had reduced GP IIb/IIIa activation during the first 24 hours after a loading dose ($P=0.025$), increased platelet inhibition ($P=0.006$), and a more optimal drug response ($P=0.003$). This polymorphism did not influence platelet aggregation profiles. No association was observed between the other polymorphisms and clopidogrel responsiveness.

Conclusions—The IVS10+12G>A polymorphism of the CYP3A4 gene modulates platelet activation in patients treated with clopidogrel and may therefore contribute to clopidogrel response variability. (*Arterioscler Thromb Vasc Biol.* 2006;26:1895-1900.)

Key Words: clopidogrel ■ platelet ■ polymorphism

Treatment with clopidogrel is associated with a broad variability in antiplatelet effects.¹⁻⁴ This may be partly attributed to the levels of clopidogrel's active metabolite.⁵ Clopidogrel in fact is an inactive pro-drug that requires oxidation by the hepatic cytochrome P450 3A4 (CYP3A4) to generate an active metabolite.⁶⁻⁷ The active metabolite of clopidogrel inhibits platelet activation through an irreversible blockage of the platelet adenosine diphosphate (ADP) P2Y₁₂ receptor. The P2Y₁₂ receptor inhibits adenylyl cyclase and in turn decreases platelet cAMP (cAMP) levels and cAMP-mediated phosphorylation of the vasodilator-stimulated phosphoprotein, critical for inhibition of glycoprotein (GP) IIb/IIIa receptor activation.⁸

active metabolite.⁹⁻¹¹ Accordingly, the metabolic activity of the CYP3A4 enzyme, which varies considerably among individuals,¹² has shown to influence platelet reactivity after a clopidogrel loading dose.¹³ Because genetic predisposition is the major determinant of heterogeneity in metabolic activity of the CYP3A4 enzyme,^{12,14-15} we hypothesized that single-nucleotide polymorphisms (SNPs) of CYP3A4 may account for interindividual variability of platelet reactivity in patients treated with clopidogrel. In the present study we examined the influence of SNPs of the CYP3A4 enzyme on platelet reactivity and responsiveness to clopidogrel in patients with coronary artery disease.

See page 1681

Drugs that are substrates or inhibit CYP3A4 can potentially interfere with the conversion of clopidogrel into its

Materials and Methods

Detailed Materials and Methods are provided in the expanded methods, available online at <http://atvb.ahajournals.org>.

Original received November 28, 2005; final version accepted April 10, 2006.

From the Division of Cardiology (D.J.A., M.A.C., T.A.B.), University of Florida, Shands Jacksonville; Cardiovascular Institute (A.F.-O., E.B., C.R., M.S., R.H., R.M., J.E., F.A., C.B., C.M.), San Carlos University Hospital, Madrid, Spain; and the Department of Mother and Child & Biology-Genetics (U.C., E.T., P.F.), University of Verona, Italy.

Correspondence to Dominick J Angiolillo, MD, PhD, FACC, FESC, Division of Cardiology, University of Florida, Shands Jacksonville, 655 West 8th St, Jacksonville, FL 32209. E-mail dominick.angiolillo@jax.ufl.edu

© 2006 American Heart Association, Inc.

Arterioscler Thromb Vasc Biol. is available at <http://www.atvbaha.org>

DOI: 10.1161/01.ATV.0000223867.25324.1a

Population

Eighty-two patients with stable coronary artery disease using combined aspirin and clopidogrel therapy were studied. Because ethnicity may contribute to variability in genotype distribution of specific SNPs, to avoid a patient selection bias only white patients homogeneous for ethnic background were included;^{12,14–15} all patients were from the central regions of Spain. In addition, because clopidogrel may take several days or weeks to achieve full antiplatelet effects, which per se is responsible for individual response variability,³ only patients who were in a steady phase (≥ 1 month) of clopidogrel (75 mg/daily) treatment were included. All patients were clinically stable from the time that they initiated dual antiplatelet therapy and were on the same aspirin dose (100 mg/daily). Platelet reactivity was assessed 2 to 4 hours after drug intake. Because platelet reactivity may vary over time, platelet function measurements were repeated at a separate time point in patients who were still using dual antiplatelet therapy (70%) to sustain our study hypothesis. Compliance to antiplatelet treatment was ensured by pill count and/or interview. Patients on other antithrombotic drugs (oral anticoagulants, dipyridamole, cilostazol) or using clopidogrel for < 1 month were not eligible for the study.

A separate cohort of 45 clopidogrel naïve patients scheduled to undergo elective coronary stenting receiving a 300-mg clopidogrel loading dose was also studied. All patients were on aspirin for at least 7 days and received clopidogrel at the time of coronary intervention. Platelet reactivity was assessed before clopidogrel administration (patients only on aspirin) and 4 and 24 hours after front loading. During intervention, unfractionated heparin (100 IU/kg) was administered according to standard practice. Only bare metal stents were used. Exclusion criteria were use of lytics or platelet GP IIb/IIIa blockers before, during, or after intervention; before use of thienopyridines; and/or aspirin intolerance/allergy.

This study was approved by the Ethical Committee of the San Carlos University Hospital and all patients gave informed consent.

Genotyping

Genomic DNA was extracted from whole blood samples according to standard protocols.^{16–17} Overall > 30 SNPs of CYP3A4 have been described.^{12,14–15} These SNPs have been identified in different regions of the CYP3A4 gene. However, SNPs of the CYP3A4 gene are not homogeneously present and vary among different ethnicities. Therefore, the selection of SNPs for this study was based on the allele heterozygosity information for the CYP3A4 gene in the CEPH population derived from the HapMap Project Public Release #20.¹⁸ All the HapMap genotyped SNPs with a minor allele frequency $> 1\%$ in the CEPH population were selected for the study: CYP3A4*1B, CYP3A4*3, IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A.

Genotyping of the CYP3A4*1B (c.-392A>G) SNP, located in the promoter region of the CYP3A4 gene, was performed using polymerase chain reaction (PCR) and incubation of the amplification product with the restriction endonuclease *Bst*NI; 2 fragments of 149 and 20 bp for the G allele were obtained, and a single fragment of 169 was obtained for the A allele. Genotyping of the CYP3A4*3 (p.M445T) SNP located in exon 12 of the CYP3A4 gene was performed using PCR and incubation of the amplification product with the restriction endonuclease *M*slI; 2 fragments of 201 and 91 bp for the M allele were obtained, whereas a single fragment of 292 bp, representing the T allele, was not found. Genotyping of the IVS7+258A>G (c.670+258A>G) SNP located in the intron 7 of the CYP3A4 gene was performed using PCR and incubation of the amplification product with the restriction endonuclease *M*nII; 2 fragments of 153 and 43 bp for the G allele were obtained, and a single fragment of 196 was obtained for the A allele. Genotyping of the IVS7+894C>T (c.670+894C>T) SNP located in the intron 7 of the CYP3A4 gene was performed using PCR and incubation of the amplification product with the restriction endonuclease *A*ciI; and 2 fragments of 149 and 172 bp for the C allele were obtained, and a single fragment of 323 was obtained for the T allele.

Genotyping of the IVS10+12G>A (c.1026+12G>A) SNP located in intron 10 of the CYP3A4 gene was performed using PCR and incubation of the amplification product with the restriction

endonuclease *R*saI; 2 fragments of 240 and 100 bp for the G allele were obtained, and a single fragment of 340 bp was obtained for the A allele.

Although these SNPs have been previously reported, their prevalence in our population composed of patients with coronary artery disease is unknown. Therefore, a priori, these SNPs were considered for functional implication analysis only if sufficiently polymorphic and in Hardy-Weinberg equilibrium. Patients were predefined to be divided into 2 groups according to a dominant model, in which subjects homozygous for the wild-type allele are compared with subjects carrying ≥ 1 variant allele.¹⁷

GP IIb/IIIa Activation

Platelet reactivity was determined by assessing platelet surface expression of activated GP IIb/IIIa after 2 μ mol/L ADP (ChronoLog Corp, Havertown, Pa) stimuli through flow cytometry, as previously described.^{1,2}

Assessment of Platelet Aggregation

ADP-induced (6 μ mol/L and 20 μ mol/L) platelet aggregation was assessed in duplicate using platelet rich plasma (PRP) by the turbidimetric method in a 2-channel aggregometer (Chrono-Log 490 Model; Chrono-Log Corp), as previously described.^{1,2,17}

Variability in Clopidogrel Responsiveness

In patients using sustained clopidogrel treatment, variability in clopidogrel induced antiplatelet effects was assessed by determining the distribution of platelet reactivity using a coefficient of variability (CV). A significant variability was defined when CV ($CV = SD/mean$) was > 0.25 in continuous variables with a normal distribution.² To assess the influence of SNPs of the CYP3A4 gene on variability of individual response to clopidogrel, this patient population was stratified into 3 groups on the basis of the tertiles of platelet reactivity (low, moderate, and high platelet reactivity). Patients in the lower tertile were considered as more optimal responders as opposed to those in the higher tertile, who were considered poor responders.

In patients treated with a 300-mg clopidogrel loading dose, clopidogrel responsiveness was defined according to the degree of inhibition of platelet reactivity 24 hours after clopidogrel administration compared with baseline values. Patients were classified as nonresponders, low responders, and responders when platelet inhibition was $< 10\%$, 10% to 29% , and $\geq 30\%$, respectively.²

Statistical Analysis

Continuous variables are expressed as mean \pm SD. Categorical variables are expressed as frequencies and percentages. The Kolmogorov-Smirnov test was performed to assess normal distribution. Comparisons between categorical variables were performed using 2-tailed Fisher exact test or the Pearson χ^2 test as appropriate. The Student *t* test was used for normally distributed continuous variables. Hardy-Weinberg equilibrium was estimated by χ^2 test for each SNP. Lewontin *D'* was used as a measure of pairwise linkage disequilibrium.¹⁹ Haplotype analysis and linkage disequilibrium calculations were performed using R statistical package with haplostats and genetic libraries.²⁰ A multivariate analysis of variance (MANOVA) was used to assess intra-subject variance of platelet function parameters in patients receiving a 300-mg loading dose and their change over the study time period.² $P < 0.05$ was considered statistically significant. Statistical analysis was performed using a SPSSv11.0 software (SPSS Inc, Chicago, Ill).

Results

Genotyping

The variant allele of the CYP3A4*1B polymorphism was present only in 3% of the overall patient population. The variant allele of the CYP3A4*3 was not observed in any patient. Therefore, the extremely low prevalence of these

Genetic Distribution of the IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A Polymorphisms in Patients Using Sustained Clopidogrel Therapy (Long-Term) and After Loading Dose Administration (Acute)

	IVS7+258A>G		IVS7+894C>T		IVS10+12G>A	
	Long-Term	Acute	Long-Term	Acute	Long-Term	Acute
wt/wt	41 (50%)	18 (39%)	64 (78%)	36 (80%)	68 (83%)	36 (80%)
wt/v+v/v	41 (50%)	27 (61%)	18 (22%)	9 (20%)	14 (17%)	9 (20%)

Patients were predefined to be divided into 2 groups according to a dominant model, in which subjects homozygous for the wild-type allele (wt) are compared with subjects carrying ≥ 1 variant allele (v).

alleles did not allow them to be considered as genetic modulators of clopidogrel induced antiplatelet effects. The IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A polymorphisms were in Hardy-Weinberg equilibrium and sufficiently polymorphic to be considered as potential modulators of platelet reactivity. Genetic distribution of the IVS7+258A>G, IVS7+894C>T, and IVS10+12G>A polymorphisms is described in the Table. Linkage disequilibrium was observed for some of the polymorphisms, particularly among the IVS7+894C>T, the IVS10+12G>A, and the CYP3A4*1B SNP (supplemental Table I, available online at <http://atvb.ahajournals.org>). Two common haplotypes constituted by 4 SNPs represent 58.3% and 28.1% of the studied population (supplemental Table II).

Platelet Reactivity in Patients Using Sustained Clopidogrel Therapy

GP IIb/IIIa receptor activation was highly variable (CV=0.65) and followed a normal distribution. The duration of clopidogrel treatment in these patients was 8.6 ± 4.9 months. Increased GP IIb/IIIa activation was observed in noncarriers compared with carriers of the IVS10+12A allele ($P=0.025$; Figure 1a). Repeated platelet function measurements were performed in 70% (n=54) of this patient population observing similar variability profile (CV=0.66), as well as differences in platelet activation measures in noncarriers compared with carriers of the IVS10+12A allele ($30.6 \pm 19.5\%$ versus $17.4 \pm 10.9\%$; $P=0.035$). The distribution of IVS10+12A allele carrier patients related to their clinical characteristics was not significantly different (supplemental Table III). There were no differences between groups, including the use of CYP3A4 and non-CYP3A4 pathway metabolized statins.

After tertile distribution analysis of platelet reactivity profiles, carriers of the variant allele were more likely to have low platelet activation ($P=0.02$; Figure 1b). Similar platelet activation distribution (lower, 0.7 to 16.7%; middle, 16.7 to 35.1%; upper, 35.1 to 74.7%) and allocation of carriers and noncarriers of the variant allele were observed in the subgroup of patients with repeated measurements ($P=0.03$). A broad variability in platelet aggregation profiles was also observed after $6 \mu\text{mol/L}$ (CV=0.44) and $20 \mu\text{mol/L}$ (CV=0.32) ADP stimuli. However, noncarriers and carriers of the variant allele of the IVS10+12G>A polymorphism had similar degrees of platelet aggregation ($31.7 \pm 14.1\%$ versus $32.9 \pm 15.7\%$, $P=0.7$ for $6 \mu\text{mol/L}$ ADP; and $45.4 \pm 14.8\%$ versus $46.5 \pm 15.9\%$, $P=0.8$ for $20 \mu\text{mol/L}$

ADP). Hence, their distribution according to platelet aggregation tertiles was also similar ($P=0.7$ for $6 \mu\text{mol/L}$ ADP and $P=0.9$ for $20 \mu\text{mol/L}$ ADP). Similar findings were observed in the subgroup of patients with repeated measurements (data not shown). No association was observed between platelet reactivity and the other genotyped SNPs in patients using sustained clopidogrel therapy. The analysis performed on the identified haplotypes showed no significant association.

Platelet Reactivity After a 300-mg Clopidogrel Loading Dose

GP IIb/IIIa receptor activation was similar in noncarriers and carriers of the variant allele of the IVS10+12G>A poly-




Figure 1. A, ADP-induced ($2 \mu\text{mol/L}$) glycoprotein IIb/IIIa receptor activation in patients carriers (gray) and noncarriers (white) of the variant allele of the IVS10+12G>A polymorphism on sustained clopidogrel therapy. Values are expressed as percentage of positive platelets (mean \pm SD) and compared using Student *t* test. B, ADP-induced ($2 \mu\text{mol/L}$) glycoprotein IIb/IIIa receptor activation in patients carriers (gray) and noncarriers (white) of the variant allele of the IVS10+12G>A polymorphism on sustained clopidogrel therapy according to tertile distribution. Probability value was calculated using χ^2 test.




Figure 2. ADP (2 μ mol/L)-induced glycoprotein IIb/IIIa receptor activation before (baseline, patients only on aspirin) and 4 and 24 hours after a 300-mg clopidogrel loading dose in patients carriers (gray) and noncarriers (black) of the variant allele of the IVS10+12G>A polymorphism. MANOVA was used for comparison between the 2 groups during the overall study ($P=0.025$). Values are expressed as percentage of positive platelets (mean \pm SD).

morphism at baseline, before clopidogrel front-loading. No significant difference was observed in the IVS10+12G>A genotype distribution regarding to their clinical characteristics (supplemental Table III). After loading dose administration, a lower reduction of GP IIb/IIIa receptor activation was observed in noncarriers of the variant allele of this polymorphism at 4 and 24 hours, with platelet activation significantly higher in these patients during the overall study time course ($P=0.025$) (Figure 2). At 24 hours the degree of platelet inhibition was significantly higher in carriers of the variant allele ($P=0.006$; Figure 3a). Noncarriers of the variant allele had a poorer response to clopidogrel as compared with carriers (non responders, 36% versus 11%; low responders, 36% versus 0%; responders, 28% versus 89%; $P=0.003$; Figure 3b). ADP-induced platelet aggregation sig-




Figure 3. A, Percentage inhibition of ADP (2 μ mol/L)-induced glycoprotein IIb/IIIa receptor activation 24 hours after a 300-mg clopidogrel loading dose in patients carriers (gray) and noncarriers (white) of the variant allele of the IVS10+12G>A polymorphism. Values are expressed as percentage of inhibition of positive platelets (mean \pm SD) and compared using Student *t* test. B, Percentage platelet inhibition and clopidogrel responsiveness. Individual patients are identified according to percentage inhibition of platelet activation and defined as clopidogrel nonresponders (\circ), low responders (\square), and responders (\triangle) when platelet inhibition was <10%, 10% to 29%, and \geq 30%, respectively, and compared using χ^2 testing. Dashed lines indicate 10% and 30% inhibition of platelet activation.

nificantly reduced after clopidogrel administration; however, it was similar in both groups (data not shown). No association was observed between platelet reactivity and the other genotyped SNPs in patients on 300 mg clopidogrel loading dose. The analysis performed on the identified haplotypes showed no significant association.

Discussion

The present study demonstrates the role of gene sequence variations of CYP3A4 in modulating platelet activation in patients treated with clopidogrel. In particular, carriers of the IVS10+12A allele had reduced GP IIb/IIIa activation and a better response to clopidogrel compared with noncarriers of the variant allele. The hypothesis of the modulating effect of this polymorphism on platelet activation in patients treated with clopidogrel is supported not only by the differences in platelet activation observed in patients already in a steady phase of clopidogrel treatment, but also in the acute phase of treatment. Although platelet reactivity may vary over time, analogous findings persisted in patients undergoing repeated measurements in the sustained phase of clopidogrel treatment, further sustaining our study hypothesis. The modulating effects of the IVS10+12G>A polymorphism as assessed by the degree of inhibition of platelet activation is noteworthy in clopidogrel naïve patients carriers of the A allele who, except for 1 patient, were all clopidogrel responders. However, this polymorphism was not associated with a modulating effect on platelet aggregation. These discrepancies with platelet activation may be attributed to the mechanism of action of clopidogrel. In fact, because clopidogrel is per se an inhibitor of platelet activation, this may explain why GP IIb/IIIa receptor activation and not platelet aggregation resulted modulated by the IVS10+12G>A polymorphism of the CYP3A4 gene. In patients treated with clopidogrel, inhibition of platelet aggregation represents a downstream effect after inhibition of platelet activation. In addition, the resulting degree of platelet aggregation is influenced by numerous stimuli, which may affect to a lesser extent platelet activation.² Because clopidogrel mediates its effects through vasodilator-stimulated phosphoprotein phosphorylation, which has a pivotal role for GP IIb/IIIa receptor inhibition, assessment of GP IIb/IIIa receptor activation may be a more sensitive platelet biomarker to assess the modulating effects of CYP3A4 gene polymorphisms.

Clopidogrel response variability has been primarily assessed during the acute phases of treatment during which clopidogrel's pharmacokinetic and pharmacodynamic profiles vary considerably.^{5,21} One may argue that such variability depends on the fact that drug-induced metabolic activity of CYP3A4 requires several days or weeks of treatment before this reaches a plateau and thus a more uniform response. This is supported by the fact that clopidogrel responsiveness ameliorates with length of treatment.³ However, a broad variability in clopidogrel induced effects continued to be observed even in patients in a steady phase of treatment.

Several factors may contribute to clopidogrel response variability.²² SNPs of targets within the clopidogrel pathway, namely CYP3A4 enzyme, P2Y₁₂ receptor, and GP IIb/IIIa

receptor, have been suggested to contribute to individual response variability to clopidogrel. A minor haplotype of the P2Y₁₂ receptor was found to be associated with increased platelet reactivity in nonmedicated healthy volunteers.²³ However, these findings could not be duplicated by several authors studying patients with coronary artery disease treated with clopidogrel.^{17,24–26} In addition to P2Y₁₂ SNP analysis, Lev et al failed to observe any impact of SNPs of the P2Y₁ and GPIIIa receptors on responsiveness to acute front-loading of clopidogrel.²⁶ We recently demonstrated the lack of association between SNPs of the GP IIIa and P2Y₁₂ receptors and platelet reactivity in patients on chronic clopidogrel therapy.^{17,27} These findings are likely related to the fact that an active metabolite and not clopidogrel per se, is responsible for inhibition of the P2Y₁₂ receptor, therefore suggesting that an upstream target within clopidogrel's metabolic pathway to have a more important modulating role of its downstream antiplatelet effects. One may argue that CYP3A5 has overlapping substrate specificity with CYP3A4 and may contribute to clopidogrel's metabolism and be a determinant of drug response as well. However, gene sequence variations of the CYP3A5 enzyme in patients with coronary artery disease have not been associated with variability in clopidogrel responsiveness.²⁵ This latter observation may be explained by the fact that CYP3A4 is the dominant CYP3A enzyme among whites¹² and therefore may be more important in clopidogrel metabolism. Drug–drug interactions and clinical status are also responsible for variability in platelet reactivity and clopidogrel responsiveness.²² However, our patient population was very homogenous not only for ethnicity but also for clinical status and treatment.

In the present study, the overall prevalence of clopidogrel nonresponders compares favorably with that previously described.^{2,3} Notably, noncarriers of the IVS10+12A allele were predominant. This may explain the high prevalence of clopidogrel nonresponders using standard clopidogrel dosing. Recent reports have shown that higher doses of clopidogrel improve responsiveness.^{2,28} However, our study did not assess if SNPs of the CYP3A4 gene continue to contribute to variability in platelet reactivity when higher clopidogrel doses are used. Nevertheless, it is noteworthy to emphasize that although increasing clopidogrel doses improves responsiveness, a broad variability in antiplatelet effects persist independent of the dose used.^{2,28} Platelet reactivity in fact may be within higher or lower ranges according to the dose used but still remain highly variable. This is related to the fact that clopidogrel's antiplatelet effects primarily depend on the metabolic activity of CYP3A4 which is per se heterogeneous. On the contrary, direct P2Y₁₂ antagonists such as AZD6140 or cangrelor, which do not require hepatic metabolism to be active, do not present such heterogeneous effects.²⁹

Although this study was not aimed to assess clinical events, it is important to underscore that increased post-treatment platelet reactivity and suboptimal response to antiplatelet drugs, including clopidogrel, which were both observed in this study, are associated with increased atherothrombotic risk.^{30–33} Larger population studies are warranted to assess if the functional findings from this study have clinical sequelae and the assessment of candidate genetic traits appears as a

promising arena of research to identify such high-risk individuals. Ultimately, the identification of relevant genetic markers may guide a more rationale and individualized use of drugs and dosages, thus optimizing therapeutic regimens which are essential for the prevention of atherothrombotic events.

Study Limitations

Measurement of CYP3A4's metabolic activity was not performed in this study. However, a correlation between CYP3A4's metabolic activity and clopidogrel response has been described¹³ and the functional findings of our study support the hypothesis that individuals genetic predisposition, a major determinant of CYP3A4's metabolic activity, contributes to this phenomenon. Although all the described SNPs with minor allele frequency >1% have been genotyped in this study, we cannot exclude the contributing role of other SNPs in modulating clopidogrel response. IVS10+12G>A polymorphism of the CYP3A4 gene, which is in linkage disequilibrium with some other SNPs, may be a marker for an haplotype which regulates the metabolic activity of CYP3A4s by either modifying catalytic activity and/or enzyme expression, possibly through other, as yet undetected, SNPs.^{12,16}

Acknowledgments

We thank Cristina Fernandez, MD, PhD, for her support with the statistical analyses of this study.

Sources of Funding

This work has been partially supported by a grant from the Spanish Ministry of Health, the Italian Ministry of University and Research and the Veneto Region Sanitary Research Project.

Disclosures

Dominick J. Angiolillo is on the speaker bureau and is a consultant for Sanofi-Aventis and Bristol Myers Squibb. The remaining authors report no conflicts.

References

1. Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Barrera-Ramirez C, Sabate M, Hernandez R, Moreno R, Escaned J, Alfonso F, Banuelos C, Costa MA, Bass TA, Macaya C. Identification of low responders to a 300 mg clopidogrel loading dose in patients undergoing coronary stenting. *Thromb Res*. 2005;115:101–108.
2. Angiolillo DJ, Fernandez-Ortiz A, Bernardo E, Ramirez C, Sabate M, Banuelos C, Hernandez-Antolin R, Escaned J, Moreno R, Alfonso F, Macaya C. High clopidogrel loading dose during coronary stenting: Effects on drug response and interindividual variability. *Eur Heart J*. 2004;25:1903–1910.
3. Gurbel PA, Bliden KP, Hiatt BL, O'Connor CM. Clopidogrel for coronary stenting: response variability, drug resistance, and the effect of pretreatment platelet reactivity. *Circulation*. 2003;107:2908–2913.
4. Serebruany VL, Steinhubl SR, Berger PB, Malinin AI, Bhatt DL, Topol EJ. Variability in platelet responsiveness to clopidogrel among 544 individuals. *J Am Coll Cardiol*. 2005;45:246–251.
5. Taubert D, Kastrati A, Harlfinger S, Gorchakova O, Lazar A, von Beckerath N, Schomig A, Schomig E. Pharmacokinetics of clopidogrel after administration of a high loading dose. *Thromb Haemost*. 2004;92:311–316.
6. Savi P, Pereillo JM, Uzabiaga MF, Combalbert J, Picard C, Maffrand JP, Pascal M, Herbert JM. Identification and biological activity of the active metabolite of clopidogrel. *Thromb Haemost*. 2000;84:891–896.
7. Pereillo JM, Maftouh M, Andrieu A, Uzabiaga MF, Fedeli O, Savi P, Pascal M, Herbert JM, Maffrand JP, Picard C. Structure and stereochemistry of the active metabolite of clopidogrel. *Drug Metab Dispos*. 2002;30:1288–1295.

8. Geiger J, Brich J, Honig-Liedl P, Eigenthaler M, Schanzenbacher P, Herbert JM, Walter U. Specific impairment of human platelet P2Y₁₂(AC) ADP receptor-mediated signaling by the antiplatelet drug clopidogrel. *Arterioscler Thromb Vasc Biol.* 1999;19:2007–2011.
9. Lau WC, Waskell LA, Watkins PB, Neer CJ, Horowitz K, Hopp AS, Tait AR, Carville DG, Guyer KE, Bates ER. Atorvastatin reduces the ability of clopidogrel to inhibit platelet aggregation: a new drug-drug interaction. *Circulation.* 2003;107:32–37.
10. Neubauer H, Gunesdogan B, Hanefeld C, Spiecker M, Mugge A. Lipophilic statins interfere with the inhibitory effects of clopidogrel on platelet function—a flow cytometry study. *Eur Heart J.* 2003;24:1744–1749.
11. Clarke TA, Waskell LA. The metabolism of clopidogrel is catalyzed by human cytochrome P450 3A and is inhibited by atorvastatin. *Drug Metab Dispos.* 2003;31:53–59.
12. Lamba JK, Lin YS, Schuetz EG, Thummel KE. Genetic contribution to variable human CYP3A-mediated metabolism. *Adv Drug Deliv Rev.* 2002;54:1271–1294.
13. Lau WC, Gurbel PA, Watkins PB, Neer CJ, Hopp AS, Carville DG, Guyer KE, Tait AR, Bates ER. Contribution of hepatic cytochrome P450 3A4 metabolic activity to the phenomenon of clopidogrel resistance. *Circulation.* 2004;109:166–171.
14. Wojnowski L. Genetics of the variable expression of CYP3A in humans. *Ther Drug Monit.* 2004;26:192–199.
15. Lamba JK, Lin YS, Thummel K, Daly A, Watkins PB, Strom S, Zhang J, Schuetz EG. Common allelic variants of cytochrome P4503A4 and their prevalence in different populations. *Pharmacogenetics.* 2002;12:121–132.
16. Quinzii C, Belpinati F, Pignatti PF. Predictive genetic testing: new possibilities in determination of risk of complex diseases. *Croat Med J.* 2001;42:458–462.
17. Angiolillo DJ, Fernández-Ortiz A, Bernardo E, Ramirez C, Cavallari U, Trabetti E, Sabate M, Jimenez-Quevedo P, Hernandez R, Moreno R, Escaned J, Alfonso F, Banuelos C, Costa MA, Bass TA, Pignatti PF, Macaya C. Lack of association between the P2Y₁₂ receptor gene polymorphism and platelet response to clopidogrel in patients with coronary artery disease. *Thromb Res.* 2005;116:491–497.
18. The International HapMap Consortium. The International HapMap Project. *Nature.* 2003;426:789–796.
19. Lewontin RC. On measures of gametic disequilibrium. *Genetics.* 1988;120:849–852.
20. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. (<http://www.R-project.org>).
21. Caplain H, Donat F, Gaud C, Necciari J. Pharmacokinetics of clopidogrel. *Semin Thromb Hemost.* 1999;25(Suppl 2):25–28.
22. Nguyen TA, Diodati JG, Pharand C. Resistance to clopidogrel: a review of the evidence. *J Am Coll Cardiol.* 2005;45:1157–1164.
23. Fontana P, Dupont A, Gandrille S, Bachelot-Loza C, Reny JL, Aiach M, Gaussem P. Adenosine diphosphate-induced platelet aggregation is associated with P2Y₁₂ gene sequence variations in healthy subjects. *Circulation.* 2003;108:989–995.
24. von Beckerath N, von Beckerath O, Koch W, Eichinger M, Schomig A, Kastrati A. P2Y₁₂ gene H2 haplotype is not associated with increased adenosine diphosphate-induced platelet aggregation after initiation of clopidogrel therapy with a high loading dose. *Blood Coagul Fibrinolysis.* 2005;16:199–204.
25. Smith SMG, Judge HM, Peters G, Wilmington, Armstrong M, Fontana P, Gaussem P, Storey RF. Common sequence variations in the P2Y₁₂ and CYP3A5 genes do not explain variability in the inhibitory effects of clopidogrel therapy. *Eur Heart J.* 2005;26:P2963 [abstract].
26. Lev EI, Patel RT, Guthikonda S, Lopez D, Bray PF, Kleiman NS. Genetic polymorphisms of the platelet receptors P2Y₁₂(12), P2Y₁(1) and GP IIIa and response to aspirin and clopidogrel. *Thromb Res.* 2006 (in press).
27. Angiolillo DJ, Bernardo E, Ramirez C, Cavallari U, Jimenez-Quevedo P, Escaned J, Trabetti E, Pignatti PF, Fernández-Ortiz A. Polymorphisms of the GP IIIa and P2Y₁₂ receptors and modulation of antiplatelet effects of combined aspirin and clopidogrel treatment. *Circulation.* 2004;110: 2013 [abstract].
28. Gurbel PA, Bliden KP, Hayes KM, Yoho JA, Herzog WR, Tantry US. The relation of dosing to clopidogrel responsiveness and the incidence of high post-treatment platelet aggregation in patients undergoing coronary stenting. *J Am Coll Cardiol.* 2005;45:1392–1396.
29. van Giezen JJ, Humphries RG. Preclinical and clinical studies with selective reversible direct P2Y₁₂ antagonists. *Semin Thromb Hemost.* 2005;31:195–204.
30. Gurbel PA, Bliden KP, Samara W, Yoho JA, Hayes KM, Tantry U. Clopidogrel Resistance and Stent Thrombosis (CREST) study. *J Am Coll Cardiol.* 2005;46:1827–1832.
31. Gurbel PA, Bliden KP, Guyer K, Cho PW, Zaman KA, Kreutz RP, Bassi AK, Tantry US. Platelet reactivity in patients and recurrent events post-stenting: results of the PREPARE POST-STENTING Study. *J Am Coll Cardiol.* 2005;46:1820–1826.
32. Matetzky S, Shenkman B, Guetta V, Shechter M, Bienart R, Goldenberg I, Novikov I, Pres H, Savion N, Varon D, Hod H. Clopidogrel resistance is associated with increased risk of recurrent atherothrombotic events in patients with acute myocardial infarction. *Circulation.* 2004;109: 3171–3175.
33. Muller I, Besta F, Schulz C, Massberg S, Schomig A, Gawaz M. Prevalence of clopidogrel non-responders among patients with stable angina pectoris scheduled for elective coronary stent placement. *Thromb Haemost.* 2003;89:783–787.

Online Supplement for:

Angiolillo DJ et al. "Contribution of Gene Sequence Variations of the Hepatic Cytochrome P450 3A4 Enzyme to Variability in Individual Responsiveness to Clopidogrel" *Arterioscler Thromb Vasc Biol.* 2006

- - - -

Materials and Methods

GP IIb/IIIa activation

Platelet reactivity was determined by assessing platelet surface expression of activated GP IIb/IIIa following 2 μ M ADP (ChronoLog Corp, Havertown, PA) stimuli through flow cytometry, as previously described (1,2) . GP IIb/IIIa activation was assessed using a polyclonal fluorescein isothiocyanate (FITC)-conjugated rabbit anti-human fibrinogen antibody (800nM, DAKO Diagnostics, Glostrup, Denmark), which detects fibrinogen binding to the activated GPIIb/IIIa receptor. An EPICS-XL PROFILE II Coulter flow cytometer (Coulter Corp. Miami, FL) was used for the assessment. After discard of the initial first millilitres of blood to avoid spontaneous platelet activation, whole blood was drawn into sterile tubes containing 3.8% trisodium citrate and then diluted with HEPES-tyrodes buffer (5mM HEPES hydroxyethylpiperazineethane-sulfonic acid-, 137mM NaCl, 2,7mM NaHCO₃, 0,36mM NaH₂PO₄, 2mM CaCl₂, 5mM glucose, 0,2% BSA) to a final volume of 1:8:1 (blood:HEPES-Tyrodes:citrate) resulting in a 1/10 dilution of whole blood during sampling. Previously HEPES-Tyrodes buffer were filtered through 0.22 μ m sterile filters to avoid interference from particles. Then, 50 μ L of diluted whole blood was stimulated in vitro with 2 μ M ADP before

immunolabelling. The corresponding antibody was then added and incubated for 20 min in dark at room temperature. After incubation, 300µl of 0.5% PBS-buffered paraformaldehyde was added for fixation. Samples were analyzed within 2 hours by flow cytometry and platelets were identified based on particle size (forward scatter) and complexity (side scatter). Light scatter and fluorescence data from 10000 platelet events were collected with all detectors in logarithmic mode. Acquisition and processing data were analyzed with XL2 software (Coulter Corp. Miami, FL). Platelet activation was expressed as the percentage of platelets positive for antibody binding.

Assessment of platelet aggregation

Blood was collected in tubes containing 3.8% trisodium citrate. Platelet aggregation was assessed in duplicate using platelet rich plasma (PRP) by the turbidimetric method in a 2-channel aggregometer (Chrono-Log 490 Model, Chrono-Log Corp., Havertown, PA), as previously described (1, 2, 17). Platelet agonists included 6 µM ADP and 20 µM ADP (Chrono-Log Corp., Havertown, PA) (1, 2, 17). PRP was obtained as a supernatant after centrifugation of citrated blood at 800 rpm for 10 minutes. Platelet poor plasma (PPP) was obtained by a second centrifugation of the blood fraction at 2500 rpm for 10 minutes. The platelet count in PRP was adjusted to the range of 250.000/µL by dilution with autologous plasma when platelet count was out of range. Light transmission was adjusted to 0% with PRP and to 100% for PPP for each measurement. Platelet aggregation was assessed within 2 hours from blood sampling. PRP was kept at 22° C prior to use and at 37°C one minute before running the aggregatory test. Aggregation was assessed in siliconized tubes at 37°C in constant

stirring conditions and curves were recorded for 5 minutes. Platelet aggregation was determined as the maximal percent change in light transmittance from baseline using PPP as reference.

Online Table I. Linkage disequilibrium observed between the studied SNPs. D' with p value in brackets are reported for each SNP pair comparison.

	<i>IVS7+258A>G</i>	<i>IVS7+894C>T</i>	<i>IVS10+12G>A</i>
CYP 3A4*1B	0.141 (0.5585)	0.685 (1.19e ⁻⁰⁸)	0.757 (1.28e ⁻¹²)
IVS7+258	-	0.200 (0.1052)	0.264 (0.0375)
IVS7+894	-	-	0.845 (<2e ⁻¹⁶)

Lewontin's D' was used as a measure of pairwise linkage disequilibrium (reference 19).

Online Table II. Frequency of the haplotypes of the CYP3A4 gene estimated in the study population.

<i>CYP 3A4*1B</i> (c.-392A>G)	<i>CYP3A4*3</i> (p.M445T)	<i>IVS7+258A>G</i> (c.670+258A>G)	<i>IVS7+894C>T</i> (c.670+894C>T)	<i>IVS10+12G>A</i> (c.1026+12G>A)	Haplotype frequency
A	M	A	C	G	0.58383
A	M	G	C	G	0.28122
A	M	A	T	A	0.03477
A	M	G	T	A	0.03320
G	M	A	T	A	0.02422
A	M	A	T	G	0.02176
A	M	G	C	A	0.01216
G	M	A	C	G	0.00346
G	M	G	C	G	0.00341
A	M	G	T	G	0.00122
G	M	G	T	A	0.00076

Haplotype analyses were performed using R statistical package with haplo.stats and genetic libraries (<http://www.R-project.org>). (reference 20)

Online Table III. Demographics of the study population according to a dominant model distribution of the IVS10+12G>A polymorphism of the CYP3A4 gene

	Long-term phase			Acute-phase		
	(n=82)		p	(n=45)		p
	Non-carriers n=68 (83%)	Carriers n=14 (17%)		Non-carriers n=36 (80%)	Carriers n=9 (20%)	
Age (y)	63 ±11	61±11	0.56	62±10	59±11	0.39
Males	51 (75%)	11 (79 %)	1.00	31 (86%)	9 (100%)	0.57
Risk factors						
Diabetes	31 (47%)	3 (21%)	0.14	11 (31%)	2 (22%)	1.00
Hypertension	36 (53%)	5 (36%)	0.38	19 (53%)	4 (44%)	0.72
Hyperlipemia	36 (53%)	8 (57%)	1.00	22 (61%)	5 (56%)	1.00
Current smokers	6 (9%)	1 (7%)	1.00	21 (58%)	8 (89%)	0.13
Treatment						
B-Blockers	43 (63%)	9 (64%)	1.00	22 (61%)	7 (78%)	0.46
Nitrates	36 (53%)	9 (54%)	0.56	23 (64%)	7 (78%)	0.69
ACE-Inhibitors	29 (43%)	3 (21%)	0.22	14 (39%)	1 (11%)	0.23
Statins						
3A4 pathway metabolized	37 (54%)	7 (50%)	0.77	9 (25%)	2 (22%)	1.00
non-3A4 pathway metabolized	4 (6%)	1 (7%)	1.00	2 (6%)	1 (11%)	0.49

Data are presented as the mean value ± SD or number (%).