

# Overview of searches for dark matter at the LHC

**Vasiliki A Mitsou**

Instituto de Física Corpuscular (IFIC), CSIC – Universitat de València,  
Parc Científic de la U.V., C/ Catedrático José Beltrán 2,  
E-46980 Paterna (Valencia), Spain

E-mail: [vasiliki.mitsou@ific.uv.es](mailto:vasiliki.mitsou@ific.uv.es)

**Abstract.** Dark matter remains one of the most puzzling mysteries in Fundamental Physics of our times. Experiments at high-energy physics colliders are expected to shed light to its nature and determine its properties. This review talk focuses on recent searches for dark-matter signatures at the Large Hadron Collider, either within specific theoretical scenarios, such as supersymmetry, or in a model-independent scheme looking for mono- $X$  events arising in WIMP-pair production.

## 1. Introduction

Both Astroparticle and Particle Physics pursue the exploration of the nature of dark matter (DM) [1]. Among the long list of well-motivated candidates, the most popular particles are *cold* and weakly interacting, typically predicting missing-energy signals at particle colliders. Supersymmetry [2] and models with extra dimensions [3] are theoretical ideas that inherently provide such a dark matter candidate. High-energy colliders, such as the Large Hadron Collider [4] at CERN, are ideal machines for producing and eventually detecting DM [5].

The structure of this paper is as follows. Section 1 provides a brief introduction to the relevance of colliders, and in particular the LHC experiments ATLAS and CMS, for the production of dark matter. In Section 2, the strategy, methods, and results of the LHC experiments as far as model-independent DM-production is concerned are discussed. In Section 3, the latest results in searches for supersymmetry at the LHC are presented. The paper concludes with a summary and an outlook in Section 4.

### 1.1. Dark matter and colliders

The nature of the dark sector of the Universe constitutes one of the major mysteries in fundamental physics. According to recent observations from anisotropies of the cosmic microwave background made by the Planck mission team [6], most of our Universe energy budget consists of unknown entities:  $\sim 26.8\%$  is dark matter and  $\sim 68.3\%$  is dark energy, a form of ground-state energy. Dark energy is believed to be responsible for the current-era acceleration of the Universe. Dark matter, on the other hand, is matter inferred to exist from gravitational effects on visible matter, being undetectable by emitted or scattered electromagnetic radiation.

Evidence from the formation of large-scale structure (galaxies and their clusters) strongly favour cosmologies where non-baryonic DM is entirely composed of cold dark matter (CDM), i.e. non-relativistic particles. CDM particles, in turn, may be weakly interacting massive particles (WIMPs), a class of DM candidates that arise naturally in models which attempt to explain the

origin of electroweak symmetry breaking. Furthermore, the typical (weak-scale) cross sections characterising these models are of the same order of magnitude as the WIMP annihilation cross section, thus establishing the so-called *WIMP miracle*; this is precisely where the connection between Cosmology and Particle Physics lies [7].

WIMP dark matter candidates include the lightest neutralino in models with weak-scale supersymmetry [2], Kaluza-Klein photons arise in scenarios with universal extra dimensions (UED) [3], while lightest  $T$ -odd particles are predicted in Little Higgs models [8] with a conserved  $T$ -parity. The common denominator in these theories is that they all predict the existence of an electrically neutral, colorless and *stable* particle, whose decay is prevented by a kind of symmetry:  $R$ -parity, connected to baryon and lepton number conservation in SUSY models; KK-parity, the four-dimensional remnant of momentum conservation in extra dimension scenarios; and a  $Z_2$  discrete symmetry called  $T$ -parity in Little Higgs models.

Weakly interacting massive particles do not interact neither electromagnetically nor strongly with matter and thus, once produced, they traverse the various detectors layers without leaving a trace, just like neutrinos. However by exploiting the hermeticity of the experiments, we can get a hint of the WIMP presence through the balance of the energy/momentum measured in the various detector components, the so-called *missing energy*. In hadron colliders, in particular, since the longitudinal momenta of the colliding partons are unknown, only the *transverse missing energy*,  $E_T^{\text{miss}}$ , can be reliably used to ‘detect’ DM particles.

### 1.2. The ATLAS and CMS experiments at the LHC

The Large Hadron Collider (LHC) [4], situated at CERN, the European Laboratory for Particle Physics, outside Geneva, Switzerland, started its physics program in 2010 colliding two counter-rotating beams of protons or heavy ions. Before the scheduled 2013–2014 long shutdown, the LHC succeeded in delivering  $\sim 5 \text{ fb}^{-1}$  of integrated luminosity at centre-of-mass energy of 7 TeV during 2010–2011 and another  $\sim 23 \text{ fb}^{-1}$  at  $\sqrt{s} = 8 \text{ TeV}$  in 2012. The LHC has already extended considerably the reach of its predecessor hadron machine, the Fermilab Tevatron, both in terms of instantaneous luminosity and energy, despite the fact that it has not arrived yet to its design capabilities.

The two general-purpose experiments, ATLAS (A Toroidal LHC ApparatuS) [9] and CMS (Compact Muon Solenoid) [10], have been constructed and operate with the aim of exploring a wide range of possible signals of New Physics that LHC renders accessible, on one hand, and performing precision measurements of Standard Model (SM) parameters, on the other. It is worth mentioning that the MoEDAL [11] experiment is specifically designed to explore high-ionisation signatures that may also arise in some theoretical scenarios of dark matter [12].

The ATLAS [9] and CMS [10] detectors are designed to overcome difficult experimental challenges: high radiation levels, large interaction rate and extremely small production cross sections of New Physics signals with respect to known SM processes. To this end, both experiments feature separate subsystems to measure charged particle momentum, energy deposited by electromagnetic showers from photons and electrons, energy from hadronic showers of strongly-interacting particles and muon-track momentum.

The most remarkable highlight of ATLAS and CMS operation so far is undoubtedly the discovery of a new particle [13] that so far seems to have all the features pinpointing to a SM(-like) Higgs boson [14]. The observation of this new boson has strong impact not only on our understanding of the fundamental interactions of Nature, as encoded in the SM, but on the proposed theoretical scenarios of Physics beyond the SM (BSM).

## 2. Model-independent DM production at the LHC

Collider searches for dark matter are highly complementary to direct [15, 16] and indirect [16, 17] DM detection methods. The main advantage of collider searches is that they do not suffer from

astrophysical uncertainties and that there is no lower limit to their sensitivity on DM masses. The leading generic diagrams responsible for DM production [18,19] at hadron colliders involve the pair-production of WIMPs plus the initial- or final-state radiation (ISR/FSR) of a gluon, photon or a weak gauge boson  $Z, W$ . The ISR/FSR particle is necessary to balance the two WIMPs' momentum, so that they are not produced back-to-back resulting in negligible  $E_T^{\text{miss}}$ . Therefore the search is based on selecting events high- $E_T^{\text{miss}}$  events, due to the WIMPs, and a single jet, photon or boson candidate.

The search results are interpreted in terms of a largely model-independent effective-field-theory framework, in which the interactions between a DM Dirac fermion  $\chi$  and SM fermions  $f$  are described by contact operators. Some of the possible operators are listed in Table 1. In this framework, the interaction between SM and DM particles is determined by only two parameters, namely the DM-particle mass,  $m_\chi$ , and the suppression scale,  $M_*$ , which is related to the mediator mass and to its coupling to SM and DM particles. The derived limits are independent of the theory behind the WIMP (SUSY, extra dimensions, etc), yet it *has* been assumed that other hypothetical particles are too heavy to be produced directly in  $pp$  collisions. Henceforth, Dirac DM fermions are considered, however conclusions for Majorana fermions can also be drawn, since their production cross section only differs by a factor of two.

**Table 1.** Effective interaction operators of WIMP pair production considered in the mono- $X$  analyses, following the formalism of Ref. [18].

Name	Initial state	Type	Operator
D1	$qq$	scalar	$\frac{m_q}{M_*^3} \bar{\chi} \chi \bar{q} q$
D5	$qq$	vector	$\frac{1}{M_*^2} \bar{\chi} \gamma^\mu \chi \bar{q} \gamma_\mu q$
D8	$qq$	axial-vector	$\frac{1}{M_*^2} \bar{\chi} \gamma^\mu \gamma^5 \chi \bar{q} \gamma_\mu \gamma^5 q$
D9	$qq$	tensor	$\frac{1}{M_*^2} \bar{\chi} \sigma^{\mu\nu} \chi \bar{q} \sigma_{\mu\nu} q$
D11	$gg$	scalar	$\frac{1}{4M_*^3} \bar{\chi} \chi \alpha_s (G_{\mu\nu}^s)^2$

### 2.1. Monojet searches


Event topologies with a single high- $E_T$  jet and large  $E_T^{\text{miss}}$ , hereafter referred to as *monojets*, constitute valuable probes of physics beyond the SM at the LHC. Both ATLAS [20] and CMS [21] experiments have performed searches for an excess of monojet events over SM expectations in a wide range of signatures. The analyses outlined here use the full 2011  $pp$  LHC dataset at a centre-of-mass energy of  $\sqrt{s} = 7$  TeV. The primary SM process yielding a true monojet final state is  $Z$ -boson production in association with a jet, where the  $Z$  decays to two neutrinos. Other known processes acting as background in this search are  $Z(\rightarrow \ell\ell)$ +jets —with  $\ell = e, \mu$ —,  $W$ +jets,  $t\bar{t}$  as well as single-top events. All electroweak backgrounds and multijet events passing the selections criteria, as well as non-collision backgrounds, are in most cases determined by data-driven methods. Top and diboson backgrounds are estimated solely from Monte Carlo (MC) simulation.

The monojet analyses for ATLAS and CMS are based on some general requirements: large  $E_T^{\text{miss}}$ , with thresholds typically ranging from 120 GeV to 500 GeV and a energetic jet with a variable  $p_T$  threshold higher than 110 GeV that fulfils high jet-reconstruction quality criteria. Moreover, events with at least one electron or muon or a third jet are vetoed. Back-to-back dijet events are suppressed by requiring the subleading jet not to be collinear with  $\mathbf{p}_T^{\text{miss}}$ . The

selected data are required to pass a trigger based on high  $E_T^{\text{miss}}$  (ATLAS) or large  $E_T^{\text{miss}}$  plus one high- $E_T$  jet (CMS).

The data, amounting to  $\sim 5 \text{ fb}^{-1}$ , are found to be in agreement with the SM expectations. The results are interpreted in a framework of WIMP production with the simulated WIMP-signal MC samples corresponding to various assumptions of the effective field theory, as discussed previously. In this framework, the interaction between SM and DM particles are defined by only two parameters, namely the DM-particle mass,  $m_\chi$ , and the suppression scale,  $M_*$ , which is related to the mediator mass and to its coupling to SM and DM particles.


Experimental and theoretical systematic uncertainties are considered when setting limits on the model parameters  $M_*$  and  $m_\chi$ . The experimental uncertainties on jet energy scale and resolution and on  $E_T^{\text{miss}}$  range from 1 – 20% of the WIMP event yield, depending on the  $E_T^{\text{miss}}$  and  $p_T$  thresholds and the considered interaction operator. Other experimental uncertainties include the ones associated with the trigger efficiency and the luminosity measurement. On the other hand, the parton-distribution-function set, the amount of ISR/FSR, and the factorisation and renormalisation scales assumed lead to theoretical uncertainties on the simulated WIMP signal.



**Figure 1.** ATLAS lower limits at 90% CL on the suppression scale,  $M_*$ , for different masses of  $\chi$  obtained with the monojet analysis for the operators D1 (top left), D5 (top right) and D8 (bottom). The region below the limit lines is excluded. All shown curves and areas are explained in the text. From Ref. [20].

From the limit on the visible cross section of new BSM physics processes, lower limits on the suppression scale as a function of the WIMP mass have been derived by the ATLAS Collaboration [20]. The 90% confidence level (CL) lower limits for the D1, D5 and D8 operators are shown in Fig. 1. The observed limit on  $M_*$  includes experimental uncertainties; the effect of theoretical uncertainties is indicated by dotted  $\pm 1\sigma$  lines above and below it. Around the expected limit,  $\pm 1\sigma$  variations due to statistical and systematic uncertainties are shown as a grey band. The lower limits are flat up to  $m_\chi \lesssim 100$  GeV and become weaker at higher mass due to the collision energy. In the bottom-right corner of the  $(m_\chi, M_*)$  plane (light-grey shaded area), the effective field theory approach is no longer valid. The rising lines correspond to couplings consistent with the measured thermal relic density [18], assuming annihilation in the early universe proceeded exclusively via the given operator. Similar exclusion limits for all operators listed in Table 1 are given in Ref. [20]. For the operator D1, the limits are much weaker ( $\sim 30$  GeV) than for other operators. Nevertheless, if heavy-quark loops are included in the analysis, much stronger bounds on  $M_*$  can be obtained [22].

The observed limit on the dark matter-nucleon scattering cross section depends on the mass of the dark matter particle and the nature of its interaction with the SM particles. The limits on the suppression scale as a function of  $m_\chi$  can be translated into a limit on the cross section using the reduced mass of  $\chi$ -nucleon system [19], which can be compared with the constraints from direct and indirect detection experiments, as we shall see at the end of this Section in conjunction with bounds acquired with other mono- $X$  analyses.




**Figure 2.** ATLAS 95% CL limits on WIMP annihilation rates  $\langle\sigma v\rangle$  versus mass  $m_\chi$ , inferred from the monojet analysis. Explanation of the shown curves is given in the text. From Ref. [20].

The ATLAS collider limits on vector (D5) and axial-vector (D8) interactions are also interpreted in terms of the relic abundance of WIMPs, using the same effective theory approach [18]. The upper limits on the annihilation rate of WIMPs into light quarks, defined as the product of the annihilation cross section  $\sigma$  and the relative WIMP velocity  $v$  averaged over the WIMP velocity distribution  $\langle\sigma v\rangle$ , are shown in Fig. 2. The results are compared to limits on WIMP annihilation to  $b\bar{b}$ , obtained from galactic high-energy gamma-ray observations, measured by the Fermi-LAT telescope [23]. Gamma-ray spectra and yields from WIMPs annihilating to  $b\bar{b}$ , where photons are produced in the hadronisation of the quarks, are expected to be very similar to those from WIMPs annihilating to light quarks [24]. Under this assumption, the ATLAS and Fermi-LAT limits can be compared, after scaling up the Fermi-LAT values by a factor of two to account for the Majorana-to-Dirac fermion adaptation. Again, the ATLAS bounds are especially important for small WIMP masses: below 10 GeV for vector couplings and below about 100 GeV for axial-vector ones. In this region, the ATLAS limits are below the annihilation cross section needed to be consistent with the thermic relic value, keeping the assumption that

WIMPs have annihilated to SM quarks only via the particular operator in question. For masses of  $m_\chi \gtrsim 200$  GeV the ATLAS sensitivity becomes worse than the Fermi-LAT one. In this region, improvements can be expected when going to larger centre-of-mass energies at the LHC.

The case in which the mediator is light enough to be accessible to the LHC has been considered too by the CMS experiment in a monojet search performed with  $\sim 20 \text{ fb}^{-1}$  at 8 TeV [25]. Figure 3 shows the observed limits on the contact interaction scale  $\Lambda$  as a function of the mass of the mediator  $M$ , assuming vector interactions and a dark matter mass of 50 GeV and 500 GeV. The width  $\Gamma$  of the mediator is varied between  $M/3$  and  $M/8\pi$  [26]. It shows the resonant enhancement in the production cross section once the mass of the mediator is within the kinematic range and can be produced on-shell. For  $m_\chi \gtrsim 100$  GeV, this approach is adequate and quite conservative in the bounds on  $\Lambda$ . For  $m_\chi \lesssim 100$  GeV, the collider bounds are considerably weaker. At large mediator masses, i.e.  $M \gtrsim 5$  TeV, the limits on  $\Lambda$  approximate to those obtained in the effective theory framework.




**Figure 3.** Observed limits on the scale  $\Lambda$  as a function of the mass  $M$ , assuming vector interactions and a dark matter mass of 50 GeV (blue) and 500 GeV (red) in a CMS monojet analysis. The width of the mediator was varied between  $M/3$ ,  $M/10$  and  $M/8\pi$ . From Ref. [25].

## 2.2. Monophoton-based probes

In the same fashion as in the monojet searches, the *monophoton* analyses aim at probing dark matter requiring large  $E_T^{\text{miss}}$  —from the  $\chi$ -pair production— and at least one ISR/FSR photon. Searches in monophoton events by ATLAS [27] and CMS [28] also show an agreement with the SM expectations. The limits are derived in a similar fashion as for the monojet search, however the monophoton search is found to be somewhat less sensitive with respect to the monojet topology.

The primary (irreducible) background for a  $\gamma + E_T^{\text{miss}}$  signal comes from  $Z\gamma \rightarrow \nu\bar{\nu}\gamma$  production. This together with other SM backgrounds, including  $W\gamma$ ,  $W \rightarrow e\nu$ ,  $\gamma + \text{jet}$  multijet, diphoton and diboson events, as well as backgrounds from beam halo and cosmic-ray muons, are taken into account in the analyses. The CMS analysis is based on single-photon triggers, whilst ATLAS relies on high- $E_T^{\text{miss}}$  triggered events. The photon candidate is required to pass tight quality and isolation criteria, in particular in order to reject events with electrons faking photons. The missing transverse momentum of the selected events should be as high as 150 GeV (130 GeV) in the ATLAS (CMS) search. In CMS, events with a reconstructed jet are vetoed, while the ATLAS analysis rejects events with an electron, a muon or a second jet.

Both analyses, observe no significant excess of events over the expected background when applied on  $\sim 5 \text{ fb}^{-1}$  of  $pp$  collision data at  $\sqrt{s} = 7 \text{ TeV}$ . Hence they set lower limits on the suppression scale,  $M_*$  versus the DM fermion mass,  $m_\chi$ , which in turn they are translated into upper limits on the nucleon-WIMP interaction cross section applying the prescription in Ref. [18]. Figure 4 shows the 90% CL upper limits on the nucleon-WIMP cross section as a function of  $m_\chi$  derived from the ATLAS search [27]. The results are compared with previous CDF [29], CMS [21, 28] and direct WIMP detection experiments [30–34] results. The CMS limit curve generally overlaps the ATLAS curve.



**Figure 4.** ATLAS 90% CL upper limits on the nucleon-WIMP cross section as a function of  $m_\chi$  for spin-dependent (left) and spin-independent (right) interactions, corresponding to D8, D9, D1, and D5 operators, derived from the monophoton analysis. Explanation of the shown curves is given in the text. From Ref. [27].


The observed limits on  $M_*$  typically decrease by 2% to 10% if the  $-1\sigma$  theoretical uncertainty, resulting from the same sources as the one cited in the monojet analysis, is considered. This translates into a 10% to 50% increase of the quoted nucleon-WIMP cross section limits. To recapitulate, the exclusion in the region  $1 \text{ GeV} < m_\chi < 1 \text{ TeV}$  ( $1 \text{ GeV} < m_\chi < 3.5 \text{ GeV}$ ) for spin-dependent (spin-independent) nucleon-WIMP interactions is driven by the results from collider experiments, always under the assumption of the validity of the effective theory, and is still dominated by the monojet results.

### 2.3. Mono- $W$ and mono- $Z$ final states

As demonstrated in the previous sections, searches for monojet or monophoton signatures have yielded powerful constraints on dark matter interactions with SM particles. Other studies propose probing DM at LHC through a  $pp \rightarrow \chi\bar{\chi} + W/Z$ , with a leptonically decaying  $W$  [35] or  $Z$  [36]. The final state in this case would be large  $E_{\text{T}}^{\text{miss}}$  and a single charged lepton (electron or muon) for the *mono- $W$*  signature (*monolepton*) or a pair of charged leptons that reconstruct to the  $Z$  mass for the *mono- $Z$*  signature. In either case, the gauge boson radiations off a  $q\bar{q}$  initial

state and an effective field theory is deployed to describe the contact interactions that couple the SM particle with the WIMP.

In Ref. [37], the existing  $W'$  searches from CMS [38] —which share a similar final state with mono- $W$  searches— are used to place a bound on mono- $W$  production at LHC, which for some choices of couplings are better than monojet bounds. This is illustrated in the left (right) panel of Fig. 5, where the spin-dependent (spin-independent) WIMP-proton cross section limits are drawn. The parameter  $\xi$  parametrises the relative strength of the coupling to down-quarks with respect to up-quarks:  $\xi = +1$  for equal couplings;  $\xi = -1$  for opposite-sign ones; and  $\xi = 0$  when there is no coupling to down-quarks. Even in cases where the monoleptons do not provide the most stringent constraints, they provide an interesting mechanism to disentangle WIMP couplings to up-type versus down-type quarks. Such an interpretation has also been performed in Ref. [35] yielding similar limits.




**Figure 5.** CMS monolepton search with  $20 \text{ fb}^{-1}$  at 8 TeV. Excluded proton-dark matter cross section for axial-vector-like, i.e. spin dependent (left), and vector-like, i.e. spin independent (right), for the combination of electron and muon channels. The CMS monojet result is for  $20 \text{ fb}^{-1}$  of 2012 data [25]. From Ref. [37].

The ATLAS Collaboration has extended the range of possible mono- $X$  probes by looking for  $pp \rightarrow \chi\bar{\chi} + W/Z$ , when the gauge boson decays to two quarks [39], as opposed to the leptonic signatures discussed so far. The analysis searches for the production of  $W$  or  $Z$  bosons decaying hadronically and reconstructed as a single massive jet in association with large  $E_{\text{T}}^{\text{miss}}$  from the undetected  $\chi\bar{\chi}$  particles. For this analysis, the jet candidates are reconstructed using a filtering procedure referred to as *large-radius jets* [40]. This search, the first of its kind, is sensitive to WIMP pair production, as well as to other DM-related models, such as invisible Higgs boson decays ( $WH$  or  $ZH$  production with  $H \rightarrow \chi\bar{\chi}$ ).

Figure 6 shows the 90% CL upper limits on the dark matter-nucleon scattering cross section as a function of the mass of DM particle for the spin-independent (left) and spin-dependent (right) models obtained by the ATLAS mono- $W/Z$  analysis [39]. The new limits are also compared to the limits set by ATLAS in the 7 TeV monojet analysis [20]. Limits from XENON100 [41], CoGent [31], CDMS II [32], SIMPLE [33], COUPP [42], Picasso [34], IceCube [43] are superimposed for comparison. For the spin-independent case with the opposite-sign up-type and





**Figure 6.** ATLAS-derived limits on  $\chi$ -nucleon cross sections as a function of  $m_\chi$  at 90% CL for spin-independent (left) and spin-dependent (right) cases, obtained with the mono- $W/Z$  analysis and compared to previous limits. From Ref. [39].

down-type couplings, the limits are improved by about three orders of magnitude. For other cases, the bounds are similar. Comparable limits have been obtained by the CMS experiment.

It is worth noting that the spin-dependent limits derived from the operator D9, give a smaller, hence better, bound on the WIMP-nucleon cross section throughout the range of  $m_\chi$ , compared to direct DM experiments. In the spin-independent case the bounds from direct detection experiments are stronger for  $m_\chi \gtrsim 10$  GeV, whereas the collider bounds, acquired with the operator D5, get important for the region of low DM masses.

### 3. Searches for supersymmetry

Supersymmetry (SUSY) [44] is an extension of the Standard Model which assigns to each SM field a superpartner field with a spin differing by a half unit. SUSY provides elegant solutions to several open issues in the SM, such as the hierarchy problem and the grand unification. In particular, SUSY predicts the existence of a stable weakly interacting particle—the lightest supersymmetric particle (LSP)—that has the pertinent properties to be a dark matter particle, thus providing a very compelling argument in favour of SUSY [45].

SUSY searches in the ATLAS [9] and CMS [10] experiments typically focus on events with high transverse missing energy, which can arise from (weakly interacting) LSPs, in the case of  $R$ -parity conserving SUSY, or from neutrinos produced in LSP decays, if  $R$ -parity is broken (c.f. Section 3.4). Hence, the event selection criteria of inclusive channels are based on large  $E_T^{\text{miss}}$ , no or few leptons ( $e$ ,  $\mu$ ), many jets and/or  $b$ -jets,  $\tau$ -leptons and photons. In addition, kinematical variables such as the transverse mass,  $M_T$ , and the effective mass,  $M_{\text{eff}}$ , assist in distinguishing further SUSY from SM events, whilst the *effective transverse energy* [46] can be useful to cross-check results, allowing a better and more robust identification of the SUSY mass scale, should a positive signal is found. Although the majority of the analysis simply look for an excess of events over the SM background, there is an increasing application of distribution

shape fitting techniques [47].

Typical SM backgrounds are top-quark production —including single-top—,  $W/Z$  in association with jets, dibosons and QCD multijet events. These are estimated using semi- or fully data-driven techniques. Although the various analyses are optimised for a specific SUSY scenario, the interpretation of the results are extended to various SUSY models or topologies.

Analyses exploring  $R$ -parity conserving SUSY models at LHC are roughly divided into inclusive searches for squarks and gluinos, for third-generation fermions, and for electroweak production (pairs of  $\tilde{\chi}^0$ ,  $\tilde{\chi}^\pm$ ,  $\tilde{\ell}$ ). Although these searches are designed and optimised to look for  $R$ -parity conserving SUSY, interpretation in terms of  $R$ -parity violating (RPV) models is also possible. Other analyses are purely motivated by oriented by RPV scenarios and/or by the expectation of long-lived sparticles. Recent summary results from each category of ATLAS and CMS searches are presented in this section.


### 3.1. Gluinos and first two generations of quarks

At the LHC, supersymmetric particles are expected to be predominantly produced hadronically, i.e. through gluino-pair, squark-pair and squark-gluino production. Each of these (heavy) sparticles is going to decay into lighter ones in a cascade decay that finally leads to an LSP, which in most of the scenarios considered is the lightest neutralino  $\tilde{\chi}_1^0$ . The two LSPs would escape detection giving rise to high transverse missing energy, hence the search strategy followed is based on the detection of high  $E_T^{\text{miss}}$ , many jets and possibly energetic leptons. The analyses make extensive use of data-driven Standard Model background measurements.

The most powerful of the existing searches are based on all-hadronic final states with large missing transverse momentum [48, 49]. In the 0-lepton search, events are selected based on a jet+ $E_T^{\text{miss}}$  trigger, applying a lepton veto, requiring a minimum number of jets, high  $E_T^{\text{miss}}$ , and large azimuthal separation between the  $E_T^{\text{miss}}$  and reconstructed jets, in order to reject multijet background. In addition, searches for squark and gluino production in a final state with one or two leptons have been performed [50, 51]. The events are categorised by whether the leptons have higher or lower momentum and are referred to as the *hard* and *soft* lepton channels respectively. The soft-lepton analysis which enhances the sensitivity of the search in the difficult kinematic region where the neutralino and gluino masses are close to each other forming the so-called *compressed spectrum*. [52] Leptons in the soft category are characterised by low lepton- $p_T$  thresholds (6 – 10 GeV) and such events are triggered by sufficient  $E_T^{\text{miss}}$ . Hard leptons pass a threshold of  $\sim 25$  GeV and are seeded with both lepton and  $E_T^{\text{miss}}$  triggers. Analyses based on the *razor* [53] variable have also been carried out by both experiments [54, 55].

The major backgrounds ( $t\bar{t}$ ,  $W$ +jets,  $Z$ +jets) are estimated by isolating each of them in a dedicated control region, normalising the simulation to data in that control region, and then using the simulation to extrapolate the background expectations into the signal region. The multijet background is determined from the data by a matrix method. All other (smaller) backgrounds are estimated entirely from the simulation, using the most accurate theoretical cross sections available. To account for the cross-contamination of physics processes across control regions, the final estimate of the background is obtained with a simultaneous, combined fit to all control regions.

In the absence of deviations from SM predictions, limits for squark and gluino production are set. Figure 7 illustrates the 95% CL limits set by ATLAS under the minimal Supergravity (mSUGRA) model in the  $(m_0, m_{1/2})$  plane [48, 56]. The remaining parameters are set to  $\tan\beta = 30$ ,  $A_0 = -2m_0$ ,  $\mu > 0$ , so as to acquire parameter-space points where the predicted mass of the lightest Higgs boson,  $h^0$ , is near 125 GeV, i.e. compatible with the recently observed Higgs-like boson [13, 14]. Exclusion limits are obtained by using the signal region with the best expected sensitivity at each point. By assumption, the mSUGRA model avoids both flavour-changing neutral currents and extra sources of  $CP$  violation. For masses in the TeV range, it



**Figure 7.** Exclusion limits at 95% CL for 8 TeV ATLAS analyses [48, 56] in the  $(m_0, m_{1/2})$  plane for the mSUGRA model. From Ref. [57].

typically predicts too much cold dark matter, however these predictions depend of the presence of stringy effects that may dilute [58] or enhance [59] the predicted relic dark matter density. In the mSUGRA case, the limit on squark mass reaches 1750 GeV and on gluino mass is 1400 GeV.


### 3.2. Third-generation squarks

The previously presented limits from inclusive channels indicate that the masses of gluinos and first/second generation squarks are expected to be above 1 TeV. Nevertheless, in order to solve the hierarchy problem in a *natural* way, the masses of the stops, sbottoms, higgsinos and gluinos have to be below the TeV-scale to properly cancel the divergences in the Higgs mass radiative corrections. Despite their production cross sections being smaller than for the first and second generation squarks, stop and sbottom may well be directly produced at the LHC and could provide the only direct observation of SUSY at the LHC in case the other sparticles are outside of the LHC energy reach. The lightest mass eigenstates of the sbottom and stop particles,  $\tilde{b}_1$  and  $\tilde{t}_1$ , could hence be produced either directly in pairs or through  $\tilde{g}$  pair production followed by  $\tilde{g} \rightarrow \tilde{b}_1 b$  or  $\tilde{g} \rightarrow \tilde{t}_1 t$  decays. Both cases will be discussed in the following.


For the aforementioned reasons, direct searches for third generation squarks have become a priority in both ATLAS and CMS. Such events are characterised by several energetic jets (some of them  $b$ -jets), possibly accompanied by light leptons, as well as high  $E_T^{\text{miss}}$ . A suite of channels have been considered, depending on the topologies allowed and the exclusions generally come with some assumptions driven by the shortcomings of the techniques and variables used, such as the requirement of 100% branching ratios into particular decay modes.

In the case of the gluino-mediated production of stops, a simplified scenario (“Gtt model”), where  $\tilde{t}_1$  is the lightest squark but  $m_{\tilde{g}} < m_{\tilde{t}_1}$ , has been considered. Pair production of gluinos is the only process taken into account since the mass of all other sparticles apart from the  $\tilde{\chi}_1^0$  are above the TeV scale. A three-body decay via off-shell stop is assumed for the gluino, yielding a

100% branching ratio for the decay  $\tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ . The stop mass has no impact on the kinematics of the decay and the exclusion limits [54, 60, 61] set by the CMS experiment are presented in the  $(m_{\tilde{g}}, m_{\tilde{\chi}_1^0})$  plane in Fig. 8. For a massless LSP, gluinos with masses from 560 GeV to 1320 GeV are excluded. Similar results are obtained if the decay  $\tilde{g} \rightarrow b\bar{b}\tilde{\chi}_1^0$  is considered instead, as shown in Fig. 9.




**Figure 8.** Summary of observed and expected limits [54, 60, 61] for gluino pair production with gluino decaying via a 3-body decay to a top, an anti-top and a neutralino. From Ref. [62].



**Figure 9.** Summary of observed and expected limits [54, 60, 61] for gluino pair production with gluino decaying via a 3-body decay to a bottom, an anti-bottom and a neutralino. From Ref. [62].

If the gluino is too heavy to be produced at the LHC, the only remaining possibility is the direct  $\tilde{t}_1\tilde{t}_1$  and  $\tilde{b}_1\tilde{b}_1$  production. If stop pairs are considered, two decay channels can be distinguished depending on the mass of the stop:  $\tilde{t}_1 \rightarrow b\tilde{\chi}_1^\pm$  and  $\tilde{t}_1 \rightarrow t\tilde{\chi}_1^0$ . CMS and ATLAS carried out a wide range of different analyses in each of these modes at both 7 TeV and 8 TeV centre-of-mass energy. In all these searches, the number of observed events has been found to be consistent with the SM expectation. Limits have been set by ATLAS on the mass of the scalar

top for different assumptions on the mass hierarchy scalar top-chargino-lightest neutralino [63], as shown in the left panel of Fig. 10. A scalar top quark of mass of up to 480 GeV is excluded at 95% CL for a massless neutralino and a 150 GeV chargino. For a 300 GeV scalar top quark and a 290 GeV chargino, models with a neutralino with mass lower than 175 GeV are excluded at 95% CL.




**Figure 10.** Summary of the dedicated ATLAS searches [63] for stop pair production based on  $20 - 21 \text{ fb}^{-1}$  of  $pp$  collision data taken at  $\sqrt{s} = 8 \text{ TeV}$ , and  $4.7 \text{ fb}^{-1}$  of  $pp$  collision data taken at  $\sqrt{s} = 7 \text{ TeV}$ . Exclusion limits at 95% CL are shown in the  $(\tilde{t}_1, \tilde{\chi}_1^0)$  mass plane for channels targeting  $\tilde{t}_1 \rightarrow b \tilde{\chi}_1^\pm, \tilde{\chi}_1^\pm \rightarrow W^\pm \tilde{\chi}_1^0$  decays (left) and  $\tilde{t}_1$  decays to  $t \tilde{\chi}_1^0$  or  $W b \tilde{\chi}_1^0$  or  $c \tilde{\chi}_1^0$  (right). The dashed and solid lines show the expected and observed limits, respectively, including all uncertainties except the theoretical signal cross section uncertainty. From Ref. [57].

For the case of a high-mass stop decaying to a top and neutralino ( $\tilde{t}_1 \rightarrow t \tilde{\chi}_1^0$ ), analyses requiring one, two or three isolated leptons, jets and large  $E_T^{\text{miss}}$  have been carried out. No significant excess of events above the rate predicted by the SM is observed and 95% CL upper limits are set on the stop mass in the stop-neutralino mass plane. The region of excluded stop and neutralino masses is shown on the right panel of Fig. 10 for the ATLAS analyses. Stop masses are excluded between 200 GeV and 680 GeV for massless neutralinos, and stop masses around 500 GeV are excluded along a line which approximately corresponds to neutralino masses up to 250 GeV. It is worth noting that a monojet analysis with  $c$ -tagging is deployed to cover part of the low- $m_{\tilde{t}_1}$ , low- $m_{\tilde{\chi}_1^0}$  region through the  $\tilde{t}_1 \rightarrow c \tilde{\chi}_1^0$  channel.

### 3.3. Electroweak gaugino production

If all squarks and gluinos are above the TeV scale, weak gauginos with masses of few hundred gigaelectronvolts may be the only sparticles accessible at the LHC. As an example, at  $\sqrt{s} = 7 \text{ TeV}$ , the cross-section of the associated production  $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$  with degenerate masses of 200 GeV is above the 1-TeV gluino-gluino production cross section by one order of magnitude. Chargino pair production is searched for in events with two opposite-sign leptons and  $E_T^{\text{miss}}$  using a jet veto, through the decay  $\tilde{\chi}_1^\pm \rightarrow \ell^\pm \nu \tilde{\chi}_1^0$ . A summary of related analyses [64] performed by CMS is shown in Fig. 11. Charginos with masses between 140 and 560 GeV are excluded for a massless LSP in the chargino-pair production with an intermediate slepton/sneutrino between the  $\tilde{\chi}_1^\pm$

and the  $\tilde{\chi}_1^0$ . If  $\tilde{\chi}_1^\pm \tilde{\chi}_2^0$  production is assumed instead, the limits range from 11 to 760 GeV. The corresponding limits involving intermediate  $W$ ,  $Z$  and/or  $H$  are significantly weaker.



**Figure 11.** Summary of observed limits for electroweak-gaugino production from CMS [64]. From Ref. [62].


### 3.4. $R$ -parity violating SUSY and meta-stable sparticles

$R$ -parity is defined as:  $R = (-1)^{3(B-L)+2S}$ , where  $B$ ,  $L$  and  $S$  are the baryon number, lepton number and spin, respectively. Hence  $R = +1$  for all Standard Model particles and  $R = -1$  for all SUSY particles. It is stressed that the conservation of  $R$ -parity is an *ad-hoc* assumption. The only firm restriction comes from the proton lifetime: non-conservation of both  $B$  and  $L$  leads to rapid proton decay.  $R$ -parity conservation has serious consequences in SUSY phenomenology in colliders: the SUSY particles are produced in pairs and the lightest SUSY particle is absolutely stable, thus providing a WIMP candidate. Here we highlight the status of RPV supersymmetry [65] searches at the LHC.

Both ATLAS and CMS experiments have probed RPV SUSY through various channels, either by exclusively searching for specific decay chains, or by inclusively searching for multilepton events. ATLAS has looked for resonant production of  $e\mu$ ,  $e\tau$  and  $\mu\tau$  [66], for multijets [67], for events with at least four leptons [68] and for excesses in the  $e\mu$  continuum [69]. Null inclusive searches in the one-lepton channel [70] have also been interpreted in the context of a model where RPV is induced through bilinear terms [71].

Recent CMS analyses are focused on studying the lepton number violating terms  $\lambda_{ijk} L_i L_j \bar{e}_k$  and  $\lambda'_{ijk} L_i Q_j \bar{d}_k$ , which result in specific signatures involving leptons in events produced in  $pp$  collisions at LHC. A search for resonant production and the following decay of  $\tilde{\mu}$  which is caused by  $\lambda'_{211} \neq 0$  has been conducted [72]. Multilepton signatures caused by LSP decays due to various  $\lambda$  and  $\lambda'$  terms in stop production have been probed [73]. Reference [74] discusses the possibility of the generic model independent search for RPV SUSY in 4-lepton events. A summary of the limits set by several CMS analyses [61, 73, 75] are listed in Fig. 12.

In view of the null results in other SUSY searches, it became mandatory to fully explore the SUSY scenario predicting meta-stable or long-lived particles. These particles, not present in the Standard Model, would provide striking signatures in the detector and rely heavily on a detailed understanding of its performance. In SUSY, non-prompt particle decay can be caused by (i) very weak RPV [76], (ii) low mass difference between a SUSY particle and the LSP [77], or (iii) very weak coupling to the gravitino in GMSB models [78, 79]. A small part of these



**Figure 12.** Best exclusion limits for the masses of the mother particles, for RPV scenarios, for each topology, for all CMS results [61, 73, 75]. In this plot, the lowest mass range is  $m_{\text{mother}} = 0$ , but results are available starting from a certain mass depending on the analyses and topologies. Branching ratios of 100% are assumed, values shown in plot are to be interpreted as upper bounds on the mass limits. From Ref. [62].

possibilities have been explored by the ATLAS [57] and CMS [62] experiments covering specific cases, difficult to summarise here. There is still a wide panorama of signatures to be explored, in view of various proposed SUSY scenarios pointing towards this direction.

As a last remark, we address the issue of (not necessarily cold) dark matter in RPV SUSY models. These seemingly incompatible concepts *can* be reconciled in models with a gravitino [80, 81] or an axino [82, 83] LSP with a lifetime exceeding the age of the Universe. In both cases, RPV is induced by bilinear terms in the superpotential that can also explain current data on neutrino masses and mixings without invoking any GUT-scale physics [71]. Decays of the next-to-lightest superparticle occur rapidly via RPV interaction, and thus they do not upset the Big-Bang nucleosynthesis, unlike the  $R$ -parity conserving case. Such gravitino DM is proposed in the context of  $\mu\nu$ SSM [84] with profound prospects for detecting  $\gamma$  rays from their decay [85].

Recent evidence on the four-year Fermi data that have found excess of a 130 GeV gamma-ray line from the Galactic Center [86] have been studied in the framework of  $R$ -parity breaking SUSY. A decaying axino DM scenario based on the SUSY KSVZ axion model with the bilinear  $R$ -parity violation explains the Fermi 130 GeV gamma-ray line excess from the GC while satisfying other cosmological constraints [83]. On the other hand, gravitino dark matter with trilinear RPV—in particular models with the  $LLE$  RPV coupling—can account for the gamma-ray line, since there is no overproduction of anti-proton flux, while being consistent with big-bang nucleosynthesis and thermal leptogenesis [81].

#### 4. Summary and outlook

The nature of dark matter remains one of the mysteries of Particle Physics and Cosmology. Mono- $X$  searches at the LHC provide strong constraints on dark matter properties in an effective field theory formalism. Colliders are superior to direct searches if dark matter is very light ( $< 10$  GeV) or if interactions are spin-dependent. Extensive efforts are currently in progress on the validity of the effective field-theory approach, the proper comparison with the results from direct detection experiments and the use of simplified models with light mediators. Analyses looking for specific models providing DM candidates, such as Supersymmetry, are ongoing. Searches continue with the full 2012 dataset but a new discovery might eventually require more energy and more data coming up in 2015.

#### Acknowledgments

The author is grateful to the XIV MWPF organisers for the kind invitation and support that gave her the opportunity to present this plenary talk. She acknowledges support by the Spanish Ministry of Economy and Competitiveness (MINECO) under the projects FPA2009-13234-C04-01 and FPA2012-39055-C02-01, by the Generalitat Valenciana through the project PROMETEO II/2013-017 and by the Spanish National Research Council (CSIC) under the JAE-Doc program co-funded by the European Social Fund (ESF).

#### References

- [1] For a pedagogical introduction, see e.g.: Profumo S 2013 TASI 2012 Lectures on Astrophysical Probes of Dark Matter *Preprint* arXiv:1301.0952 [hep-ph]
- [2] Ellis J R, Falk T, Ganis G and Olive K A 2000 Supersymmetric dark matter in the light of LEP and the Tevatron collider *Phys. Rev. D* **62** 075010 (*Preprint* hep-ph/0004169)  
Drees M, Kim Y G, Nojiri M M, Toya D, Hasuko K and Kobayashi T 2001 Scrutinizing LSP dark matter at the CERN LHC *Phys. Rev. D* **63** 035008 (*Preprint* hep-ph/0007202)  
Chattopadhyay U and Nath P 2001 Upper limits on sparticle masses from  $g - 2$  and the possibility for discovery of SUSY at colliders and in dark matter searches *Phys. Rev. Lett.* **86** 5854 (*Preprint* hep-ph/0102157)  
Lahanas A B, Mavromatos N E and Nanopoulos D V 2003 WMAPing the universe: Supersymmetry, dark matter, dark energy, proton decay and collider physics *Int. J. Mod. Phys. D* **12** 1529 (*Preprint* hep-ph/0308251)  
Baer H, Belyaev A, Krupovnickas T and O’Farrill J 2004 Indirect, direct and collider detection of neutralino dark matter *JCAP* **0408** 005 (*Preprint* hep-ph/0405210)  
Allanach B C, Belanger G, Boudjema F and Pukhov A 2004 Requirements on collider data to match the precision of WMAP on supersymmetric dark matter *JHEP* **0412** 020 (*Preprint* hep-ph/0410091)  
Roszkowski L, Ruiz de Austri R and Choi K-Y 2005 Gravitino dark matter in the CMSSM and implications for leptogenesis and the LHC *JHEP* **0508** 080 (*Preprint* hep-ph/0408227)  
Baer H, Mustafayev A, Profumo S, Belyaev A and Tata X 2005 Direct, indirect and collider detection of neutralino dark matter in SUSY models with non-universal Higgs masses *JHEP* **0507** 065 (*Preprint* hep-ph/0504001)  
Arbey A, Battaglia M and Mahmoudi F 2012 Light Neutralino Dark Matter in the pMSSM: Implications of LEP, LHC and Dark Matter Searches on SUSY Particle Spectra *Eur. Phys. J. C* **72** 2169 (*Preprint* arXiv:1205.2557 [hep-ph])
- [3] Hooper D and Profumo S 2007 Dark matter and collider phenomenology of universal extra dimensions *Phys. Rept.* **453** 29 (*Preprint* hep-ph/0701197)
- [4] Evans L and Bryant P 2008 LHC Machine *JINST* **3** S08001
- [5] Mitsou V A 2013 Shedding Light on Dark Matter at Colliders *Int. J. Mod. Phys. A* **28** 1330052 (*Preprint* arXiv:1310.1072 [hep-ex])
- [6] Ade P A R *et al.* [Planck Collab.] 2013 Planck 2013 results. XVI. Cosmological parameters *Preprint* arXiv:1303.5076 [astro-ph.CO]
- [7] Mavromatos N E 2012 Some Aspects of String Cosmology and the LHC *Preprint* arXiv:1210.0211 [hep-ph] in *Proc. 1<sup>st</sup> Int. Conf. on New Frontiers in Physics*, to appear in *Eur. Phys. J. Web Conf.*
- [8] Birkedal A, Noble A, Perelstein M and Spray A 2006 Little Higgs dark matter *Phys. Rev. D* **74** 035002 (*Preprint* arXiv:hep-ph/0603077)



- [9] Aad G *et al.* [ATLAS Collab.] 2008 The ATLAS Experiment at the CERN Large Hadron Collider *JINST* **3** S08003
- [10] Adolphi R *et al.* [CMS Collab.] 2008 The CMS experiment at the CERN LHC *JINST* **3** S08004
- [11] MoEDAL Collaboration 2009 Technical Design Report of the MoEDAL Experiment *CERN Preprint* CERN-LHC-2009-006, MoEDAL-TDR-1.1
- [12] Acharya B *et al.* 2014 The Physics Program of the MoEDAL Experiment at the LHC, to appear in *Int. J. Mod. Phys. A*
- [13] Aad G *et al.* [ATLAS Collab.] 2012 Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC *Phys. Lett. B* **716** 1 (*Preprint* arXiv:1207.7214 [hep-ex])  
Chatrchyan S *et al.* [CMS Collab.] 2012 Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC *Phys. Lett. B* **716** 30 (*Preprint* arXiv:1207.7235 [hep-ex])
- [14] Chatrchyan S *et al.* [CMS Collab.] 2013 Study of the Mass and Spin-Parity of the Higgs Boson Candidate Via Its Decays to Z Boson Pairs *Phys. Rev. Lett.* **110** 081803 (*Preprint* arXiv:1212.6639 [hep-ex])  
Aad G *et al.* [ATLAS Collab.] 2013 Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC *Phys. Lett. B* **726** 88 (*Preprint* arXiv:1307.1427 [hep-ex])  
Aad G *et al.* [ATLAS Collab.] 2013 Evidence for the spin-0 nature of the Higgs boson using ATLAS data *Phys. Lett. B* **726** 120 (*Preprint* arXiv:1307.1432 [hep-ex])
- [15] Baer H and Brhlik M 1998 Neutralino dark matter in minimal supergravity: Direct detection versus collider searches *Phys. Rev. D* **57** 567 (*Preprint* hep-ph/9706509)  
Pato M, Baudis L, Bertone G, Ruiz de Austri R, Strigari L E and Trotta R 2011 Complementarity of Dark Matter Direct Detection Targets *Phys. Rev. D* **83** 083505 (*Preprint* arXiv:1012.3458 [astro-ph.CO])
- [16] Rajaraman A, Shepherd W, Tait T M P and Wijangco A M 2011 LHC Bounds on Interactions of Dark Matter *Phys. Rev. D* **84** 095013 (*Preprint* arXiv:1108.1196 [hep-ph])  
Cheung K, Tseng P-Y, Tsai Y-L S and Yuan T-C 2012 Global Constraints on Effective Dark Matter Interactions: Relic Density, Direct Detection, Indirect Detection, and Collider *JCAP* **1205** 001 (*Preprint* arXiv:1201.3402 [hep-ph])  
Bauer D *et al.* 2013 Dark Matter in the Coming Decade: Complementary Paths to Discovery and Beyond *Preprint* arXiv:1305.1605 [hep-ph]
- [17] Bertone G, Cerdeno D G, Fornasa M, Pieri L, Ruiz de Austri R and Trotta R 2012 Complementarity of Indirect and Accelerator Dark Matter Searches *Phys. Rev. D* **85** 055014 (*Preprint* arXiv:1111.2607 [astro-ph.HE])
- [18] Goodman J, Ibe M, Rajaraman A, Shepherd W, Tait T M P and Yu H-B 2010 Constraints on Dark Matter from Colliders *Phys. Rev. D* **82** 116010 (*Preprint* arXiv:1008.1783 [hep-ph])
- [19] Bai Y, Fox P J and Harnik R 2010 The Tevatron at the Frontier of Dark Matter Direct Detection *JHEP* **1012** 048 (*Preprint* arXiv:1005.3797 [hep-ph])  
Beltran M, Hooper D, Kolb E W, Krusberg Z A C and Tait T M P 2010 Maverick dark matter at colliders *JHEP* **1009** 037 (*Preprint* arXiv:1002.4137 [hep-ph])
- [20] Aad G *et al.* [ATLAS Collab.] 2013 Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector *JHEP* **1304** 075 (*Preprint* arXiv:1210.4491 [hep-ex])
- [21] Chatrchyan S *et al.* [CMS Collab.] 2012 Search for dark matter and large extra dimensions in monojet events in  $pp$  collisions at  $\sqrt{s} = 7$  TeV *JHEP* **1209** 094 (*Preprint* arXiv:1206.5663 [hep-ex])
- [22] Haisch U, Kahlhoefer F and Unwin J 2013 The impact of heavy-quark loops on LHC dark matter searches *JHEP* **1307** 125 (*Preprint* arXiv:1208.4605 [hep-ph])
- [23] Ackermann M *et al.* [Fermi-LAT Collab.] 2011 Constraining Dark Matter Models from a Combined Analysis of Milky Way Satellites with the Fermi Large Area Telescope *Phys. Rev. Lett.* **107** 241302 (*Preprint* arXiv:1108.3546 [astro-ph.HE])
- [24] Bergstrom L, Ullio P and Buckley J H 1998 Observability of gamma-rays from dark matter neutralino annihilations in the Milky Way halo *Astropart. Phys.* **9** 137 (*Preprint* astro-ph/9712318)  
Cirelli M *et al.* 2011 PPPC 4 DM ID: A Poor Particle Physicist Cookbook for Dark Matter Indirect Detection *JCAP* **1103** 051 [Erratum-ibid. 2012 **1210** E01] (*Preprint* arXiv:1012.4515 [hep-ph])
- [25] CMS Collaboration 2012 Search for new physics in monojet events in  $pp$  collisions at  $\sqrt{s} = 8$  TeV *CMS Note* CMS-PAS-EXO-12-048
- [26] Fox P J, Harnik R, Kopp J and Tsai Y 2012 Missing Energy Signatures of Dark Matter at the LHC *Phys. Rev. D* **85** 056011 (*Preprint* arXiv:1109.4398 [hep-ph])
- [27] Aad G *et al.* [ATLAS Collab.] 2013 Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in  $pp$  collision data at  $\sqrt{s} = 7$  TeV with the ATLAS detector *Phys. Rev. Lett.* **110** 011802 (*Preprint* arXiv:1209.4625 [hep-ex])
- [28] Chatrchyan S *et al.* [CMS Collab.] 2012 Search for Dark Matter and Large Extra Dimensions in  $pp$  Collisions

- Yielding a Photon and Missing Transverse Energy *Phys. Rev. Lett.* **108** 261803 (*Preprint* arXiv:1204.0821 [hep-ex])
- [29] Aaltonen T *et al.* [CDF Collab.] 2012 A Search for dark matter in events with one jet and missing transverse energy in  $p\bar{p}$  collisions at  $\sqrt{s} = 1.96$  TeV *Phys. Rev. Lett.* **108** 211804 (*Preprint* arXiv:1203.0742 [hep-ex])
- [30] Aprile E *et al.* [XENON100 Collab.] 2011 Dark Matter Results from 100 Live Days of XENON100 Data *Phys. Rev. Lett.* **107** 131302 (*Preprint* arXiv:1104.2549 [astro-ph.CO])
- [31] Aalseth C E *et al.* [CoGeNT Collab.] 2011 Results from a Search for Light-Mass Dark Matter with a P-type Point Contact Germanium Detector *Phys. Rev. Lett.* **106** 131301 (*Preprint* arXiv:1002.4703 [astro-ph.CO])
- [32] Ahmed Z *et al.* [CDMS-II Collab.] 2010 Dark Matter Search Results from the CDMS II Experiment *Science* **327** 1619 (*Preprint* arXiv:0912.3592 [astro-ph.CO])  
 Ahmed Z *et al.* [CDMS-II Collab.] 2011 Results from a Low-Energy Analysis of the CDMS II Germanium Data *Phys. Rev. Lett.* **106** 131302 (*Preprint* arXiv:1011.2482 [astro-ph.CO])
- [33] Felizardo M *et al.* 2012 Final Analysis and Results of the Phase II SIMPLE Dark Matter Search *Phys. Rev. Lett.* **108** 201302 (*Preprint* arXiv:1106.3014 [astro-ph.CO])
- [34] Archambault S *et al.* [PICASSO Collab.] 2012 Constraints on Low-Mass WIMP Interactions on  $^{19}\text{F}$  from PICASSO *Phys. Lett. B* **711** 153 (*Preprint* arXiv:1202.1240 [hep-ex])
- [35] Bai Y and Tait T M P 2013 Searches with Mono-Leptons *Phys. Lett. B* **723** 384 (*Preprint* arXiv:1208.4361 [hep-ph])
- [36] Bell N F, Dent J B, Galea A J, Jacques T D, Krauss L M and Weiler T J 2012 Searching for Dark Matter at the LHC with a Mono-Z *Phys. Rev. D* **86** 096011 (*Preprint* arXiv:1209.0231 [hep-ph])  
 Carpenter L M, Nelson A, Shimmin C, Tait T M P and Whiteson D 2013 Collider searches for dark matter in events with a Z boson and missing energy *Phys. Rev. D* **87** 074005 (*Preprint* arXiv:1212.3352 [hep-ex])
- [37] CMS Collaboration 2013 Search for dark matter in the mono-lepton channel with  $pp$  collision events at  $\sqrt{s} = 8$  TeV *CMS Note* CMS-PAS-EXO-13-004
- [38] CMS Collaboration 2012 Search for new physics in the final states with a lepton and missing transverse energy at  $\sqrt{s} = 8$  TeV *CMS Note* CMS-PAS-EXO-12-060
- [39] Aad G *et al.* [ATLAS Collab.] 2014 Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in  $pp$  collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector *Phys. Rev. Lett.* **112** 041802 (*Preprint* arXiv:1309.4017 [hep-ex])
- [40] Aad G *et al.* [ATLAS Collab.] 2013 Performance of jet substructure techniques for large- $R$  jets in proton-proton collisions at  $\sqrt{s} = 7$  TeV using the ATLAS detector *JHEP* **1309** 076 (*Preprint* arXiv:1306.4945 [hep-ex])
- [41] Aprile E *et al.* [XENON100 Collab.] 2012 Dark Matter Results from 225 Live Days of XENON100 Data *Phys. Rev. Lett.* **109** 181301 (*Preprint* arXiv:1207.5988 [astro-ph.CO])
- [42] Behnke E *et al.* [COUPP Collab.] 2012 First Dark Matter Search Results from a 4-kg  $\text{CF}_3\text{I}$  Bubble Chamber Operated in a Deep Underground Site *Phys. Rev. D* **86** 052001 (*Preprint* arXiv:1204.3094 [astro-ph.CO])
- [43] Aartsen M G *et al.* [IceCube Collab.] 2013 Search for dark matter annihilations in the Sun with the 79-string IceCube detector *Phys. Rev. Lett.* **110** 131302 (*Preprint* arXiv:1212.4097 [astro-ph.HE])
- [44] Nilles H P 1984 Supersymmetry, Supergravity and Particle Physics *Phys. Rept.* **110** 1
- [45] Mitsou V A 2013 Experimental status of particle and astroparticle searches for supersymmetry *J. Phys. Conf. Series* **447** 012019 (*Preprint* arXiv:1304.1414 [hep-ph])
- [46] Cabrera M E and Casas J A 2012 Understanding and improving the Effective Mass for LHC searches *Preprint* arXiv:1207.0435 [hep-ph]
- [47] Cabrera M E, Casas J A, Mitsou V A, Ruiz de Austri R and Terron J 2012 Histogram comparison as a powerful tool for the search of new physics at LHC. Application to CMSSM *JHEP* **1204** 133 (*Preprint* arXiv:1109.3759 [hep-ph])
- [48] Aad G *et al.* [ATLAS Collab.] 2013 Search for new phenomena in final states with large jet multiplicities and missing transverse momentum at  $\sqrt{s} = 8$  TeV proton-proton collisions using the ATLAS experiment *JHEP* **1310** 130 (*Preprint* arXiv:1308.1841 [hep-ex])
- [49] Chatrchyan S *et al.* [CMS Collab.] 2012 Search for supersymmetry in hadronic final states using MT2 in  $pp$  collisions at  $\sqrt{s} = 7$  TeV *JHEP* **1210** 018 (*Preprint* arXiv:1207.1798 [hep-ex])  
 Chatrchyan S *et al.* [CMS Collab.] 2012 Search for new physics in the multijet and missing transverse momentum final state in proton-proton collisions at  $\sqrt{s} = 7$  TeV *Phys. Rev. Lett.* **109** 171803 (*Preprint* arXiv:1207.1898 [hep-ex])
- [50] Aad G *et al.* [ATLAS Collab.] 2012 Further search for supersymmetry at  $\sqrt{s} = 7$  TeV in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector *Phys. Rev. D* **86** 092002 (*Preprint* arXiv:1208.4688 [hep-ex])
- [51] Chatrchyan S *et al.* [CMS Collab.] 2013 Search for supersymmetry in  $pp$  collisions at  $\sqrt{s} = 7$  TeV in events with a single lepton, jets, and missing transverse momentum *Eur. Phys. J. C* **73** 2404 (*Preprint*

- arXiv:1212.6428 [hep-ex])  
 Chatrchyan S *et al.* [CMS Collab.] 2013 Search for supersymmetry in events with opposite-sign dileptons and missing transverse energy using an artificial neural network *Phys. Rev. D* **87** 072001 (*Preprint* arXiv:1301.0916 [hep-ex])
- [52] Baer H, Box A, Park E-K and Tata X 2007 Implications of compressed supersymmetry for collider and dark matter searches *JHEP* **0708** 060 (*Preprint* arXiv:0707.0618 [hep-ph])  
 LeCompte T J and Martin S P 2012 Compressed supersymmetry after  $1 \text{ fb}^{-1}$  at the Large Hadron Collider *Phys. Rev. D* **85** 035023 (*Preprint* arXiv:1111.6897 [hep-ph])
- [53] Rogan C 2010 Kinematical variables towards new dynamics at the LHC *Preprint* arXiv:1006.2727 [hep-ph]
- [54] Chatrchyan S *et al.* [CMS Collab.] 2013 Inclusive search for supersymmetry using the razor variables in  $pp$  collisions at  $\sqrt{s} = 7 \text{ TeV}$  *Phys. Rev. Lett.* **111** 081802 (*Preprint* arXiv:1212.6961 [hep-ex])
- [55] Aad G *et al.* [ATLAS Collab.] 2013 Multi-channel search for squarks and gluinos in  $\sqrt{s} = 7 \text{ TeV}$   $pp$  collisions with the ATLAS detector *Eur. Phys. J. C* **73** 2362 (*Preprint* arXiv:1212.6149 [hep-ex])
- [56] ATLAS Collaboration 2013 Search for strongly produced superpartners in final states with two same sign leptons with the ATLAS detector using  $21 \text{ fb}^{-1}$  of proton-proton collisions at  $\sqrt{s} = 8 \text{ TeV}$  *ATLAS Note* ATLAS-CONF-2013-007  
 ATLAS Collaboration 2013 Search for Supersymmetry in Events with Large Missing Transverse Momentum, Jets, and at Least One Tau Lepton in  $21 \text{ fb}^{-1}$  of  $\sqrt{s} = 8 \text{ TeV}$  Proton-Proton Collision Data with the ATLAS Detector *ATLAS Note* ATLAS-CONF-2013-026  
 ATLAS Collaboration 2013 Measurement of multi-jet cross-section ratios and determination of the strong coupling constant in proton-proton collisions at  $\sqrt{s} = 7 \text{ TeV}$  with the ATLAS detector *ATLAS Note* ATLAS-CONF-2013-041  
 ATLAS Collaboration 2013 Search for strong production of supersymmetric particles in final states with missing transverse momentum and at least three  $b$ -jets using  $20.1 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 8 \text{ TeV}$  with the ATLAS Detector *ATLAS Note* ATLAS-CONF-2013-061  
 ATLAS Collaboration 2013 Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at  $\sqrt{s} = 8 \text{ TeV}$  with the ATLAS detector *ATLAS Note* ATLAS-CONF-2013-062
- [57] ATLAS Supersymmetry Searches Public Results:  
<https://twiki.cern.ch/twiki/bin/view/AtlasPublic/SupersymmetryPublicResults>
- [58] Lahanas A B, Mavromatos N E and Nanopoulos D V 2007 Smoothly evolving supercritical-string dark energy relaxes supersymmetric-dark-matter constraints *Phys. Lett. B* **649** 83 (*Preprint* hep-ph/0612152)
- [59] Mavromatos N E, Sarkar S and Vergou A 2011 Stringy Space-Time Foam, Finsler-like Metrics and Dark Matter Relics *Phys. Lett. B* **696** 300 (*Preprint* arXiv:1009.2880 [hep-th])  
 Mavromatos N E, Mitsou V A, Sarkar S and Vergou A 2012 Implications of a Stochastic Microscopic Finsler Cosmology *Eur. Phys. J. C* **72** 1956 (*Preprint* arXiv:1012.4094 [hep-ph])
- [60] Chatrchyan S *et al.* [CMS Collab.] 2013 Search for gluino mediated bottom- and top-squark production in multijet final states in  $pp$  collisions at  $8 \text{ TeV}$  *Phys. Lett. B* **725** 243 (*Preprint* arXiv:1305.2390 [hep-ex])  
 CMS Collaboration 2013 Search for supersymmetry in  $pp$  collisions at a center-of-mass energy of  $8 \text{ TeV}$  in events with a single lepton, multiple jets and  $b$ -tags *CMS Note* CMS-PAS-SUS-13-007  
 CMS Collaboration 2013 Search for supersymmetry in  $pp$  collisions at  $\sqrt{s} = 8 \text{ TeV}$  in events with three leptons and at least one  $b$ -tagged jet *CMS Note* CMS-PAS-SUS-13-008
- [61] CMS Collaboration 2013 Search for new physics in events with same-sign dileptons and jets in  $pp$  collisions at  $\sqrt{s} = 8 \text{ TeV}$  *CMS Note* CMS-PAS-SUS-13-013
- [62] CMS Supersymmetry Physics Results:  
<https://twiki.cern.ch/twiki/bin/view/CMSPublic/PhysicsResultsSUS>
- [63] Aad G *et al.* [ATLAS Collab.] 2012 Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at  $\sqrt{s} = 7 \text{ TeV}$  with the ATLAS detector *Phys. Rev. Lett.* **109** 211802 (*Preprint* arXiv:1208.1447 [hep-ex])  
 Aad G *et al.* [ATLAS Collab.] 2012 Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in  $\sqrt{s} = 7 \text{ TeV}$   $pp$  collisions using  $4.7 \text{ fb}^{-1}$  of ATLAS data *Phys. Rev. Lett.* **109** 211803 (*Preprint* arXiv:1208.2590 [hep-ex])  
 Aad G *et al.* [ATLAS Collab.] 2012 Search for light scalar top quark pair production in final states with two leptons with the ATLAS detector in  $\sqrt{s} = 7 \text{ TeV}$  proton-proton collisions *Eur. Phys. J. C* **72** 2237 (*Preprint* arXiv:1208.4305 [hep-ex])  
 Aad G *et al.* [ATLAS Collab.] 2012 Search for a heavy top-quark partner in final states with two leptons with the ATLAS detector at the LHC *JHEP* **1211** 094 (*Preprint* arXiv:1209.4186 [hep-ex])  
 Aad G *et al.* [ATLAS Collab.] 2013 Search for light top squark pair production in final states with leptons and  $b$ -jets with the ATLAS detector in  $\sqrt{s} = 7 \text{ TeV}$  proton-proton collisions *Phys. Lett. B* **720** 13 (*Preprint* arXiv:1209.2102 [hep-ex])

- Aad G *et al.* [ATLAS Collab.] 2013 Search for direct third-generation squark pair production in final states with missing transverse momentum and two  $b$ -jets in  $\sqrt{s} = 8$  TeV  $pp$  collisions with the ATLAS detector *JHEP* **1310** 189 (*Preprint* arXiv:1308.2631 [hep-ex])
- ATLAS Collaboration 2013 Search for direct production of the top squark in the all-hadronic  $t\bar{t} + E_T^{\text{miss}}$  final state in  $21 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector *ATLAS Note* ATLAS-CONF-2013-024
- ATLAS Collaboration 2013 Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in  $\sqrt{s} = 8$  TeV  $pp$  collisions using  $21 \text{ fb}^{-1}$  of ATLAS data *ATLAS Note* ATLAS-CONF-2013-037
- ATLAS Collaboration 2013 Search for direct top squark pair production in final states with two leptons in  $\sqrt{s} = 8$  TeV  $pp$  collisions using  $20 \text{ fb}^{-1}$  of ATLAS data *ATLAS Note* ATLAS-CONF-2013-048
- ATLAS Collaboration 2013 Searches for direct scalar top pair production in final states with two leptons using the transverse mass variable and a multivariate analysis technique in  $\sqrt{s} = 8$  TeV  $pp$  collisions using  $20.3 \text{ fb}^{-1}$  of ATLAS data *ATLAS Note* ATLAS-CONF-2013-065
- ATLAS Collaboration 2013 Search for pair-produced top squarks decaying into charm quarks and the lightest neutralinos using  $20.3 \text{ fb}^{-1}$  of  $pp$  collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector at the LHC *ATLAS Note* ATLAS-CONF-2013-068
- [64] CMS Collaboration 2013 Search for electroweak production of charginos, neutralinos, and sleptons using leptonic final states in  $pp$  collisions at  $\sqrt{s} = 8$  TeV *CMS Note* CMS-PAS-SUS-13-006
- CMS Collaboration 2013 Search for electroweak production of charginos and neutralinos in final states with a Higgs boson in  $pp$  collisions at  $\sqrt{s} = 8$  TeV *CMS Note* CMS-PAS-SUS-13-017
- [65] Barbier R *et al.* 2005  $R$ -parity violating supersymmetry *Phys. Rept.* **420** 1 (*Preprint* hep-ph/0406039)
- [66] Aad G *et al.* [ATLAS Collab.] 2011 Search for a heavy particle decaying into an electron and a muon with the ATLAS detector in  $\sqrt{s} = 7$  TeV  $pp$  collisions at the LHC *Phys. Rev. Lett.* **106** 251801 (*Preprint* arXiv:1103.5559 [hep-ex])
- Aad G *et al.* [ATLAS Collab.] 2011 Search for a heavy neutral particle decaying into an electron and a muon using  $1 \text{ fb}^{-1}$  of ATLAS data *Eur. Phys. J. C* **71** 1809 (*Preprint* arXiv:1109.3089 [hep-ex])
- Aad G *et al.* [ATLAS Collab.] 2013 Search for a heavy narrow resonance decaying to  $e\mu$ ,  $e\tau$ , or  $\mu\tau$  with the ATLAS detector in  $\sqrt{s} = 7$  TeV  $pp$  collisions at the LHC *Phys. Lett. B* **723** 15 (*Preprint* arXiv:1212.1272 [hep-ex])
- [67] Aad G *et al.* [ATLAS Collab.] 2012 Search for pair production of massive particles decaying into three quarks with the ATLAS detector in  $\sqrt{s} = 7$  TeV  $pp$  collisions at the LHC *JHEP* **1212** 086 (*Preprint* arXiv:1210.4813 [hep-ex])
- [68] Aad G *et al.* [ATLAS Collab.] 2012 Search for  $R$ -parity-violating supersymmetry in events with four or more leptons in  $\sqrt{s} = 7$  TeV  $pp$  collisions with the ATLAS detector *JHEP* **1212** 124 (*Preprint* arXiv:1210.4457 [hep-ex])
- [69] Aad G *et al.* [ATLAS Collab.] 2012 Search for lepton flavour violation in the  $e\mu$  continuum with the ATLAS detector in  $\sqrt{s} = 7$  TeV  $pp$  collisions at the LHC *Eur. Phys. J. C* **72** 2040 (*Preprint* arXiv:1205.0725 [hep-ex])
- [70] Aad G *et al.* [ATLAS Collab.] 2012 Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in  $\sqrt{s} = 7$  TeV  $pp$  collisions using  $1 \text{ fb}^{-1}$  of ATLAS data *Phys. Rev. D* **85** 012006 (*Preprint* arXiv:1109.6606 [hep-ex])
- ATLAS Collaboration 2012 Search for supersymmetry at  $\sqrt{s} = 7$  TeV in final states with large jet multiplicity, missing transverse momentum and one isolated lepton with the ATLAS detector *ATLAS Note* ATLAS-CONF-2012-140
- [71] Romao J C, Diaz M A, Hirsch M, Porod W and Valle J W F 2000 A Supersymmetric solution to the solar and atmospheric neutrino problems *Phys. Rev. D* **61** 071703 (*Preprint* hep-ph/9907499)
- Hirsch M, Diaz M A, Porod W, Romao J C and Valle J W F 2000 Neutrino masses and mixings from supersymmetry with bilinear  $R$  parity violation: A Theory for solar and atmospheric neutrino oscillations *Phys. Rev. D* **62** 113008 [Erratum-ibid. 2002 D **65** 119901] (*Preprint* hep-ph/0004115)
- Porod W, Hirsch M, Romao J and Valle J W F 2001 Testing neutrino mixing at future collider experiments *Phys. Rev. D* **63** 115004 (*Preprint* hep-ph/0011248)
- Diaz M A, Hirsch M, Porod W, Romao J C and Valle J W F 2003 Solar neutrino masses and mixing from bilinear  $R$  parity broken supersymmetry: Analytical versus numerical results *Phys. Rev. D* **68** 013009 [Erratum-ibid. 2005 D **71** 059904] (*Preprint* hep-ph/0302021)
- Hirsch M and Valle J W F 2004 Supersymmetric origin of neutrino mass *New J. Phys.* **6** 76 (*Preprint* hep-ph/0405015)
- [72] CMS Collaboration 2013 Search for RPV SUSY resonant second generation slepton production in same-sign dimuon events at  $\sqrt{s} = 7$  TeV *CMS Note* CMS-PAS-SUS-13-005

- [73] Chatrchyan S *et al.* [CMS Collab.] 2013 Search for top squarks in  $R$ -parity-violating supersymmetry using three or more leptons and  $b$ -tagged jets *Phys. Rev. Lett.* **111** 221801 (*Preprint* arXiv:1306.6643 [hep-ex])
- [74] CMS Collaboration 2013 Search for RPV SUSY in the four-lepton final state *CMS Note* CMS-PAS-SUS-13-010
- [75] Chatrchyan S *et al.* [CMS Collab.] 2013 Search for pair production of third-generation leptoquarks and top squarks in  $pp$  collisions at  $\sqrt{s} = 7$  TeV *Phys. Rev. Lett.* **110** 081801 (*Preprint* arXiv:1210.5629 [hep-ex])  
 CMS Collaboration 2012 Search for RPV supersymmetry with three or more leptons and  $b$ -tags *CMS Note* CMS-PAS-SUS-12-027  
 CMS Collaboration 2012 Search for Light- and Heavy-flavour Three-Jet Resonances In Multijet Final States at 8 TeV *CMS Note* CMS-PAS-EXO-12-049
- [76] Aad G *et al.* [ATLAS Collab.] 2013 Search for long-lived, heavy particles in final states with a muon and multi-track displaced vertex in proton-proton collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector *Phys. Lett. B* **719** 280 (*Preprint* arXiv:1210.7451 [hep-ex])
- [77] Aad G *et al.* [ATLAS Collab.] 2013 Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in  $pp$  collisions at  $\sqrt{s} = 7$  TeV with the ATLAS detector *JHEP* **1301** 131 (*Preprint* arXiv:1210.2852 [hep-ex])
- [78] Aad G *et al.* [ATLAS Collab.] 2013 Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in  $pp$  collisions at  $\sqrt{s} = 7$  TeV *Phys. Lett. B* **720** 277 (*Preprint* arXiv:1211.1597 [hep-ex])  
 Chatrchyan S *et al.* [CMS Collab.] 2013 Searches for long-lived charged particles in  $pp$  collisions at  $\sqrt{s} = 7$  and 8 TeV *JHEP* **1307** 122 (*Preprint* arXiv:1305.0491 [hep-ex])
- [79] Aad G *et al.* [ATLAS Collab.] 2013 Search for non-pointing photons in the diphoton and  $E_{\text{T}}^{\text{miss}}$  final state in  $\sqrt{s} = 7$  TeV proton-proton collisions using the ATLAS detector *Phys. Rev. D* **88** 012001 (*Preprint* arXiv:1304.6310 [hep-ex])  
 Chatrchyan S *et al.* [CMS Collab.] 2013 Search for long-lived particles decaying to photons and missing energy in proton-proton collisions at  $\sqrt{s} = 7$  TeV *Phys. Lett. B* **722** 273 (*Preprint* arXiv:1212.1838 [hep-ex])
- [80] Takayama F and Yamaguchi M 2000 Gravitino dark matter without  $R$ -parity *Phys. Lett. B* **485** 388 (*Preprint* hep-ph/0005214)  
 Hirsch M, Porod W and Restrepo D 2005 Collider signals of gravitino dark matter in bilinearly broken  $R$ -parity *JHEP* **0503** 062 (*Preprint* hep-ph/0503059)  
 Buchmuller W, Covi L, Hamaguchi K, Ibarra A and Yanagida T 2007 Gravitino Dark Matter in  $R$ -Parity Breaking Vacua *JHEP* **0703** 037 (*Preprint* hep-ph/0702184)
- [81] Liew S P 2013 Gamma-ray line from radiative decay of gravitino dark matter *Phys. Lett. B* **724** 88 (*Preprint* arXiv:1304.1992 [hep-ph])
- [82] Chun E J and Kim H B 2006 Axino Light Dark Matter and Neutrino Masses with  $R$ -parity Violation *JHEP* **0610** 082 (*Preprint* hep-ph/0607076)
- [83] Endo M, Hamaguchi K, Liew S P, Mukaida K and Nakayama K 2013 Axino dark matter with  $R$ -parity violation and 130 GeV gamma-ray line *Phys. Lett. B* **721** 111 (*Preprint* arXiv:1301.7536 [hep-ph])
- [84] Lopez-Fogliani D E and Munoz C 2006 Proposal for a new minimal supersymmetric standard model *Phys. Rev. Lett.* **97** 041801 (*Preprint* hep-ph/0508297)  
 Escudero N, Lopez-Fogliani D E, Munoz C and Ruiz de Austri R 2008 Analysis of the parameter space and spectrum of the  $\mu\nu$ SSM *JHEP* **0812** 099 (*Preprint* arXiv:0810.1507 [hep-ph])  
 Ghosh P, Lopez-Fogliani D E, Mitsou V A, Munoz C and Ruiz de Austri R 2013 Probing the  $\mu$  from  $\nu$  supersymmetric standard model with displaced multileptons from the decay of a Higgs boson at the LHC *Phys. Rev. D* **88** 015009 (*Preprint* arXiv:1211.3177 [hep-ph])
- [85] Choi K-Y, Lopez-Fogliani D E, Munoz C and Ruiz de Austri R 2010 Gamma-ray detection from gravitino dark matter decay in the  $\mu\nu$ SSM *JCAP* **1003** 028 (*Preprint* arXiv:0906.3681 [hep-ph])
- [86] Bringmann T, Huang X, Ibarra A, Vogl S and Weniger C 2012 Fermi LAT Search for Internal Bremsstrahlung Signatures from Dark Matter Annihilation *JCAP* **1207** 054 (*Preprint* arXiv:1203.1312 [hep-ph])  
 Weniger C 2012 A Tentative Gamma-Ray Line from Dark Matter Annihilation at the Fermi Large Area Telescope *JCAP* **1208** 007 (*Preprint* arXiv:1204.2797 [hep-ph])  
 Tempel E, Hektor A and Raidal M 2012 Fermi 130 GeV gamma-ray excess and dark matter annihilation in sub-haloes and in the Galactic centre *JCAP* **1209** 032 [Addendum-ibid. 2012 **1211** A01] (*Preprint* arXiv:1205.1045 [hep-ph])