

Iodine status worldwide

WHO Global Database
on Iodine Deficiency

Editors

Bruno de Benoist

Maria Andersson

Ines Egli

Bahi Takkouche

Henrietta Allen



Department of Nutrition for Health and Development
World Health Organization

Geneva

2004

WHO Library Cataloguing-in-Publication Data

Iodine status worldwide : WHO Global Database on Iodine Deficiency / editors: Bruno de Benoist ... [et al.]

I.Iodine 2.Deficiency diseases – epidemiology 3.Goiter – epidemiology 4.Nutrition surveys 5.Databases, Factual
I.World Health Organization II.De Benoist, Bruno.

ISBN 92 4 159200 1

(NLM classification: WD 105)

© **World Health Organization 2004**

All rights reserved. Publications of the World Health Organization can be obtained from Marketing and Dissemination, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel: +41 22 791 2476; fax: +41 22 791 4857; email: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to Publications, at the above address (fax: +41 22 791 4806; email: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

The World Health Organization does not warrant that the information contained in this publication is complete and correct and shall not be liable for any damages incurred as a result of its use.

The named editors alone are responsible for the views expressed in this publication.

In issuing this document, the World Health Organization welcomes comments from experts and institutions for further advice for future updating of the document.

Front cover photos from WHO photo library.

Photographers: Kelly T, Julio Vizcarra, James Nalty, Carlos Gaggero, Jean-Marc Giboux.

Designed by minimum graphics

Printed in Malta

Contents

Preface	v
Acknowledgements	vi
Abbreviations	vii
1. Introduction	1
1.1 Iodine deficiency disorders: a public health problem	1
1.1.1 Etiology	1
1.1.2 Health consequences	1
1.1.3 Indicators for assessment and monitoring	1
1.2 Control of IDD	2
1.2.1 Correcting iodine deficiency	2
1.2.2 Monitoring and evaluating the IDD control programmes	3
1.2.3 Increasing awareness of public health authorities and the general public	3
1.2.4 Reinforcing the collaboration between sectors	3
1.2.5 Sustaining IDD control programmes	4
2. Methods	5
2.1 Data sources – The WHO Global Database on Iodine Deficiency	5
2.2 Selection of survey data	5
2.2.1 Administrative level	5
2.2.2 Population groups	6
2.3 Classification of iodine nutrition	6
2.4 Population coverage, proportion of population and the number of individuals with insufficient iodine intake	7
2.4.1 Population coverage	7
2.4.2 Proportion of population and the number of individuals with insufficient iodine intake	7
2.5 TGP	8
3. Results and discussion	9
3.1 Results	9
3.1.1 Population coverage	9
3.1.2 Classification of countries by degree of public health significance of iodine nutrition based on median UI	10
3.1.3 Proportion of population and number of individuals with insufficient iodine intake	12
3.1.4 TGP	12
3.2 Discussion	13
3.2.1 Population coverage	13
3.2.2 Limitations of data sources	13
3.2.3 Classification of countries by degree of public health significance of iodine nutrition based on median UI	13
3.2.4 Proportion of population and the number of individuals with insufficient iodine intake	14
3.2.5 TGP	14
3.3 Conclusion	14
References	16

Annexes		17
Annex 1.	WHO Member States grouped by WHO and UN regions	17
	Table A1.1 WHO Member States grouped by WHO region	17
	Table A1.2 WHO Member States grouped by UN region and subregion	18
Annex 2.	Results by UN region	20
	Table A2.1 Population coverage by UI surveys carried out between 1993 and 2003, by UN region	20
	Table A2.2 Type of UI survey data by UN region	20
	Table A2.3 Population coverage by TGP surveys carried out between 1993 and 2003, by UN region	21
	Table A2.4 Type of TGP survey data by UN region	21
	Table A2.5 Number of countries classified by degrees of public health significance of iodine nutrition based on median UI in school-age children, by UN region, 2003	22
	Table A2.6 Proportion of population, and number of individuals with, insufficient iodine intake in school-age children (6–12 years) and the general population, by UN region, 2003	23
	Table A2.7 TGP in the general population by UN region, 2003	23
Annex 3.	National estimates of iodine status	25
	Table A3.1 Country data on UI and national estimate of iodine nutrition	26
	Table A3.2 Country data on TGP	33
Tables		
Table 1.1	The spectrum of IDD across the life-span	1
Table 1.2	Criteria for monitoring progress towards sustaining elimination of IDD	4
Table 2.1	Epidemiological criteria for assessing iodine nutrition based on median UI concentrations in school-age children	6
Table 3.1	Population coverage by UI surveys carried out between 1993 and 2003, by WHO region	9
Table 3.2	Type of UI survey data by WHO region	9
Table 3.3	Population coverage by TGP surveys carried out between 1993 and 2003, by WHO region	10
Table 3.4	Type of TGP survey data by WHO region	10
Table 3.5	Number of countries classified by degrees of public health significance of iodine nutrition based on median UI in school-age children by WHO region, 2003	12
Table 3.6	Proportion of population, and number of individuals with insufficient iodine intake in school-age children (6–12 years), and in the general population (all age groups), by WHO region, 2003	12
Table 3.7	Change in total goitre prevalence between 1993 and 2003, by WHO region	12
Figures		
Figure 2.1	Relation between median UI ($\mu\text{g/l}$) and mean UI ($\mu\text{g/l}$) with linear regression line	7
Figure 2.2	Relation between median UI ($\mu\text{g/l}$) and proportion (%) of UI values below 100 $\mu\text{g/l}$ with quadratic regression curve	7
Figure 2.3	Relation between general population TGP and school-age children TGP with linear regression line	8
Figure 3.1	Type of UI survey data	9
Figure 3.2	Type of TGP survey data	10
Figure 3.3	Degree of public health significance of iodine nutrition based on median UI	11

Preface

In 1960, the World Health Organization (WHO) published the first global review on the extent of endemic goitre. This review, covering 115 countries, was instrumental in focusing attention on the scale of the public health problem of Iodine Deficiency Disorders (IDD). It was only in the mid 1980s that the international community committed themselves to the elimination of IDD, through a number of declarations and resolutions.

WHO subsequently established a global database on iodine deficiency which now holds surveys dating back from the 1940s to the present day. Its objective is to assess the global magnitude of iodine deficiency, to evaluate the strategies for its control and to monitor each country's progress towards achieving the international community's goal of IDD elimination.

In 1993, WHO published the first version of the WHO Global Database on Iodine Deficiency with global estimates on the prevalence of iodine deficiency based on total goitre prevalence (TGP), using data from 121 countries.

Since then the international community and the authorities in most countries where IDD was identified as a public health problem have taken measures to control iodine deficiency, in particular through salt iodization programmes – the WHO recommended strategy to prevent and control IDD. As a result, it is assumed that the iodine status of populations throughout the world has improved over the past decade. The WHO Global Database on Iodine Deficiency is therefore being revised and updated to reflect the current situation of iodine deficiency worldwide.

Until the 1990s TGP was the recommended indicator for assessing iodine status. However, goitre responds slowly to a change in iodine status and today urinary iodine (UI)

is recommended as a more sensitive indicator of recent changes in iodine nutrition. The methodology used for this updated version of global iodine status thus rests on UI data and only uses TGP to make a comparison with the data published in 1993.

This report provides general information on iodine deficiency, its health consequences and current control interventions (Chapter 1). The methodology used to generate estimates at national, regional and global levels is described in Chapter 2. The estimates of iodine deficiency at national, regional and worldwide levels are given in Chapter 3 followed by a critical analysis of the methodology used. Annex 1 provides detailed information on the status of iodine deficiency, UI and TGP, for each country for which data are available.

The objective of this report is to provide an updated analysis of the iodine deficiency situation in the world at the beginning of the 21st century. It forms part of WHO's work to track the progress made by each country to meet the goal of IDD elimination. We hope that this report will help governments recognize the progress made in improving iodine nutrition over the past decade, and also to be aware that iodine deficiency is still a public health problem in some countries. In order to reach the goal of IDD elimination continued efforts are needed on the part of health authorities. It will also require that control programmes are sustained and strengthened.

Bruno de Benoist, MSc, MD

Focal Point, Micronutrient Programme

Department of Nutrition for Health and Development

World Health Organization, Geneva

Acknowledgements

The WHO Global Database on Iodine Deficiency is maintained at the Department of Nutrition for Health and Development. Update of the database and overall revision of iodine status worldwide was made possible through financial support by UNICEF. WHO gratefully acknowledges the contribution of UNICEF towards the publication of this report.

The database was developed and managed by Henrietta Allen and Maria Andersson under the coordination of Bruno de Benoist. Grace Rob was assisting in data management. Bruno de Benoist, Maria Andersson, Bahi Takkouche and Ines Egli were engaged in data analysis and preparation of the report.

WHO wishes to thank the numerous individuals, institutions, governments, and non-governmental and international organizations who provided data to the global database. Without continual international collaboration in keeping the global database up-to-date, this compilation on worldwide patterns and trends in iodine deficiency would not have been possible. Special thanks are due to ministries of health of the WHO Member States; WHO Regional Offices; WHO Country Offices; the Nutrition section, UNICEF, New York, NY, USA; UNICEF Regional Of-

ices; UNICEF Country Offices; the International Council for the Control of Iodine Deficiency Disorders.

The following individuals provided technical comments and valuable advice to improve the clarity of the text:

Fereidoun Azizi, Shahid Beheshti University of Medical Sciences, Teheran, Islamic Republic of Iran; Zu-pei Chen, Tianjin Medical University, Tianjin, China; Nita Dalmiya, Nutrition section, UNICEF, New York, NY, USA; Ian Darnton-Hill, Nutrition Section, UNICEF, New York, NY, USA; François Delange, ICCIDD, Brussels, Belgium; John T Dunn^a, ICCIDD, Charlottesville, VA, USA; Pieter Jooste, Medical Research Council, Cape Town, South Africa; Aldo Pinchera, ICCIDD, Pisa, Italy; Eduardo Pretell, ICCIDD, Lima, Peru; Claudia Stein, WHO, Geneva, Switzerland; Kevin Sullivan, Rollins School of Public Health of Emory University, Atlanta, GA, USA; Charles Todd, European Commission, Brussels, Belgium.

Editorial assistance was provided by Kai Lashley. The support provided by the staff of the Department of Nutrition for Health and Development, WHO, Geneva, especially Trudy Wijnhoven and Anna Wolter, is also much appreciated.

^a deceased

Abbreviations

CDC	Centers for Disease Control and Prevention
FAO	Food and Agricultural Organization of the United Nations
ICCIDD	International Council for Control of Iodine Deficiency Disorders
IDD	Iodine deficiency disorders. The spectrum of clinical, social and intellectual consequences of iodine deficiency.
IIH	Iodine-induced hyperthyroidism
MI	The Micronutrient Initiative
ppm	Parts per million
SAC	School-age children (6–12 years)
TGP	Total goitre prevalence. Prevalence of enlarged goitres in a population (usually school-age children).
TSH	Thyroid stimulating hormone
UI	Urinary iodine
UL	Upper limit
UN	United Nations
UNICEF	The United Nations Children's Fund
USI	Universal salt iodization
WHO	World Health Organization

1. Introduction

1.1 Iodine deficiency disorders: a public health problem

Iodine deficiency is a major public health problem for populations throughout the world, particularly for pregnant women and young children. They are a threat to the social and economic development of countries. The most devastating outcomes of iodine deficiency are increased perinatal mortality and mental retardation – iodine deficiency is the greatest cause of preventable brain damage in childhood which is the primary motivation behind the current world-wide drive to eliminate it.

1.1.1 Etiology

The main factor responsible for iodine deficiency is a low dietary supply of iodine (1). It occurs in populations living in areas where the soil has a low iodine content as a result of past glaciation or the repeated leaching effects of snow, water and heavy rainfall. Crops grown in this soil, therefore, do not provide adequate amounts of iodine when consumed.

1.1.2 Health consequences

Iodine is present in the body in minute amounts, mainly in the thyroid gland. Its main role is in the synthesis of thyroid hormones. When iodine requirements are not met, thyroid hormone synthesis is impaired, resulting in hypothyroidism and a series of functional and developmental abnormalities grouped under the heading of “Iodine Deficiency Disorders (IDD)” as shown in Table 1.1.

Goitre is the most visible manifestation of IDD. Endemic goitre results from increased thyroid stimulation by thyroid stimulating hormone (TSH) to maximize the utilization of available iodine and thus represents maladaptation to iodine deficiency (5, 6). However, the most damaging disorders induced by iodine deficiency are irreversible mental retardation and cretinism (2, 7, 8). If iodine deficiency occurs during the most critical period of brain development (from the fetal stage up to the third month after birth), the resulting thyroid failure will lead to irreversible alterations in brain function (9, 10). In severely endemic areas, cretinism may affect up to 5–15% of the population. A meta-analysis

Table 1.1 *The spectrum of IDD across the life-span*

Fetus	Abortions Stillbirths Congenital anomalies Increased perinatal mortality Endemic cretinism Deaf mutism
Neonate	Neonatal goitre Neonatal hypothyroidism Endemic mental retardation Increased susceptibility of the thyroid gland to nuclear radiation
Child and adolescent	Goitre (Subclinical) hypothyroidism (Subclinical) hyperthyroidism Impaired mental function Retarded physical development Increased susceptibility of the thyroid gland to nuclear radiation
Adult	Goitre, with its complications Hypothyroidism Impaired mental function Spontaneous hyperthyroidism in the elderly Iodine-induced hyperthyroidism Increased susceptibility of the thyroid gland to nuclear radiation

Source: Adapted with permission of the publisher, from Hetzel (2), Laurberg et al. (3) Stanbury et al. (4).

of 19 studies conducted in severely iodine deficient areas showed that iodine deficiency is responsible for a mean IQ loss of 13.5 points in the population (10). While cretinism is the most extreme manifestation, of considerably greater significance are the more subtle degrees of mental impairment leading to poor school performance, reduced intellectual ability and impaired work capacity (7).

1.1.3 Indicators for assessment and monitoring

Several indicators are used to assess the iodine status of a population: thyroid size by palpation and/or by ultrasonography, urinary iodine (UI) and the blood constituents, TSH or thyrotropin, and thyroglobulin.

Until the 1990s total goitre prevalence (TGP)¹ was recommended as the main indicator to assess IDD prevalence. However, TGP is of limited utility in assessing the impact of salt iodization. In endemic areas, TGP may not return to normal for months or years after correction of iodine deficiency. During this period, TGP is a poor indicator because it reflects a population's history of iodine nutrition but not its present iodine status. TGP is still useful to assess the severity of IDD at baseline and has a role in evaluating the long term impact of control programmes.

As UI is a more sensitive indicator to recent changes in iodine intake, it is now recommended over TGP (12). Most countries have started to implement IDD control programmes, and a growing number of countries are consequently monitoring iodine status using UI.

TSH levels in neonates are particularly sensitive to iodine deficiency however difficulties in interpretation remain and the cost of implementing a TSH screening programme is high. The value of thyroglobulin as an indicator of global IDD status has yet to be fully explored.

While IDD affects the entire population, a school-based sampling method is recommended for UI and TGP as the most efficient and practical approach to monitor IDD as this group is usually easily accessible and can be used as a proxy for the general population (12). School-age refers to children aged 6–12 years, hereafter referred to as school-age children unless otherwise noted. Iodine deficiency is considered to be a public health problem in populations of school-age children where the median UI is below 100 µg/l (see table 2.1) or goitre prevalence is above 5% (12).

1.2 Control of IDD

The recommended strategy for IDD control is based on correcting the deficiency by increasing iodine intake through supplementation or food fortification. Four main components are required to implement the strategy: correction of iodine deficiency, surveillance including monitoring and evaluation, inter-sectorial collaboration and advocacy and communication to mobilize public health authorities and educate the public.

1.2.1 Correcting iodine deficiency

1.2.1.1 Iodine supplementation

The first iodine supplements were in the form of an oral solution of iodine such as Lugol, which was given daily. After the Second World War, considerable progress was made in reducing IDD with iodized oil – initially using the intramuscular form and in the 1990s, using the oral form. For example, iodized oil was used with success in Papua New Guinea and thereafter in China, several countries in Africa

¹ In this report “total goitre prevalence” is used instead of “total goitre rate” (TGR) to be in agreement with the terminology that is universally used in epidemiology (11).

and Latin America and in other severely endemic areas.

The oral form of iodized oil has several advantages over the intramuscular form: it does not require special storage conditions or trained health personnel for the injection and it can be given once a year. Compared to iodized salt, however, it is more expensive and coverage can be limited since it requires direct contact with each person. With the introduction of iodized salt on a large scale, iodized oil is now only recommended for populations living in severely endemic areas with no access to iodized salt.

1.2.1.2 Food fortification with iodine

Over the past century, many food vehicles have been fortified with iodine: bread, milk (13), water (14) and salt. Salt is the most commonly used vehicle. It was first introduced in the 1920s in the United States (15) and in Switzerland (16). However, this strategy was not widely replicated until the 1990s when the World Health Assembly adopted universal salt iodization (USI) (the iodization of salt for both human and livestock consumption) as the method of choice to eliminate IDD. In 2002, at the Special Session on Children of the United Nations (UN) General Assembly, the goal to eliminate IDD by the year 2005 was set.

USI was chosen as the best strategy based on the following facts: (i) salt is one of the few commodities consumed by everyone; (ii) salt consumption is fairly stable throughout the year; (iii) salt production is usually in the hands of few producers; (iv) salt iodization technology is easy to implement and available at a reasonable cost (0.4 to 0.5 US cents/kg, or 2 to 9 US cents per person/year); (v) the addition of iodine to salt does not affect its colour, taste or odour; (vi) the quality of iodized salt can be monitored at the production, retail and household levels; and (vii) salt iodization programmes are easy to implement.

In order to meet the iodine requirements of a population it is recommended to add 20 to 40 parts per million (ppm) of iodine to salt (assuming an average salt intake of 10 g per capita/day) (17). There are two forms of iodine fortificants, potassium iodate and potassium iodide. Because iodate is more stable under extreme climatic conditions it is preferred to iodide, especially in hot and humid climates (17). For historical reasons, North America and some European countries use potassium iodide while most tropical countries use potassium iodate.

1.2.1.3 Safety in approaches to control iodine deficiency

Iodine fortification and supplementation are safe if the amount of iodine administered is within the recommended range. For more than 50 years iodine has been added to salt and bread without noticeable toxic effects (18). However, a rapid increase in iodine intake can increase the risk of iodine toxicity in individuals who have previously had chronic iodine deficiency.

Iodine-induced hyperthyroidism (IIH) is the most common complication of iodine prophylaxis and it has been reported in almost all iodine supplementation programmes in their early phases (4). For programmes using iodized salt, there is less information. IIH occurs in the early phase of the iodine intervention and primarily affects the elderly who have longstanding thyroid nodules. However, it is transient and its incidence reverts to normal after one to ten years. Monitoring of salt quality and iodine status of populations, and training of health staff in identification and treatment of IIH are the most effective means for preventing IIH and its health consequences (19).

1.2.2 Monitoring and evaluating the IDD control programmes

1.2.2.1 Monitoring iodine levels of salt

Governments usually set the level at which salt should be iodized. Monitoring aims to ensure that the salt industry complies with the regulations set by the government and that the iodine levels are re-adjusted if necessary. Iodine levels are monitored (at a minimum) at the factory and household levels, and if possible at the retail level. If iodized salt is imported it is monitored at the point of entry into the country. The monitoring process at the factory level is the salt producer's or importer's responsibility and is regularly supervised by the relevant public authorities. In most cases the Ministry of Health carries out the monitoring at the household level.

Iodine content in salt is best measured by titration. Field test kits have been developed. They only give qualitative results, indicating if iodine is present or not. Because of this, they are of limited use, moreover their reliability has recently been questioned. However, they are can still be useful for training educating and for advocacy purposes for the public and staff.

1.2.2.2 Monitoring of iodine status

When salt is adequately iodized, it is likely that a population's iodine status will improve and the thyroid function of that population will normalize. Monitoring the population's iodine status is nevertheless necessary since dietary habits may change in some segments of the population or the iodine level of salt may not be sufficient to meet the requirements of some groups, in particular pregnant women.

Indicators used to monitor iodine status are described in section 1.1.3.

1.2.3 Increasing awareness of public health authorities and the general public

WHO has played a pioneer role in mobilizing the international community and public health authorities by providing strategic guidance and technical support. In 1990, the World Health Assembly adopted a resolution urging Mem-

ber States to take the appropriate measures to eliminate IDD. This goal was reaffirmed in a series of subsequent international fora including: the 1990 World Summit for Children (New York), the Joint WHO/Food and Agricultural Organization of the United Nations (FAO) and the International Conference on Nutrition in 1992 (Rome) and the Special Session on Children of the UN General Assembly in 2002 (New York).

This commitment catalysed the involvement of a large number of additional actors. The United Nations Children's Fund (UNICEF) was one of the first organizations to assist countries in establishing salt iodization programmes and still now plays a leading role in this regard. The International Council for the Control of Iodine Deficiency Disorders (ICCIDD) played an instrumental role in providing technical support. Other important actors were the bilateral co-operation agencies, non-governmental organizations (NGOs) such as the Micronutrient Initiative, the salt industry, and donor foundations such as Kiwanis International and the Bill and Melinda Gates Foundation.

1.2.4 Reinforcing the collaboration between sectors

1.2.4.1 Network for sustainable elimination of iodine deficiency

Effective IDD control demands collaboration and clearly the salt industry has a major role to play by iodizing salt and ensuring its delivery to regions worldwide. To facilitate the participation and co-ordination of the salt industry as well as other sectors in IDD control, the Global Network for Sustained Elimination of Iodine Deficiency was established in 2002. Further information on this network can be obtained on the Internet: <http://www.sph.emory.edu/iodinenetwork/>.

In most countries where iodine deficiency is a public health problem, a national multi-sectoral IDD body has been established, usually chaired by the Ministry of Health (20). Its main roles are to design and supervise the implementation of an IDD control plan and to coordinate the activities of the various sectors and partners involved. It acts in concert with the national and international partners involved in IDD control.

1.2.4.2 International resource laboratory network

Where iodine deficiency is a public health problem, laboratory facilities to measure the indicators required to monitor the programme are often insufficient or lacking altogether. To overcome this problem, the International Resource Laboratories for Iodine (IRLI) Network has been created, under the coordination of the Centers for Disease Control and Prevention (CDC), WHO, UNICEF, the Micronutrient Initiative and ICCIDD. The main role of this network is to provide technical support to national laboratories which may need assistance through regional or subregional

resource laboratories in monitoring their IDD control programmes. In every WHO region at least one resource laboratory has been identified. Further information can be obtained at IRLI's web site: <http://www.cdc.gov/nceh/dls/iodinelabnetwork.htm>.

1.2.5 Sustaining IDD control programmes

In order to achieve the global goal set for 2005, IDD control programmes and monitoring need to be constantly

sustained due to the fact that IDD simply re-appears if salt iodization is interrupted. This may happen when the responsible public health authorities are demobilized or if the salt industry fails to effectively monitor iodine content. In order to assess the sustainability of control programmes and track their progress towards the IDD elimination goal, criteria have been established by WHO (Table 1.2).

Table 1.2 **Criteria for monitoring progress towards sustainable IDD elimination**

Indicators	Goals
<p><i>Salt iodization coverage</i></p> <ul style="list-style-type: none"> ■ Proportion of households consuming adequately iodized salt^a 	>90%
<p><i>Urinary iodine</i></p> <ul style="list-style-type: none"> ■ Proportion of population with urinary iodine levels below 100 µg/l ■ Proportion of population with urinary iodine levels below 50 µg/l 	<p><50%</p> <p><20%</p>
<p><i>Programmatic indicators</i></p> <ul style="list-style-type: none"> ■ National body responsible to the government for IDD elimination. It should be multidisciplinary, involving the relevant fields of nutrition, medicine, education, the salt industry, the media, and consumers, with a chairman appointed by the Minister of Health; ■ Evidence of political commitment to USI and elimination of IDD; ■ Appointment of a responsible executive officer for the IDD elimination programme; ■ Legislation or regulation of USI; ■ Commitment to regular progress in IDD elimination, with access to laboratories able to provide accurate data on salt and urinary iodine; ■ A programme of public education and social mobilization on the importance of IDD and the consumption of iodized salt; ■ Regular data on iodized salt at the factory, retail and household levels; ■ Regular laboratory data on urinary iodine in school-age children, with appropriate sampling for higher-risk areas; ■ Co-operation from the salt industry in maintenance of quality control; and ■ A database for recording results or regular monitoring procedures particularly for salt iodine, urinary iodine and, if available, neonatal thyroid stimulating hormone (TSH), with mandatory public reporting. 	At least 8 of the 10

Source: WHO et al. (12).

^a Adequately iodized salt refers to at least 15 ppm at household level

2. Methods

This report provides estimates of the current worldwide situation of iodine nutrition based on UI data collected between 1993 and 2003. It is a continuation of the previous report on the global prevalence of IDD published by WHO in 1993 (21). For comparison purposes with the IDD estimates in 1993 (21), the present report provides estimates of IDD based on TGP in addition to UI.

2.1 Data sources – The WHO Global Database on Iodine Deficiency

The estimates presented are based on the data available in the WHO Global Database on Iodine Deficiency, accessible on the Internet: <http://www3.who.int/whosis/micronutrient/>. This database compiles country data on UI and TGP and presents it in a standardized and easily accessible format.

Data are collected from the scientific literature and through a broad network of collaborators, including WHO regional and country offices, United Nations organizations, non-governmental organizations, ministries of health, other national institutions, and research and academic institutions. MEDLINE and regional databases (African Index Medicus, Index Medicus for the WHO Eastern Mediterranean Region, Latin American and Caribbean Center on Health Sciences Information, Pan American Health Organization Library Institutional Memory Database, Index Medicus for South-East Asia Region) are systematically searched. Articles published in non-indexed medical and professional journals and reports from principal investigators are also systematically looked for. Data are extracted from reports written in any language.

For inclusion in the database, a complete original survey report providing details of the sampling method used is necessary. Studies must have a population-based sample frame and must use standard UI and TGP measuring techniques (21). Only TGP data measuring goitre by palpation are included. Until recently no international reference values for thyroid size measured by ultrasonography were available, and thus results from surveys using this technique have not yet been included (22).

When a potentially relevant survey is identified and

the full report obtained, all data are checked for consistency as part of routine quality control. When necessary, the authors are contacted for clarification or additional information. Final data are extracted and entered into a standard data form. The full archived documentation and correspondence are available on request.

As of June 2003, the database contained 389 UI surveys and 409 goitre surveys. Surveys received at WHO after this date were not included in this analysis but are available in the online database and will be included in future analysis.

2.2 Selection of survey data

Data collected between 1993 and 2003, available to WHO in June 2003, were reviewed for WHO's 192 Member States. Data on UI and TGP were selected for each country using two variables: the administrative level for which the population sample is representative (national or sub-national) and the population groups surveyed (school-age children or other).

2.2.1 Administrative level

Surveys were first selected according to the administrative level. Surveys are considered as national level when they are carried out on a nationally representative sample of the population group surveyed, or as sub-national level when they are carried out on a sample representative of a given administrative level: region, state, province, district or local.

Whenever available, data from the most recent national survey were used in preference to sub-national surveys. WHO recommends that iodine status is regularly assessed (12). Thus, if a national survey was 5 years old or more, and more recent sub-national data were available, preference was given to the sub-national data.

In the absence of national data, sub-national data were used. When two or more sub-national surveys of the same sub-national level had been carried out in different locations in a country during the analysis period, the survey results were pooled into a single summary measure, using a weighted sample size for each survey. When, in a few cases,

sample size information was missing for one sub-national survey, it was assumed to have a number of subjects equal to the average sample size of the other surveys included in the pooling. For one country, sample size information was missing from all data pooled, and thus unweighted average was computed. Exceptionally, data from different sub-national levels were pooled, for example a survey carried in the capital city, classified as local, with a district level survey.

2.2.2 Population groups

WHO recommends that iodine deficiency surveys examine school-age children from 6 to 12 years (12). When data for this age group were not available, data of the next closest age group were used in the following order of priority: data from the children closest to school age, adults, the general population, preschool-age children, other population groups.

2.3 Classification of iodine nutrition

Median UI of the distribution was used to classify countries into different degrees of public health significance. Since UI values from populations are usually not normally distributed, the median rather than the mean is used as a measure of central tendency (12). The WHO cut-off points applied for classifying iodine nutrition into different degrees of public health significance are shown in Table 2.1. Median UI below 100 µg/l define a population which has iodine deficiency.

If a national median UI was not available for the severity classification the following methods were applied to derive median UI from various UI data.

1. When UI means were the only available data, UI medians were derived through simple linear regression using the equation:

$$\text{Median} = 1.128 + 0.864 * \text{Mean}$$

This equation was obtained from a model based on surveys available in the database which presented both median UI and mean UI. To perform this linear regression, disaggregated data were introduced for each survey, i.e. sub-samples of the same survey stratified by age, sex or region. A total of 351 regression points were identified. The relation is shown in Figure 2.1.

2. When only disaggregated UI medians were presented (e.g. UI medians for each age, sex or region), aggregated total median UI was estimated using the following procedure:

Step 1: disaggregated UI means from disaggregated UI medians were derived, assuming linearity through linear regression using the equation:

$$\text{Mean} = 7.447 + 1.081 * \text{Median}$$

These were obtained from the 351 pairs of points used in paragraph 1.

Step 2: the average of the means thus obtained was computed, weighted by the sample size of each group.

Step 3: median UI corresponding to the total survey population through linear regression was calculated using the equation:

$$\text{Median} = 1.128 + 0.864 * \text{Mean}$$

explained in paragraph 1 above.

Proceeding in this manner was necessary because medians, unlike means, when pooled directly, give rise to erroneous results when the distribution of data is not normal but skewed as in the case of UI.

3. When the proportion of UI values below 100 µg/l (% UI <100 µg/l) was the only available information on UI, median UI was derived through a quadratic regression using the following equation:

Table 2.1 **Epidemiological criteria for assessing iodine nutrition based on median UI concentrations in school-age children**

Median UI (µg/l)	Iodine intake	Iodine nutrition
< 20	Insufficient	Severe iodine deficiency
20–49	Insufficient	Moderate iodine deficiency
50–99	Insufficient	Mild iodine deficiency
100–199	Adequate	Optimal iodine nutrition
200–299	More than adequate	Risk of iodine-induced hyperthyroidism within 5–10 years following introduction of iodized salt in susceptible groups
≥ 300	Excessive	Risk of adverse health consequences (iodine induced hyperthyroidism, auto-immune thyroid diseases)

Source: WHO et al. (12).

$$\text{Median} = 277.670 - 4.96 * (\% \text{ UI } < 100 \text{ } \mu\text{g/l}) + 0.0254 * (\% \text{ UI } < 100 \text{ } \mu\text{g/l})^2$$

The quadratic regression model was computed from 408 pairs of points obtained from surveys, performed all over the world and included in the database, where data were available on both medians and proportions of UI values below 100 µg/l for the same population group. All populations groups used in the computation above were mutually exclusive. The quadratic regression was the model that best described the relation between median UIs and proportions of UI values below 100 µg/l ($r^2 = 0.83$) and was preferred to the simple linear regression model ($r^2 = 0.76$). The relation is shown in Figure 2.2.

Note that when the equation above is used, a proportion of UI values below 100 µg/l of 50% yield a median figure of 102 µg/l, instead of the expected value of 100 µg/l. In this case, no attempt to modify the intercept or the slope of the equation was made in order to make it fit predicted values. Instead, this caveat is mentioned in the “notes” section of Table 4.1, where applicable.

Countries with high medians (>300 µg/l) were given zero per cent as a proportion of UI values below 100 µg/l, and not the value predicted by the equation.

2.4 Population coverage, proportion of population and the number of individuals with insufficient iodine intake

2.4.1 Population coverage

The coverage of the estimates for a given WHO region was calculated as the sum of the populations of countries with data divided by the total population of the region. The same procedure was used to calculate global coverage.

2.4.2 Proportion of population and the number of individuals with insufficient iodine intake

National, regional and global populations (school-age children and general population) with insufficient iodine intake was estimated based on each country’s proportion of population with UI below 100 µg/l. The following method was used:

1. The number of subjects with insufficient iodine intake at the country level was calculated by applying the proportion of population with UI below 100 µg/l to the national population of both children aged 6–12 years and general population (all age groups including children aged 6–12 years). The population figures are based on the year 2002 (23).

If the proportion of population with UI values below 100 µg/l was not presented, it was computed from

Figure 2.1 Relation between median UI (µg/l) and mean UI (µg/l) with linear regression line





Figure 2.2 Relation between median UI (µg/l) and proportion (%) of UI values below 100 µg/l with quadratic regression curve



median UI, using the simple linear regression equation based on the data points presented in Figure 2.2:

$$\% \text{ UI } < 100 \text{ } \mu\text{g/l} = 86.3 - 0.324 * \text{Median}$$

The 95% confidence intervals of for the proportion of a population with UI below 100 µg/l for each country are presented as a measure of uncertainty (Table A3.1).

2. The number of subjects with insufficient iodine intake at the regional level was calculated by summing the number of individuals with UI below 100 µg/l in each country of the region and dividing the sum by the total population of all countries with available data. The calculations were made for both WHO and UN regions.
3. The global estimate was calculated by summing the number of individuals with insufficient iodine intake in each region and dividing the sum by the total population of all countries with data available.

2.5 TGP

TGP was computed from data in school-age children. In order to compare present TGP data, with the 1993 TGP estimates which were generated for the general population (21), it was necessary to calculate current TGP estimates for the general population. To that end, an algorithm was developed from surveys available in the database that measured prevalence in both population groups.

Eight countries were found to have carried out such surveys between 1993 and 2003: Burkina Faso, Ethiopia, France (the island of Réunion), Guinea-Bissau, India, Islamic Republic of Iran, Italy and the Philippines. A total of 23 pairs of points corresponding to different population subgroups were included.


Assuming linearity in the data range, the equation of the linear regression model that predicts TGP in the general population from school age children TGP is:

$$\text{General population TGP (\%)} = 0.954 + 0.742 * \text{school-age children TGP (\%)}$$

The graph that displays the relation between the two sets of prevalences is shown in Figure 2.3 above.

Goitre prevalences by country, region (both WHO and

Figure 2.3 Relation between general population TGP and school-age children TGP with linear regression line



UN) and worldwide were derived for the general population, applying the algorithm described above at country level, following the same procedure as described for UI calculations (section 2.5.2).

Along with the point estimates of TGP, 95% confidence intervals of TGP for each country are presented as a measure of uncertainty (Table A3.2).

3. Results and discussion

3.1 Results

Regional and worldwide estimates of iodine status are based on data from 192 WHO Member States (Annex 1). Estimates by WHO region are presented in this chapter; estimates by UN region appear in Annex 2. National estimates of iodine status for each WHO Member State are presented in Annex 3.

3.1.1 Population coverage

3.1.1.1 UI surveys

Data on UI collected between 1993 and 2003 were available from 126 countries. Sixty-six countries have no data on UI.

Table 3.1 presents the population coverage for the age group 6–12 years based on UI data by WHO region. The number of countries with national and, if not available, sub-national UI survey data is shown in Table 3.2. Figure 3.1 shows the worldwide coverage of national and sub-national UI surveys.

Overall, the available UI data covers 92.1% of the world's 6–12 year old population. Regional population coverage varies from 83.4% in the Eastern Mediterranean to 98.8% in South-East Asia.

Table 3.1 Population coverage^a by UI surveys carried out between 1993 and 2003, by WHO region

WHO region ^b	Total number of school-age children (millions) ^c	School-age children covered (millions)	Coverage (%)
Africa	128.9	116.9	90.7
Americas	109.0	98.8	90.6
South-East Asia	242.4	239.4	98.8
Europe	81.2	70.5	86.8
Eastern Mediterranean	87.1	72.6	83.4
Western Pacific	199.4	183.0	91.8
Total	848.0	781.2	92.1

^a School-age children (6–12 years).

^b 192 WHO Member States.

^c Based on population estimates for the year 2002 (23).

Table 3.2 Type of UI survey data by WHO region

WHO region ^a	National	Sub-national	No data
Africa	17	17	12
Americas	13	7	15
South-East Asia	7	2	2
Europe	21	17	14
Eastern Mediterranean	11	4	6
Western Pacific	6	4	17
Total	75	51	66

^a 192 WHO Member States.

Figure 3.1 Type of UI survey data




Figure 3.2 Type of TGP survey data




Table 3.3 Population coverage^a by TGP surveys carried out between 1993 and 2003, by WHO region

WHO region ^b	School-age children (millions) ^c	School-age children covered (millions)	Coverage (%)
Africa	128.9	117.6	91.2
Americas	109.0	50.7	46.5
South-East Asia	242.4	232.1	95.7
Europe	81.2	46.9	57.8
Eastern Mediterranean	87.1	76.5	87.8
Western Pacific	199.4	184.0	92.3
Total	848.0	707.7	83.5

^a School-age children (6–12 years).

^b 192 WHO Member States.

^c Based on population estimates for the year 2002 (23).

Table 3.4 Type of TGP survey data by WHO region

WHO region ^a	National	Sub-national	No data
Africa	18	17	11
Americas	4	3	28
South-East Asia	7	1	3
Europe	12	15	25
Eastern Mediterranean	10	5	6
Western Pacific	6	2	19
Total	57	43	92

^a 192 WHO Member States.

Of the 126 countries with data available on UI, 75 have nationally representative surveys covering 45.7% of the school-age children population.

3.1.1.2 TGP surveys

Data on goitre collected between 1993 and 2003 were available from 100 countries. Table 3.3 presents the population coverage for the age group 6–12 years based on TGP data by WHO region. Table 3.4 presents the number of countries with national and, if not available, sub-national surveys. Figure 3.2 shows the worldwide coverage of national and sub-national TGP surveys.

Population coverage for TGP surveys is 83.5%, ranging from 46.5% in the Americas to 95.7% in South-East Asia. Of the 100 countries with data available on TGP, 57 had nationally representative surveys, covering 43.4% of the school-age children population.

3.1.2 Classification of countries by degree of public health significance of iodine nutrition based on median UI

In Figure 3.3, countries are classified into six different degrees of public health significance with respect to their iodine nutrition estimated from median UI. Table 3.5 shows the number of countries classified by degree and by WHO region.

In 54 countries the population has insufficient iodine intake as indicated by a median UI below 100 µg/l. These countries are classified as iodine deficient: one country is severely deficient, 13 are moderately deficient and 40 mildly deficient. In 43 countries, the population have adequate iodine intake with a median UI between 100 and 199 µg/l. Iodine nutrition of these countries is considered as optimal. In 24 countries, median UI is between 200

Figure 3.3 Degree of public health significance of iodine nutrition based on median UI




Table 3.5 Number of countries classified by degrees of public health significance of iodine nutrition based on median UI in school-age children by WHO region, 2003

WHO region ^a	Classification of iodine nutrition						No data
	Severe iodine deficiency (Median UI <20 µg/l)	Moderate iodine deficiency (Median UI 20–49 µg/l)	Mild iodine deficiency (Median UI 50–99 µg/l)	Optimal iodine nutrition (Median UI 100–199 µg/l)	Risk of IIH in susceptible groups (Median UI 200–299 µg/l)	Risk of adverse health consequences (Median UI ≥300 µg/l)	
Africa	0	6	8	11	7	2	12
Americas	0	1	1	3	12	3	15
South-East Asia	0	0	3	5	1	0	2
Europe	0	4	19	15	0	0	14
Eastern Mediterranean	1	0	5	6	3	0	6
Western Pacific	0	2	4	3	1	0	17
Total	1	13	40	43	24	5	66

^a 192 WHO Member States.

and 299 µg/l indicating that the population has more than adequate iodine intake. In these countries, there is a risk of iodine-induced hyperthyroidism in susceptible groups. In 5 countries, there is excessive iodine intake as shown by a median UI above 300 µg/l. In these countries, there is a risk of iodine-induced hyperthyroidism and other adverse health consequences.

3.1.3 Proportion of population and number of individuals with insufficient iodine intake

The proportion of the population and the number of individuals (school-age children and general population) with insufficient iodine intake (defined as proportion of population with UI below 100 µg/l) by WHO region is presented in Table 3.6.

Table 3.6 Proportion of population, and number of individuals with insufficient iodine intake in school-age children (6–12 years), and in the general population (all age groups) by WHO region, 2003

WHO region ^a	Insufficient iodine intake (UI <100 µg/l)			
	School-age children		General population	
	Proportion (%)	Total number (millions) ^b	Proportion (%)	Total number (millions) ^b
Africa	42.3	49.5	42.6	260.3
Americas	10.1	10.0	9.8	75.1
South-East Asia	39.9	95.6	39.8	624.0
Europe	59.9	42.2	56.9	435.5
Eastern Mediterranean	55.4	40.2	54.1	228.5
Western Pacific	26.2	48.0	24.0	365.3
Total	36.5	285.4	35.2	1988.7

^a 192 WHO Members States.

^b Based on population estimates in the year 2002 (23).

It is estimated that the iodine intake of 36.5% (285 million) school-age children worldwide is insufficient. Extrapolating the proportion of school-age children to the general population, it is estimated that nearly two billion individuals have insufficient iodine intake.

The most affected region is South-East Asia where 96 million children have a low iodine intake. Africa and the Western Pacific follow, both with an estimated 50 million children with a low iodine intake. Europe and the Eastern Mediterranean harbour about 40 million children each, and the Americas have 10 million. The highest proportions are found in Europe (59.9%) and South-East Asia (39.9%) while the lowest are found in the Americas (10.1%) and the Western Pacific (26.2%).

3.1.4 TGP

Goitre prevalence in the general population is presented with the purpose of comparing the current estimate with that of 1993 (21) (Table 3.7).

Globally, the TGP in the general population is estimated

Table 3.7 Change in total goitre prevalence between 1993 and 2003, by WHO region

WHO region ^a	TGP (%)		
	General population		% change
	1993	2003	
Africa	15.6	28.3	+ 81.4
Americas	8.7	4.7	- 46.0
South-East Asia	13.0	15.4	+ 18.5
Europe	11.4	20.6	+ 80.7
Eastern Mediterranean	22.9	37.3	+ 62.9
Western Pacific	9.0	6.1	- 32.2
Total	12.0	15.8	+ 31.7

^a 192 WHO Member States.

to be 15.8%, varying between 4.7% in the Americas to 28.3% in Africa. When comparing current TGP estimates with the 1993 estimates, TGP has increased by 31.7% worldwide. This masks a decrease in two regions of 46.0% in the Americas and 32.2% in the Western Pacific. All other regions experienced an increase in TGP ranging from 18.5% in South-East Asia to 81.4% in Africa.

3.2 Discussion

Data gathered in the WHO Global Database on Iodine Deficiency permit a description to be made of the magnitude, severity and distribution of iodine deficiency worldwide and facilitates decisions on the most effective strategy to eliminate iodine deficiency.

3.2.1 Population coverage

Estimates of iodine nutrition were calculated based on UI data available from 126 countries representing 92.1% of the world's population of school-age children (Table 3.1). The current estimates are thus believed to be a true reflection of the situation. The remaining 66 countries lacking data, or lacking recent data, represent only 7.9% of the world's school-age population. However, the risk of iodine deficiency is unlikely to be a public health problem in many of these countries.

3.2.2 Limitations of data sources

Estimates presented are subject to several limitations. Sixty percent of the 126 countries with data have nationally representative surveys; the remainder have only one or several sub-national surveys. The lack of nationally representative surveys may lead to substantial bias. Under estimation may occur when parts of a country which may have an inadequate iodine intake, have not been surveyed. Over estimation may occur when the population of one or more endemic regions is over-sampled. The data for some countries are still weak which makes their classification and accurate analysis of their national situation difficult. For example, for India and Spain the only available data are sub-national, which pooled show optimal iodine nutrition. In the absence of national representative data the entire country has therefore been classified accordingly, when in fact the situation might be very different. Thus, the methods used for pooling sub-national survey results into one summary measure are not perfect. Nevertheless, they are regarded as the best estimate in the absence of nationally representative data.

The data compiled in the database are extracted from final publications and reports, which present data in various formats and with varying degrees of analysis. The models developed to standardize the data and derive one measure from another are a potential source of error. Raw data sets are not available in the database and thus render any further verification impossible.

When the proportion of population with UI below 100 µg/l was missing for a particular country, the equation presented in section 2.3 was used to produce a value for this variable from the value of the median. Because the model is based on real data, no manipulation (changes in slope or intercept) was carried out in the equation that predicts the proportion of population below 100 µg/l from the median UI, to adapt it to expected predictions. The results may then slightly depart from expected values. In accordance with current knowledge on modelling, in the case that the predicted results lead to mistaken classification, this was mentioned in the "notes" section of Annex 3, Table A3.1. This problem only arose for one country, for which the predicted value of the proportion of population with UI below 100 µg/l was 50.1% (hence classifying this country as iodine deficient) while the median was 109 µg/l.

These limitations highlight the need to improve data quality. It is important for countries to conduct nationally representative surveys on a regular basis and ensure representative samples. Standardized data collection and presentation will also aid the comparison of countries and regions, allow for more precise monitoring and a lower level of uncertainty around future global estimates of iodine nutrition.

3.2.3 Classification of countries by degree of public health significance of iodine nutrition based on median UI

Iodine nutrition is optimal in 43 countries (Table 3.5). The number of countries with iodine deficiency as a public health problem decreased from 110 to 54 between 1993 (using TGP as an indicator) and 2003 (using UI). Nevertheless, in 54 countries, located in all regions of the world, the iodine intake of the population is insufficient and iodine deficiency with its impact on health and development is still a public health concern. In these countries USI needs to be strengthened and fully implemented.

Iodine intake is more than adequate, with a median UI between 200 and 299 µg/l, in 24 countries. Here attention should be drawn to the emerging risk of iodine-induced hyperthyroidism in susceptible groups following introduction of iodized salt.

Five countries have a median UI equal to or above 300 µg/l indicating an excessive iodine intake and are therefore exposed to the risk of hyperthyroidism and iodine toxicity. Elevated median UI is most likely due to high levels of iodine added to salt. Salt quality monitoring should be re-enforced to ensure that the level of salt fortification with iodine is not too high but is adequate to ensure optimal iodine nutrition.

Sixty-six countries have no data on UI and iodine nutrition can therefore not be classified. Even though iodine deficiency is unlikely in many of these countries, urinary

iodine surveys should be performed in order to investigate the level of iodine intake and evaluate the effectiveness of pre-existing salt iodization programmes. There is evidence that iodine deficiency may be re-emerging in countries that were previously thought to be iodine sufficient, like Australia and New Zealand (Annex 3, Table A3.1).

3.2.4 Proportion of population and the number of individuals with insufficient iodine intake

Overall, one third of the world's school-age children population has UI below 100 µg/l indicating insufficient iodine intake (Table 3.6). This group is therefore exposed to the risk of iodine deficiency.

For the six WHO regions the proportion of the population with UI below 100 µg/l ranges from 10% (in the Americas) to 60% (in Europe).

Noteworthy is the correlation between household coverage of iodized salt and prevalence of low iodine intake. The proportion of households consuming iodized salt increased from 10% in the 1990s (20) to 66% in the year 2003 (24).

The Americas has the highest number of households consuming iodized salt (90%) and the lowest proportion of its population with an insufficient iodine intake. In contrast, the European Region which has the lowest household consumption of iodized salt (27%), has the highest proportion of its population with an insufficient iodine intake (20). These results, however, should not mask the fact that there are large variations both between countries within regions, and within countries themselves.

WHO recommends that school-age children are surveyed to assess iodine status because they are readily accessible and their iodine status is an acceptable proxy for the iodine status of the general population. Results of surveys of school-age children were thus extrapolated to the general population (Table 3.6). However, it has recently been recognized that national systems to monitor the impact of USI also need to include other vulnerable groups, especially pregnant women. Data for this population group may be considered for future global analysis as more data become available.

3.2.5 TGP

The worldwide TGP of 15.8% is above the 5% cut-off used to signal a public health problem (12). Its increase of 31.7% between 1993 and 2003 is inconsistent with current iodine status based on UI. This has several possible explanations.

First, there is a time lag between the implementation of a salt iodization programme and the disappearance of clinically detectable goitre (25). This time-lag may be further increased when USI is only partially implemented.

Second, 70% of the TGP surveys in the analysis period 1993–2003 were carried out between 1993 to 1998, which

was prior to extensive implementation of USI programmes. In fact, when analysis is restricted to surveys carried out in the last five years, TGP shows a decrease of 28.9% compared to 1993. Analysis of data available in the WHO database also shows that between 1993 and 1998 TGP was the main indicator used to assess iodine deficiency, while UI was measured only in a few countries. The shift in indicators from TGP to UI over the last decade resulted in less TGP data covering the last five years since many countries only measured UI in their most recent surveys.

Third, in areas affected by mild iodine deficiency, the sensitivity and specificity of TGP measured by palpation are poor (26). Ultrasonography is a promising method to overcome the inherent limitations of the clinical assessment of thyroid volume as iodine status improves. New international reference values are now available allowing comparison between countries (22). In spite of its limitations for global trend analysis TGP measured by palpation remains a practical indicator for baseline assessment, especially in severely endemic areas (12, 27).

3.3 Conclusion

In conclusion, there has been substantial progress in the last decade towards the elimination of iodine deficiency. Improved iodine nutrition reflects the validity of the strategy adopted by WHO based on salt iodization complemented with iodine supplementation in remote areas not reached by iodized salt or in population groups who are severely deficient. It reflects the efforts made by countries to implement effective IDD control programmes and is proof of the successful collaboration between all the partners in IDD control, in particular the health authorities and the salt industry.

Having said that, every effort needs to be made to ensure that programmes continue to cover at-risk populations if the goal of eliminating IDD is to be reached. Current iodine deficiency estimates based on UI provide the baseline for future global estimates. The challenge now is to improve the quality of the data in order to trigger appropriate and timely interventions and to track progress more accurately and rapidly.

With regard to the WHO Global Database on Iodine Deficiency and the measurement of iodine nutrition, attention must be drawn to the following issues.

- It is important that each country carries out nationally representative surveys on a regular basis. Efforts should be made to ensure that samples are representative (i.e. to make sure that no region of a given country is deliberately excluded from the sampling procedure). Countries where evaluation data are missing introduce considerable uncertainty as to the impact of the iodization efforts.

- National data may not reflect the presence of pockets of iodine deficiency in some parts of the country. Additional closer monitoring might be required.
- UI is the most reliable indicator to assess, monitor and evaluate iodine status in a population. At the assessment stage, clinical detection of goitre may be useful but it should always be associated with the measurement of urinary iodine. Neonatal TSH screening may be useful if a reliable system is already in place and the resources are available.
- To improve the reliability of goitre data, thyroid volume can be measured by ultrasonography. For comparison between countries and regions results from surveys measuring thyroid size by ultrasound should apply the new international reference values for thyroid volume measured by ultrasonography (22).
- Efforts to enforce a standardized approach in presenting data are important for cross-comparisons between countries and regions. This will reduce the level of uncertainty of future global estimates of iodine deficiency. It will also aid in the monitoring of IDD and management of USI programmes.
- The quality of data also depends on the capacity of national laboratories to carry out reliable measurements of UI. The IRLI network provides country support.

References

1. Delange F. The disorders induced by iodine deficiency. *Thyroid*, 1994, 4:107–128.
2. Hetzel BS. Iodine deficiency disorders (IDD) and their eradication. *Lancet*, 1983, 2:1126–1129.
3. Laurberg P et al. Thyroid disorders in mild iodine deficiency. *Thyroid*, 2000, 10:951–963.
4. Stanbury JB et al. Iodine-induced hyperthyroidism: occurrence and epidemiology. *Thyroid*, 1998, 8: 83–100.
5. Delange F. Adaptation to iodine deficiency during growth: Etiopathogenesis of endemic goiter and cretinism. In: Delange F, Fisher D, Malvaux P eds. *Pediatric Thyroidology*. Basel, S Karger, 1985:295–326.
6. Dumont JE et al. Large goiter as a maladaptation to iodine deficiency. *Clinical Endocrinology*, 1995,43:1–10.
7. Stanbury JB. *The damaged brain of iodine deficiency*. New York, Cognizant Communication, 1994.
8. Delange F. Iodine deficiency as a cause of brain damage. *Postgraduate Medical Journal*, 2001, 77:217–220.
9. Boyages SC. Primary pediatric hypothyroidism and endemic cretinism. *Current therapy in endocrinology and metabolism*, 1994, 5:94–8.
10. Bleichrodt N, Born MP. A meta-analysis of research on iodine and its relationship to cognitive development. In: Stanbury JB, ed. *The damaged brain of iodine deficiency*. New York, Cognizant Communication, 1994: 195–200.
11. Elandt-Johnson RC. Definition of rates: some remarks on their use and misuse. *American Journal of Epidemiology*, 1975, 102:267–71.
12. WHO, UNICEF, ICCIDD. *Assessment of iodine deficiency disorders and monitoring their elimination*. Geneva, World Health Organization, 2001 (WHO/NHD/01.1).
13. Phillips DIW. Iodine, milk, and the elimination of endemic goitre in Britain: the story of an accidental public health triumph. *Journal of Epidemiology and Community Health*, 1997, 51:391–393.
14. Anonymous. Iodized water to eliminate iodine deficiency. *IDD Newsletter*, 1997, 13:33–39.
15. Marine D, Kimball OP. Prevention of simple goiter in man. *Archives of Internal Medicine*, 1920, 25:661–672.
16. Bürgi H, Supersaxo Z, Selz B. Iodine deficiency diseases in Switzerland one hundred years after Theodor Kocher's survey: a historical review with some new goitre prevalence data. *Acta Endocrinology (Copenhagen)*, 1990, 123:577–590.
17. WHO, UNICEF, ICCIDD. *Recommended iodine levels in salt and guidelines for monitoring their adequacy and effectiveness*. Geneva, World Health Organization, 1996 (WHO/NUT/96.13).
18. Bürgi H, Schaffner T, Seiler JP. The toxicology of iodate: A review of the literature. *Thyroid*, 2001, 11: 449–456.
19. Todd CH. *Hyperthyroidism and other thyroid disorders. A practical handbook for recognition and management*. Geneva, World Health Organization, 1999 (WHO/AFRO/NUT/99.1, WHO/NUT/99.1).
20. WHO, UNICEF, ICCIDD. *Progress towards the elimination of iodine deficiency disorders (IDD)*. Geneva, World Health Organization, 1999 (WHO/NHD/99.4).
21. WHO, UNICEF, ICCIDD. *Global prevalence of iodine deficiency disorders*. Micronutrient Deficiency Information System working paper 1. Geneva, World Health Organization, 1993.
22. Zimmermann MB et al. New reference values for thyroid volume by ultrasound in iodine-sufficient schoolchildren: a World Health Organization/Nutrition for Health and Development Iodine Deficiency Study Group Report. *American Journal of Clinical Nutrition*, 2004, 79: 231–237.
23. UN Population Division. *World population prospects: the 2002 revision*. New York, United Nations, 2003.
24. UNICEF. *The state of the world's children 2004*. New York, United Nations Children's Fund, 2004.
25. Delange F. Iodine deficiency in the world: Where do we stand at the turn of the century? *Thyroid* 2001, 11: 437–447.
26. Zimmermann MB et al. Thyroid ultrasound compared with World Health Organization 1996 and 1994 palpation criteria for determination of goitre prevalence in regions of mild and severe iodine deficiency. *European Journal of Endocrinology* 2000, 143:727–31.
27. Peterson S et al. Classification of thyroid size by palpation and ultrasonography in field surveys. *Lancet* 2000, 355:106–110.

ANNEX 1

WHO Member States grouped by WHO and UN regions

Table A1.1 WHO Member States grouped by WHO region

Africa	Sierra Leone	Saint Vincent and the Grenadines	Germany
Algeria	South Africa	Suriname	Greece
Angola	Swaziland	Trinidad and Tobago	Hungary
Benin	Togo	United States of America	Iceland
Botswana	Uganda	Uruguay	Ireland
Burkina Faso	United Republic of Tanzania	Venezuela	Israel
Burundi	Zambia	South-East Asia	Italy
Cameroon	Zimbabwe	Bangladesh	Kazakhstan
Cape Verde	Americas	Bhutan	Kyrgyzstan
Central African Republic	Antigua and Barbuda	Democratic People's Republic of Korea	Latvia
Chad	Argentina	India	Lithuania
Comoros	Bahamas	Indonesia	Luxemburg
Congo	Barbados	Maldives	Malta
Côte d'Ivoire	Belize	Myanmar	Monaco
Democratic Republic of the Congo	Bolivia	Nepal	Netherlands
Equatorial Guinea	Brazil	Sri Lanka	Norway
Eritrea	Canada	Thailand	Poland
Ethiopia	Chile	Timor Leste	Portugal
Gabon	Colombia	Europe	Republic of Moldova
Gambia	Costa Rica	Albania	Romania
Ghana	Cuba	Andorra	Russian Federation
Guinea	Dominica	Armenia	San Marino
Guinea-Bissau	Dominican Republic	Austria	Serbia and Montenegro
Kenya	Ecuador	Azerbaijan	Slovakia
Lesotho	El Salvador	Belarus	Slovenia
Liberia	Grenada	Belgium	Spain
Madagascar	Guatemala	Bosnia and Herzegovina	Sweden
Malawi	Guyana	Bulgaria	Switzerland
Mali	Haiti	Croatia	Tajikistan
Mauritania	Honduras	Cyprus	The former Yugoslav Republic of Macedonia
Mauritius	Jamaica	Czech Republic	Turkey
Mozambique	Mexico	Denmark	Turkmenistan
Namibia	Nicaragua	Estonia	Ukraine
Niger	Panama	Finland	United Kingdom of Great Britain and Northern Ireland
Nigeria	Paraguay	France	
Rwanda	Peru	Georgia	
Sao Tome and Principe	Saint Kitts and Nevis		
Senegal	Saint Lucia		
Seychelles			

Eastern Mediterranean

Afghanistan
Bahrain
Djibouti
Egypt
Iran (Islamic Republic of)
Iraq
Jordan
Kuwait
Lebanon
Libyan Arab Jamahiriya
Morocco
Oman
Pakistan

Qatar
Saudi Arabia
Somalia
Sudan
Syrian Arab Republic
Tunisia
United Arab Emirate
Yemen

Western Pacific

Australia
Brunei Darussalam
Cambodia
China

Cook Islands
Fiji
Japan
Kiribati
Lao People's Democratic
Republic
Malaysia
Marshall Islands
Micronesia (Federated
States of)
Mongolia
Nauru
New Zealand
Niue

Palau
Papua New Guinea
Philippines
Republic of Korea
Samoa
Singapore
Solomon Islands
Tonga
Tuvalu
Vanuatu
Viet Nam

Table A1.2 WHO Member States grouped by UN region and subregion

Africa**Eastern Africa**

Burundi
Comoros
Djibouti
Eritrea
Ethiopia
Kenya
Madagascar
Malawi
Mauritius
Mozambique Rwanda
Seychelles
Somalia
Uganda
United Republic of
Tanzania
Zambia
Zimbabwe

Middle Africa

Angola
Cameroon
Central African Republic
Chad
Congo
Democratic Republic of
The Congo
Equatorial Guinea Gabon
Sao Tome and Principe

Northern Africa

Algeria
Egypt
Libyan Arab Jamahiriya

Morocco
Sudan
Tunisia

Southern Africa

Botswana
Lesotho
Namibia
South Africa
Swaziland

Western Africa

Benin
Burkina Faso
Cape Verde
Côte d'Ivoire
Gambia
Ghana
Guinea
Guinea-Bissau
Liberia
Mali
Mauritania
Niger
Nigeria
Senegal
Sierra Leone
Togo

Asia**Eastern Asia**

China
Democratic People's
Republic of Korea
Japan

Mongolia
Republic of Korea

South-central Asia

Afghanistan
Bangladesh
Bhutan
India
Iran (Islamic Republic of)
Kazakhstan
Kyrgyzstan
Maldives
Nepal
Pakistan
Sri Lanka
Tajikistan
Turkmenistan
Uzbekistan

South-eastern Asia

Brunei Darussalam
Cambodia
Timor Leste
Indonesia
Lao People's Democratic
Republic
Malaysia
Myanmar
Philippines
Singapore
Thailand
Viet Nam

Western Asia

Armenia

Azerbaijan
Bahrain
Cyprus
Georgia
Iraq
Israel
Jordan
Kuwait
Lebanon
Oman
Qatar
Saudi Arabia
Syrian Arab Republic
Turkey
United Arab Emirates
Yemen

Europe**Eastern Europe**

Belarus
Bulgaria
Czech Republic
Hungary
Poland
Republic of Moldova
Romania
Russian Federation
Slovakia
Ukraine

Northern Europe

Denmark
Estonia
Finland
Iceland

Ireland
Latvia
Lithuania
Norway
Sweden

Southern Europe

Albania
Andorra
Bosnia and Herzegovina
Croatia
Greece
Italy
Malta
Portugal
San Marino

Western Europe

Austria
Belgium
France
Germany
Luxembourg
Monaco
Netherlands
Switzerland

Latin America and the Caribbean

Caribbean
Antigua and Barbuda
Bahamas
Barbados
Cuba
Dominica
Dominican Republic
Grenada
Haiti
Jamaica
Saint Kitts and Nevis
Saint Lucia
Saint Vincent and the
Grenadines
Trinidad and Tobago

Central America

Belize
Costa Rica
El Salvador
Guatemala
Honduras
Mexico
Nicaragua
Panama

South America

Argentina
Bolivia
Brazil
Chile
Colombia
Ecuador
Guyana
Paraguay
Peru
Suriname
Uruguay
Venezuela

Northern America

Canada
United States of America

Oceania

Australia-New Zealand

Australia
New Zealand

Melanesia

Fiji
Papua New Guinea
Solomon Islands
Vanuatu

Micronesia

Kiribati
Marshall Islands
Micronesia (Federated
States of)
Nauru
Palau

Polynesia

Cook Islands
Samoa
Tonga
Tuvalu
Niue

ANNEX 2

Results by UN region

Table A2.1 Population coverage^a by UI surveys carried out between 1993 and 2003, by UN region

UN region ^b	School-age children (millions) ^c	School-age children surveyed (millions)	Coverage (%)
Africa	154.6	139.8	90.4
Eastern Africa	52.1	43.1	82.7
Middle Africa	19.1	15.6	82.0
Northern Africa	28.7	27.9	97.3
Southern Africa	8.3	7.9	95.4
Western Africa	46.4	45.2	97.5
Asia	521.7	488.8	93.7
Eastern Asia	164.8	148.8	90.3
South-central Asia	246.5	241.2	97.8
South-eastern Asia	79.0	75.8	95.9
Western Asia	31.5	23.0	73.2
Europe	59.1	50.3	85.2
Eastern Europe	25.2	25.2	100.0
Northern Europe	8.6	2.0	23.1
Southern Europe	10.6	8.5	79.7
Western Europe	14.7	14.7	100.0
Latin America and the Caribbean	76.6	69.2	53.7
Caribbean	4.5	2.4	53.0
Central America	22.4	22.4	100.0
South America	49.7	44.5	89.5
Northern America	32.4	29.6	91.2
Oceania	3.7	3.5	94.0
Australia–New Zealand	2.3	2.3	100.0
Melanesia	1.3	1.2	90.2
Micronesia	0.04	0	0.0
Polynesia	0.05	0	0.0
Total	848.0	781.2	92.1

^a School-age children (6–12 years).

^b Based on 192 WHO Member States.

^c Based on population estimates for the year 2002 (23).

Table A2.2 Type of UI survey data by UN region

UN region ^a	National	Sub-national	No data
Africa	20	18	15
Eastern Africa	7	3	7
Middle Africa	3	2	4
Northern Africa	3	2	1
Southern Africa	3	1	1
Western Africa	4	10	2
Asia	26	8	13
Eastern Asia	2	0	3
South-central Asia	8	4	2
South-eastern Asia	6	1	4
Western Asia	10	3	4
Europe	16	14	11
Eastern Europe	6	4	0
Northern Europe	3	4	3
Southern Europe	3	3	7
Western Europe	4	3	1
Latin America and the Caribbean	12	7	14
Caribbean	1	1	11
Central America	7	1	0
South America	4	5	3
Northern America	1	0	1
Oceania	0	4	12
Australia–New Zealand	0	2	0
Melanesia	0	2	2
Micronesia	0	0	5
Polynesia	0	0	5
Total	75	51	66

^a Based on 192 WHO Member States.

Table A2.3 **Population coverage^a by TGP surveys carried out between 1993 and 2003, by UN region**

UN region ^b	School-age children (millions) ^c	School-age children surveyed (millions)	Coverage (%)
Africa	154.6	140.5	90.9
Eastern Africa	52.1	46.1	88.4
Middle Africa	19.1	16.4	85.8
Northern Africa	28.7	27.9	97.3
Southern Africa	8.3	7.9	95.4
Western Africa	46.4	42.2	91.1
Asia	521.7	487.4	93.4
Eastern Asia	164.8	148.8	90.3
South-central Asia	246.5	246.5	100.0
South-eastern Asia	79.0	69.9	88.6
Western Asia	31.5	22.2	70.6
Europe	59.1	26.1	44.2
Eastern Europe	25.2	9.2	36.4
Northern Europe	8.6	0.5	5.5
Southern Europe	10.6	9.4	88.2
Western Europe	14.7	7.1	48.4
Latin America and the Caribbean	76.6	50.7	66.2
Caribbean	4.5	1.3	28.4
Central America	22.4	16.1	72.0
South America	49.7	33.3	67.0
Northern America	32.4	0	0.0
Oceania	3.7	2.9	79.3
Australia–New Zealand	2.3	1.9	81.7
Melanesia	1.3	1.0	80.7
Micronesia	0.040	0	0.0
Polynesia	0.054	0	0.0
Total	848.0	707.7	83.5

^a School-age children (6–12 years).

^b Based on 192 WHO Member States.

^c Based on population estimates for the year 2002 (UN 2003).

Table A2.4 **Type of TGP survey data by UN region**

UN region ^a	National	Sub-national	No data
Africa	21	18	14
Eastern Africa	6	5	6
Middle Africa	3	3	3
Northern Africa	3	2	1
Southern Africa	3	1	1
Western Africa	6	7	3
Asia	24	9	14
Eastern Asia	2	0	3
South-central Asia	7	7	0
South-eastern Asia	6	0	5
Western Asia	8	3	6
Europe	9	10	22
Eastern Europe	3	4	3
Northern Europe	0	1	9
Southern Europe	5	3	5
Western Europe	1	2	5
Latin America and the Caribbean	4	3	26
Caribbean	1	0	12
Central America	1	1	6
South America	2	2	8
Northern America	0	0	2
Oceania	0	2	14
Australia–New Zealand	0	1	1
Melanesia	0	1	3
Micronesia	0	0	5
Polynesia	0	0	5
Total	57	43	92

^a Based on 192 WHO Member States.

Table A2.5 **Number of countries classified by degrees of public health significance of iodine nutrition based on median UI in school-age children, by UN region, 2003**

UN region ^a	Classification of iodine nutrition						No data
	Severe iodine deficiency (Median UI <20 µg/l)	Moderate iodine deficiency (Median UI 20–49 µg/l)	Mild iodine deficiency (Median UI 50–99 µg/l)	Optimal iodine nutrition (Median UI 100–199 µg/l)	Risk of IIH in susceptible groups (Median UI 200–299 µg/l)	Risk of adverse health consequences (Median UI ≥300 µg/l)	
Africa	0	6	10	13	7	2	15
Eastern Africa	0	0	3	4	2	1	7
Middle Africa	0	2	1	1	1	0	4
Northern Africa	0	1	2	2	0	0	1
Southern Africa	0	1	0	2	1	0	1
Western Africa	0	2	4	4	3	1	2
Asia	1	4	12	12	5	0	13
Eastern Asia	0	0	0	1	1	0	3
South-central Asia	1	2	4	3	2	0	2
South-eastern Asia	0	1	3	3	0	0	4
Western Asia	0	1	5	5	2	0	4
Europe	0	1	15	14	0	0	11
Eastern Europe	0	1	6	3	0	0	0
Northern Europe	0	0	5	2	0	0	3
Southern Europe	0	0	1	5	0	0	7
Western Europe	0	0	3	4	0	0	1
Latin America and the Caribbean	0	1	1	3	11	3	14
Caribbean	0	1	1	0	0	0	11
Central America	0	0	0	2	6	0	0
South America	0	0	0	1	5	3	3
Northern America	0	0	0	0	1	0	1
Oceania	0	1	2	1	0	0	12
Australia–New Zealand	0	0	2	0	0	0	0
Melanesia	0	1	0	1	0	0	2
Micronesia	0	0	0	0	0	0	5
Polynesia	0	0	0	0	0	0	5
Total	1	13	40	43	24	5	66

^a Based on 192 WHO Member States.

Table A2.6 **Proportion of population, and number of individuals with, insufficient iodine intake in school-age children (6–12 years) and the general population, by UN region, 2003**

UN region ^a	Insufficient iodine intake (UI <100 µg/l)			
	School-age children		General population	
	Prevalence (%)	Total number (millions) ^b	Prevalence (%)	Total number (millions) ^b
Africa	42.7	59.7	43.0	324.2
Eastern Africa	45.1	19.4	45.2	98.2
Middle Africa	32.4	5.1	32.7	26.3
Northern Africa	50.7	14.1	50.6	88.2
Southern Africa	31.6	2.5	31.2	15.4
Western Africa	41.1	18.6	41.4	96.2
Asia	38.3	187.0	35.6	1239.3
Eastern Asia	16.3	24.2	16.3	212.2
South-central Asia	43.2	104.1	41.9	631.9
South-eastern Asia	61.2	46.4	60.5	312.6
Western Asia	53.2	12.2	55.8	82.6
Europe	53.1	26.7	52.7	330.8
Eastern Europe	60.0	15.1	59.9	180.6
Northern Europe	59.3	1.2	59.2	13.0
Southern Europe	47.8	4.1	49.2	58.8
Western Europe	43.6	6.4	42.6	78.5
Latin America and the Caribbean	10.3	7.1	10.0	47.4
Caribbean	69.8	1.7	66.2	13.2
Central America	9.9	2.2	9.7	13.5
South America	7.3	3.3	6.6	20.8
Northern America	9.5	2.8	9.5	27.6
Oceania	59.4	2.1	64.5	19.2
Australia–New Zealand	73.0	1.7	72.8	17.0
Melanesia	32.7	0.4	33.9	2.2
Micronesia	–	–	–	–
Polynesia	–	–	–	–
Total	36.5	285.4	35.2	1988.7

– No data.

^a Based on 192 WHO Member States.

^b Based on population estimates for the year 2002 (UN 2003).

Table A2.7 **TGP in the general population^a by UN region, 2003**

UN region ^b	Total goitre prevalence (%)
Africa	26.8
Eastern Africa	29.5
Middle Africa	23.3
Northern Africa	25.3
Southern Africa	29.1
Western Africa	25.9
Asia	14.5
Eastern Asia	5.3
South-central Asia	23.8
South-eastern Asia	8.4
Western Asia	20.4
Europe	16.3
Eastern Europe	27.2
Northern Europe	12.1
Southern Europe	10.9
Western Europe	10.7
Latin America and the Caribbean	4.7
Caribbean	4.9
Central America	8.7
South America	2.9
Northern America	–
Oceania	12.9
Australia–New Zealand	15.3
Melanesia	4.4
Micronesia	–
Polynesia	–
Total	15.8

– No data.

^a Estimated from TGP data from school-age children (6–12 years).

^b Based on 192 WHO Member States.

ANNEX 3

National estimates of iodine status

Table A3.1 Country data on UI and national estimate of iodine nutrition

Member State	Population year 2002 ^a		Survey data										Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)
Afghanistan	4197	22 930	No data												
Albania	434	3141	No data												
Algeria	4993	31 266	1994 P	Local	SAC (6-11)	169	27	77.7	71.4-84.0	Survey in endemic area. Baseline data before intervention. % <100 µg/l calculated from median.	1348	Insufficient	Moderate iodine deficiency	3879	24294
Andorra	6	69	No data												
Angola	2594	13 184	No data												
Antigua and Barbuda	10	73	No data												
Argentina	4792	37 981	No data												
Armenia	346	3072	1998	National	Pre-SAC (0-5)	2596	146	31.8	30.0-33.6	Median calculated from % <100 µg/l.	3329	Adequate	Optimal iodine nutrition	110	977
Australia	1872	19 544	2000, 2001	State, local	SAC (4-18)	802	77	71.5	68.4-74.6	Medians and % <100 µg/l from two surveys pooled.	3379, 3598	Insufficient	Mild iodine deficiency	1339	13974
Austria	650	8111	1994	Local	SAC (6-15)	589	111	49.4	46.3-54.3	Thyromobile study. % <100 µg/l calculated from median.	1319	Adequate	Optimal iodine nutrition	327	4080
Azerbaijan	1268	8297	2001 P	Regional	SAC (8-14)	347	54	74.4	69.8-79.0	% <100 µg/l calculated from median.	3413	Insufficient	Mild iodine deficiency	943	6173
Bahamas	42	310	No data												
Bahrain	97	709	1999	National	SAC (8-12)	749	204	16.2	13.6-18.8	Median calculated from % <100 µg/l.	1142	More than adequate	Risk of IIH in susceptible groups	16	115
Bangladesh	25 239	143 809	1993	National	SAC (5-11)	2054	54	70.7	68.7-72.7	Median calculated from % <100 µg/l.	642	Insufficient	Mild iodine deficiency	17844	101673
Barbados	26	269	No data												
Belarus	862	9940	1995-1998	National	SAC (6-18)	11 562	45	80.9	80.2-81.6		3181	Insufficient	Moderate iodine deficiency	697	8041
Belgium	846	10 296	1998	National	SAC (6-12)	2585	80	66.9	65.1-68.7		1336	Insufficient	Mild iodine deficiency	566	6888
Belize	44	251	1994-1995	National	SAC (7-14)	1656	184	26.7	24.6-28.8	% <100 µg/l calculated from median.	3133	Adequate	Optimal iodine nutrition	12	67
Benin	1302	6558	1999	Local	SAC (6-12)	433	289	8.3	5.7-10.9	Thyromobile study. % <100 µg/l calculated from median.	2535	More than adequate	Risk of IIH in susceptible groups	108	544
Bhutan	409	2190	1996	National	SAC (6-11)	333	230	24.0	19.4-28.6		2649	More than adequate	Risk of IIH in susceptible groups	98	526
Bolivia	1546	8645	1996	National	Women 15-49 and children <5	508	250	19.0	15.6-22.4		3339	More than adequate	Risk of IIH in susceptible groups	294	1643
Bosnia and Herzegovina	357	4126	1999	National	SAC (7-14)	1945	111	52.4	50.2-54.6	Medians and % <100 µg/l pooled separately from two surveys (Republica Sipska and Federation of Bosnia & Herzegovina). Pooled % <100 µg/l higher than 50% in spite of pooled median higher than 100 µg/l.	2994, 3453	Adequate	Optimal iodine nutrition	187	2162
Botswana	321	1770	1994	National	SAC (8-10)	287	219	15.3	11.1-19.5	Medians from disaggregated data by districts pooled. % <100 µg/l calculated from median.	2805	More than adequate	Risk of IIH in susceptible groups	49	271
Brazil	23 198	176 257	2000	State	SAC (6-12)	1013	360	0.0	0.0-0.0		3350	Excessive	Risk of adverse health consequences	0	0
Brunei Darussalam	49	350	No data												
Bulgaria	599	7965	1996	National	SAC (6-14)	1028	111	42.9	39.9-45.9		3017	Adequate	Optimal iodine nutrition	257	3417
Burkina Faso	2608	12 624	1999	Local	SAC (6-12)	391	114	47.5	42.6-52.4	Thyromobile study.	2535	Adequate	Optimal iodine nutrition	1239	5997

Table A3.1

Member State	Population year 2002 ^a		Survey data										Classification of iodine intake	Classification of iodine nutrition	Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	6-12 years (000)			General population (000)		
Burundi	1361	6602	No data														
Cambodia	2585	13 810	No data														
Cameroon	3007	15 729	1993	National	SAC (6-18)	757	52	91.7	89.7-93.7			1431	Insufficient	Mild iodine deficiency	2757	14 424	
Canada	2840	31 271	No data														
Cape Verde	85	454	1996	National	SAC (6-12)	302	52	77.4	72.7-82.1			1622	Insufficient	Mild iodine deficiency	66	351	
Central African Republic	729	3819	1993-1994	Province	General population	319	21	79.5	75.1-83.9	% <100 µg/l calculated from median.		1351	Insufficient	Moderate iodine deficiency	579	3036	
Chad	1638	8348	1993-1994	National	SAC (10-20)	1141	29	99.6	99.2-100.0			390	Insufficient	Moderate iodine deficiency	1632	8314	
Chile	2046	15 613	2001	Local urban	SAC (6-18)	371	984	0.2	0.0-0.7	Median calculated from mean.		3335	Excessive	Risk of adverse health consequences	4	31	
China	148 422	1 302 307	2002	National	SAC (8-10)	11 766	241	16.2	15.5-16.9	% <100 µg/l calculated from median.		3579	More than adequate	Risk of IIH in susceptible groups	24 044	210 974	
Colombia	6514	43 526	1994-1998	National urban	SAC (8-12)	7363	249	6.4	5.8-7.0	Median calculated from % <100 µg/l.		3296	More than adequate	Risk of IIH in susceptible groups	417	2786	
Comoros	137	747	No data														
Congo	727	3633	No data														
Cook Islands	2	18	No data														
Costa Rica	595	4094	1996	National	SAC	538	233	8.9	6.5-11.3			1634	More than adequate	Risk of IIH in susceptible groups	53	364	
Cote d'Ivoire	3067	16 365	1999-2000	Local	SAC (4-16)	400	162	33.8	29.2-38.4	% <100 µg/l calculated from median.		3239	Adequate	Optimal iodine nutrition	1036	5531	
Croatia	351	4439	2002	National	SAC (6-12)	927	140	28.8	25.9-31.7			3429	Adequate	Optimal iodine nutrition	101	1278	
Cuba	1119	11 271	1995	National rural	SAC (6-12)	3027	95	51.0	49.2-52.8			1512	Insufficient	Mild iodine deficiency	571	5748	
Cyprus	87	796	No data														
Czech Republic	810	10 246	2000 P	Regional	SAC (6, 10, 13)	714	119	47.7	44.0-51.4	Median calculated from mean. % <100 µg/l calculated from median.		515	Adequate	Optimal iodine nutrition	386	4887	
Democratic People's Republic of Korea	2825	22 541	No data														
Democratic Republic of the Congo	10 011	51 201	1995	Local	SAC (6-14)	305	267	0.0	0.0-0.0	Medians from disaggregated data by zones pooled. % <100 µg/l calculated from median.		3601	More than adequate	Risk of IIH in susceptible groups	0	0	
Denmark	473	5351	1997-1998	Regional	Adults (18-65)	4616	61	70.8	69.5-72.1	% <100 µg/l calculated from median.		3205	Insufficient	Mild iodine deficiency	335	3789	
Djibouti	129	693	No data														
Dominica	11	78	No data														
Dominican Republic	1292	8616	1993	National	SAC (6-14)	837	39	86.0	83.7-88.4	Median calculated from % <100 µg/l.		771	Insufficient	Moderate iodine deficiency	1111	7410	
Ecuador	1978	12810	1999	Local	SAC	630	420	0.0	0.0-0.0	Thyromobile study. % <100 µg/l calculated from median.		3615	Excessive	Risk of adverse health consequences	0	0	
Egypt	11 373	70 507	1998	State	SAC (6-10)	706	148	31.2	27.8-34.6	Median calculated from % <100 µg/l.		2639	Adequate	Optimal iodine nutrition	3548	21 998	
El Salvador	1033	6415	1996-1997	National	SAC (6-14)	2394	>150	4.6	3.8-5.4			3108	Adequate	Optimal iodine nutrition	48	295	
Equatorial Guinea	90	481	No data														
Eritrea	794	3991	1998	National	SAC (6-12)	2100	168	25.3	23.4-27.2	Median calculated from % <100 µg/l.		3122	Adequate	Optimal iodine nutrition	201	1010	
Estonia	112	1338	1995	National	SAC (8-10)	1840	65	67.0	64.9-69.2			1225	Insufficient	Mild iodine deficiency	75	896	

Table A3.1

Member State	Population year 2002 ^a		Survey data										Classification of iodine intake		Population with insufficient iodine intake	
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)	
Ethiopia	13 685	68 961	2000 P	District	SAC	512	58	68.4	64.4-72.4	Means from disaggregated data by regions pooled. Median calculated from mean. % <100 µg/l calculated from median.	3199	Insufficient	Mild iodine deficiency	9360	47 169	
Fiji	123	831	1994	District	SAC	479	34	75.4	71.5-79.3	Medians from disaggregated data by site pooled. % <100 µg/l calculated from median.	3236	Insufficient	Moderate iodine deficiency	93	626	
Finland	448	5197	1997	Local	Adults (30-42)	342	164	35.5	30.4-40.6	% <100 µg/l calculated from median.	3605	Adequate	Optimal iodine nutrition	159	1845	
France	5128	59 850	1996	National	Adults (35-60)	12 014	85	60.4	59.5-61.3	Medians from disaggregated data by sex pooled. % <100 µg/l calculated from median.	1269	Insufficient	Mild iodine deficiency	3097	36 149	
Gabon	245	1306	2001	National	SAC (6-12)	NS	190	38.3	% <100 µg/l calculated from median.	3611	Adequate	Optimal iodine nutrition	94	500		
Gambia	248	1388	1999	National	SAC (8-12)	594	42	72.8	69.2-76.4	% <100 µg/l calculated from median.	2595	Insufficient	Moderate iodine deficiency	180	1011	
Georgia	494	5177	1998	National	NS	NS	62	80.0	Median calculated from % <100 µg/l.	3699	Insufficient	Mild iodine deficiency	395	4142		
Germany	6022	82414	1999	National	SAC (6-12)	3065	148	27.0	25.4-28.6		3126	Adequate	Optimal iodine nutrition	1626	22 252	
Ghana	3712	20471	1994	District		292	54	71.3	66.1-76.5		1772	Insufficient	Mild iodine deficiency	2647	14 596	
Greece	759	10970	No data													
Grenada	12	80	No data													
Guatemala	2309	12036	1995	National	SAC, women	814	222	14.4	12.0-16.8	% <100 µg/l calculated from median.	3091	More than adequate	Risk of IIH in susceptible groups	333	1733	
Guinea	1567	8359	1999	Region	SAC (8-19)	1234	91	63.6	60.9-66.3	Medians and % <100 µg/l from disaggregated data by prefectures pooled.	2617	Insufficient	Mild iodine deficiency	997	5316	
Guinea-Bissau	283	1449	No data													
Guyana	104	764	1997 P	National	SAC (5-14)	342	162	26.9	22.2-31.6	% <100 µg/l from disaggregated data by sex pooled. Median calculated from % <100 µg/l.	3094	Adequate	Optimal iodine nutrition	28	205	
Haiti	1459	8218	No data													
Honduras	1262	6781	1999	Local	SAC	609	240	31.3	27.6-35.0	Thyromobile study. % <100 µg/l calculated from median.	3394	More than adequate	Risk of IIH in susceptible groups	395	2122	
Hungary	817	9923	1994-1997	National	SAC (7-11)	2814	80	65.2	63.4-67.0	% <100 µg/l from disaggregated data by county pooled.	3041, 3684, 3683, 3682, 3681	Insufficient	Mild iodine deficiency	533	6470	
Iceland	32	287	1998 P	Local	Elderly (66-70)	89	150	37.7	27.6-47.8	% <100 µg/l calculated from median.	1251	Adequate	Optimal iodine nutrition	12	108	
India	16 1973	1 049 549	1993-1993, 1995, 1996, 1996 P, 1997, 1997 P, 1998, 1998 P, 1999, 2000 P, 2001 P, 2001, 2002	State, district	SAC	17 321	133	31.3	30.6-32.0	Medians and % <100 µg/l from 20 state and district surveys, pooled at district level.	1159, 1164, 1161, 1160, 1215, 1166, 1163, 3538, 1165, 3539, 3456, 3584, 3585, 3578, 3577, 3565, 3545, 3534	Adequate	Optimal iodine nutrition	50 698	328 509	
Indonesia	30 322	217 131	1996 P	District	SAC (8-10)	544	65	63.7	59.7-67.7		1183	Insufficient	Mild iodine deficiency	19 315	138 313	
Iran (Islamic Republic of)	11 208	68 070	1996	National	SAC (8-10)	2917	205	14.9	13.6-16.2		3317	More than adequate	Risk of IIH in susceptible groups	1670	10 142	

Table A3.1

Member State	Population year 2002 ^a		Survey data										Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)
Iraq	4498	24 510	No data												
Ireland	375	3911	1999	Local	Adults (22-61)	132	82	60.8	52.5-69.1	% <100 µg/l calculated from median.	3608	Insufficient	Mild iodine deficiency	228	2378
Israel	795	6304	No data												
Italy	3866	57 482	1992-1994, 1993-1995, 1994 P, 1997 P, 1998 P, 1999 P	Region, local	SAC (6-15)	11 226	94	55.7	54.8-56.6	Medians from nine local and regional surveys pooled. % <100 µg/l calculated from median.	2058, 1291, 3419, 1273, 1287, 1286, 2059, 1272	Insufficient	Mild iodine deficiency	2154	32 018
Jamaica	381	2627	No data												
Japan	8482	127 478	No data												
Jordan	917	5329	2000	National	SAC (8-10)	2601	154	24.4	22.8-26.1		2534	Adequate	Optimal iodine nutrition	224	1300
Kazakhstan	1991	15 469	1999	National	Women (15-49)	951	97	53.1	49.9-56.3	Medians from disaggregated data by region pooled.	3056	Insufficient	Mild iodine deficiency	1057	8214
Kenya	6050	31540	1994	National	SAC (8-10)	3042	115	36.7	35.0-38.4	% <100 µg/l from disaggregated data by district pooled.	391	Adequate	Optimal iodine nutrition	2220	11575
Kiribati	10	87	No data												
Kuwait	278	2443	1997	National	SAC (6-9)	341	147	31.4	26.5-36.3	Median calculated from % <100 µg/l.	3135	Adequate	Optimal iodine nutrition	87	767
Kyrgyzstan	787	5067	1994	Region	SAC (7-11)	221	30	88.1	83.8-92.4	Survey in 4 out of 7 States (Oblasts). The 4 States considered endemic. % <100 µg/l calculated from median.	3230	Insufficient	Moderate iodine deficiency	693	4464
Lao People's Democratic Republic	1038	5529	2000	National	SAC (8-12)	900	162	26.9	24.0-29.8	Median calculated from % <100 µg/l.	770	Adequate	Optimal iodine nutrition	279	1487
Latvia	200	2329	2000	National	SAC (8-10)	599	59	76.8	73.4-80.2		3058	Insufficient	Mild iodine deficiency	154	1789
Lebanon	510	3596	1997	National	SAC (7-15)	586	95	55.5	51.5-59.5	% <100 µg/l calculated from median.	3222	Insufficient	Mild iodine deficiency	283	1996
Lesotho	331	1800	1999	National	SAC (8-12)	500	26	100.0	—		3481	Insufficient	Moderate iodine deficiency	331	1800
Liberia	631	3239	1999	National	SAC (6-11)	2060	321	3.5	2.7-4.3		1242	Excessive	Risk of adverse health consequences	22	113
Libyan Arab Jamahiriya	765	5445	No data												
Lithuania	334	3465	1995	National	SAC	2087	75	62.0	59.9-64.1	% <100 µg/l calculated from median.	3613	Insufficient	Mild iodine deficiency	207	2148
Luxembourg	40	447	1994	Local	SAC (6-15)	124	90	57.4	48.7-66.1	Thyromobile study. % <100 µg/l calculated from median.	1319	Insufficient	Mild iodine deficiency	23	257
Madagascar	3231	16916	No data												
Malawi	2315	11871	No data												
Malaysia	3717	23965	1995	National	SAC (8-10)	11 362	91	57.0	56.1-57.9	Medians from three state surveys pooled. % <100 µg/l calculated from median.	2637, 2637, 2840	Insufficient	Mild iodine deficiency	2118	13660
Maldives	58	309	1995	National	SAC (6-12)	316	67	65.5	60.3-70.7		2650	Insufficient	Mild iodine deficiency	38	202
Mali	2636	12 623	1999	Local	SAC (6-12)	352	203	34.1	29.2-39.1	Thyromobile study.	2535	More than adequate	Risk of IIH in susceptible groups	899	4304
Malta	37	393	No data												
Marshall Islands	6	52	No data												
Mauritania	514	2807	1995	National	SAC (6-14)	240	55	69.8	64.0-75.6	Median calculated from % <100 µg/l.	392	Insufficient	Mild iodine deficiency	359	1959
Mauritius	147	1210	1995	National	Adults	225	154	4.4	1.7-7.1	Median calculated from mean.	395	Adequate	Optimal iodine nutrition	6	53

Table A3.1

Member State	Population year 2002 ^a		Survey data										Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)
Mexico	15 671	101 965	1999	National	SAC (5-12)	585	235	8.5	6.2-10.8	Median calculated from % <100 µg/l.	2997	More than adequate	Risk of IHI in susceptible groups	1332	8667
Micronesia (Federated States of)	19	108	No data												
Monaco	3	34	No data												
Mongolia	407	2559	2001	National	SAC (4-16)	2748	102	48.9	47.0-50.8		3227	Adequate	Optimal iodine nutrition	199	1252
Morocco	4385	30 072	1993	National	SAC (6-12)	281	75	63.0	57.4-68.7	Median calculated from mean.	491	Insufficient	Mild iodine deficiency	2762	18945
Mozambique	3587	18 537	1998	Province	SAC	567	69	65.4	61.5-69.3		2872	Insufficient	Mild iodine deficiency	2346	12123
Myanmar	7352	48 852	2001	National	SAC (6-11)	3345	136	38.2	36.6-39.9		3076	Adequate	Optimal iodine nutrition	2809	18662
Namibia	385	1961	No data												
Nauru	2	13	No data												
Nepal	4430	24 609	1997-1998	National	SAC (6-11)	1450	144	35.1	32.6-37.6		1083	Adequate	Optimal iodine nutrition	1555	8638
Netherlands	1395	16 067	1995-1996	Local	SAC (6-18)	937	154	37.5	34.4-40.6	Thyromobile study, % <100 µg/l calculated from median.	3204	Adequate	Optimal iodine nutrition	523	6025
New Zealand	419	3846	1996-1999	Local	SAC (8-10)	282	66	79.7	75.0-84.4		3597	Insufficient	Mild iodine deficiency	334	3065
Nicaragua	1007	5335	2000	National	SAC (6-9)	886	271	0.0	0.0-0.0	% <100 µg/l calculated from median.	3109	More than adequate	Risk of IHI in susceptible groups	0	0
Niger	2357	11 544	1998	Region	SAC	944	270	0.0	0.0-0.0	% <100 µg/l calculated from median.	3611	More than adequate	Risk of IHI in susceptible groups	0	0
Nigeria	23 677	120 911	1998	State	SAC (8-12)	537	147	38.8	34.7-42.9	% <100 µg/l calculated from median.	3604	Adequate	Optimal iodine nutrition	9187	46914
Niue	0	2	No data												
Norway	427	4514	No data												
Oman	450	2768	1993-1994	National	SAC (8-11)	951	91	49.8	46.6-53.0		481	Insufficient	Mild iodine deficiency	224	1379
Pakistan	27 695	149911	1993-1994	Region	SAC (8-10)	1500	16	90.4	88.9-91.9		492	Insufficient	Severe iodine deficiency	25037	13 5519
Palau	2	20	No data												
Panama	436	3064	1999	National	SAC (6-12)	604	235	8.6	6.4-10.8	Median calculated from % <100 µg/l.	3098	More than adequate	Risk of IHI in susceptible groups	37	263
Papua New Guinea	1048	5586	1996	Local	SAC (8-10)	627	181	27.7	24.2-31.2	Medians from disaggregated data by site pooled, % <100 µg/l calculated from median.	723	Adequate	Optimal iodine nutrition	290	1547
Paraguay	1013	5740	1999	National	SAC (6-12)	5864	294	13.4	12.5-14.3		3167	More than adequate	Risk of IHI in susceptible groups	136	769
Peru	4201	26767	1999	National	SAC	4936	230	11.8	10.9-12.7	% <100 µg/l calculated from median.	3147	More than adequate	Risk of IHI in susceptible groups	496	3158
Philippines	13 388	78580	1998	National	SAC (6-12)	10 616	71	65.3	64.4-66.2		2536	Insufficient	Mild iodine deficiency	8742	51313
Poland	3437	38622	1999	Local	SAC (6-15)	873	84	64.0	60.8-67.2	Thyromobile study, Median calculated from mean.	2574	Insufficient	Mild iodine deficiency	2200	24718
Portugal	774	10049	No data												
Qatar	72	601	1996	Local	SAC (6-15)	59	203	30.0	26.3-33.7	Median calculated from mean.	3614	More than adequate	Risk of IHI in susceptible groups	22	180
Republic of Korea	4640	47430	No data												
Republic of Moldova	455	4270	1996	National	SAC (8-10)	516	78	62.0	57.8-66.2		3332	Insufficient	Mild iodine deficiency	282	2648
Romania	1836	22387	2000-2001	National	SAC (6-16)	7358	68	64.2	63.1-65.3	Median from disaggregated data by county pooled, % <100 µg/l calculated from median.	3544	Insufficient	Mild iodine deficiency	1178	14373

Table A3.1

Member State	Population year 2002 ^a		Survey data										Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)
Russian Federation	11 779	144082	1999, 2000, 2001, 2002	Local	SAC (7-12)	3401	93	56.2	54.5-57.9	Medians from eight local surveys pooled. % <100 µg/l calculated from median.	1191, 743 1222, 1143 1139, 3610 3610, 3610	Insufficient	Mild iodine deficiency	6620	80974
Rwanda	1593	8272	1996	National	SAC (5-19)	1246	298	0.0	0.0-0.0	% <100 µg/l calculated from median.	2558	More than adequate	Risk of IIH in susceptible groups	0	0
Saint Kitts and Nevis	6	42	No data												
Saint Lucia	22	148	No data												
Saint Vincent and the Grenadines	18	119	No data												
Samoa	33	176	No data												
San Marino	2	27	No data												
Sao Tome and Principe	28	157	No data												
Saudi Arabia	4063	23520	1994-1995	National	SAC (8-10)	4590	180	23.0	21.8-24.2		490	Adequate	Optimal iodine nutrition	934	5410
Senegal	1898	9855	1996-1997	Region	SAC (10-14)	1054	45	75.7	73.1-78.3		1633	Insufficient	Moderate iodine deficiency	1437	7460
Serbia and Montenegro	976	10535	1998-1999	Region	SAC (7-15)	1515	158	20.8	18.8-22.8		3062	Adequate	Optimal iodine nutrition	203	2191
Seychelles	15	80	No data												
Sierra Leone	876	4764	No data												
Singapore	450	4183	No data												
Slovakia	496	5398	2002	National	SAC	1744	183	15.0	13.3-16.7		558	Adequate	Optimal iodine nutrition	74	810
Slovenia	144	1986													
Solomon Islands	89	463	No data												
Somalia	1850	9480	No data												
South Africa	7070	44759	1998	National	SAC (7-11)	8254	177	29.0	28.0-30.0	% <100 µg/l calculated from median.	2618	Adequate	Optimal iodine nutrition	2050	12980
Spain	2712	40977	1995, 2000, 2000 P, 2001 P, 2002 P	Regional, province	SAC	3154	109	50.1	49.3-52.7	Medians from five regional and provincial surveys pooled. % <100 µg/l calculated from median. % <100 µg/l borderline, while median is 109 µg/l due to the equation used for the estimation (see section 2.3).	2091, 3607 3581, 3606 3580	Adequate	Optimal iodine nutrition	1383	20898
Sri Lanka	2236	18910	2000-2001	National	SAC (8-10)	2630	145	30.6	28.8-32.4	Medians from disaggregated data by zones pooled. % <100 µg/l calculated from median.	404	Adequate	Optimal iodine nutrition	684	5786
Sudan	5797	32878	1997	National	SAC	3544	75	62.0	60.4-63.6		2937	Insufficient	Mild iodine deficiency	3594	20385
Suriname	60	432	No data												
Swaziland	213	1069	1998	Local	SAC (6-18)	170	170	34.5	27.4-41.7	Medians from disaggregated data by sites pooled.	2589a	Adequate	Optimal iodine nutrition	73	369
Sweden	807	8867	No data												
Switzerland	576	7171	1999	National	SAC (6-12)	600	115	39.5	35.6-43.4		2662	Adequate	Optimal iodine nutrition	228	2833
Syrian Arab Republic	3039	17381	No data												
Tajikistan	1108	6195	No data												
Thailand	7414	62193	2000	National	Pregnant women	3557	150	34.9	33.3-36.5		3554	Adequate	Optimal iodine nutrition	2588	21705
The former Yugoslav Republic of Macedonia	213	2046	2002	National	SAC (7-11)	1216	199	11.8	10.0-13.6		3609	Adequate	Optimal iodine nutrition	25	241
Timor Leste	147	739	No data												
Togo	927	4801	1999	Local	SAC (6-12)	381	116	42.8	37.8-47.8	Thyromobile study.	2535	Adequate	Optimal iodine nutrition	397	2055

Table A3.1

Member State	Population year 2002 ^a		Survey data										Population with insufficient iodine intake		
	6-12 years (000)	General (000)	Date of survey (years)	Level of survey	Population group and age (years)	Sample size	Median UI (µg/l)	Proportion of population with UI <100 µg/l (%)	95% CI of proportion of population with UI <100 µg/l (%)	Notes	Bibliographic references ^b	Classification of iodine intake	Classification of iodine nutrition	6-12 years (000)	General population (000)
Tonga	17	103	No data												
Trinidad and Tobago	147	1298	No data												
Tunisia	1364	9728	1996-1997	National	SAC (6-9)	94	164	26.4	17.5-35.3	Median calculated from % <100 µg/l.	2485	Adequate	Optimal iodine nutrition	360	2568
Turkey	10 119	70318	1997-1999	National	SAC (9-11)	5948	36	74.6	73.5-75.7	% <100 µg/l calculated from median.	3426	Insufficient	Moderate iodine deficiency	7549	52457
Turkmenistan	821	4794	1999	Local	SAC (8-10)	65	64	65.6	54.1-77.2	% <100 µg/l calculated from median.	3620	Insufficient	Mild iodine deficiency	539	3145
Tuvalu	1	10	No data												
Uganda	5226	25004	1999	National	SAC (6-12)	293	310	11.9	8.2-15.6	% <100 µg/l from disaggregated data by district pooled.	2582	Excessive	Risk of adverse health consequences	622	2975
Ukraine	4114	48902	1991-1996, 1996-1999	Local	SAC (5-20)	3506	50	70.1	68.6-71.6	Medians and % <100 µg/l from two local surveys pooled.	1238, 3600	Insufficient	Mild iodine deficiency	2884	34280
United Arab Emirates	361	2937	1994	Region	SAC (9-13)	258	91	56.6	50.6-62.7	Survey in six endemic zones.	483	Insufficient	Mild iodine deficiency	205	1662
United Kingdom of Great Britain and Northern Ireland	5349	59068	No data												
United Republic of Tanzania	7282	36276	1996	State	SAC (8-9)	586	127	37.7	33.8-41.6		1772	Adequate	Optimal iodine nutrition	2745	13676
United States of America	29 589	291038	1988-1994	National	SAC (6-11)	3058	237	9.5	8.5-10.5	% <100 µg/l calculated from median.	1523	More than adequate	Risk of IIH in susceptible groups	2811	27649
Uruguay	388	3391	No data												
Uzbekistan	4316	25705	1998	National	SAC (7-10)	800	36	97.4	96.3-98.5	Median calculated from % <100 µg/l.	570	Insufficient	Moderate iodine deficiency	4204	25037
Vanuatu	38	207	No data												
Venezuela	3869	25226	2000, 2001	State	SAC	1040	286	0.0	0.0-0.0	Medians from disaggregated data by district from three state surveys pooled. % <100 µg/l calculated from median.	3168	More than adequate	Risk of IIH in susceptible groups	0	0
Viet Nam	12 520	80278	1993	National	SAC (8-12)	3062	40	84.0	82.7-85.3	Median calculated from % <100 µg/l.	1076	Insufficient	Moderate iodine deficiency	10517	67434
Yemen	4065	19315	1998	National	SAC (6-12)	974	173	30.2	27.3-33.1	% <100 µg/l calculated from median.	1561	Adequate	Optimal iodine nutrition	1228	5833
Zambia	2176	10698	1993	National	SAC	2505	60	72.0	70.2-73.8		394	Insufficient	Mild iodine deficiency	1567	7703
Zimbabwe	2555	12835	1999	National	SAC (6-14)	847	245	14.8	12.4-17.2		2641	More than adequate	Risk of IIH in susceptible groups	378	1900

^a (JUN 2003)^b Numeric references correspond to those on the WHO web site [<http://www3.who.int/whosis/micronutrient/>].

CI: Confidence interval; NS: Non specified; P: Published; SAC: School-age children

Table A3.2 Country data on TGP

Member State	Survey data							Notes
	Date of survey (yrs)	Level of survey	Population group and age (yrs)	Sample size	TGP (%)	95% confidence interval of TGP (%)	Bibliographic references ^a	
Afghanistan	1995	Province	Pre-SAC (1-5)	1553	9.9	8.4-11.4	493	
Albania	No data							
Algeria	1994 P	Local	SAC (6-11)	169	65.0	57.8-72.2	1348	Survey in endemic area. Baseline data before intervention.
Andorra	No data							
Angola	No data							
Antigua and Barbuda	No data							
Argentina	No data							
Armenia	1998	National	Women (15-45)	2569	30.2	28.4-32.0	3329	
Australia	2001	Local	SAC (11-18)	577	19.4	16.2-22.6	3598	
Austria	No data							
Azerbaijan	1996	National	Adults	904	11.1	9.1-13.2	827	
Bahamas	No data							
Bahrain	1999	National	SAC (8-12)	1600	1.7	1.1-2.3	1142	
Bangladesh	1993	National	SAC (5-11)	12 862	49.9	49.0-50.8	642a	
Barbados	No data							
Belarus	1995-98	National	SAC (6-18)	11 562	33.4	32.5-34.3	3181	
Belgium	No data							
Belize	No data							
Benin	1995	Province	SAC (7-13)	10 560	19.6	18.8-20.4	393	
Bhutan	1996	National	SAC (6-11)	1200	14.0	12.0-16.0	2649	
Bolivia	1994	National	NS	360	4.5	2.4-6.6	3612	
Bosnia and Herzegovina	1999	National	SAC (7-14)	9183	25.7	24.8-26.6	2994, 3453	TGP from two surveys (Republica Srpska and Federation of Bosnia & Herzegovina) pooled.
Botswana	1994	National	SAC (8-10)	4268	16.5	15.4-17.6	2805	
Brazil	1994-1996	National	SAC (6-14)	178 774	4.0	3.9-4.1	3599	
Brunei Darussalam	No data							
Bulgaria	1998, 2001	Region, local	SAC (6-11)	9045	27.4	26.5-28.3	1086, 3016	TGP from one regional (endemic) and one local survey pooled.
Burkina Faso	1995	Local	General population (0-45)	210	55.2	48.5-61.9	1349	
Burundi	No data							
Cambodia	1996-1997	National	SAC (8-12)	35 418	17.0	16.6-17.4	1076	
Cameroon	1995	National	SAC (8-18)	NS	10.3		401	
Canada	No data							
Cape Verde	1996	National	SAC (6-12)	1200	25.5	23.0-28.0	1622	
Central African Republic	1993-1994	Province	General population	3090	60.9	59.2-62.6	1351	
Chad	1993-94	National	SAC (10-20)	1171	63.0	60.2-65.8	390	
Chile	2001	Local urban	SAC (6-18)	3712	6.4	5.6-7.2	3335	
China	2002	National	SAC (8-10)	38 894	5.8	5.6-6.0	3579	
Colombia	1994-1998	National urban	SAC (8-12)	19 530	6.5	6.2-6.9	3296	
Comoros	1994	National	SAC (8-11)	2249	13.6	12.2-15.0	396	
Congo	1995	District	General population	2620	12.0	10.8-13.2	2279	
Cook Islands	No data							
Costa Rica	No data							
Cote d'Ivoire	1997	Local	SAC (6-15)	419	44.8	40.0-49.6	3120	
Croatia	1999	National	SAC (7-11)	2054	16.1	14.5-17.7	2941	
Cuba	No data							
Cyprus	No data							

Table A3.2

Member State	Survey data							Notes
	Date of survey (yrs)	Level of survey	Population group and age (yrs)	Sample size	TGP (%)	95% confidence interval of TGP (%)	Bibliographic references ^a	
Czech Republic	No data							
Democratic People's Republic of Korea	No data							
Democratic Republic of the Congo	1995	Local	SAC (6-14)	313	28.4	23.4-33.4	3601b	TGP from disaggregated data by region pooled.
Denmark	1997-1998	Region	Adults (18-65)	4649	12.1	11.2-13.0	3003	TGP from disaggregated data by region pooled.
Djibouti	No data							
Dominica	No data							
Dominican Republic	1993	National	SAC (6-14)	837	5.3	3.8-6.8	771	
Ecuador	No data							
Egypt	1995	National	Women	1629	21.4	19.4-23.4	486	
El Salvador	No data							
Equatorial Guinea	No data							
Eritrea	1998	National	SAC (6-12)	3270	36.7	35.1-38.4	3122	
Estonia	No data							
Ethiopia	2000 P	District	SAC	2485	53.3	51.3-55.3	3199	
Fiji	No data							
Finland	No data							
France	1996	National	Adults (35-60)	12 014	12.9	12.3-13.5	1269	TGP from disaggregated data by sex pooled.
Gabon	2001	National	SAC (6-12)	3280	17.1	15.8-18.4	3611	
Gambia	1999	National	SAC (8-12)	3010	16.3	15.0-17.6	2595	
Georgia	No data							
Germany	No data							
Ghana	1991-1994	National	NS	27 000	23.8	23.3-24.3	388	TGP from disaggregated data by district pooled.
Greece	1995 P 1999 P	Local, region	SAC (9-15)	5129	17.5	16.5-18.5	2539, 3430	TGP from one local and one regional survey pooled.
Grenada	No data							
Guatemala	No data							
Guinea	1999	Region	SAC (8-19)	1234	59.0	56.3-61.7	2617	
Guinea-Bissau	1995	National	SAC (6-12)	5480	32.6	31.4-33.8	397	
Guyana	No data							
Haiti	No data							
Honduras	No data							
Hungary	1994-1997	National	SAC (7-11, male)	299 351	11.6	11.5-11.7	3027	TGP from disaggregated data by county pooled.
Iceland	No data							
India	1992-1993, 1993, 1995 P, 1995, 1996, 1996 P, 1997, 1997 P, 1997-1998, 1998, 1998-1999, 1999, 2000 P, 2001 P, 2001, 2002	State, district	SAC	84 407	17.9	17.6-18.2	1434, 1158, 1169, 1168, 1162, 1159, 1164, 1161, 1160, 1215, 1166, 1163, 3586, 3538, 1165, 3576, 3539, 3456, 3584, 1432, 3585, 3578, 3566, 3534	TGP from 24 state and district surveys pooled at district level.
Indonesia	1995-1998	National	SAC (6-12)	1 156 367	9.8	9.8-9.9	1085	
Iran (Islamic Republic of)	1996	National	SAC (8-10)	36 178	54.6	54.1-55.1	3317	TGP from disaggregated data by province pooled.
Iraq	No data							
Ireland	No data							
Israel	No data							
Italy	1992-1994, 1993-1995, 1994, 1998 P, 1999 P	Region, local	SAC	12 744	13.9	13.3-14.5	2058, 1291, 3419, 3419, 1319, 3435, 1287, 1286, 2059	TGP from nine local and regional surveys pooled.

Table A3.2

Member State	Survey data							Notes
	Date of survey (yrs)	Level of survey	Population group and age (yrs)	Sample size	TGP (%)	95% confidence interval of TGP (%)	Bibliographic references ^a	
Jamaica	No data							
Japan	No data							
Jordan	2000	National	SAC (8-10)	2601	33.5	31.7-35.3	2534	
Kazakhstan	2000-2001	District	SAC (9-15)	3369	36.7	35.1-38.3	3055	TGP from disaggregated data by site pooled.
Kenya	1994	National	SAC (8-10)	20 916	15.5	15.0-16.0	391	
Kiribati	No data							
Kuwait	1997	National	SAC (6-9)	799	0.0	—	3135	
Kyrgyzstan	1994	Region	SAC (7-11)	440	49.1	44.4-53.8	3230	Survey in 4 out of 7 States (Oblasts). (The 4 States considered endemic.)
Lao People's Democratic Republic	No data							
Latvia	No data							
Lebanon	1993	National	SAC (7-15)	7319	25.7	24.7-26.7	485	
Lesotho	1999	National	SAC (8-12)	500	4.9	3.0-6.8	3481	
Liberia	No data							
Libyan Arab Jamahiriya	No data							
Lithuania	No data							
Luxembourg	No data							
Madagascar	1995	Local	SAC (6-11)	3635	22.8	21.4-24.2	398	
Malawi	1996	Region	SAC	9434	28.1	27.2-29.0	400	Survey in seven endemic districts.
Malaysia	1995	National	SAC (8-10)	2814	4.0	3.3-4.7	2637, 2840	TGP from two state surveys pooled.
Maldives	1995	National	SAC (6-12)	2834	23.6	22.0-25.2	2650	
Mali	No data							
Malta	No data							
Marshall Islands	No data							
Mauritania	1995	National	SAC (6-14)	4820	30.9	29.6-32.2	392	
Mauritius	No data							
Mexico	2002 P	Region	SAC (6-14)	673	10.4	8.1-12.7	12	
Micronesia (Federated States of)	No data							
Monaco	No data							
Mongolia	2001	National	SAC (5-16)	2455	23.0	21.3-24.7	3227	
Morocco	1993	National	SAC (6-12)	1594	22.0	20.0-24.0	491	
Mozambique	1998	Province	SAC	5684	14.3	13.4-15.2	2872	
Myanmar	No data							
Namibia	No data							
Nauru	No data							
Nepal	1997-1998	National	SAC (6-11)	15 542	40.0	38.8-40.2	1083	
Netherlands	1995-1996	Local	SAC (6-18)	937	1.8	1.0-2.7	3204	Thyromobile study.
New Zealand	No data							
Nicaragua	No data							
Niger	1994	National	SAC (10-15)	8933	35.8	34.8-36.8	384	
Nigeria	1995	State	SAC (6-14)	590	29.1	25.4-32.8	3601	TGP from disaggregated data by state pooled.
Niue	No data							
Norway	No data							
Oman	1993-1994	National	SAC (8-11)	2996	10.0	8.9-11.1	481	
Pakistan	1993-1994	Region	SAC (8-10)	6000	84.9	84.0-85.8	492	
Palau	No data							
Panama	1999	National	SAC (6-12)	2959	10.2	9.1-11.3	3098	
Papua New Guinea	1996	Local	SAC (8-10)	627	4.6	3.0-6.2	723	

Table A3.2

Member State	Survey data							Notes
	Date of survey (yrs)	Level of survey	Population group and age (yrs)	Sample size	TGP (%)	95% confidence interval of TGP (%)	Bibliographic references ^a	
Paraguay	No data							
Peru	No data							
Philippines	1993	National	SAC (7-14)	4579	5.4	4.8-6.1	715	
Poland	No data							
Portugal	No data							
Qatar	No data							
Republic of Korea	No data							
Republic of Moldova	1996	National	SAC (8-10)	3313	36.7	35.1-38.3	3332	
Romania	1995	Local	SAC	1010	12.8	10.7-14.9	1093	
Russian Federation	No data							
Rwanda	1996	National	SAC (5-19)	6886	25.9	24.9-26.9	2558	
Saint Kitts and Nevis	No data							
Saint Lucia	No data							
Saint Vincent and the Grenadines	No data							
Samoa	No data							
San Marino	No data							
Sao Tome and Principe	No data							
Saudi Arabia	2001, P	Region	SAC	940	24.0	21.3-26.7	567	
Senegal	1995-1996, 1996-1997	Region	SAC (10-14)	2346	28.7	26.9-30.5	1633, 1562	TGP from two regional surveys pooled.
Serbia and Montenegro	1998	National	SAC (9-18)	1421	1.3	0.7-1.9	763	Refugees survey.
Seychelles	No data							
Sierra Leone	No data							
Singapore	No data							
Slovakia	1989-1995	Region	SAC (6-15)	1923	4.4	3.5-5.3	1099	TGP from disaggregated data by age and sex pooled.
Slovenia	1994	National	SAC (13)	1740	79.0	77.1-80.9	3391	TGP measured by ultrasonography: 6.3%.
Solomon Islands	No data							
Somalia	No data							
South Africa	1998	National	SAC	2377	40.9	38.9-42.9	2618	
Spain	1995, 2000, 2002 P	Province	SAC (6-16)	2745	10.4	9.3-11.5	2091, 3581	
Sri Lanka	2001	National	SAC (8-10)	6733	20.9	19.9-21.9	404	
Sudan	1997	National	SAC	40 922	22.0	21.6-22.4	2937	
Suriname	No data							
Swaziland	1998	Local	SAC (6-18)	778	5.4	3.8-7.0	2589	Survey in one endemic site within each of Swaziland's 4 regions.
Sweden	No data							
Switzerland	1994, 1995	Local	SAC (6-17)	514	14.5	11.5-17.5	1212, 1207	TGP from two local surveys pooled.
Syrian Arab Republic	No data							
Tajikistan	1999	Local	SAC	NS	68.7		3705	
Thailand	2000	National	SAC (6-12)	3 953 730	2.2	2.19-2.21	2555	
The former Yugoslav Republic of Macedonia	2002	National	SAC (7-11)	1222	5.8	4.5-7.1	3609	
Timor Leste	No data							
Togo	1995	Region	SAC (10-19)	791	32.6	29.3-35.9	387	
Tonga	No data							
Trinidad and Tobago	No data							

Table A3.2

Member State	Survey data							Notes
	Date of survey (yrs)	Level of survey	Population group and age (yrs)	Sample size	TGP (%)	95% confidence interval of TGP (%)	Bibliographic references ^a	
Tunisia	1993	District	SAC (8–10)	1675	36.3	34.0–38.6	1559	
Turkey	1995	Province	SAC (6–12)	6906	30.3	29.2–31.4	3428	
Turkmenistan	1999	Local urban	SAC (8–10)	65	25.0	14.5–35.5	3620	
Tuvalu	No data							
Uganda	1999	National	SAC (6–12)	2870	60.2	58.4–62.0	2582	
Ukraine	1996–1999	Local	SAC (8–12)	2675	55.6	53.7–57.5	3600	
United Arab Emirates	1994	Region	SAC (9–13)	4778	40.4	39.0–41.8	483	Survey in six endemic zones.
United Kingdom of Great Britain and Northern Ireland	No data							
United Republic of Tanzania	1995	Local	SAC (6–14)	713	44.0	40.4–47.6	3601	TGP from disaggregated data by district pooled.
United States of America	No data							
Uruguay	No data							
Uzbekistan	1998	National	Pre-SAC, SAC (0–17)	19 895	73.0	72.4–73.6	570	TGP from disaggregated data by province plus Tashkent city pooled.
Vanuatu	No data							
Venezuela	No data							
Viet Nam	1993	National	SAC (8–12)	3062	21.9	20.4–23.4	1076	TGP from disaggregated data by province pooled.
Yemen	1998	National	SAC (6–12)	2984	16.8	15.5–18.1	1561	
Zambia	1993	National	SAC	2505	31.6	29.8–33.4	394	
Zimbabwe	No data							

^a Numeric references correspond to those on the WHO web site [<http://www3.who.int/whosis/micronutrient/>].

CI: Confidence interval; NS: Non specified; P: Published; SAC: School-age children

Annex 3.3 References

Afghanistan

Kakar F, Kakar S. *Indicators of child morbidity and mortality in three Afghan provinces*. Afghanistan, RACA, Afghanistan Program Office, United Nations Children's Fund, 1996. Ref 493

Algeria

Benmiloud M et al. Oral iodized oil for correcting iodine deficiency: optimal dosing and outcome indicator selection. *Journal of Clinical Endocrinology and Metabolism*, 1994, 79: 20–24. Ref 1348

Armenia

Branca F et al. *The health and nutritional status of children and women in Armenia*. Rome, National Institute of Nutrition, 1998. Ref 3329

Australia

Guttikonda K et al. Recurrent iodine deficiency in Tasmania, Australia: a salutary lesson in sustainable iodine prophylaxis and its monitoring. *Journal of Clinical Endocrinology and Metabolism*, 2002, 87:2809–2815. Ref 3379

McDonnell CM, Harris M, Zacharin MR. Iodine deficiency and goitre in schoolchildren in Melbourne, 2001. *Medical Journal of Australia*, 2003, 178:159–162. Ref 3598

Austria

Delange F et al. Thyroid volume and urinary iodine in European schoolchildren: standardization of values for assessment of iodine deficiency. *European Journal of Endocrinology*, 1997, 136:180–187. Ref 1319

Azerbaijan

Markou KB et al. Iodine deficiency in Azerbaijan after the discontinuation of an iodine prophylaxis program: reassessment of iodine intake and goiter prevalence in schoolchildren. *Thyroid*, 2001, 11:141–146. Ref 3413

Branca F et al. *Health and nutrition survey of internally displaced and resident population of Azerbaijan – April 1996*. US Agency for International Development, World Health Organization, United Nations Children's Fund, 1996. Ref 827

Bahrain

Moosa K et al. *National study on the prevalence of iodine deficiency disorders among school children aged 8–12 years old in Bahrain*. Manama, Ministry of Health, 2000. Ref 1142

Bangladesh

Yusuf HKM et al. *Report of the National Iodine Deficiency Disorders Survey in Bangladesh – 1993*. Dhaka, Dhaka University, 1993. Ref 642

Belarus

Arinchin A et al. Goiter prevalence and urinary iodine excretion in Belarus children born after the Chernobyl accident. *IDD Newsletter*, 2000, 16:7–9. Ref 3181

Belgium

Delange F et al. Silent iodine prophylaxis in Western Europe only partly corrects iodine deficiency: the case of Belgium. *European Journal of Endocrinology*, 2000, 143: 89–196. Ref 1336

Belize

National Iodine Survey Belize 1994–1995. Belmopan, Ministry of Health, 1995. Ref 3133

Benin

Kibambe N et al. *Projet Thyromobil en Afrique de l'Ouest (première phase). Rapport final. [Thyromobil project in West Africa (first phase). Final report.]* Lomé, Conseil International de Lutte contre les Troubles Dus à la Carence en Iode, 2000. Ref 2535

Ategbo EA. *Résultats de l'enquête transversale sur la prévalence du goitre effectuée dans la partie nord du Bénin [Results of cross-sectional goiter prevalence study in the northern part of Benin] – draft*. Cotonou, Université Nationale du Bénin, 1995. Ref 393

Bhutan

Royal Government of Bhutan et al. *Tracking Progress towards sustainable elimination of IDD in Bhutan*. Thimphu, Royal Government, 1996. Ref 2649

Bolivia

Gutierrez Sardan M. *Encuesta Nacional de Múltiples Indicadores 1996 [National Multiple Indicator Cluster Survey 1996]*. La Paz, Dirección Nacional de Epidemiología – Fondo de las Naciones Unidas para la Infancia, 1997. Ref 3339

[Anonymous]. Bolivia conquers iodine deficiency. *IDD Newsletter*, 1996, 12:33–34. Ref 3612

Bosnia and Herzegovina

Lolic A, Lolic B. *Iodine deficiency*. Banja Luka, United Nations Children's Fund, 1999. Ref 2994

Tahirovic H et al. Assessment of the current status of iodine prophylaxis in Bosnia and Herzegovina Federation. *Journal of Clinical Endocrinology and Metabolism*, 2001, 14: 139–144. Ref 3453

Botswana

Micronutrient malnutrition in Botswana. A National survey to assess the status of iodine, iron, and vitamin A. Gaborone, Ministry of Health, 1996. Ref 2805

Brazil

Pretell EA. *Thyromobil project in Latin America; Report of the study in Brazil*. Brasilia, Ministry of Health, 2002. Ref 3350

Correa Filho HR et al. Inquérito sobre a prevalência de bócio endêmico no Brasil em escolares de 6 a 14 anos: 1994 a 1996. [Endemic goiter prevalence survey in Brazilian schoolchildren 6 to 14 years old, 1994–1996]. *Revista Panamericana de Salud Pública/Pan American Journal of Public Health*, 2002, 12:317–326. Ref 3599

Bulgaria

Ivanova L et al. Urinary Iodine in IDD monitoring in Bulgaria, In: Geertmann RM, ed. *8th World Salt Symposium*, Amsterdam, Elsevier, 2000:1249–1250. Ref 3017

Timtcheva T. *Information on IDD situation in Bulgaria* [unpublished report]. Sofia, Ministry of Health, 1999. Ref 1086

Kovatcheva R et al. Thyroid volume and urinary iodine in Bulgarian school children from Sofia's region [in Bulgarian]. *Endocrinologia*, 2001, 6. Ref 3016

Burkina Faso

Kibambe N et al. *Projet Thyromobil en Afrique de l'Ouest (première phase). Rapport final. [Thyromobil project in West Africa (first phase). Final report]*. Lumé, Conseil International de Lutte contre les Troubles Dus à la Carence en Iode, 2000. Ref 2535

Thiebaut R et al. Prévalence du goitre endémique dans le secteur sanitaire de Zitenga (Burkina Faso) [Prevalence of endemic goiter in the health sector of Zitenga (Burkina Faso)]. *Santé*, 1998, 8:269–274. Ref 1349

Cambodia

Ministry of Health of Cambodia, National Maternal and Child Health Center. *Workshop on regional strategy for control of IDD*. Cambodia, The National Sub Committee for control of IDD (NSCIDD), 1997. Ref 1076

Cameroon

Lantum DN. *Monitoring and evaluation of IDD elimination programme in Cameroon 1992–1993*. Yaoundé, University of Yaoundé I, 1995. Ref 1431

Lantum DN. *Action towards elimination of iodine deficiency disorders 1990–1995*. Yaoundé, University of Yaoundé I, 1995. Ref 401

Cape Verde

Ntambwe-Kibambe. *Enquête nationale sur la carence en iode et la consommation du sel au Cap Vert – rapport final [National survey on iodine deficiency and salt consumption*

in Cape Verde – final report]. Cape Verde, Ministère de la Santé et Promotion Sociale, 1997. Ref 1622

Central African Republic

Yazipo D et al. Effectiveness of a water iodination system for preventing iodine deficiency diseases in Central Africa. *Santé*, 1995, 5:9–17. Ref 1351

Chad

Ministère de la Santé Publique Tchad, WHO, UNICEF. *Résultats de l'enquête nationale de prévalence des TDCI en République du Tchad – du 05/11/93 au 06/01/94 [Results of the national survey on IDD prevalence in the Republic of Chad – from 05/11/93 to 06/01/94]*. N'Djaména, Ministère de la Santé Publique, 1994. Ref 390

Chile

Muzzo SB, Leiva L, Mora JR. *Nutricion de yodo del escolar Chileno despues del cambio del reglamento sanitario de los alimentos [Iodine nutrition in Chilean schoolchildren after the revision of the sanitary regulation on foods]*. Santiago de Chile, Universidad de Chile, 2002. Ref 3335

China

Center for Endemic Disease Control, Chinese Center for Disease Control and Prevention. Specialist Group of the Fourth National Survey on IDD. *China national iodine deficiency disorders surveillance report 2002*. Beijing, Chinese Center for Disease Control and Prevention, 2003. Ref 3579

Colombia

Ruiz H, Jimenez G. *Prevalencia de los desórdenes por deficiencia de yodo e ingestión promedio de sal Colombia 1994–1998 [Prevalence of iodine deficiency disorders and salt consumption, Colombia 1994–1998]*. Bogotá, Instituto Nacional de Salud, 2001. Ref 3296

Comoros

Rapport des résultats de l'enquête sur la prévalence du goître et l'avitaminose A – Enquête organisée du 8 au 14 décembre 1994 [Report of the results from the survey on goiter prevalence and vitamin A deficiency – Survey from 8th to 14th December 1994]. Moroni, Ministère de la Santé Publique République Fédérale Islamique des Comores, 1995. Ref 396

Congo

Doumtabé NL. *Rapports de mission de la distribution de l'huile iodée dans le district de Dongou [Mission report of iodized oil distribution in the Dongou district]*. Brazzaville, WHO, 1996. Ref 2279

Costa Rica

Encuesta Nacional de Nutrición: 2 Fascículo Micronutrientes [National nutrition survey: Part 2 micronutrients]. San José, Ministerio de Salud, 1996. Ref 1634

Côte d'Ivoire

Hess SY et al. Treatment of iron deficiency in goitrous children improves the efficacy of iodized salt in Cote d'Ivoire. *American Journal of Clinical Nutrition*, 2002, 75:743–748. Ref 3239

Zimmermann M et al. Persistence of goiter despite oral iodine supplementation in goitrous children with iron deficiency anemia in Cote d'Ivoire. *American Journal of Clinical Nutrition*, 2000, 71:88–93. Ref 3120

Croatia

Kusic Z et al. Croatia has reached iodine sufficiency. *Journal of Endocrinological Investigation*, 2003, 26:738–742. Ref 3429

Croatian National Institute of Public Health, UNICEF Office for Croatia, Kiwanis International. *Public Health Significance of iodine deficiency disorders in Croatia: Results of the 1997–99 Eradication Program*. Zagreb, Croatian National Institute of Public Health, 2000. Ref 2941

Cuba

Rodriguez-Ojea A et al. Low levels of urinary iodine excretion in schoolchildren of rural areas in Cuba. *European Journal of Clinical Nutrition*, 1998, 52:372–375. Ref 1512

Czech Republic

Zamrazil V. Iodine deficiency. In: Levinski A, ed. *Abstracts of the 1st EFES Regional Polish-Slovak-Czech-Hungarian Postgraduate course in clinical endocrinology*. Polanczyk, Polish Society of Endocrinology, 2000. Ref 515

Democratic Republic of the Congo

WHO, UNICEF, ICCIDD. *Review of findings from 7-country study in Africa on levels of salt iodization in relation to iodine deficiency disorders, including iodine induced hyperthyroidism*. Geneva, World Health Organization, 1997 (WHO/AFRO/NUT/97.2). Ref 3601

Denmark

Rasmussen LB et al. Dietary iodine intake and urinary iodine excretion in a Danish population: effect of geography, supplements and food choice. *British Journal of Nutrition*, 2002, 87:61–69. Ref 3205

Knudsen N et al. Goitre prevalence and thyroid abnormalities at ultrasonography: a comparative epidemiological study in two regions with slightly different iodine status. *Clinical Endocrinology*, 2000, 53:479–485. Ref 3003

Dominican Republic

Noguera A, Gueri M. *Analisis de la situacion de deficiencia de yodo en America Latina – sus tendencias y estrategias de accion [Analysis of the situation of iodine deficiency in Latin America – trends and strategies]*. Washington, D.C., Pan American Health Organization, 1994. Ref 771

Ecuador

[Anonymous]. The western hemisphere nears iodine sufficiency. *IDD Newsletter*, 2001,17:1–9. Ref. 3394

Egypt

Shaheen FM, Hassan H. *Nutritional deficiencies among primary school children in Egypt*. Cairo, Ministry of Health and Population Egypt. 1998. Ref 2639

Ministry of Health and Population Egypt, UNICEF. *National survey for assessment of vitamin A status in Egypt*. Cairo, Ministry of Health and Population Egypt, 1995. Ref 486

El Salvador

Primer estudio nacional sobre excrecion urinaria de yodo en escolares en El Salvador [First national survey on urinary iodine excretion in schoolchildren in El Salvador]. San Salvador, Ministerio de Salud Publica y Asistencia Social El Salvador, 1998. Ref 3108

Eritrea

National IDD Survey 1998. Draft report. Asmara, Ministry of Health, 1998. Ref 3122

Estonia

Veinpalu M et al. Urinary iodine excretion in Estonian children. *European Journal of Endocrinology*, 1996, 135: 248. Ref 1225

Ethiopia

Cherinet A, Kelbessa U. Determinants of iodine deficiency in school children in different regions of Ethiopia. *East African Medical Journal*, 2000, 77:133–137. Ref 3199

Fiji

UNICEF-Pacific Office. *IDD in Fiji: Outcome and follow up to a consultancy visit by Dr Rainer Gutekunst (August 1994)*. Suva, United Nations Children's Fund, 1997. Ref 3236

Finland

Valsta M et al. Iodine status of middle-aged subjects in Finland [abstract]. In: *Spring meeting of the Finnish Society for Nutrition Research 2003*. Kuopio, Finnish Society for Nutrition Research, 2003. Ref 3605

France

Valeix P et al. Iodine deficiency in France. *Lancet*, 1999, 353:1766–1767. Ref 1269

Gabon

[Anonymous]. IDD in some Sub-Saharan Countries. *IDD Newsletter*, 2002, 18:4. 2002. Ref 3611

Gambia

Egbuta JO. *Consultancy report on iodine deficiency disorders survey in The Gambia*. Banjul, The National Nutrition Agency, 2001. Ref 2595

Georgia

Gerasimov G. IDD in Eastern Europe and Central Asia. *IDD Newsletter*, 2002, 18:33–37. Ref 3699

Germany

Hampel R et al. Jodidurie bei Schulkindern in Deutschland 1999 im Normbereich [Urinary iodine levels within normal range in German school-age children]. *Medizinische Klinik*, 2001, 96:125–128. Ref 3126

Ghana

The Partnership for Child Development. The health and nutritional status of schoolchildren in Africa: evidence from school-based health programmes in Ghana and Tanzania. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1998, 92:254–261. Ref 1772

Asibey-Berko E, Orroca-Tetteh R, eds. *Proceedings of the National Workshop on Iodine Deficiency Disorders in Ghana, July 1994*. Accra, University of Ghana, 1995. Ref 388

Greece

Tsatsoulis A et al. An epidemiological survey on the prevalence of goiter among schoolchildren in Northwestern Greece. *European Journal of Internal Medicine*, 1996, 7: 35–39. Ref 2539

Doufas AG et al. The predominant form of non-toxic goiter in Greece is now autoimmune thyroiditis. *European Journal of Endocrinology*, 1999, 140:505–511. Ref 3430

Guatemala

Encuesta nacional de micronutrientes [National micronutrient survey]. Guatemala City, Ministerio de Salud Publica y Asistencia Social Guatemala, 1996. Ref 3091

Guinea

Ingenbleek Y. *Evaluation des troubles dus à la carence iodée dans les préfectures de Kindia, Labé, Lola et Siguirí [Evaluation of iodine deficiency disorders in the prefectures of Kindia, Labé, Lola and Siguirí]*. United Nations Children's Fund, 1999. Ref 2617

Guinea-Bissau

Doh A. *Rapport d'enquête nationale sur les manifestations de la carence en iode en Guinée-Bissau [Report of the national iodine deficiency survey in Guinea-Bissau]*. Lomé, Organisation de Coordination et de Coopération pour la lutte contre les Grandes Endémies, Centre régional de Recherches en Alimentation et Nutrition, 1995. Ref 397

Guyana

Ministry of Health of Guyana, Caribbean Food and Nutrition Institute, PAHO. *Executive summary micronutrient study report – Guyana. An assessment of the vitamin A, beta-Carotene, iron and iodine status in the population*. Georgetown, Ministry of Health, 1997. Ref 3094

Honduras

[Anonymous]. The western hemisphere nears iodine sufficiency. *IDD Newsletter*, 2001, 17:1–9. Ref 3394

Hungary

Farkas I et al. Golyvagyakorisság, jodellátottság az általános iskolák I–IV. Oszályos fiú tanulóinak körében Budapesten 1996-ban. In *Budapesti Közegészségügy*. Budapest, 1997. Ref 3041

Farkas I, Sajgo M, Fehér Z. *Jelentés az 1997 évben I–IV osztályos fiútanulókon végzett tapintásos golyvagyakorisság és vizeletjód vizsgálatok eredményeiről [Report of 1997 survey of primary school boys, I–IV grade: goitre prevalence and urinary iodine excretion]*. Budapest, “Johan Bela” National Public Health Institute, 1997. Ref 3684

Farkas I, Sajgo M, Fehér Z. *Jelentés az 1996 évben I–IV osztályos fiútanulókon végzett tapintásos golyvagyakorisság és vizeletjód vizsgálatok eredményeiről [Report of 1996 survey of primary school boys, I–IV grade: goitre prevalence and urinary iodine excretion]*. Budapest, “Johan Bela” National Public Health Institute, 1996. Ref 3683

Farkas I, Sajgo M, Fehér Z. *Jelentés az 1995/1996. ös tanévben I–IV osztályos fiútanulókon végzett tapintásos golyvagyakorisság és vizeletjód vizsgálatok eredményeiről [Report of 1995/1996 survey of primary school boys, I–IV grade: goitre prevalence and urinary iodine excretion]*. Budapest, “Johan Bela” National Public Health Institute, 1995. Ref 3682

Farkas I, Sajgo M, Fehér Z. *Jelentés az 1994/1995. ös tanévben I–IV osztályos fiútanulókon végzett tapintásos golyvagyakorisság és vizeletjód vizsgálatok eredményeiről [Report of 1994/1995 survey of primary school boys, I–IV grade: goitre prevalence and urinary iodine excretion]*. Budapest, “Johan Bela” National Public Health Institute, 1994. Ref 3681

Farkas I, Sajgo M. Goitre frequency among boys, primary school class I–IV. Reports to the Ministry of Health 1994/95, 1995/96, 1996/97. *Public Health of Budapest* 1997, 1: 15–19. Ref 3027

Iceland

Laurberg P et al. Iodine intake and the pattern of thyroid disorders: A comparative epidemiological study of thyroid abnormalities in the elderly in Iceland and in Jutland, Denmark. *Journal of Clinical Endocrinology and Metabolism*, 1998, 83:765–769. Ref 1251

India

Mallik AK et al. Iodine deficiency disorders in Car Nicobar (Andaman and Nicobar Islands). *The National Medical Journal of India*, 1998, 11:9–11. Ref 1158

Kapil U et al. Status of iodine deficiency in selected blocks of Kangra district, Himachal Pradesh. *Indian Pediatrics*, 1997, 34:338–340. Ref 1162

Bhardwaj AK et al. Assessment of iodine deficiency in district Bikaner, Rajasthan. *Indian Journal of Maternal and Child Health*, 1997, 8:18–20. Ref 1159

Sohal KS et al. Assessment of iodine deficiency disorders in district Hamirpur, Himachal Pradesh. *Indian Pediatrics*, 1998, 35:1008–1011. Ref 1164

Kapil U et al. Assessment of iodine deficiency disorders using the 30 cluster approach in the national capital territory of Delhi. *Indian Pediatrics*, 1996, 33:1013–1017. Ref 1161

Kapil U et al. Iodine deficiency in district Kinnaur, Himachal Pradesh. *Indian Journal of Pediatrics*, 1998, 65:451–453. Ref 1160

Kapil U, Ramachandran S, Tandon M. Assessment of iodine deficiency in Andaman district of Union Territory of Andaman and Nicobar. *Indian Journal of Maternal and Child Health*, 1998, 9:19–20. Ref 1215

Kapil U et al. Assessment of iodine deficiency in selected blocks of east and west Champaran districts of Bihar. *Indian Pediatrics*, 1997, 34:1087–1091. Ref 1166

Kapil U, Ramachandran S, Tandon M. Assessment of iodine deficiency in Pondicherry. *Indian Pediatrics*, 1998, 35:357–359. Ref 1163

Kapil U et al. Status of iodine deficiency in selected hill districts of Uttar Pradesh – a pilot study. *Indian Journal of Maternal and Child Health*, 1999, 10:24–27. Ref 3538

Kapil U, Tandon M, Pathak P. Assessment of iodine deficiency in Ernakulam district, Kerala state. *Indian Pediatrics*, 1999, 36:178–180. Ref 1165

Kapil U et al. Assessment of iodine deficiency disorders using the 30 cluster approach in district Kangra, Himachal Pradesh, India. *Journal of Tropical Pediatrics*, 2000, 46:264–266. Ref 3539

Kapil U et al. Assessment of iodine deficiency in Kottayam district, Kerala State: a pilot study. *Asia Pacific Journal of Clinical Nutrition*, 2002, 11:33–35. Ref 3456

Mittal M, Tandon M, Raghuvanshi RS. Iodine status of children and use of iodized salt in Tarai region of North India. *Journal of Tropical Pediatrics*, 2000, 46:300–302. Ref 3584

Bhasin SK, Kumar P, Dubey KK. Comparison of urinary iodine excretion and goiter survey to determine the prevalence of iodine deficiency. *Indian Pediatrics*, 2001, 38:901–905. Ref 3585

Chandra AK, Ray I. Dietary supplies of iodine and thiocyanate in the etiology of endemic goiter in Tripura. *Indian Journal of Pediatrics*, 2001, 68:399–404. Ref 3578

Brahmbhatt SR et al. Biochemical assessment of iodine deficiency disorders in Baroda and Dang districts of Gujarat State. *Indian Pediatrics*, 2001, 38:247–255. Ref 3577

Brahmbhatt SR et al. Study of biochemical prevalence indicators for the assessment of iodine deficiency disorders in adults at field conditions in Gujarat (India). *Asia Pacific Journal of Clinical Nutrition*, 2001, 10:51–57. Ref 3565

All India Institute of Medical Sciences, Government of India, Salt Department. *Monitoring of quality of iodised salt to prevent iodine deficiency disorders and increase production of iodised salt through networking of medical colleges in Andhra Pradesh, Karnataka, Kerala, Tamil Nadu and Pondicherry states of India*. Dew Delhi, All India Institute of Medical Sciences, 2002. Ref 3545

Kapil U et al. Assessment of iodine deficiency disorders in district Bharatpur, Rajasthan. *Indian Pediatrics*, 2003, 40:147–149. Ref 3534

Patowary AC et al. Iodine deficiency disorders (IDD) and iodised salt in Assam: a few observations. *Indian Journal of Public Health*, 1995, 39:135–140. Ref 1434

Zargar AH et al. Prevalence of goitre in school children in Baramulla (Kashmir Valley). *Indian Journal of Pediatrics*, 1997, 64:225–230. Ref 1169

Zargar AH et al. Prevalence of goiter in schoolchildren in Kashmir Valley, India. *American Journal of Clinical Nutrition*, 1995, 62:1020–1021. Ref 1168

Gakkhar RK et al. Prevalence of goitre in school-going children in Jabalpur. *Indian Journal of Pediatrics*, 2001, 68:229–233. Ref 3586

Rao RS et al. Prevalence of goitre among school children in coastal Karnataka. *Indian Journal of Pediatrics*, 2002, 69:477–479. Ref 3576

Brahmbhatt S, Brahmbhatt RM, Boyages SC. Thyroid ultrasound is the best prevalence indicator for assessment of iodine deficiency disorders: a study in rural/tribal school-children from Gujarat (Western India). *European Journal of Endocrinology*, 2000, 143:37–46. Ref 1432

Biswas AB et al. Iodine deficiency disorders among school children of Malda, West Bengal, India. *Journal of Health, Population and Nutrition*, 2002, 20:180–183. Ref 3566

Indonesia

Pardede LVH et al. Urinary iodine excretion is the most appropriate outcome indicator for iodine deficiency at field conditions at district level. *Journal of Nutrition, Growth and Cancer*, 1998, 128:1122–1126. Ref 1183

WHO, World Bank, Ministry of Health Indonesia. *Iodine deficiency in Indonesia: A detailed nationwide map of goitre prevalence*. Geneva, World Health Organization, 2001, (WHO/NHD/01.4). (Available on request from Department of Nutrition for Health and Development, World Health Organization, 1211 Geneva 27, Switzerland). Ref 1085

Iran (Islamic Republic of)

Azizi F et al. Sustainable control of iodine deficiency in Iran: Beneficial results of the implementation of the mandatory law on salt iodization. *Journal of Endocrinological Investigation*, 2002, 25:409–413. Ref 3317

Ireland

Smyth PPA. *IDD status in Ireland* [unpublished data]. Dublin, University College of Dublin, 2002. Ref 3608

Italy

Frigato F et al. Epidemiological survey of goiter and iodine deficiency in Veneto region. *Journal of Endocrinological Investigation*, 1996, 19:734–738. Ref 2058

Pagliara S et al. Diffusione del gozzo endemico e della carenza iodica in provincia di Avellino [Widespread endemic goiter and iodine deficiency in the province of Avellino]. *Annali dell Istituto Superiore di Sanita*, 1998, 34:417–421. Ref 1291

Vitti P et al. Thyroid volume measurement by ultrasound in children as a tool for the assessment of mild iodine deficiency. *Journal of Clinical Endocrinology and Metabolism*, 1994, 79:600–603. Ref 3419

Aghini-Lombardi F. Effect of iodized salt on thyroid volume of children living in an area previously characterized by moderate iodine deficiency. *Journal of Clinical Endocrinology and Metabolism*, 1997, 82:1136–1139. Ref 1273

Zini M et al. Indagine epidemiologica sulla prevalenza del gozzo e sulla escrezione urinaria di iodio nella popolazione scolastica della provincia di Reggio Emilia [Epidemiologic investigation on the prevalence of goiter and urinary excretion of iodine in the school population of the province of Reggio Emilia]. *Annali dell Istituto Superiore di Sanita*, 1998, 34:383–387. Ref 1287

Cassio A et al. Prevalenza di gozzo e ioduria nella popolazione della scuola dell'obbligo in un'area dell'Appennino bolognese [Prevalence of goiter and urinary iodine in a school population in an area of the Bolognese Apennines]. *Annali dell Istituto Superiore di Sanita*, 1998, 34:389–391. Ref 1286

Lupoli G et al. Evaluation of goiter endemia by ultrasound in schoolchildren in Val Sarmiento (Italy). *Journal of Endocrinological Investigation*, 1999, 22:503–507. Ref 2059

Rapa A et al. Iodine deficiency in Italy. *Lancet*, 1999, 354:596–597. Ref 1272

Delange F et al. Thyroid volume and urinary iodine in European schoolchildren: standardization of values for assessment of iodine deficiency. *European Journal of Endocrinology*, 1997, 136:180–187. Ref 1319

Panunzi C et al. Prevalenza di gozzo ed escrezione urinaria di iodio in un campione di bambini in età scolare della città di Roma [Goiter prevalence and urinary excretion of iodine in a sample of school age children in the city of Rome]. *Annali dell Istituto Superiore di Sanita*, 1998, 34:409–412. Ref 3435

Jordan

Kharabsheh SA, Belbesi A, Qarqash W. *Assessment of iodine deficiency status among Jordanian children after introduction of iodized salt*. Jordan, Ministry of Health, 2000. Ref 2534

Kazakhstan

Ospanova F. Iodine urine excretion as estimation of +A29 status of reproductive age women. *Astana Medical Journal*, 2000, 1:109–111. Ref 3056

Zeltser ME et al. *Summary of 2000–2001 survey* [unpublished data]. Almaty, Kazakh Academy of Nutrition. Ref 3055

Kenya

Gitau W. *Report of the National Micronutrient Survey – February to August 1994, iodine deficiency disorders*. Nairobi, University of Nairobi, 1994. Ref 391

Kuwait

Toward a national plan for prevention and control of iron deficiency and its anemia. Kuwait, Ministry of Health, 2001. Ref 3135

Kyrgyzstan

Houston R, Rashid B, Kalanzi J. *Rapid assessment of iodine deficiency in Kyrgyzstan*. United Nations Children's Fund, 1994. Ref 3230

Lao People's Democratic Republic

Report on National Health Survey. Health status of the people of Lao PDR. Vientiane, Ministry of Health, 2001. Ref 770

Latvia

Selga G, Sauka M, Gerasimov G. Status of iodine deficiency in Latvia reconsidered: results of nation-wide survey of 587 school children in the year 2000. *IDD Newsletter*, 2000, 16:54. Ref 3058

Lebanon

Ministry of Public Health of Lebanon, UNICEF. *Evaluation Studies of the IDD programme in Lebanon 1997*. Beirut, Ministry of Public Health, 1997. Ref 3222

Ministry of Health of Lebanon, UNICEF, WHO, American University of Beirut. *IDD survey in Lebanon, 1993 – final report*. Beirut, Ministry of Health, 1994. Ref 485

Lesotho

Sebotsa MLD et al. Prevalence of goitre and urinary iodine status of primary school children in Lesotho. *Bulletin of the World Health Organization*, 2003, 81:28–34. Ref 3481

Liberia

Mulder-Sibanda M et al. *National Micronutrient Survey. A national prevalence study on vitamin A deficiency, iron deficiency anemia, iodine deficiency*. Monrovia, Ministry of Health and Social Welfare, 1999. Ref 1242

Lithuania

Knowles J. IDD in the Baltics. *IDD Newsletter*, 2000, 16: 53. Ref 3613

Luxembourg

Delange F et al. Thyroid volume and urinary iodine in European schoolchildren: standardization of values for assessment of iodine deficiency. *European Journal of Endocrinology*, 1997, 136:180–187. Ref 1319

Madagascar

Hantaniaina R. *Rapport de l'évaluation des 7 postes sentinelles TDCI [Report of IDD evaluation in 7 sentinel sites]*. Antananarivo, Service de Nutrition et Alimentation, 1995. Ref 398

Malawi

Mdebwe HJ, Banda TW. *A survey report on the trend of prevalence of goitre in junior primary school pupils and coverage of iodized salt at household level in seven districts in northern and central regions of Malawi*. Lilongwe, Ministry of Health and Population, 1996. Ref 400

Malaysia

Institute for Medical Research, Ministry of Health of Malaysia. *Survey of IDD in Peninsula Malaysia*. Kuala Lumpur, Ministry of Health, 1995. Ref 2637

Tayong IB et al. *Iodine deficiency disorders in Sabab (a prevalence survey)*. Sabah, State Health Department, 2001. Ref 2840

Maldives

Pandav CS (ICCIDD), Government of Maldives, Department of Public Health, UNICEF Malé. *Iodine deficiency disorders in Maldives*. Malé, International Council for Control of Iodine Deficiency Disorders, 1995. Ref 2650

Mali

Kibambe N et al. *Projet Thyromobil en Afrique de l'Ouest (première phase). Rapport final. [Thyromobil project in West Africa (first phase). Final report.]* Lomé, Conseil International de Lutte contre les Troubles Dus à la Carence en Iode, 2000. Ref 2535

Mauritania

Kibambe N. *Rapport traitement et analyse – enquête TDCI Mauritanie [Data and analysis report – IDD survey Mauritania]*. Nouakchott, Ministère de la Santé et des Affaires Sociales, 1995. Ref 392

Mauritius

A survey of nutrition in Mauritius and Rodrigues (1995) – final report. Port Louis, Ministry of Health, 1995. Ref 395

Mexico

Encuesta Nacional de Nutrición 1999 [National nutrition survey 1999]. Mexico City, Instituto Nacional de Salud Publica, 1999. Ref 2997

Martinez-Salgado H et al. Deficiencia de yodo y otros posibles bociógenos en la persistencia del bocio endémico en México [Iodine deficiency and other potential goitrogens in the persistence of endemic goiter in Mexico]. *Gaceta Medica de Mexico*, 2002, 138:149–156. Ref 12

Mongolia

Bolormaa et al. *Report on national survey of iodine deficiency disorders in Mongolia in 2001*. Ulaanbaatar, Ministry of Health, 2001. Ref 3227

Morocco

Chaouki N et al. Etude de prévalence des troubles dus à la carence iodée chez les enfants âgés de 6 à 12 ans au Maroc [Study of prevalence of iodine deficiency disorders in children aged 6 to 12 years in Morocco]. *Bulletin Épidémiologique*, 1996, 17–20:2–15. Ref 491

Mozambique

Fildalgo L, Ismael C, Khan S. *Avaliação da deficiência em micronutrientes a nível das províncias de C. Delgado, Manica, Gaza e Maputo* [Evaluation of micronutrient deficiency in the provinces of C. Delgado, Manica, Gaza and Maputo]. Maputo, Ministerio de Saude, 1999. Ref 2872

Myanmar

Naing KM. *Assessment of urinary iodine levels among school age children in Myanmar, 2001 (the second national urinary iodine survey)*. Report to UNICEF. Myanmar, United Nations Children's Fund, 2001. Ref 3076

Nepal

Ministry of Health of Nepal, New ERA, The Micronutrient Initiative, UNICEF Nepal, WHO. *Nepal Micronutrient Status Survey 1998*. Kathmandu, Ministry of Health, 1999. Ref 1083

Netherlands

Wiersinga WM et al. A survey of iodine intake and thyroid volume in Dutch schoolchildren: reference values in an iodine-sufficient area and the effect of puberty. *European Journal of Endocrinology*, 2001, 144:595–603. Ref 3204

New Zealand

Skeaff SA, Thomson CD, Gibson RS. Mild iodine deficiency in a sample of New Zealand schoolchildren. *European Journal of Clinical Nutrition*, 2002, 56:1169–1175. Ref 3597

Nicaragua

Segunda encuesta nacional de micronutrientes (II ENM 2000) [Second national micronutrient survey (II ENM 2000)]. Managua, Ministerio de Salud, 2000. Ref 3109

Niger

[Anonymous]. IDD in some Sub-Saharan countries. *IDD Newsletter*, 2002, 18:2–6. Ref 3611

Hamani D. *Enquête nationale sur la prévalence du goitre au Niger – Rapport No.1* [National survey on goiter prevalence in Niger – Report no. 1]. Niamey, United Nations Children's Fund, 1994. Ref 384

Nigeria

Egbuta J, Onyezili F, Vanormelingen K. Impact evaluation of efforts to eliminate iodine deficiency disorders in Nigeria. *Public Health Nutrition*, 2003, 6:169–173. Ref 3604

WHO, UNICEF, ICCIDD. *Joint WHO/UNICEF/ICCIDD Consultation. Review of findings from 7-country study in Africa on levels of salt iodization in relation to iodine deficiency disorders, including iodine induced hyperthyroidism*. Geneva, World Health Organization, 1997. (Available on request from Department of Nutrition for Health and Development, World Health Organization, 1211 Geneva 27, Switzerland). Ref 3601

Oman

Ministry of Health of Oman, WHO, UNICEF, Sultan Qaboos University. *National study on prevalence of iodine deficiency disorders (IDD) in Oman*. Muscat, Ministry of Health, 1993. Ref 481

Pakistan

Gardezi SMA. *Iodine deficiency disorders prevalence survey in Azad Jammu & Kashmir, Pakistan 1994*. Kashmir, United Nations Children's Fund, Directorate of Health Services Azad Jammu & Kashmir, 1996. Ref 492

Panama

Encuesta nacional de bocio y anemia en escolares de 6 a 12 años [National survey of goiter and anemia in school children 6 to 12 years old]. Panama City, Ministerio de Salud, 1999. Ref 3098.

Papua New Guinea

Amoa B, Attah Johnson F, Saweri W. *Iodine deficiency disorders in Huon and Menyamya districts of Morobe province*. United Nations Children's Fund, 1997. Ref 723

Paraguay

Sanchez S, Bianchetto F. *Deficiencia de micronutrientes* [Micronutrient deficiencies]. Asunción, Instituto de Nacional de Alimentación y Nutrición, 2002. Ref 3167

Peru

Evaluación de la situación de los desórdenes por deficiencia de yodo en las áreas de riesgo de 1999 [Evaluation of iodine deficiency disorders in risk areas in 1999]. Lima, Dirección General de Salud de Las Personas Peru, 1999. Ref 3147

Philippines

Food and Nutrition Research Institute, Department of Science and Technology. *Philippine Nutrition Facts & Figures*. Metro Manila, Food and Nutrition Research Institute, 2001. Ref 2536

Food and Nutrition Research Institute, Department of Science and Technology. *Fourth national nutrition survey*. Metro Manila, Food and Nutrition Research Institute, 1993. Ref 715

Poland

Szybinski Z et al. A programme of iodine supplementation using only iodised household salt is efficient – the case of Poland. *European Journal of Endocrinology*, 2001, 144: 331–337. Ref 2574

Qatar

Verster A. Country activities: Qatar. *IDD Newsletter*, 1996, 12:23. Ref 3614

Republic of Moldova

UNICEF Bureau pour la Moldavie. Rapport de mission de consultation (17–27 septembre 1996) [Mission report (17–27 September 1996)]. In: Chauliac M et al. eds. *Situation alimentaire et nutritionnelle des mères et des enfants Moldaves*, 1996. Ref 3332

Romania

Simescu M, Dimitriu I, Sava M. *Iodine concentration in spot urine samples of school children from 30 counties 2000–2001*. Bucharest, Ministry of Health, 2002. Ref 3544

Circo E et al. Study of the nontoxic goiter prevalence in a paraendemic territory. *Romanian Journal of Endocrinology*, 1996, 34:25–39. Ref 1093

Russian Federation

Endocrinology Research Center, Russian Academy of Medical Science, National IDD Center, Ministry of Health of Russian Federation. *Report on IDD survey in the Tyva Republic*. Moscow, Endocrinology Research Center, Russian Academy of Medical Science, 2000. Ref 1191

Endocrinology Research Center, Russian Academy of Medical Science, National IDD Center, Ministry of Health. *Report on IDD survey in the Belgorod Oblast*. Moscow, Endocrinology Research Center, Russian Academy of Medical Science, 2001. Ref 743

Endocrinology Research Center, Russian Academy of Medical Science, National IDD Center, Ministry of Health, Health Department of Volgograd Oblast. *IDD survey in the Volgogr Oblast*. Moscow, Endocrinology Research Center, Russian Academy of Medical Science, 2001. Ref 1222

Endocrinology Research Center, Russian Academy of Medical Science, National IDD Center, Ministry of Health. *IDD survey in the city of Electrostal, Moscow Oblast*. Moscow, Endocrinology Research Center, Russian Academy of Medical Science, 2001. Ref 1143

Endocrinology Research Center, Russian Academy of Medical Sciences. *Evaluation of iodine status and goitre prevalence in children population of Kirov Oblast*. Moscow, Endocrinology Research Center, Russian Academy of Medical Sciences, 2001. Ref 1139

Endocrinology Research Center, Ministry of Health. *Urinary iodine and thyroid volume in Astrachan, Volgogradsky and Krasnodarsky Regions [unpublished report]*. Moscow, Endocrinology Research Center, Ministry of Health, 2002. Ref 3610

Rwanda

Ministère de la Santé Rwanda, UNICEF, WHO. *National nutrition survey of women and children in Rwanda in 1996: final report*. Kigali, Ministère de la Santé, 1997. Ref 2558

Saudi Arabia

Al-Nuim A et al. *National iodine deficiency disorders survey – Saudi Arabia, 1994–1995*. Riyadh, Ministry of Health, 1995. Ref 490

Abu-Eshy SA, Abolfotouh MA, Al-Naggar YM. Endemic goitre in school children in high and low altitude areas of Asir region, Saudi Arabia. *Saudi Medical Journal*, 2001, 22: 146–149. Ref 567

Senegal

SANAS, ORSTOM Nutrition. *Enquête sur la prévalence des troubles dus à la carence en iode dans la région de Tambacounda, novembre 1995–avril 1996 [Survey on the prevalence of iodine deficiency disorders in the Tambacounda region, November 1995–April 1996]*. Dakar, 1997. Ref 1633

Dillon JC, Milliez J. Reproductive failure in women living in iodine deficient areas of West Africa. *British Journal of Obstetrics and Gynaecology*, 2000, 107:631–636. Ref 1562

Serbia and Montenegro

Simic M et al. *Republic of Serbia is free of iodine deficiency: results of comprehensive survey of school children* [Unpublished report]. Belgrade, Institute of Public Health of Serbia, 2001. Ref 3062

Institute of Public Health of Serbia et al. *The health and nutrition of the refugee population in the Federal Republic of Yugoslavia*. Rome, National Institute of Nutrition Italy, 1998. Ref 763

Slovakia

Kostalova L. *Urinary iodine assessment in children in Slovakia* [Unpublished report]. Bratislava, Children's University Hospital, 2003. Ref 558

Podoba J Jr et al. Účinnost jódovej profylaxie endemickej strumy na Slovensku z hľadiska fyzikálneho vyšetrenia štítnej žľazy [The effectiveness of iodine prophylaxis of endemic goiter in Slovakia from the viewpoint of physical and ultrasonographic examinations of the thyroid gland]. *Bratislavske Lekarske Listy*, 1995, 96:622–626. Ref 1099

Slovenia

Zaletel K, Hojker S. *IDD control in Slovenia. Iodine nutrition in Slovenia* [unpublished data]. Ljubljana, University Medical Center Ljubljana, 2002. Ref 3391

South Africa

Department of Health and South African Institute for Medical Research. *Report of the South African Institute for Medical Research (SAIMR) on the IDD Survey of Primary School learners for the DOH, South Africa*. Johannesburg, South African Institute for Medical Research, 2000. Ref 2618

Spain

Garcia-Mayor RV et al. Effect of iodine supplementation on a pediatric population with mild iodine deficiency. *Thyroid*, 1999, 9:1089–1093. Ref 2091

Serna Arnaiz MC et al. The prevalence of antithyroid antibodies in Lleida. *Anales de Medicina Interna*, 2000, 17: 62–66. Ref 3607

Delgado Álvarez E et al. Nutrición de yodo en los escolares Asturianos tras 18 años de yodoprofilaxis con sal. Erradicación del bocio endémico? [Iodine nutrition in Asturian schoolchildren after 18 years of iodized salt prophylaxis. Eradication of endemic goiter?] *Endocrinología Y Nutrición*, 2001, 48:4. Ref 3581

Maudeno Car AJ et al. Prevalencia de bocio y deficiencia de yodo en población escolar de una zona básica de salud tradicionalmente endémica [Prevalence of goitre and iodine deficiency in a school population from a traditionally endemic health area]. *Atencion Primaria*, 2001, 27: 258–262. Ref 3606

Santiago P et al. Prevalencia de déficit de yodo en la provincia de Jaén [Prevalence of iodine deficiency in Jaén province]. *Endocrinología Y Nutrición*, 2001, 49:77–78. Ref 3580

Sri Lanka

Medical Research Institute, UNICEF. *Iodine deficiency status of children in Sri Lanka, 2000–2001*. Colombo, Medical Research Institute, 2001. Ref 404

Sudan

IDD baseline survey report. Khartoum, Ministry of Health, 1999. Ref 2937

Swaziland

Todd CH. *Report on a consultancy to review Swaziland's iodine deficiency disorders control programme*. Mbabane, National Nutrition Council, 1998. Ref 2589

Switzerland

Hess SY et al. Monitoring the adequacy of salt iodization in Switzerland: a national study of school children and pregnant women. *European Journal of Clinical Nutrition*, 2001, 55:162–166. Ref 2662

Fleury Y et al. Iodine nutrition and prevalence of goiter in adolescents in the Canton of Vaud. *Swiss Medical Weekly*, 1999, 129:1831–1838. Ref 1207

Hoang Truong T et al. Iodine supply at various periods in life and ultrasonographic thyroid volume in school children in a region of Switzerland. *Swiss Medical Weekly*, 1997, 127: 715–721. Ref 1212

Tajikistan

Kasymova S. Prevalence of iodine deficiency disorders in Tajikistan. *WHO CAR News*, 2000, 6(23):5. Ref 3705

Thailand

Surveillance system for 'Tracking progress towards the sustainable elimination of IDD in Thailand': survey 2000 presented in table form. Bangkok, Ministry of Public Health, 2000. Ref 3554

IDD prevalence rate of school children in Thailand: annual surveys 1992–2000 presented in table form. Bangkok, Ministry of Public Health, 2001. Ref 2555

The former Yugoslav Republic of Macedonia

UNICEF. *Report on the activities of the National Committee for iodine deficiency in the year 2002*. Skopje, United Nations Children's Fund, 2002. Ref 3609

Togo

Kibambe N et al. *Projet Thyromobil en Afrique de l'Ouest (première phase). Rapport final. [Thyromobil project in West Africa (first phase). Final report.]* Lomé, Conseil International de Lutte contre les Troubles Dus à la Carence en Iode, 2000. Ref 2535

Ministère de la Santé Togo, WHO. *Troubles neurologiques dans une région d'endemie goitreuse – enquête exhaustive dans une population rurale de 4182 habitants [Neurologic disorders in an endemic goitre region – exhaustive survey in a rural population of 4182 inhabitants]*. Lomé, Ministère de la Santé, 1995. Ref 387

Tunisia

Ministère de la Santé Publique Tunisie. *Rapport National: évaluation de l'état nutritionnel de la population tunisienne [National Report: evaluation of the nutritional status of the Tunisian population]*. Tunis, Ministère de la Santé Publique, 1996. Ref 2485

Hsairi M et al. Prévalence du goitre endémique dans une région du nord ouest de la Tunisie, 1993 [Prevalence of endemic goiter in the north western region of Tunisia, 1993]. *Tunisie Médicale*, 1994, 72:663–669. Ref 1559

Turkey

Erdogan G et al. Iodine status and goiter prevalence in Turkey before mandatory iodization. *Journal of Endocrinological Investigation*, 2002, 25:224–228. Ref 3426

Hacettepe Universitesi. 15 il'de beslenme eğitimi ve araştırması projesi, 1995 [Training and research project on nutrition in 15 provinces, 1995]. Turkey, 1995. Ref 3428

Turkmenistan

Akmuradova G. Prevention of iodine deficiency disorders in Turkmenistan [in Russian]. *WHO CAR News*, 2000, 6(23):6–7. Ref 3620

Uganda

Olico-Okui. *Monitoring progress towards control of IDD through universal salt iodization in Uganda: study report*. Kampala, Makerere University, 2000. Ref 2582

Ukraine

Ashizawa K et al. Prevalence of goiter and urinary iodine excretion levels in children around Chernobyl. *Journal of Clinical Endocrinology and Metabolism*, 1997, 82:3430–3433. Ref 1238

Pankiv V. Epidemiological survey of goiter and iodine deficiency in the Ukrainian Carpathians. *IDD Newsletter*, 2000, 16:56. Ref 3600

United Arab Emirates

Pandav CS. *Survey of prevalence of iodine deficiency disorders in the United Arab Emirates – 7 February through 15 March 1994*. Cairo, World Health Organization, 1994. Ref 483

United Republic of Tanzania

The Partnership for Child Development. The health and nutritional status of schoolchildren in Africa: evidence from school-based health programmes in Ghana and Tanzania. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 1998, 92:254–261. Ref 1772

WHO, UNICEF, ICCIDD. *Joint WHO/UNICEF/ICCIDD Consultation. Review of findings from 7-country study in Africa on levels of salt iodization in relation to iodine deficiency disorders, including iodine induced hyperthyroidism*. Geneva, World Health Organization, 1997. (Available on request from Department of Nutrition for Health and Development, World Health Organization, 1211 Geneva 27, Switzerland). Ref 3601

United States of America

Hollowell JG et al. Iodine nutrition in the United States. Trends and public health implications: iodine excretion data from National Health and Nutrition Examination Surveys I and III (1971–1974 and 1988–1994). *Journal of Clinical Endocrinology and Metabolism*, 1998, 83:3401–3408. Ref 1523

Uzbekistan

Ismailov SI. *Iodine deficiency disorders in Uzbekistan*. Tashkent, Ministry of Health, 1998. Ref 570

Venezuela

Situación actual de los DDY en Venezuela [Current situation of IDD in Venezuela]. Caracas, Republica Bolivariana de Venezuela, 2002. Ref 3168

Viet Nam

Ministry of Health of Cambodia, National Maternal and Child Health Center. *Workshop on regional strategy for control of IDD*. Cambodia, The National Sub Committee for control of IDD (NSCIDD), 1997. Ref 1076

Yemen

Zein A et al. The epidemiology of iodine deficiency disorders (IDD) in Yemen. *Public Health Nutrition*, 2000, 3: 245–252. Ref 1561

Zambia

Lumbwe CM et al. *Iodine deficiency disorders in Zambia – 1993 survey draft report*. Lusaka, National Food and Nutrition Commission, 1995. Ref 394

Zimbabwe

Zimbabwe National Micronutrient Survey: 1999. Harare, Ministry of Health and Child Welfare, 2001. Ref 2641