Scope Dominance with Upward Monotone Quantifiers

Alon Altman Ya'acov Peterzil Yoad Winter

Abstract

We give a complete characterization of the class of upward monotone generalized quantifiers Q_1 and Q_2 over countable domains that satisfy the scheme $Q_1x\ Q_2y\ \phi \rightarrow Q_2y\ Q_1x\ \phi$. This generalizes the characterization of such quantifiers over *finite* domains, according to which the scheme holds iff either Q_1 is $\exists$ or Q_2 is $\forall$ (excluding trivial cases). Our result shows that in infinite domains, there are more general types of quantifiers that support these entailments.

1 Introduction

A *generalized quantifier* over a domain E is a set $Q \subseteq \wp(E)$. For instance, over a domain E and some $X \subseteq E$, the following are the generalized quantifiers that are more traditionally written as $\exists x \in X$ and $\forall x \in X$, respectively:

$$\mathsf{some}(X) \stackrel{def}{=} \{A \subseteq E : X \cap A \neq \emptyset\}.$$

$$\mathsf{every}(X) \stackrel{def}{=} \{A \subseteq E : X \subseteq A\}.$$

We call such quantifiers *EXIST* ("existential") and *UNIV*, respectively.[1] The quantifiers Q that are both *EXIST* and *UNIV* are of the form $\{A \subseteq E : x \in A\}$ for some $x \in E$, which are precisely the principal ultrafilters over E.

When Q_1 and Q_2 are generalized quantifiers and R a binary relation, the formula $Q_1x\ \ Q_2y\ \ R(x,y)$ is often written Q_1Q_2R, which is interpreted in E as follows.

(1) $\{x \in E : R_x \in Q_2\} \in Q_1$,

[1] We do not simply say that $\mathsf{some}(X)$ is *existential* to avoid confusion with the larger class of quantifiers that Keenan and Westerståhl (1996) call *intersective*, and which are often referred to as *existential*.

where $R_x = \{y \in E : R(x,y)\}$. Henceforth we will also use the notation R^y for $\{x \in E : R(x,y)\}$, considering the following equivalence:

(2) $Q_2Q_1R^{-1} \Leftrightarrow \{y \in E : R^y \in Q_1\} \in Q_2$.

Previous studies of generalized quantifiers have characterized various *scope commutativity* properties of quantifiers in constructions with multiple quantification. Notably, Westerståhl (1996) characterizes the class of *self-commuting* quantifiers – those quantifiers Q that satisfy the following equivalence:

(3) For all $R \subseteq E^2$: $QQR \Leftrightarrow QQR^{-1}$.

Zimmermann (1993) characterizes the class of *scopeless* quantifiers – those quantifiers Q that satisfy the following equivalence.

(4) For all $Q_1 \subseteq \wp(E)$, for all $R \subseteq E^2$: $QQ_1R \Leftrightarrow Q_1QR^{-1}$.

He shows that the scopeless quantifiers over E are precisely the ultrafilters over E that are closed under intersections of length $\leq |E|$.

Westerståhl (1986) studies the more general problem of characterizing the generalized quantifiers Q_1, Q_2 that satisfy the following unidirectional entailment.

(5) For all $R \subseteq E^2$: $Q_1Q_2R \Rightarrow Q_2Q_1R^{-1}$.

When this entailment holds, we say that Q_1 is (scopally) *dominant* over Q_2.

For the sake of completeness we give in section 2 a simple proof of Westerståhl's characterization of domination between quantifiers in *finite* domains. The main part of the paper is section 3, where this characterization is extended to *countable* domains. Section 4 concludes with some remarks about scope commutativity, finiteness and monotonicity in natural language semantics.

2 Finite domains

Westerståhl's characterization is restricted to *upward monotone* quantifiers over *finite domains*. Standardly, by saying that a quantifier Q over E is upward monotone we mean that Q is closed under supersets: $A \in Q$ and $A \subseteq B$ implies $B \in Q$. Under upward monotonicity and finiteness of the domain, Westerståhl's claim can be stated as follows.[2]

Fact 1 *Let Q_1 and Q_2 be upward monotone quantifiers over a finite domain E. Q_1 is dominant over Q_2 iff these quantifiers fall under at least one of the following cases.*

[2]Westerståhl characterizes scope entailments for *determiners* – functions from sets to generalized quantifiers. The following is a simpler statement of the result for generalized quantifiers.

(i) Q_1 *is EXIST.*

(ii) Q_2 *is UNIV.*

(iii) $Q_1 = \wp(E)$ *and* $Q_2 \neq \emptyset$.

(iv) $Q_2 = \emptyset$ *and* $Q_1 \neq \wp(E)$.

Proof The 'if" direction of the proof is easy. For the "only if" direction, assume that Q_1 is dominant over Q_2. First it is easy to see that if $Q_1 = \wp(E)$ then $Q_2 \neq \emptyset$ and that if $Q_2 = \emptyset$ then $Q_1 \neq \wp(E)$. Assume for contradiction that no one of the conditions *(i)-(iv)* holds. Then by finiteness of E there is a minimal set $A \in Q_1$ such that $|A| \geq 2$ (otherwise by upward monotonicity, $Q_1 = \wp(E)$ or $Q_1 = \mathsf{some}(\bigcup_{\{x\} \in Q_1} \{x\})$). Similarly, by finiteness of E there are $B_1, B_2 \in Q_2$ such that $B_1 \cap B_2 \notin Q_2$ (otherwise by upward monotonicity, $Q_2 = \emptyset$ or $Q_2 = \mathsf{every}(\cap Q_2)$). Given the sets A, B_1 and B_2, and an arbitrary $a \in A$, it is easy to verify that the relation $(\{a\} \times B_1) \cup ((A \setminus \{a\}) \times B_2)$ contradicts our assumption that Q_1 is dominant over Q_2. □

Westerståhl (1996) calls two quantifiers $Q_1, Q_2 \subseteq \wp(E)$ *independent* if they satisfy the following equivalence.

(6) For all $R \subseteq E^2$: $Q_1 Q_2 R \Leftrightarrow Q_2 Q_1 R^{-1}$.

Using Fact 1 it is easy to establish the following corollary.

Corollary 2 *Let* Q_1 *and* Q_2 *be upward monotone quantifiers over a finite domain* E. *Then* Q_1 *and* Q_2 *are independent iff* Q_1 *and* Q_2 *fall under at least one of the following cases.*

(i) Both Q_1 *and* Q_2 *are EXIST.*

(ii) Both Q_1 *and* Q_2 *are UNIV.*

(iii) Either Q_1 *or* Q_2 *(or both) are principal ultrafilters.*

(iv) Either Q_1 *or* Q_2 *are trivial, but not both.*

By saying that a quantifier Q is *trivial* we mean that $Q = \wp(E)$ or $Q = \emptyset$.

Examples For illustrating scope domination in simple natural language sentences, consider a well-known type of example.

(7) Some priest visited every city.

Let us assume that the nouns *priest* and *city* are denoted by the sets $P, C \subseteq E$ respectively, and that the verb *visited* is denoted by the binary relation $V \subseteq E^2$. Sentence (7) has two readings, depending on the order in which the quantifiers operate on the arguments of the relation V:

(8) a. $\mathsf{some}(P)\ \mathsf{every}(C)\ V$

b. $\mathsf{every}(C)\ \mathsf{some}(P)\ V^{-1}$

The statement in (8a) is called the *object narrow scope* (ONS) reading of sentence (7), whereas the the statement in (8b) is called the *object wide scope* (OWS) reading of the sentence. As a matter of first-order logic, (8a) entails (8b) but not vice versa. Thus, the quantifier $\mathsf{some}(P)$ is dominant over the quantifier $\mathsf{every}(C)$ for any $P, C \subseteq E$, but the opposite does not hold.

The situation is similar in cases where (exactly) one of the existential/universal quantifiers is replaced by another upward monotone quantifier, not necessarily first-order. The sentences in (9) below illustrate some cases like that, where the ONS reading entails the OWS reading. The corresponding quantifiers we assume are given in (10).

(9) a. At least half/at least two/all but at most five of the priests visited every city.

b. Some priest visited at least half/at least two/all but at most five of the cities.

$$
\begin{array}{lcl}
\mathsf{at\_least\_half\_of\_the}(X) & = & \{A \subseteq E : |X \cap A| \geq |X \setminus A|\} \\
\mathsf{at\_least\_n}(X) & = & \{A \subseteq E : |X \cap A| \geq n\} \\
\mathsf{all\_but\_at\_most\_n\_of\_the}(X) & = & \{A \subseteq E : |A \setminus X| \leq n\}
\end{array} \tag{10}
$$

Note that the generalized quantifier $\mathsf{at\_least\_half\_of\_the}(X)$ is not first-order definable.

Westerståhl's result shows that over finite domains, the *EXIST* quantifiers (for Q_1) and the *UNIV* quantifiers (for Q_2) are the only non-trivial upward monotone quantifiers that lead to entailments as in (5). Thus, the sentences in (7) and (9) are representative of the cases where upward monotone quantifiers lead to an entailment from the ONS reading to the OWS reading on finite domains.

3 Countable domains

As Westerståhl observes, his characterization of scope dominance over finite domains in Fact 1 does not hold for infinite domains. Thus, over infinite domains there are non-trivial upward monotonic quantifiers besides the *EXIST* and *UNIV* quantifiers that give rise to scope entailments. Consider the following example (following Westerståhl), where E is assumed to be countable.

(11) Infinitely many dots are contained in at least one of the three circles.

$$Q_1 = \{A \subseteq E : |D \cap A| = \aleph_0\}$$
$$Q_2 = \{A \subseteq E : C \cap A \neq \emptyset\}, \text{ where } |C| = 3$$

It is easy to verify that Q_1 is dominant over Q_2, but Q_1 and Q_2 are upward monotone and the conditions in Fact 1 do not hold. Incidentally, since Q_2 is *EXIST*, it is dominant over Q_1.

In this section we characterize scope dominance in the class of upward monotone quantifiers over countable domains. First, we observe the close relation between scope dominance and *duality* of GQs. The *dual* Q^d (cf. Barwise and Cooper (1981)) of a generalized quantifier Q over E is defined by:

$$Q^d \stackrel{def}{=} \{A \subseteq E : E \setminus A \notin Q\}.$$

Any *EXIST* quantifier is the dual of a *UNIV* quantifier and vice versa (note that $(Q^d)^d = Q$ for any quantifier Q). It is also easy to observe the following fact.

Fact 3 *Q_1 is dominant over Q_2 iff Q_2^d is dominant over Q_1^d.*

This fact follows directly from the definition of scope dominance and duality, and the observation that for any $R \subseteq E^2$ we have:

$$Q_1\ Q_2\ R \Leftrightarrow \neg(Q_1^d\ Q_2^d\ (E^2 \setminus R)).$$

Let us define some properties of quantifiers that will be useful for characterizing scope dominance. First, we say that a quantifier Q satisfies the *union property* (U) if for all $A_1, A_2 \subseteq E$: if $A_1 \cup A_2 \in Q$ then $A_1 \in Q$ or $A_2 \in Q$. For example, any *EXIST* quantifier satisfies (U), while a *UNIV* quantifier $\mathsf{every}(X)$ satisfies (U) if and only if X is either a singleton or the empty set. The set of all infinite subsets of E satisfies (U) as well.

Further, we say that a quantifier Q satisfies the *Descending Chain Condition* (DCC) if for every descending sequence $A_1 \supseteq A_2 \supseteq \cdots A_n \supseteq \cdots$ in Q, the intersection $\cap A_i$ is in Q as well. For example, any *UNIV* quantifier satisfies (DCC). An *EXIST* quantifier $\mathsf{some}(X)$ satisfies (DCC) if and only if X is finite. Another quantifier that satisfies (DCC) is the following, where the domain $E = \mathbb{N}$ is the set of natural numbers:

$\{A \subseteq \mathbb{N} : \forall n \in \mathbb{N}\,[2n \in A \vee 2n+1 \in A]\}$.

If every set in a quantifier Q contains a finite subset that is also in Q, we say that Q satisfies (FIN). The following fact uses (FIN) in characterizing dual properties to (U) and (DCC) over the class of upward monotone quantifiers.

Fact 4 *For any upward monotone generalized quantifier Q over a domain E:*

(i) Q satisfies (U) *iff Q^d is a filter.*

(ii) If E is countable: Q satisfies (DCC) *iff Q^d satisfies* (FIN).

Proof The proof of (i) is easy.
For (ii), assume that Q is (DCC) and assume for contradiction that there is $A \in Q^d$ such that for all $B \subseteq A$: if $B \in Q^d$ then B is infinite. Let $B_0 \subset A$ be a finite set. Hence $B_0 \notin Q^d$, and $E \setminus B_0 \in Q$. By countability of E, we can denote $A \setminus B_0 = \{a_i\}_{i=1}^{\infty}$. Let $B_{i+1} = B_i \cup \{a_{i+1}\}$, for any $i \geq 0$. By our assumption on A we have, $B_i \notin Q^d$ for any $i \geq 0$, hence $E \setminus B_i \in Q$ for any $i \geq 0$. But $\bigcap_i (E \setminus B_i) = E \setminus A \notin Q$, in contradiction to Q being (DCC).
Conversely, assume that Q satisfies (FIN). Let $B_1 \supseteq B_2 \supseteq \ldots$ be a descending chain in Q^d, thus $E \setminus B_i \notin Q$ for any $i \geq 1$. Assume for contradiction $B = \bigcap_i B_i \notin Q^d$, thus $E \setminus B = \bigcup_i (E \setminus B_i) \in Q$. By assumption on Q, there exists $A' \subseteq \bigcup_i (E \setminus B_i)$ such that $A' \in Q$ and A' is finite. Hence, there is n such that $A' \subseteq E \setminus B_n$, and from upward monotonicity of Q, it follows that $E \setminus B_n \in Q$, in contradiction to the assumption. □

Using these two pairs of dual properties, our main Theorem can be stated as follows.

Theorem 5 *Let Q_1 and Q_2 be upward monotone quantifiers over a countable domain E. Then the following are equivalent:*

I. Q_1 is dominant over Q_2.

II. Q_1 and Q_2 fall under at least one of the following cases.

(i) Q_1 is EXIST.
(ii) Q_2 is UNIV.
(iii) Q_1 satisfies (U), *$Q_2 \neq \emptyset$ and Q_2 satisfies* (DCC).
(iv) Q_2 is a filter, $Q_1 \neq \wp(E)$ and Q_1 satisfies (FIN).
(v) $Q_1 = \wp(E)$ and $Q_2 \neq \emptyset$.
(vi) $Q_2 = \emptyset$ and $Q_1 \neq \wp(E)$.

Note that, in contradistinction to Fact 1, clauses (iii) and (iv) in this theorem imply that in countable domains *EXIST* and *UNIV* quantifiers are not the only non-trivial upward monotone quantifiers that give rise to scope entailments. Each of these two clauses contains a different generalization of *EXIST* and *UNIV* quantifiers. For (iii), over finite domains the upward monotone quantifiers that satisfy (U) are exactly the *EXIST* quantifiers and $\wp(E)$, and the (DCC) requirement for Q_2 is trivially satisfied. For (iv), over finite domains the filters are the *UNIV* quantifiers and $\emptyset$, and the "finiteness" requirement for Q_2 is trivially satisfied. Note also that, in any domain, the *UNIV* quantifiers are (DCC) and the *EXIST* quantifiers satisfy the "finiteness" requirement. Thus, Fact 1 is a (simple) special case of Theorem 5.

We note that any filter over countable domain that satisfies (DCC) is *UNIV*. From this and from Fact 4 it follows that when Q_1 and Q_2 are upward monotone quantifiers over a countable domain, statement (II) is equivalent to statement (II') below.

(II') *All* of the following requirements hold:

(i') Q_1^d or Q_2 (or both) are filters;

(ii') Q_1^d or Q_2 (or both) satisfy (DCC);

(iii') Q_1^d or Q_2 (or both) are not empty.

In the proof of Theorem 5, we use both formulations (II) and (II').

Proof of Theorem 5

(II) $\Rightarrow$ (I):

We prove only the entailment from clause (iv) to (I). The entailment from clause (iii) to (I) then follows directly from Facts 3 and 4. The entailments from the other clauses in (II) to (I) are easy to prove.

Assume that (iv) holds, hence every set in Q_1 contains a finite set in Q_1, $Q_1 \neq \wp(E)$ and Q_2 is closed under finite intersections. Assume that Q_1Q_2R holds, and denote $A = \{x \in E : R_x \in Q_2\} \in Q_1$. Let $F \in Q_1$ be a minimal finite subset of A in Q_1. By monotonicity of Q_1 and $Q_1 \neq \wp(E)$ it follows that $F \neq \emptyset$. Let $B = \bigcap_{x \in F} R_x$. Since Q_2 is closed under finite intersections, B itself is in Q_2. For every $y \in B$, $R^y \supseteq F$ and therefore $R^y \in Q_1$. It follows that the set $\{y : R^y \in Q_1\}$ contains B and hence is in Q_2. We thus showed that $Q_2Q_1R^{-1}$ holds, as required.

(I) $\Rightarrow$ (II'):

We assume that for every $R \subseteq E^2$: $Q_1Q_2R \Rightarrow Q_2Q_1R^{-1}$.

To show that (i') holds, assume that Q_1^d is not a filter, so from Fact 4 it follows

that Q_1 does not satisfy (U). Hence, by upward monotonicity of Q_1, it contains a set $A = A_1 \cup A_2$ where A_1 and A_2 are disjoint and neither of them is in Q_1. To show that Q_2 is closed under finite intersections, hence a filter, let us denote for any $B_1, B_2 \in Q_2$:

$$R = (A_1 \times B_1) \cup (A_2 \times B_2).$$

We have $\{x : R_x \in Q_2\} \in Q_1$ and therefore $B \stackrel{def}{=} \{y : R^y \in Q_1\} \in Q_2$. But, since $A_1, A_2 \notin Q_1$, we have $B = B_1 \cap B_2$, hence $B_1 \cap B_2 \in Q_2$.

To show that (ii') holds, assume that Q_1^d does not satisfy (DCC), so from Fact 4 it follows that there is a set $A \in Q_1$ such that every subset of A that is also in Q_1 is infinite. To show that Q_2 satisfies (DCC), let $B_1 \supseteq B_2 \ldots \supseteq B_n \supseteq \ldots$ be a sequence of sets in Q_2, and let B be their intersection. We again assume that E is the set of natural numbers, and enumerate $A = \{a_1, a_2, \ldots, \}$. Consider the relation

$$R = \bigcup_{n=1}^{\infty} (\{a_n\} \times B_n).$$

Then the set $\{x : R_x \in Q_2\} = A \in Q_1$, or $Q_1 Q_2 R$, and by our assumption it follows that $Q_2 Q_1 R^{-1}$ holds, or $\{y : R^y \in Q_1\} \in Q_2$. We claim that this last set equals B, so Q_2 satisfies (DCC). Clearly, for every $y \in B$: $R^y = A \in Q_1$. However, if $y \notin B$ then there is n such that $y \notin B_m$ for all $m \geq n$, and therefore $R^y \subseteq A$ is finite. By our previous observation, such R^y is not in Q_1.

Clause (iii') trivially holds. □

As in the case of scope dominance over finite domains, the following corollary of Theorem 5 allows us to characterize the pairs of *independent* quantifiers.

Corollary 6 *Let Q_1 and Q_2 be upward monotone quantifiers over a countable domain E. Then Q_1 and Q_2 are independent iff Q_1 and Q_2 fall under at least one of the following cases, with Q_1 and Q_2 interchangeable.*

- *(i) Q_1 is EXIST and non-trivial, hence equal to $\mathsf{some}(X)$ for some $X \neq \emptyset$. Either X is finite and Q_2 satisfies* (U)*, or X is infinite and Q_2 is EXIST.*
- *(ii) Q_1 is UNIV and non-trivial, hence equal to $\mathsf{every}(X)$ for some $X \neq \emptyset$. Either X is finite and Q_2 is a filter, or X is infinite and Q_2 is UNIV.*
- *(iii) Q_1 or Q_2 are principal ultrafilters.*
- *(iv) For some finite collection $\mathcal{X} \subseteq \wp(E)$ of finite sets, $Q_1 = \bigcup_{X \in \mathcal{X}} \mathsf{every}(X)$, and Q_2 is an ultrafilter.*

(v) Either Q_1 or Q_2 are trivial, but not both.

Remark: Since we assume here the Axiom of Choice, non-principal ultrafilters exist over E, so (iv) is not subsumed by (iii).

Proof The "if" direction easily follows from Theorem 5. For the "only if" direction, we use the following two claims, where Q is an upward monotone quantifier over a countable domain E.

Claim 1. Q satisfies both (U) and (DCC) iff $Q = \wp(E)$ or $Q = \mathsf{some}(X)$ for some finite $X \subseteq E$.

Proof of claim 1. The proof of the "if" direction is easy. For the "only if" dirrection, assume that Q satisfies (U) and $Q \neq \wp(E)$. Then by monotonicity there are no minimal sets in Q other than singletons. Let A be some arbitrary set in Q. If A is finite then it must contain a singleton in Q. If A is infinite, then either it contains a singleton in Q, or by (U) and the countability of E, we can form a descending chain of subsets of A, all in Q, whose intersection is empty. From (DCC) it follows that $\emptyset \in Q$ and by monotonicity $Q = \wp(E)$, in contradiction to our assumption. Thus, every set in Q contains a singleton in Q, and if $X = \{x \in E : \{x\} \in Q\}$ then by monotonicity $Q = \mathsf{some}(X)$. Suppose for contradiction that X is infinite, then again by (DCC), we conclude that $\emptyset \in Q$, contradiction.

Claim 2. If Q satisfies (DCC) and (FIN) then there are finitely many finite minimal sets in Q.

Proof of claim 2. Assume for contradiction that there are infinitely many finite minimal sets in Q, and denote this collection of sets by $\mathcal{X}$. It follows that for any $A, B \in \mathcal{X}$ such that $A \neq B$, $A \cap B$ is a proper subset of both A and B. Let F_1 be in $\mathcal{X}$. Because F_1 is finite and $\mathcal{X}$ is infinite, there must be some $F_1' \subsetneq F_1$ such that the collection of sets $\mathcal{X}_1 = \{F \in \mathcal{X} : F \cap F_1 = F_1'\}$ is infinite. We can continue this process by defining F_i, F_i' and $\mathcal{X}_i$ for every $i \geq 1$ as follows:

> F_{i+1}' is some proper subset of F_i such that $\{F \in \mathcal{X}_i : F \cap F_{i+1} = F_{i+1}'\}$ is infinite.
>
> $\mathcal{X}_{i+1} \stackrel{def}{=} \{F \in \mathcal{X}_i : F \cap F_i = F_i'\}$
>
> F_{i+1} is some set in $\mathcal{X}_i$.

We obtain an infinite sequence $F_1, \ldots, F_i, \ldots$ of finite minimal sets in Q, together with $F_1', \ldots, F_i', \ldots$, such that for every i, $F_i' \subsetneq F_i$ and for every $m > n$, $F_m \cap F_n = F_n'$.

We now let $A_n = \bigcup_{m \geq n} F_m$. This is a decreasing sequence of sets, all in Q, so by (DCC), $A = \bigcap_{n=1}^{\infty} A_n$ is in Q. We claim that $A = \bigcup_{n=1}^{\infty} F_n'$. Indeed, note

that A consists of all elements which belong to infinitely many sets F_n. Let x be some element in $\bigcup_{n=1}^{\infty} F'_n$, thus $x \in F'_n$ for some n. For every $m > n$, $x \in F_m$ because $F_m \in \mathcal{X}_n$. Thus x is in infinitely many sets F_m, and therefore $x \in A$. For the opposite direction, assume that $x \in A$, thus belongs to infinitely many F_n. In particular it belongs to some F_m, F_n for $m > n$. But then $x \in (F_m \cap F_n) = F'_n$, thus belonging to $\bigcup_{n=1}^{\infty} F'_n$.

By our assumption on Q, the set A contains a finite subset $B \in Q$. The set B is then contained in the union of finitely many sets F'_n, which implies that for some m (larger than all these n's): $B \subsetneq F_m$. Because both F_m and B are in Q, this contradicts the minimality of F_m.

We now assume that Q_1 and Q_2 are independent and prove that one of the conditions (i) to (v) holds. If one of the quantifiers Q_1 and Q_2 is trivial, it easily follows that the other one is not trivial, so (v) holds. We thus assume that both Q_1 and Q_2 are not trivial.

Assume that Q_1 is *EXIST*. By Theorem 5, Q_1 is dominant over Q_2, and for Q_2 to be dominant over Q_1, one of the following must hold:

- Q_2 too is *EXIST*, hence (i) holds;
- Q_1 is *UNIV*, hence it is a principal ultrafilter and (iii) holds;
- Q_2 satisfies (U) and Q_1 satisfies (DCC) hence Q_1 must be $\mathsf{some}(X)$ for some finite X, and thus condition (i) holds;
- Q_1 is a filter, hence it is a principal ultrafilter and (iii) holds.

Symmetrically, if Q_2 is *EXIST* then (i) or (iii) holds. Similar considerations show that if Q_1 (or Q_2) is *UNIV*, then (ii) or (iii) must hold.

Assume now that both Q_1 and Q_2 are neither *EXIST* nor *UNIV*. Consider the two possibilities according to Theorem 5 that Q_1 is dominant over Q_2:

- Q_1 satisfies (U) and Q_2 satisfies (DCC). By claim 1 and the assumption that Q_1 is not *EXIST* it follows that Q_1 does not satisfy (DCC). Therefore, by Theorem 5 and dominance of Q_2 over Q_1, it follows that Q_1 is a filter, and every set in Q_2 contains a finite set in Q_2. Since Q_1 satisfies (U), it directly follows that it is furthermore an ultrafilter. By claim 2, the set $\mathcal{X}$ of minimal sets in Q_2 is finite, and all the sets it contains are finite too. Hence, $Q_2 = \bigcup_{X \in \mathcal{X}} \mathsf{every}(X)$, so (iv) holds.
- Q_1 satisfies (FIN) and Q_2 is a filter. If Q_1 is a filter then it follows that it contains only one minimal set, in contradiction to the assumption that it is not *UNIV*. Hence, by to Theorem 5 and dominance of Q_2 over Q_1, Q_2 satisfies (U) and Q_1 satisfies (DCC), so (iv) holds again.

□

Examples First let us note that in example (11) above, $Q_2 = \mathsf{some}(C)$ for a finite C ($|C| = 3$). Q_1 satisfies (U), hence Q_1 and Q_2 fall under clause (i) in Corollary 6, and the ONS reading of the sentence is equivalent to the OWS reading. The following example illustrates quantifiers Q_1 and Q_2 that fall under the dual clause (ii) in Corollary 6.

(12) Each of the three circles contains all but finitely many dots.

$$\begin{aligned} Q_1 &= \{A \subseteq E : C \subseteq A\}, \text{ where } |C| = 3 \\ Q_2 &= \{A \subseteq E : |D \setminus A| < \aleph_0\} \end{aligned}$$

To illustrate non-trivial usages of clause (iv) in Corollary 6, we would have to use non-principal ultrafilters, which we here omit.

As for dominance between quantifiers without independence, the quantifiers in (13) and (14) below satisfy clauses (iii) and (iv) of Theorem 5 respectively. Hence, in these cases the ONS reading entails the OWS reading, but not vise versa (assuming a finite $n > 0$).

(13) Infinitely many dots are contained in all but at most n circles.

(14) At least n circles contain all but finitely many dots.

Similarly, consider the following examples.

(15) Infinitely many dots are contained in circle 1 or [circles 2 and 3].

(16) Circle 1 and [circles 2 or 3] contain all but finitely many dots.

We assume that the object of sentence (15) and the subject of sentence (16) denote the following quantifiers respectively, for three different circles c_1, c_2 and c_3.

$$\{A \subseteq E : c_1 \in A \vee (c_2 \in A \wedge c_3 \in A)\}$$

$$\{A \subseteq E : c_1 \in A \wedge (c_2 \in A \vee c_3 \in A)\}$$

Also the quantifiers in these sentences satisfy respectively clauses (iii) and (iv) of Theorem 5, hence the scope dominance, but the two quantifiers in each sentence are not independent.

4 Concluding remarks

In this paper we have characterized scope dominance and independence for upward monotone generalized quantifiers over countable domains. This is a natural extension of the results by Westerståhl and Zimmermann about self-commuting and scopeless quantifiers. This characterization directly extends a previous result by Altman et al. (2001), which concentrated on a subclass of quantifiers on countable domains, called *finitely based* quantifiers. Our results are still partial in some obvious respects. First, we did not characterize scope dominance for uncountable domains. Theorem 5 does not hold for such domains, for the similar reasons that Fact 1 about finite domains does not hold for countable domains. Consider for instance the following sentence and quantifiers, parallel to (11) above over countable domains.

(17) Uncountably many dots are contained in at least one of the countably many circles.

$$Q_1 = \{A \subseteq E : |D \cap A| = \aleph_1\}$$
$$Q_2 = \{A \subseteq E : C \cap A \neq \emptyset\}, \text{ where } |C| = \aleph_0$$

The quantifier Q_1 is dominant over Q_2, but these quantifiers do not satisfy the conditions of Theorem 5. Thus, a further generalization of our result is called for.

It is also natural to look for a characterization of dominance with *non-upward monotone* quantifiers. One can also add further requirements on the relation R in (5), and obtain more quantifiers Q_1 and Q_2 that exhibit scope dominance for this restricted class of relations. Such more refined characterizations are relevant for natural language, where there are often logical restrictions on the possible denotations of binary relations. For instance, in the sentence *every priest is taller than some peasant*, where the relation *be taller than* is transitive, the ONS reading and the OWS reading are equivalent over finite domains, by contrast to the case with general R's.

Characterizations of scope independence are useful for reducing ambiguity in computational representations of natural language sentences. A system that goes in this direction, using the results that were obtained in the present paper, appears in Altman and Winter (2002).

Acknowledgments
The first and third authors were partly supported by grant no. 1999210 ("Extensions and Implementations of Natural Logic") from the United States-Israel Binational Science Foundation (BSF), Jerusalem, Israel.

References

Altman, A., Keenan, E., and Winter, Y. (2001). Monotonicity and relative scope relations. In *Proceedings of the 14th Amsterdam Colloquium*.

Altman, A. and Winter, Y. (2002). Computing dominant readings in scopally ambiguous sentences. Unpublished ms., Technion – Israel Institute of Technology. Downloadable at http://www.cs.technion.ac.il/~winter/.

Barwise, J. and Cooper, R. (1981). Generalized quantifiers and natural language. *Linguistics and Philosophy*, 4:159–219.

Keenan, E. and Westerståhl, D. (1996). Generalized quantifiers in linguistics and logic. In van Benthem, J. and ter Meulen, A., editors, *Handbook of Logic and Language*. Elsevier, Amsterdam.

Westerståhl, D. (1986). On the order between quantifiers. In Furberg, M. et al., editors, *Acta Universitatis Gothoburgensis*, pages 273–285.

Westerståhl, D. (1996). Self-commuting quantifiers. *The Journal of Symbolic Logic*, 61:212–224.

Zimmermann, T. E. (1993). Scopeless quantifiers and operators. *Journal of Philosophical Logic*, 22:545–561.