

jnm/ CONCISE COMMUNICATION**PULMONARY AND SYSTEMIC BLOOD PRESSURE RESPONSES****TO LARGE DOSES OF ALBUMIN MICROSPHERES**

Fred S. Mishkin and Richard E. Brashear

Indiana University School of Medicine, Indianapolis, Indiana

Toxicity of intravenously injected particles for lung scanning depends directly upon the degree of pulmonary vascular-bed occlusion (1). Although change in diffusing capacity has been used as an indication of such toxicity, pulmonary artery pressure may double without any change in diffusing capacity and hence provides a more sensitive index of toxicity (2,3). We assessed the effect of the newly introduced lung scanning agent, albumin microspheres (4), on pulmonary artery pressure and systemic blood pressure in dogs.

METHODS

Eight adult mongrel dogs weighing between 18 and 26 kg were anesthetized using 30 mg/kg sodium pentobarbital administered intravenously; the dogs were ventilated with a cuffed endotracheal tube using a constant-volume ventilator adjusted to maintain the end tidal CO₂ at 5.0–5.5 volumes % measured by an infrared CO₂ analyzer. Mean pulmonary artery and aortic pressures were measured on an Electronics for Medicine recorder using catheters fluoroscopically placed in the main pulmonary artery and aortic arch

and connected to strain gages. An electrocardiogram recorded heart rate. One gram of albumin microspheres (3M albumin microspheres) with a mean diameter of 22 microns (approximately 1.4×10^8 particles) was suspended by sonicating for 5 min in 247.5 ml of saline with 2.5 ml of 5% tween 80 added to make a suspension containing approximately 560,000 particles/ml. Before infusion the particles were suspended by manual shaking. Control pressures were recorded and each of five separate increments was infused over 30 sec. Each increment contained 7.15 mg/kg of microspheres in 32.2–46.5 ml, which was 500 times the usual human dose. Pressures were measured 1 min after infusing each increment.

RESULTS

The data in Table 1 subjected to Dunnett's t-test (5) shows that although there was a significant rise in pulmonary artery pressure after the infusion of

Received Nov. 9, 1970; revision accepted Jan. 14, 1971.

For reprints contact: Fred S. Mishkin, Indiana University Medical Center, 1100 W. Michigan St., Indianapolis, Ind. 46202.

TABLE 1. CORRELATION OF MEAN PULMONARY AND AORTIC PRESSURES WITH AMOUNT OF MICROSPHERES INFUSED

Dog weight (kg)	Control pressure		After increment I*		After increment II		After increment III		After increment IV		After increment V	
	Pulmonary	Aortic	Pulmonary	Aortic	Pulmonary	Aortic	Pulmonary	Aortic	Pulmonary	Aortic	Pulmonary	Aortic
18	15	124	16	134	17	141	19	150	21	128	14	81
19	13	131	13	127	15	122	18	95	18	70	15	54
19	14	110	14	133	14	82	12	42	25	51	31	48
20	13	115	14	119	14	118	18	140	20	105	11	47
26	8	117	2	34	2	29	3	33	6	44	8	49
19	11	111	14	129	15	134	16	122	17	64	15	61
18	10	131	11	136	12	136	13	137	14	126	15	90
25	11	111	12	121	14	137	15	118	14	92	13	60
Mean ± s.d.	12±2	119±9	12±4	117±34	13±5	117±39	14±5	105±45	17±6	85±3	15±7	62±16
Level of significance of change from control			None	None	None	None	None	None	0.05	0.05	None	0.05

* Increment contained 7.15 mg/kg body weight.

MISHKIN AND BRASHEAR

28.6 mg/kg or 8.2×10^7 of the microspheres, the pulmonary artery pressure in only a single instance reached twice the control value. In contrast to this, mean aortic pressure decreased significantly after the injection of 28.6 mg/kg of the microspheres, while there was no significant change in heart rate.

CONCLUSION

The dose required to elevate pulmonary artery pressure exceeds the usual clinical dose of 0.0143 mg/kg of body weight by a factor of 2,000, showing that there is a wide margin of safety between the clinically recommended dose and toxicity in dogs. The anomalous fall in aortic pressure without undue rise in pulmonary artery pressure suggests that there may be toxic effects of the particles other than mechanical blockade of pulmonary vessels since the pulmonary artery pressure usually increases by a factor of two or more preceding systemic hypotension when inert particles such as barium or glass beads are injected. These findings deserve further investigation and are reminiscent of histamine infusion into the pulmonary artery which causes pulmonary vasoconstriction and systemic hypotension (6).

ACKNOWLEDGMENT

Supported in part by research grants HE-06228, Program Project Grant HE-06308, and Postgraduate Research Training Grant 05363, National Heart Institute, United States Public Health Service.

REFERENCES

1. TAPLIN GV, JOHNSON DE, KENNADY JC, et al: Aggregated albumin labeled with various radioisotopes. In *Radioactive Pharmaceuticals*, Andrews GA, Kniseley RM, Wagner HN, USAEC Division of Technical Information, 1966, p. 525
2. BRASHEAR RE, MISHKIN FS, ROSS JC: Effects of iron (^{113}In) hydroxide particles on pulmonary diffusing capacity and pulmonary artery pressure. *Invest Radiol* 4: 19-23, 1969
3. DALY WJ, WALDHAUSEN JA: Physiologic studies of the pulmonary capillary bed after barium sulfate embolization. *J Clin Invest* 46: 1617-1624, 1967
4. RHODES BA, ZOLLE I, BUCHANAN JW, et al: Radioactive albumin microspheres for studies of the pulmonary circulation. *Radiology* 92: 1453-1460, 1969
5. WINER BJ: *Statistical Principles in Experimental Design*. New York: McGraw-Hill Book Co., 1962, p. 89
6. DELAUNOIS AL, LORDECKI R, POLET H, et al: Cardiac output, arterial blood pressure and pulmonary arterial pressure in histamine shock. *Arch Int Pharmacodyn* 120: 114-119, 1959

CALL FOR PAPERS

**3rd Annual Meeting, Northern and Southern California Chapters of
The Society of Nuclear Medicine**

**Town and Country Hotel
San Diego, California
September 17 and 18, 1971**

Members or sponsored guests of the Northern and Southern California Chapters of The Society of Nuclear Medicine are invited to submit abstracts of papers that they may wish to present at the above meeting. The abstracts should be 400 words or less in length and accompanied by no more than two pages of supporting data. The original and five copies of the abstract and accompanying data should be mailed to Dr. Samuel E. Halpern, Division of Nuclear Medicine, University Hospital, 225 W. Dickinson Street, San Diego, California 92103. The deadline for submission of abstracts is June 1, 1971.



The Journal of
NUCLEAR MEDICINE

Pulmonary and Systemic Blood Pressure Responses to Large Doses of Albumin Microspheres

Fred S. Mishkin and Richard E. Brashear

J Nucl Med. 1971;12:251-252.

This article and updated information are available at:
<http://jnm.snmjournals.org/content/12/5/251.citation>

Information about reproducing figures, tables, or other portions of this article can be found online at:
<http://jnm.snmjournals.org/site/misc/permission.xhtml>

Information about subscriptions to JNM can be found at:
<http://jnm.snmjournals.org/site/subscriptions/online.xhtml>

The Journal of Nuclear Medicine is published monthly.
SNMMI | Society of Nuclear Medicine and Molecular Imaging
1850 Samuel Morse Drive, Reston, VA 20190.
(Print ISSN: 0161-5505, Online ISSN: 2159-662X)

© Copyright 1971 SNMMI; all rights reserved.