

(*R*)-Histidinium (*2R,3R*)-tartrate**M. N. Johnson* and N. Feeder**Pfizer Ltd, IPC 049, Ramsgate Road, Sandwich,
Kent CT13 9NJ, EnglandCorrespondence e-mail:
matthew.johnson@sandwich.pfizer.com**Key indicators**Single-crystal X-ray study
 $T = 291$ K
Mean $\sigma(\text{C}-\text{C}) = 0.003$ Å
 R factor = 0.030
 wR factor = 0.076
Data-to-parameter ratio = 7.3For details of how these key indicators were
automatically derived from the article, see
<http://journals.iucr.org/e>.

The crystal structure of (*R*)-histidinium (*2R,3R*)-tartrate, $\text{C}_6\text{H}_{10}\text{N}_3\text{O}_2^+ \cdot \text{C}_4\text{H}_5\text{O}_6^-$, has been determined as part of an ongoing study into the fundamental effects of chirality on salt formation and hydration. Repeating layers of (*R*)-histidinium and (*2R,3R*)-tartrate interlink to form a three-dimensional network through simple translational symmetry of the unit cell.


Received 23 July 2004

Accepted 29 July 2004

Online 7 August 2004

Comment

This study was undertaken to identify the effects of chirality on the formation of salts, specifically the way chirality may affect hydration, as a result of interactions between a chiral drug and a chiral counterion. The absolute configuration can be considered determined as the chiral properties of the starting materials are well characterized. (*R*)-Histidine and (*2R,3R*)-tartaric acid samples were purchased from Fluka and used in the crystallization. The asymmetric unit and unit cell contain one histidine as a monocation (protonated at the amine and imidazole ring N atoms and deprotonated at the carboxylic acid group), and one tartrate as a monoanion (see scheme below and Fig. 1).



The (*R*)-histidinium layer is formed by chains of linked histidinium cations hydrogen bonded from the imidazole ring (N2) to carboxyl oxygen (O1). The two-dimensional layer is created by a hydrogen bond from N3 (ammonium group) to the adjacent (*R*)-histidinium carboxyl group (O2). The (*2R,3R*)-tartrate anions also form chains, with each link created by hydrogen bonding between the carboxyl OH (O3) group to an oxygen (O8) of the carboxylate group (Fig. 2). The (*2R,3R*)-tartrate two-dimensional layer is maintained by a single hydrogen bond from a hydroxyl group (O6) to a neighbouring carboxyl oxygen (O7). The three-dimensional crystal structure is formed of repeating layers of (*R*)-histidinium and (*2R,3R*)-tartrate ions, as shown in Fig. 2. The (*R*)-histidinium layer is linked to one (*2R,3R*)-tartrate layer through two hydrogen bonds from the ammonium group (N3) to O7 (carboxylate). The next layer of tartrates is hydrogen bonded to the imidazole ring (N1–H1···O5) and the carboxylate atom O1. Atom O1 is bifurcated, maintaining both the histidine chains and the tartrate/histidine interaction with O5.

Experimental

A saturated aqueous solution (5 ml) of (*R*)-histidine was mixed with a saturated aqueous solution (5 ml) of (*2R,3R*)-tartaric acid and the vial was covered with a pierced film. This was placed in a larger glass vial containing methanol (25 ml), sealed and allowed to stand for three weeks at room temperature. Crystals of a suitable size for use for single-crystal X-ray diffraction analysis were removed and mounted on glass fibres.

Crystal data

$C_6H_{10}N_3O_2^+ \cdot C_4H_5O_6^-$	$Z = 1$
$M_r = 305.25$	$D_x = 1.602 \text{ Mg m}^{-3}$
Triclinic, $P1$	Mo $K\alpha$ radiation
$a = 5.3712 (18) \text{ \AA}$	Cell parameters from 1902 reflections
$b = 7.637 (3) \text{ \AA}$	$\theta = 2.6\text{--}28.1^\circ$
$c = 8.460 (3) \text{ \AA}$	$\mu = 0.14 \text{ mm}^{-1}$
$\alpha = 72.025 (5)^\circ$	$T = 291 (2) \text{ K}$
$\beta = 73.872 (5)^\circ$	Tablet, colourless
$\gamma = 81.144 (5)^\circ$	$0.5 \times 0.3 \times 0.1 \text{ mm}$
$V = 316.2 (2) \text{ \AA}^3$	

Data collection

Bruker SMART APEX CCD diffractometer	1407 independent reflections
Thin-slice ω scans	1382 reflections with $I > 2\sigma(I)$
Absorption correction: multi-scan (Blessing, 1995)	$R_{\text{int}} = 0.011$
$T_{\text{min}} = 0.802$, $T_{\text{max}} = 0.990$	$\theta_{\text{max}} = 28.4^\circ$
2658 measured reflections	$h = -6 \rightarrow 7$
	$k = -10 \rightarrow 10$
	$l = -11 \rightarrow 11$

Refinement

Refinement on F^2	$w = 1/[\sigma^2(F_o^2) + (0.0511P)^2 + 0.0492P]$
$R[F^2 > 2\sigma(F^2)] = 0.030$	where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.076$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.07$	$\Delta\rho_{\text{max}} = 0.20 \text{ e \AA}^{-3}$
1407 reflections	$\Delta\rho_{\text{min}} = -0.23 \text{ e \AA}^{-3}$
194 parameters	
H-atom parameters constrained	

Table 1

Hydrogen-bonding geometry (\AA , $^\circ$).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1\cdots O8^i$	0.86	1.90	2.760 (2)	173
$N1-H1\cdots O5^i$	0.86	2.55	2.993 (3)	113
$N2-H2A\cdots O1^{ii}$	0.86	2.09	2.810 (3)	142
$N2-H2A\cdots O6^{ii}$	0.86	2.47	2.961 (3)	117
$N3-H3A\cdots O7^{iii}$	0.89	1.91	2.799 (2)	176
$N3-H3C\cdots O2^{iv}$	0.89	1.96	2.822 (3)	162
$O5-H500\cdots O1^v$	0.82	1.95	2.741 (3)	163
$O6-H600\cdots O7^{iv}$	0.82	2.08	2.744 (2)	138
$O3-H300\cdots O8^{ii}$	0.82	1.72	2.533 (2)	172

Symmetry codes: (i) $x - 1, 1 + y, 1 + z$; (ii) $x, 1 + y, z$; (iii) $1 + x, 1 + y, z$; (iv) $1 + x, y, z$; (v) $x, y, z - 1$.

All H atoms were positioned geometrically ($N-H = 0.86\text{--}0.89$, $O-H = 0.82$ and $C-H = 0.93\text{--}0.98 \text{ \AA}$) and refined using a riding model, with $U_{\text{iso}}(\text{H}) = 1.2$ or 1.5 times $U_{\text{eq}}(\text{N})$, $1.5U_{\text{eq}}(\text{O})$ and $1.2U_{\text{eq}}(\text{C})$. In the absence of significant anomalous dispersion effects Friedel pairs were merged prior to refinement.




Figure 1
View of the title compound, with atomic displacement ellipsoids drawn at the 50% probability level.




Figure 2
Hydrogen-bonding (dashed lines) motifs in the (*R*)-histinium (blue) and (*2R,3R*)-tartrate (green) ions of (I).

Data collection: *SMART* (Bruker, 2002); cell refinement: *S SAINT* (Bruker, 2002); data reduction: *S SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *XP* (Sheldrick, 1996) and *Materials Studio* (Accelrys, 2001); software used to prepare material for publication: *SHELXL97*.

References

- Accelrys (2001). *Materials Studio*. Accelrys Ltd, 334 Cambridge Science Park, Cambridge CB4 0WN, England.
- Blessing, R. H. (1995). *Acta Cryst. A* **51**, 33–38.
- Bruker (2002). *S SAINT* (Version 6.02) and *SMART* (Version 5.622). Bruker AXS Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997). *XP*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.