



UNIVERSITY  
OF WOLLONGONG  
AUSTRALIA

University of Wollongong  
Research Online

---

Faculty of Engineering and Information Sciences -  
Papers

Faculty of Engineering and Information Sciences

---

2014

# KMS states on the C-algebras of reducible graphs

Astrid An Huef

*University of Wollongong*, [astrid@uow.edu.au](mailto:astrid@uow.edu.au)

Marcelo Laca

*University of Wollongong*, [marcelo@uow.edu.au](mailto:marcelo@uow.edu.au)

Iain Raeburn

*University of Otago*, [raeburn@uow.edu.au](mailto:raeburn@uow.edu.au)

Aidan Sims

*University of Wollongong*, [asims@uow.edu.au](mailto:asims@uow.edu.au)

---

## Publication Details

An Huef, A., Laca, M., Raeburn, I. & Sims, A. (2014). KMS states on the C-algebras of reducible graphs. *Ergodic Theory and Dynamical Systems*, 1-24.

Research Online is the open access institutional repository for the University of Wollongong. For further information contact the UOW Library:  
[research-pubs@uow.edu.au](mailto:research-pubs@uow.edu.au)

---

# KMS states on the C-algebras of reducible graphs

## Abstract

We consider the dynamics on the C-algebras of finite graphs obtained by lifting the gauge action to an action of the real line. Enomoto, Fujii and Watatani [KMS states for gauge action on OA. Math. Japon. 29 (1984), 607-619] proved that if the vertex matrix of the graph is irreducible, then the dynamics on the graph algebra admits a single Kubo-Martin-Schwinger (KMS) state. We have previously studied the dynamics on the Toeplitz algebra, and explicitly described a finite-dimensional simplex of KMS states for inverse temperatures above a critical value. Here we study the KMS states for graphs with reducible vertex matrix, and for inverse temperatures at and below the critical value. We prove a general result which describes all the KMS states at a fixed inverse temperature, and then apply this theorem to a variety of examples. We find that there can be many patterns of phase transition, depending on the behaviour of paths in the underlying graph.

## Keywords

reducible, algebras, graphs, c, kms, states

## Disciplines

Engineering | Science and Technology Studies

## Publication Details

An Huef, A., Laca, M., Raeburn, I. & Sims, A. (2014). KMS states on the C-algebras of reducible graphs. Ergodic Theory and Dynamical Systems, 1-24.

# KMS states on the $C^*$ -algebras of reducible graphs

ASTRID AN HUEF<sup>†</sup>, MARCELO LACA<sup>‡</sup>, IAIN RAEBURN<sup>†</sup> and AIDAN SIMS<sup>§</sup>

<sup>†</sup> *Department of Mathematics and Statistics, University of Otago, PO Box 56,  
Dunedin 9054, New Zealand*

*(e-mail: astrid@maths.otago.ac.nz, iraeburn@maths.otago.ac.nz)*

<sup>‡</sup> *Department of Mathematics and Statistics, University of Victoria, Victoria,  
Canada BC V8W 3P4*

*(e-mail: laca@math.uvic.ca)*

<sup>§</sup> *School of Mathematics and Applied Statistics, University of Wollongong,  
NSW 2522, Australia*

*(e-mail: asims@uow.edu.au)*

*(Received 22 February 2014 and accepted in revised form 8 May 2014)*

*Abstract.* We consider the dynamics on the  $C^*$ -algebras of finite graphs obtained by lifting the gauge action to an action of the real line. Enomoto, Fujii and Watatani [KMS states for gauge action on  $\mathcal{O}_A$ . *Math. Japon.* **29** (1984), 607–619] proved that if the vertex matrix of the graph is irreducible, then the dynamics on the graph algebra admits a single Kubo–Martin–Schwinger (KMS) state. We have previously studied the dynamics on the Toeplitz algebra, and explicitly described a finite-dimensional simplex of KMS states for inverse temperatures above a critical value. Here we study the KMS states for graphs with reducible vertex matrix, and for inverse temperatures at and below the critical value. We prove a general result which describes all the KMS states at a fixed inverse temperature, and then apply this theorem to a variety of examples. We find that there can be many patterns of phase transition, depending on the behaviour of paths in the underlying graph.

## 1. Introduction

Composing the gauge action of  $\mathbb{T}$  with the map  $t \mapsto e^{it}$  gives a natural dynamics on any Cuntz–Krieger algebra or graph algebra. Enomoto, Fujii and Watatani proved thirty years ago that for a simple Cuntz–Krieger algebra  $\mathcal{O}_A$ , this dynamics admits a unique Kubo–Martin–Schwinger (KMS) state, and that this state has inverse temperature the natural logarithm  $\ln \rho(A)$  of the spectral radius  $\rho(A)$  (which is also the Perron–Frobenius eigenvalue of  $A$ ) [3]. Recently Kajiwara and Watatani revisited this question for

the  $C^*$ -algebras of finite graphs with sources, and found many more KMS states [9]. Other authors are currently interested in KMS states on the  $C^*$ -algebras of infinite graphs [2, 19] or on the  $C^*$ -algebras of higher-rank graphs [7, 20].

We recently studied KMS states on the Toeplitz algebra  $\mathcal{TC}^*(E)$  of a finite graph  $E$  [6]. For inverse temperatures  $\beta$  larger than a critical value  $\beta_c$ , we described a simplex of  $\text{KMS}_\beta$  states whose dimension is determined by the number of vertices in the graph [6, Theorem 3.1]. This gave a concrete implementation of an earlier result of Exel and Laca [4, Theorem 18.4], at least as it applies to the gauge dynamics. The critical inverse temperature  $\beta_c$  in [6] is  $\ln \rho(A)$  where  $A$  is the vertex matrix of the graph  $E$ . When  $A$  is irreducible in the sense of Perron–Frobenius theory (and in particular if  $C^*(E)$  is simple), we showed that there is a unique  $\text{KMS}_{\ln \rho(A)}$  state on  $\mathcal{TC}^*(E)$ , and that this state factors through  $C^*(E)$ .

Here we consider a finite graph  $E$  whose vertex matrix  $A$  is reducible, and aim to find all the KMS states on  $\mathcal{TC}^*(E)$  and  $C^*(E)$ . We have organized our results so that we can describe the KMS states at each fixed inverse temperature. From [6, Theorem 3.1], we already have a concrete description of the simplex of  $\text{KMS}_\beta$  states on  $\mathcal{TC}^*(E)$  for  $\beta > \ln \rho(A)$ , and we know exactly which ones factor through  $C^*(E)$  [6, Corollary 6.1].

Our first main theorem concerns the critical value  $\beta = \ln \rho(A)$  (Theorem 4.3). It identifies two different families of extreme  $\text{KMS}_{\ln \rho(A)}$  states. The first family  $\{\psi_C\}$  is parameterized by a set of strongly connected components  $C$  of  $E$  such that the matrix  $A_C := A|_{C \times C}$  satisfies  $\beta = \ln \rho(A_C)$  (in the theorem we say exactly which components belong to this set). The states  $\psi_C$  all factor through  $C^*(E)$ . Then we consider the hereditary closure  $H$  in  $E^0$  of the components  $C$  with  $\beta = \ln \rho(A_C)$ , and the complementary graph  $E \setminus H$  with vertex set  $E^0 \setminus H$ . The second family  $\{\phi_v\}$  of extremal  $\text{KMS}_{\ln \rho(A)}$  consists of states which factor through a natural quotient map of  $\mathcal{TC}^*(E)$  onto  $\mathcal{TC}^*(E \setminus H)$  (see Proposition 2.1), and is parameterized by  $E^0 \setminus H$ . The convex hull of  $\{\psi_C\} \cup \{\phi_v\}$  is the full simplex of  $\text{KMS}_{\ln \rho(A)}$  states. The proof of Theorem 4.3 involves some rather intricate computations using the Perron–Frobenius theory for the matrices  $A_C$ .

In §5 we describe the  $\text{KMS}_\beta$  states for a fixed inverse temperature  $\beta$  satisfying  $\beta < \ln \rho(A)$ . In Theorem 5.3, we consider the hereditary closure  $H_\beta$  of the connected components  $C$  with  $\ln \rho(A_C) > \beta$ . If  $\beta > \ln \rho(A_{E^0 \setminus H_\beta})$ , the  $\text{KMS}_\beta$  states all factor through the quotient  $\mathcal{TC}^*(E \setminus H_\beta)$ , and an application of [6, Theorem 3.1] gives a concrete description of these states. If  $\beta = \ln \rho(A_{E^0 \setminus H_\beta})$ , then applying Theorem 4.3 to  $E \setminus H_\beta$  shows that there are two families  $\{\psi_C\}$  and  $\{\phi_v\}$  of extremal  $\text{KMS}_\beta$  states. Theorem 5.3 also identifies the states which factor through  $C^*(E)$ , where there are some tricky subtleties involving the saturations of the sets  $H_\beta$  and  $K_\beta$ .

By applying Theorem 5.3 as  $\beta$  decreases, we can in principle find all KMS states on  $\mathcal{TC}^*(E)$  and  $C^*(E)$  for every finite graph  $E$ . In §6 we show how this works on a variety of examples, and find in particular that there are graphs for which our dynamics has many phase transitions. These examples shed considerable light on the possible behaviour of KMS states, and in particular on what happens between the various critical inverse temperatures discussed in [4, §14]. We close with a section of concluding remarks in

which we discuss the range of possible inverse temperatures, and the connections with the results of [2, 4].

## 2. Background

**2.1. Directed graphs and their Toeplitz algebras.** Suppose that  $E = (E^0, E^1, r, s)$  is a directed graph. We use the conventions of [15] for paths, so that, for example,  $ef$  is a path when  $s(e) = r(f)$ . We write  $E^n$  for the set of paths of length  $n$ , and  $E^* := \bigcup_{n \in \mathbb{N}} E^n$ . For vertices  $v, w$ , we write  $vE^n w$  for the set  $\{\mu \in E^n : r(\mu) = v \text{ and } s(\mu) = w\}$  (and we allow variations on this theme).

A Toeplitz–Cuntz–Krieger family  $(P, S)$  consists of mutually orthogonal projections  $\{P_v \mid v \in E^0\}$  and partial isometries  $\{S_e \mid e \in E^1\}$  such that  $S_e^* S_e = P_{s(e)}$  for every  $e \in E^1$  and

$$P_v \geq \sum_{e \in F} S_e S_e^* \quad \text{for every } v \in E^0 \text{ and finite subset } F \text{ of } vE^1 = r^{-1}(v). \quad (2.1)$$

Here we consider only finite graphs, and then it suffices to impose inequality (2.1) for  $F = vE^1$ . The Toeplitz algebra  $\mathcal{TC}^*(E)$  is generated by a universal Toeplitz–Cuntz–Krieger family  $(p, s)$ ; the existence of such an algebra was proved in [5, Theorem 4.1]. For  $\mu \in E^n$ , we define  $s_\mu := s_{\mu_1} s_{\mu_2} \cdots s_{\mu_n}$ . Then each  $s_\mu$  is also a partial isometry, and we have

$$\mathcal{TC}^*(E) := \overline{\text{span}}\{s_\mu s_\nu^* \mid \mu, \nu \in E^*, s(\mu) = s(\nu)\}.$$

We shall work mostly in the Toeplitz algebra  $\mathcal{TC}^*(E)$  rather than the usual graph algebra  $C^*(E)$ , and it is therefore convenient to view  $C^*(E)$  as the quotient of  $\mathcal{TC}^*(E)$  by the ideal generated by

$$\left\{ p_v - \sum_{r(e)=v} s_e s_e^* \mid v \in E^0 \right\}.$$

We write  $\pi_E$  for the quotient map of  $\mathcal{TC}^*(E)$  onto  $C^*(E)$ , and  $\bar{p}_v := \pi_E(p_v)$ ,  $\bar{s}_e := \pi_E(s_e)$ . The pair  $(\bar{p}, \bar{s})$  is then universal for Cuntz–Krieger families in the usual way.

**2.2. Ideals in Toeplitz algebras.** We are interested in graphs whose  $C^*$ -algebras  $C^*(E)$  are not simple. The standard theory (as in [1, 11] or [15, §4]) says that ideals in  $C^*(E)$  are determined by subsets  $H$  of  $E^0$  which are both hereditary ( $v \in H$  and  $vE^* w \neq \emptyset$  imply  $w \in H$ ) and saturated ( $s(vE^1) \subset H$  implies  $v \in H$ ). In the Toeplitz algebra, there are more ideals, and in particular every hereditary subset determines one. We need to know what the quotient is.

**PROPOSITION 2.1.** *Suppose that  $H$  is a hereditary set of vertices in a directed graph  $E$  and that  $H$  is not all of  $E^0$ . Then  $E \setminus H := (E^0 \setminus H, s^{-1}(E^0 \setminus H), r, s)$  is a directed graph, and there is a homomorphism  $q_H : \mathcal{TC}^*(E) \rightarrow \mathcal{TC}^*(E \setminus H) = C^*(p^{E \setminus H}, s^{E \setminus H})$  such that*

$$q_H(p_v) = \begin{cases} p_v^{E \setminus H} & \text{if } v \in E^0 \setminus H, \\ 0 & \text{if } v \in H, \end{cases} \quad \text{and} \quad q_H(s_e) = \begin{cases} s_e^{E \setminus H} & \text{if } s(e) \in E^0 \setminus H, \\ 0 & \text{if } s(e) \in H. \end{cases} \quad (2.2)$$

*The homomorphism is surjective, and its kernel is the ideal  $J_H$  generated by  $\{p_v \mid v \in H\}$ .*

*Proof.* Since  $s$  maps  $(E \setminus H)^1 := s^{-1}(E^0 \setminus H)$  to  $(E \setminus H)^0 := E^0 \setminus H$ , and since  $r(e) \in H$  implies  $s(e) \in H$ ,  $r$  maps  $(E \setminus H)^1$  into  $(E \setminus H)^0$  also. Thus  $E \setminus H$  is a directed graph. The formulas on the right-hand sides of (2.2) define a Toeplitz–Cuntz–Krieger  $E$ -family in  $\mathcal{TC}^*(E \setminus H)$ , and hence the universal property of  $\mathcal{TC}^*(E)$  gives the existence of the homomorphism  $q_H$ . It is surjective because its range contains all the generators of  $\mathcal{TC}^*(E \setminus H)$ . The kernel of  $q_H$  contains all the generators of  $J_H$ , so  $J_H \subset \ker q_H$ , and hence  $q_H$  factors through the quotient map  $q : \mathcal{TC}^*(E) \rightarrow \mathcal{TC}^*(E)/J_H$ . We write  $\bar{q}_H$  for the homomorphism on  $\mathcal{TC}^*(E)/J_H$  such that  $q_H = \bar{q}_H \circ q$ .

To see that  $J_H$  is all of  $\ker q_H$ , we construct a left inverse for  $\bar{q}_H$ . A quick check shows that the elements  $\{q(p_v), q(s_e) \mid v \in E^0 \setminus H, e \in s^{-1}(E^0 \setminus H)\}$  form a Toeplitz–Cuntz–Krieger  $(E \setminus H)$ -family in  $\mathcal{TC}^*(E)/J_H$ . (It is crucial that we are not trying to impose a Cuntz–Krieger relation at vertices in  $E^0 \setminus H$  which receive edges from  $H$ .) Thus there is a homomorphism  $\rho : \mathcal{TC}^*(E \setminus H) \rightarrow \mathcal{TC}^*(E)/J_H$  such that  $\rho(p_v^{E \setminus H}) = q(p_v)$  and  $\rho(s_e^{E \setminus H}) = q(s_e)$ . Since  $s(e) \in H$  implies that  $q(s_e) = 0$ , the range of  $\rho$  contains the images of all the generators of  $\mathcal{TC}^*(E)$ , and hence  $\rho$  is surjective. A quick check shows that  $\bar{q}_H \circ \rho$  fixes the generators of  $\mathcal{TC}^*(E \setminus H)$ , and hence is the identity on  $\mathcal{TC}^*(E \setminus H)$ . Now the surjectivity of  $\rho$  implies that  $\rho \circ \bar{q}_H$  is the identity on  $\mathcal{TC}^*(E)/J_H$ , so  $\bar{q}_H$  is injective, and we have  $\ker q_H = J_H$ .  $\square$

**2.3. Decompositions of the vertex matrix.** Let  $E$  be a finite directed graph. The vertex matrix of  $E$  is the  $E^0 \times E^0$  matrix  $A$  with entries  $A(v, w) = |vE^1w|$ ; the powers of  $A$  then have entries  $A^n(v, w) = |vE^n w|$ . We will do computations using block decompositions of the vertex matrix  $A$ . For subsets  $C, D \subset E^0$ , we write  $A_{C,D}$  for the  $C \times D$  subblock of  $A$ , and  $A_C := A_{C,C}$ . We usually choose decompositions of  $E^0 = C_1 \sqcup C_2 \sqcup \cdots \sqcup C_n$  such that the associated block decomposition of  $A$  is upper-triangular.

For  $v, w \in E^0$ , we write  $v \leq w \iff vE^*w \neq \emptyset$ , and  $v \sim w \iff v \leq w$  and  $w \leq v$ . It is easy to check that  $\sim$  is an equivalence relation on  $E^0$  (we have  $v \sim v$  for all  $v \in E^0$  because  $E^0 \subset E^*$ ). We write  $E^0/\sim$  for the set of equivalence classes, and refer to these equivalence classes as the *strongly connected components* of  $E$ . When  $C \in E^0/\sim$ , the matrix  $A_C$  is either a  $1 \times 1$  zero matrix (if  $C = \{v\}$  is a single vertex with no loops, in which case we say  $C$  is a trivial component), or an irreducible matrix in the sense of Perron–Frobenius theory (so that for every  $v, w \in C$ , there exists  $n$  such that  $A^n(v, w) > 0$ ).

We next order the vertex set  $E^0$  to ensure that the vertex matrix takes a convenient block upper-triangular form. The relation  $\leq$  descends to a well-defined partial order on  $E^0/\sim$ ; when  $C \leq D$ , we say that  $D$  talks to  $C$ . We list first the trivial components for which  $A_C = (0)$  and which do not talk to non-trivial components; we list them in an order such that  $w$  appears after  $v$  when  $v \leq w$ . Next we list the components which are minimal for the order  $\leq$  on the remaining components, grouping the vertices in the same component together. Then we list the trivial components which talk only to the components we have listed so far, and so on. This decomposes  $A$  as a block upper-triangular matrix in which the diagonal components  $A_C$  are either  $1 \times 1$  zero matrices or irreducible. We will refer to such a decomposition as a *Seneta decomposition* of  $A$ . (Though since Seneta uses different conventions in [16, §1.2], the decomposition he discusses

there is a block lower-triangular matrix and our minimal components would become maximal<sup>†</sup>.)

2.4. *KMS states.* We denote the gauge actions of  $\mathbb{T}$  on  $\mathcal{TC}^*(E)$  and  $C^*(E)$  by  $\gamma$ . We are interested in the dynamics  $\alpha$  given, on both  $\mathcal{TC}^*(E)$  and  $C^*(E)$ , by  $\alpha_t = \gamma_{e^{it}}$ . For KMS states, we use the conventions of our previous paper [6]. Thus we know from [6, Proposition 2.1] that a state  $\phi$  of  $\mathcal{TC}^*(E)$  is a  $\text{KMS}_\beta$  state for some  $\beta \in \mathbb{R}$  if and only if

$$\phi(s_\mu s_\nu^*) = \delta_{\mu,\nu} e^{-\beta|\mu|} \phi(p_{s(\mu)}) \quad \text{for all } \mu, \nu \in E^*. \quad (2.3)$$

For fixed  $\beta$  the  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  form a simplex, which we shall refer to as the *KMS $_\beta$  simplex* of  $(\mathcal{TC}^*(E), \alpha)$ . The  $\text{KMS}_0$  states are the invariant traces.

Since the results of [6, §3] already describe all the KMS states for large inverse temperatures, we do not have anything new to say about ground states or  $\text{KMS}_\infty$  states.

### 3. KMS states and quotients

When the vertex matrix  $A$  of  $E$  is irreducible, there are no  $\text{KMS}_\beta$  states on the Toeplitz algebra  $\mathcal{TC}^*(E)$  when  $\beta < \ln \rho(A)$ . So it seems reasonable that if  $C$  is a strongly connected component with  $\ln \rho(A_C) > \beta$ , then every  $\text{KMS}_\beta$  state must vanish on vertex projections  $p_v$  with  $v \in C$ . The key to our analysis of reducible graphs is that KMS states must also vanish on any projections  $p_v$  for vertices  $v$  that connect to such components  $C$ . The next result makes this precise.

**PROPOSITION 3.1.** *Suppose that  $H$  is a hereditary subset of  $E^0$ , and  $q_H : \mathcal{TC}^*(E) \rightarrow \mathcal{TC}^*(E \setminus H)$  is the surjection of Proposition 2.1. Then for every  $\beta \in [0, \infty)$ ,  $q_H^* : \psi \mapsto \psi \circ q_H$  is an affine injection of the  $\text{KMS}_\beta$  simplex of  $(\mathcal{TC}^*(E \setminus H), \alpha)$  into the  $\text{KMS}_\beta$  simplex of  $(\mathcal{TC}^*(E), \alpha)$ . If  $\{C \in H/\sim \mid \ln \rho(A_C) > \beta\}$  generates  $H$  as a hereditary subset of  $E^0$ , then  $\phi(p_v) = 0$  for every  $\text{KMS}_\beta$  state  $\phi$  on  $\mathcal{TC}^*(E)$  and every  $v \in H$ ; if in addition  $H$  is not all of  $E^0$ , then  $q_H^*$  is surjective.*

**LEMMA 3.2.** *Suppose that  $H$  is the hereditary subset of  $E^0$  generated by  $\mathcal{C}_1 \subset E^0/\sim$ , and that  $\beta \leq \ln \rho(A_C)$  for all  $C \in \mathcal{C}_1$ . Suppose that  $\phi$  is a  $\text{KMS}_\beta$  state on  $(\mathcal{TC}^*(E), \alpha)$ , and that  $v$  belongs to the complement of  $\bigcup\{C \in \mathcal{C}_1 \mid \beta = \ln \rho(A_C)\}$  in  $H$ . Then  $\phi(p_v) = 0$ .*

When  $\{C \in H/\sim \mid \ln \rho(A_C) > \beta\}$  generates  $H$ , as in Proposition 3.1, Lemma 3.2 applies to every  $v \in H$  with  $\mathcal{C}_1 = \{C \in H/\sim \mid \ln \rho(A_C) > \beta\}$ . The extra generality in the lemma will be useful in the proof of Proposition 4.1 below.

*Proof.* For every path  $\mu$  with  $s(\mu) = v$ , (2.1) implies that  $p_{r(\mu)} \geq s_\mu s_\mu^*$ , and hence

$$0 \leq \phi(p_v) = \phi(s_\mu^* s_\mu) = e^{\beta|\mu|} \phi(s_\mu s_\mu^*) \leq e^{\beta|\mu|} \phi(p_{r(\mu)}). \quad (3.1)$$

<sup>†</sup> Unfortunately, there is no universal convention as to whether  $A(v, w)$  should refer to edges from  $w$  to  $v$  or edges from  $v$  to  $w$ . Our convention arises from viewing directed edges as arrows in a category, in which case one expects  $ef := e \circ f$  to have source  $s(f)$  and range  $r(e)$ . This convention is standard in many places: for example, in the substantial literature on higher-rank graphs, which have strong links to higher-dimensional subshifts [10, 14], and in studying equivalences for categories of modules over path algebras of quivers [17, 18]; see especially the discussion in [17, §5.4].

Proposition 2.1(c) of [6] implies that the vector  $m^\phi := (\phi(p_w))$  in  $[0, 1]^{E^0}$  satisfies the subinvariance relation  $Am^\phi \leq e^\beta m^\phi$ , and for every  $C \in \mathcal{C}_1$  we have

$$A_C(m^\phi|_C) \leq A_C(m^\phi|_C) + A_{C, H \setminus C}(m^\phi|_{H \setminus C}) = (Am^\phi)|_C \leq e^\beta m^\phi|_C. \quad (3.2)$$

Since  $v \in H$  and  $H$  is generated by  $\mathcal{C}_1$ , there exists  $C \in \mathcal{C}_1$  such that  $CE^*v \neq \emptyset$ . Then either  $\beta < \ln \rho(A_C)$  or  $\beta = \ln \rho(A_C)$ . Suppose that  $\beta < \ln \rho(A_C)$ . Then (3.2) and the last sentence in Theorem 1.6 of [16] imply that  $m^\phi|_C = 0$ . We can therefore apply (3.1) to any  $\mu \in CE^*v$ , and deduce that  $\phi(p_v) = 0$ .

Now suppose that  $\beta = \ln \rho(A_C)$ . Then by hypothesis  $v \notin C$ , and there exists  $\lambda \in CE^*v$  of the form  $\lambda = e\mu$ , where  $e \in E^1$ ,  $r(e) \in C$  and  $s(e) \notin C$ . If  $m^\phi|_C = 0$ , then we can apply (3.1) to  $\mu$  and deduce that  $\phi(p_v) = 0$ . So we suppose that  $m^\phi|_C \neq 0$ . Then (3.2) and [16, Theorem 1.6] imply that  $m^\phi|_C$  is a multiple of the Perron–Frobenius eigenvector for  $A_C$ . Since

$$\begin{aligned} (A_C(m^\phi|_C))_{r(e)} &\leq (A_C(m^\phi|_C))_{r(e)} + A(r(e), s(e))m_{s(e)}^\phi \\ &\leq ((Am^\phi)|_C)_{r(e)} \leq e^\beta m_{r(e)}^\phi = \rho(A_C)(m^\phi|_C)_{r(e)}, \end{aligned} \quad (3.3)$$

and the left and right ends of (3.3) are equal, we deduce that  $A(r(e), s(e))m_{s(e)}^\phi = 0$  and  $\phi(p_{r(\mu)}) = m_{s(e)}^\phi = 0$ ; now (3.1) implies that  $\phi(p_v) = 0$ .  $\square$

*Proof of Proposition 3.1.* Since  $(E \setminus H)^* = \{\mu \in E^* \mid s(\mu) \notin H\}$ , we can deduce from [6, Proposition 2.1(a)] that  $\psi \circ q_H$  is a KMS state if and only if  $\psi$  is. Since  $q_H$  is surjective,  $q_H^*$  is injective, and it is clearly weak\* continuous and affine. To see the assertion about surjectivity, suppose that  $\{C \in H/\sim \mid \ln \rho(A_C) > \beta\}$  generates  $H$  and  $\phi$  is a  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E), \alpha)$ . Lemma 3.2 implies that  $\phi(p_v) = 0$  for all  $v \in H$ . Now we can apply [6, Lemma 2.2] with  $\mathcal{F} = \{s_\mu s_\nu^* \mid \mu, \nu \in E^*\}$  and  $P = \{p_v \mid v \in H\}$ , and deduce that  $\phi$  factors through a state of  $\mathcal{TC}^*(E)/J_H = \mathcal{TC}^*(E)/\ker q_H$ . Thus if  $H \neq E^0$ , there is a state  $\psi$  of  $\mathcal{TC}^*(E \setminus H)$  such that  $\phi = \psi \circ q_H$ . Since  $q_H$  is surjective and is equivariant for the various actions  $\alpha$ ,  $\psi$  is a  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E \setminus H), \alpha)$ .  $\square$

The analogue of Proposition 3.1 for the graph algebra  $C^*(E)$  has a slightly different hypothesis: it suffices that  $\{C \mid \ln \rho(A_C) > \beta\}$  generates  $H$  as a *saturated* hereditary set. This happens because the identification of  $C^*(E)/I_H$  with  $C^*(E \setminus H)$  only works when  $H$  is saturated (compare Proposition 2.1 with [1, Theorem 4.1] or [15, Theorem 4.9]).

**PROPOSITION 3.3.** *Suppose that  $H$  is a saturated hereditary subset of  $E^0$ , and write  $\bar{q}_H$  for the canonical surjection of  $C^*(E)$  onto  $C^*(E \setminus H)$ . Then for every  $\beta \in [0, \infty)$ ,  $\bar{q}_H^* : \psi \mapsto \psi \circ \bar{q}_H$  is an affine injection of the  $\text{KMS}_\beta$  simplex of  $(C^*(E \setminus H), \alpha)$  into the  $\text{KMS}_\beta$  simplex of  $(C^*(E), \alpha)$ . If  $\{C \in H/\sim \mid \ln \rho(A_C) > \beta\}$  generates  $H$  as a saturated hereditary subset of  $E^0$ , then  $\phi(p_v) = 0$  for every  $\text{KMS}_\beta$  state  $\phi$  on  $C^*(E)$  and every  $v \in H$ ; if in addition  $H$  is not all of  $E^0$ , then  $\bar{q}_H^*$  is surjective.*

For the proof we need a simple lemma. Recall from the proof of [6, Corollary 6.1], for example, that the saturation  $\Sigma H$  of a hereditary set  $H$  can be viewed as  $\bigcup_{k=0}^\infty S_k H$ , where  $S_k H$  are the subsets of  $E^0$  defined recursively by

$$S_0 H = H \quad \text{and} \quad S_{k+1} H = S_k H \cup \{v \mid s(vE^1) \subset S_k H\}. \quad (3.4)$$



LEMMA 3.4. Suppose that  $H$  is a hereditary subset of  $E^0$ ,  $\beta \in [0, \infty)$ , and  $\phi, \psi$  are  $\text{KMS}_\beta$  states on  $(C^*(E), \alpha)$ .

- (a) If  $\phi(p_v) = \psi(p_v)$  for all  $v \in H$ , then  $\phi = \psi$  on the ideal  $I_H$  of  $C^*(E)$  generated by  $\{p_v \mid v \in H\}$ .
- (b) If  $\phi(p_v) = 0$  for all  $v \in H$ , then  $\phi(p_v) = 0$  for all  $v$  in the saturation  $\Sigma H$ .

*Proof.* For (a), we first claim that  $\phi(p_v) = \psi(p_v)$  for all  $v \in \Sigma H$ . We are given that  $\phi(p_v) = \psi(p_v)$  for  $v \in S_0 H$ . Suppose that  $\phi(p_v) = \psi(p_v)$  for  $v \in S_k H$ . Then for  $v \in S_{k+1} H$  and  $e \in vE^1$ , we have  $s(e) \in S_k H$ , and

$$\begin{aligned} \phi(p_v) &= \phi\left(\sum_{e \in vE^1} s_e s_e^*\right) = \sum_{e \in vE^1} e^{-\beta} \phi(p_{s(e)}) \\ &= \sum_{e \in vE^1} e^{-\beta} \psi(p_{s(e)}) = \psi(p_v). \end{aligned} \tag{3.5}$$

Thus by induction we have  $\phi(p_v) = \psi(p_v)$  for all  $v \in S_k H$  and all  $k$ , and hence for all  $v \in \Sigma H$ , as claimed.

Next, we recall that

$$I_H = \overline{\text{span}}\{s_\mu s_\nu^* \mid s(\mu) = s(\nu) \in \Sigma H\}$$

(see [1, Lemma 4.3], for example). For a typical spanning element  $s_\mu s_\nu^*$ , [6, equation (2.1)] says that

$$\phi(s_\mu s_\nu^*) = \delta_{\mu,\nu} e^{-\beta|\mu|} \phi(p_{s(\mu)}) = \delta_{\mu,\nu} e^{-\beta|\mu|} \psi(p_{s(\mu)}) = \psi(s_\mu s_\nu^*),$$

and it follows from linearity and continuity that  $\phi = \psi$  on  $I_H$ .

For (b), we repeat the induction argument of the first paragraph, and in particular the computation in the first line of (3.5).  $\square$

*Proof of Proposition 3.3.* As in the proof of Proposition 3.1,  $\bar{q}_H^* : \psi \mapsto \psi \circ \bar{q}_H$  is an affine injection of the  $\text{KMS}_\beta$  simplex of  $C^*(E \setminus H)$  into the  $\text{KMS}_\beta$  simplex of  $C^*(E)$ . Suppose that  $\phi$  is a  $\text{KMS}_\beta$  state of  $C^*(E)$ . Then applying Proposition 3.1 to the hereditary closure  $H_0$  of  $\{C \in H/\sim \mid \beta < \ln \rho(A_C)\}$  shows that  $\phi(\bar{p}_v) = \phi \circ \pi_E(p_v) = 0$  for  $v \in H_0$ . Thus  $\{v \in E^0 \mid \psi(\bar{p}_v) = 0\}$  contains  $H_0$ , and hence by Lemma 3.4 contains  $\Sigma H_0 = H$ . Now [6, Lemma 2.2] implies that  $\phi$  factors through a state of  $C^*(E)/I_H$ , and hence there is a state  $\psi$  of  $C^*(E \setminus H)$  such that  $\phi = \psi \circ \bar{q}_H$ . Then the surjectivity of  $\bar{q}_H$  implies that  $\psi$  is a  $\text{KMS}_\beta$  state, and  $\bar{q}_H^*(\psi) = \phi$ .  $\square$

#### 4. KMS states on Toeplitz algebras

We suppose that  $E$  has at least one cycle, so that  $\rho(A) \geq 1$  (by [6, Lemma A.1]), and the critical inverse temperature  $\ln \rho(A) \geq 0$ . Since a Seneta decomposition of  $A$  is upper triangular as a block matrix, we have

$$\rho(A) = \max\{\rho(A_C) \mid C \in E^0/\sim \text{ is a non-trivial strongly connected component}\}.$$

We therefore focus on the set

$$\{C \in E^0/\sim \mid \rho(A_C) = \rho(A)\} \tag{4.1}$$

of *critical components* of  $E$ , and in particular on the set  $\text{mc} = \text{mc}(E)$  of *minimal critical components* that are minimal in the induced partial order on the set (4.1).

The results of the previous section imply that if  $\beta < \ln \rho(A)$ , then every  $\text{KMS}_\beta$  state on  $\mathcal{TC}^*(E)$  vanishes on the hereditary closure of  $\{C \mid \ln \rho(A_C) = \ln \rho(A)\}$ . This hereditary closure is the same as that of  $\text{mc}(E)$ . So the location of the minimal critical components in the graph plays an important role in our analysis. Because the minimal critical components are minimal in (4.1), they cannot talk to each other. Thus in a Seneta decomposition of the vertex matrix  $A$ , our conventions ensure that the diagonal blocks  $\{A_C \mid C \in \text{mc}(E)\}$  associated to the minimal critical components appear in the decomposition above other critical components  $A_D$ .

The next result is a new version of [6, Theorem 2.1(a)].

**PROPOSITION 4.1.** *Suppose that  $E$  has at least one cycle. Let  $K = \bigcup_{C \in \text{mc}(E)} C$ , let  $H$  be the hereditary closure of  $K$ , and let  $L$  be the union of the non-trivial strongly connected components. Let  $\beta \in \mathbb{R}$ . Then:*

- (a)  $\rho(A_{E^0 \setminus H}) < \rho(A)$ ;
- (b) if  $\phi$  is a  $\text{KMS}_{\ln \rho(A)}$  state of  $(\mathcal{TC}^*(E), \alpha)$ , then  $\phi(p_v) = 0$  for  $v \in H \setminus K$ ;
- (c) if  $E^0$  is the hereditary closure of  $K$  and  $\phi$  is a  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E), \alpha)$ , then  $\ln \rho(A) \leq \beta$ ;
- (d) if  $E^0$  is the hereditary closure of  $L$  and  $\phi$  is a  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E), \alpha)$ , then there is a non-trivial component  $C$  with  $\ln \rho(A_C) \leq \beta$ ;
- (e) if  $E^0$  is the saturated hereditary closure of  $L$  and  $\phi$  is a  $\text{KMS}_\beta$  state of  $(C^*(E), \alpha)$ , then there is a non-trivial component  $C$  with  $\ln \rho(A_C) \leq \beta$ .

*Proof.* Since every minimal element of (4.1) is contained in  $H$ , so is every other strongly connected component  $C$  in (4.1). Thus  $\rho(A_C) < \rho(A)$  for every strongly connected component  $C$  that is contained in  $E^0 \setminus H$ , and

$$\rho(A_{E^0 \setminus H}) = \max\{\rho(A_C) \mid C \in E^0 / \sim, C \subset E^0 \setminus H\} < \rho(A),$$

which is (a). Next suppose that  $\phi$  is a  $\text{KMS}_{\ln \rho(A)}$  state on  $(\mathcal{TC}^*(E), \alpha)$ . We set things up so  $\text{mc}(E) = \{C \in \text{mc}(E) \mid \rho(A_C) = \rho(A)\}$ , so we can apply Lemma 3.2 with  $\beta = \ln \rho(A)$  and  $\mathcal{C}_1 = \text{mc}(E)$ , and (b) follows.

For (c), we suppose that  $\ln \rho(A) > \beta$ . Then  $\text{mc}(E) \subset \{C \in E^0 / \sim \mid \ln \rho(A_C) > \beta\}$ , and hence the hypothesis implies that  $\{C \mid \ln \rho(A_C) > \beta\}$  generates  $E^0$ . So Proposition 3.1 applies with  $H = E^0$ . Thus  $\phi(p_v) = 0$  for all  $v \in E^0$ , and  $1 = \phi(1) = \sum_{v \in E^0} \phi(p_v) = 0$ , which is a contradiction. A similar argument gives (d). For (e), we repeat the argument yet again, using Proposition 3.3 instead of Proposition 3.1.  $\square$

*Remark 4.2.* If the hereditary closure  $G$  of  $L$  is not all of  $E^0$ , then  $\rho(A \setminus G) = 0$ , and [6, Theorem 3.1] applies to  $E \setminus G$  and every  $\beta \in \mathbb{R}$ . Thus if  $\beta < \ln \rho(A_C)$  for every non-trivial component  $C$ , there is a  $(|E^0 \setminus G| - 1)$ -dimensional simplex of  $\text{KMS}_\beta$  states on  $\mathcal{TC}^*(E \setminus G)$ . It follows from Proposition 3.1 that there is also a  $(|E^0 \setminus G| - 1)$ -dimensional simplex of  $\text{KMS}_\beta$  states on  $\mathcal{TC}^*(E)$ . Whether any of these factor through  $C^*(E)$  will depend on whether  $E \setminus \Sigma G$  has sources (see [6, Corollary 6.1]), and Example 6.4 shows that  $E \setminus \Sigma G$  can have sources.

Proposition 4.1 implies that the  $\text{KMS}_{\ln \rho(A)}$  simplex does not see the set  $H \setminus K$ , and hence (via [6, Lemma 2.2]) that the  $\text{KMS}_{\ln \rho(A)}$  states vanish on the ideal  $J_{H \setminus K}$  generated by  $\{p_v \mid v \in H \setminus K\}$ . Our next result describes how the minimal critical components give rise to  $\text{KMS}_{\ln \rho(A)}$  states.

**THEOREM 4.3.** *Suppose that  $E$  is a directed graph with at least one cycle. Let  $K = \bigcup_{C \in \text{mc}(E)} C$ , and let  $H := \{v \in E^0 \mid KE^*v \neq \emptyset\}$  be the hereditary closure of  $K$ .*

- (a) *Let  $C \in \text{mc}(E)$  be a minimal critical component, and let  $x^C$  be the unimodular Perron–Frobenius eigenvector of  $A_C$  (that is, the one with  $\|x^C\|_1 = 1$ ). Define a vector  $z^C \in [0, \infty)^{E^0 \setminus H}$  by*

$$z^C := \rho(A)^{-1} (1 - \rho(A)^{-1} A_{E^0 \setminus H})^{-1} A_{E^0 \setminus H, C} x^C. \quad (4.2)$$

*Then there is a  $\text{KMS}_{\ln \rho(A)}$  state  $\psi_C$  of  $(\mathcal{T}C^*(E), \alpha)$  such that*

$$\psi_C(s_\mu s_\nu^*) = \delta_{\mu, \nu} \rho(A)^{-|\mu|} (1 + \|z^C\|_1)^{-1} \begin{cases} z_{s(\mu)}^C & \text{if } s(\mu) \in E^0 \setminus H, \\ x_{s(\mu)}^C & \text{if } s(\mu) \in C, \\ 0 & \text{if } s(\mu) \in H \setminus C. \end{cases} \quad (4.3)$$

*The state  $\psi_C$  factors through a  $\text{KMS}_{\ln \rho(A)}$  state  $\bar{\psi}_C$  of  $(C^*(E), \alpha)$ .*

- (b) *The map  $t \mapsto \sum_{C \in \text{mc}(E)} t_C \psi_C$  is an affine isomorphism of*

$$S_E := \left\{ t \in [0, 1]^{\text{mc}(E)} \mid \sum_{C \in \text{mc}(E)} t_C = 1 \right\}$$

*onto a simplex  $\Sigma_{\text{mc}(E)}$  of  $\text{KMS}_{\ln \rho(A)}$  states of  $(\mathcal{T}C^*(E), \alpha)$ . Every  $\text{KMS}_{\ln \rho(A)}$  state of  $(\mathcal{T}C^*(E), \alpha)$  is a convex combination of a state of the form  $q_H^*(\phi) = \phi \circ q_H$  and a state in  $\Sigma_{\text{mc}(E)}$ .*

The idea in part (a) is that the values of a KMS state on vertices in  $C$  contribute to the values  $\phi(p_v)$  for  $v \in E^0 \setminus H$  when there are paths  $\lambda$  from  $C$  to  $v$ . As discussed at the beginning of [6, §3], for  $\beta > \ln \rho(A_{E^0 \setminus H})$  the series  $\sum_{n=0}^{\infty} e^{-\beta n} A_{E^0 \setminus H}^n$  converges in operator norm to  $(1 - e^{-\beta} A_{E^0 \setminus H})^{-1}$ , and so

$$(1 - e^{-\beta} A_{E^0 \setminus H})^{-1}(v, w) = \sum_{n=0}^{\infty} e^{-\beta n} A_{E^0 \setminus H}^n(v, w) = \sum_{\lambda \in vE^*w} e^{-\beta|\lambda|}. \quad (4.4)$$

Since  $\rho(A) > \rho(A_{E^0 \setminus H})$ , the  $(E^0 \setminus H) \times C$  matrix  $(1 - \rho(A)^{-1} A_{E^0 \setminus H})^{-1} A_{E^0 \setminus H, C}$  in equation (4.2) has entries

$$((1 - \rho(A)^{-1} A_{E^0 \setminus H})^{-1} A_{E^0 \setminus H, C})(v, w) = \sum_{e \in (E^0 \setminus H)E^1w} \sum_{\mu \in vE^*r(e)} \rho(A)^{-|\mu|}. \quad (4.5)$$

We use (4.4) in the proof of part (a) and (4.5) in the proof of part (b), and again in Lemma 5.7 and Theorem 5.3(c).

*Proof of Theorem 4.3(a).* We partition  $E^0$  as  $(E^0 \setminus H) \cup C \cup (H \setminus C)$ , and claim that the vector  $(z^C, x^C, 0)$  satisfies

$$A(z^C, x^C, 0) = \rho(A)(z^C, x^C, 0). \quad (4.6)$$

Since  $C$  is minimal, it does not talk to any of the other components in  $H \setminus C$ , and we have

$$A(z^C, x^C, 0) = (A_{E^0 \setminus H} z^C + A_{E^0 \setminus H, C} x^C, A_C x^C, 0). \quad (4.7)$$

We know that  $A_C x^C = \rho(A) x^C$ , so we concentrate on the first term. Proposition 4.1 implies that  $\rho(A_{E^0 \setminus H}) < \rho(A)$ . Since  $e^{-\rho(A)} = \rho(A)^{-1}$ , (4.4) gives

$$z^C = \sum_{n=0}^{\infty} \rho(A)^{-n-1} A_{E^0 \setminus H}^n A_{E^0 \setminus H, C} x^C,$$

and we have

$$\begin{aligned} A_{E^0 \setminus H} z^C + A_{E^0 \setminus H, C} x^C &= A_{E^0 \setminus H} \left( \sum_{n=0}^{\infty} \rho(A)^{-n-1} A_{E^0 \setminus H}^n A_{E^0 \setminus H, C} x^C \right) + A_{E^0 \setminus H, C} x^C \\ &= \left( \sum_{m=1}^{\infty} \rho(A)^{-m} A_{E^0 \setminus H}^m A_{E^0 \setminus H, C} x^C \right) + A_{E^0 \setminus H, C} x^C \\ &= \sum_{m=0}^{\infty} \rho(A)^{-m} A_{E^0 \setminus H}^m A_{E^0 \setminus H, C} x^C \\ &= \rho(A) z^C. \end{aligned}$$

From this and (4.7), we deduce that  $(z^C, x^C, 0)$  satisfies (4.6), as claimed.

Since  $x^C$  is unimodular,  $m := (1 + \|z^C\|_1)^{-1} (z^C, x^C, 0)$  satisfies  $\|m\|_1 = 1$ , and hence is a probability measure on  $E^0$ . Equation (4.6) implies that  $Am = \rho(A)m$ . Thus [6, Proposition 4.1] implies that there is a  $\text{KMS}_{\ln \rho(A)}$  state  $\psi_C$  on  $(\mathcal{T}C^*(E), \alpha)$  satisfying (4.3), and that  $\psi_C$  factors through a  $\text{KMS}_{\ln \rho(A)}$  state of  $(C^*(E), \alpha)$ .  $\square$

The double sum appearing on the right-hand side of (4.5) is parameterized by paths in  $vE^*w$  of the form  $\mu e$ , where  $r(e)$  is in  $E^0 \setminus C$  and  $\mu$  is a path in  $E \setminus H$ . We say that such paths *make a quick exit from  $C$* . For a minimal critical component  $C$ , we write  $\text{QE}(C)$  for the set

$$\text{QE}(C) := \{\mu e \mid e \in E^1 C, r(e) \notin C, \mu \in E^* r(e)\}$$

of paths which start in  $C$  and make a quick exit from  $C$ , and  $\text{QE}(K) := \bigcup_{C \in \text{emc}(E)} \text{QE}(C)$ . With this notation, the right-hand side of (4.5) becomes

$$\sum_{\lambda \in v \text{QE}(C) w} \rho(A)^{-(|\lambda|-1)}.$$

LEMMA 4.4. *The projections  $\{s_\lambda s_\lambda^* \mid \lambda \in \text{QE}(K)\}$  are mutually orthogonal.*

*Proof.* Suppose that  $\mu, \nu \in \text{QE}(K)$  and  $\mu \neq \nu$ . If  $|\mu| = |\nu|$ , then  $(s_\mu s_\mu^*)(s_\nu s_\nu^*) = s_\mu (s_\mu^* s_\nu) s_\nu^* = 0$ . So suppose that one path is longer, say  $|\mu| > |\nu|$ . Then  $s(\nu)$  is in  $K$  and  $s(\mu_{|\nu|})$  is not in  $K$  because the different minimal critical components do not talk to each other. Thus  $\mu$  does not have the form  $\nu \mu'$ , and we have  $s_\mu^* s_\nu = 0$ , which implies that  $(s_\mu s_\mu^*)(s_\nu s_\nu^*) = 0$ .  $\square$

*Proof of Theorem 4.3(b).* Suppose that  $\phi$  is a  $\text{KMS}_{\ln \rho(A)}$  state of  $(\mathcal{TC}^*(E), \alpha)$ , and consider  $m^\phi = (\phi(p_v))$ , which by [6, Proposition 2.1(c)] satisfies the subinvariance relation  $Am^\phi \leq \rho(A)m^\phi$ . Suppose that  $C \in \text{mc} = \text{mc}(E)$ . Proposition 4.1 implies that  $m_v^\phi = 0$  for  $v \in H \setminus K$ , which since the minimal critical components do not talk to each other implies that  $(Am^\phi)|_C = A_C(m^\phi|_C)$ . So subinvariance implies that

$$A_C(m^\phi|_C) = (Am^\phi)|_C \leq \rho(A)m^\phi|_C = \rho(A_C)m^\phi|_C;$$

now [16, Theorem 1.6] implies that we have equality throughout, and that  $m^\phi|_C$  is a multiple of the unimodular Perron–Frobenius eigenvector  $x^C$  for  $A_C$ . We define  $t_C \in [0, \infty)$  by  $m^\phi|_C = t_C(1 + \|z^C\|_1)^{-1}x^C$ .

We claim that  $\sum_{C \in \text{mc}} t_C \leq 1$ . For  $v \in E^0 \setminus H$ , Lemma 4.4 implies that  $\phi(p_v) \geq \sum_{\lambda \in v \text{ QE}(K)} \phi(s_\lambda s_\lambda^*)$ . Now we calculate, using [6, Proposition 2.1(a)] and (4.5):

$$\begin{aligned} \phi(p_v) &\geq \sum_{\lambda \in v \text{ QE}(K)} \phi(s_\lambda s_\lambda^*) = \sum_{\lambda \in v \text{ QE}(K)} \rho(A)^{-|\lambda|} \phi(p_{s(\lambda)}) \\ &= \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} \left( \sum_{\lambda \in v \text{ QE}(C)} \rho(A)^{-|\lambda|} x_{s(\lambda)}^C \right) \\ &= \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} \left( \sum_{w \in C} \sum_{\lambda \in v \text{ QE}(C)w} \rho(A)^{-|\lambda|} x_w^C \right) \\ &= \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} \\ &\quad \times \left( \sum_{w \in C} \rho(A)^{-1} ((1 - \rho(A))^{-1} A_{E^0 \setminus H})^{-1} A_{E^0 \setminus H, C} \right) (v, w) x_w^C \\ &= \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} z_v^C. \end{aligned} \tag{4.8}$$

For  $v \in C$  we have  $\phi(p_v) = t_C(1 + \|z^C\|_1)^{-1}x_v^C$  by definition of  $t_C$ . Thus

$$\begin{aligned} 1 = \phi(1) &= \sum_{v \in E^0} \phi(p_v) \geq \sum_{v \in E^0 \setminus H} \phi(p_v) + \sum_{C \in \text{mc}} \sum_{v \in C} \phi(p_v) \\ &\geq \sum_{v \in E^0 \setminus H} \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} z_v^C + \sum_{C \in \text{mc}} \sum_{v \in C} t_C (1 + \|z^C\|_1)^{-1} x_v^C \\ &= \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} \|z^C\|_1 + \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} \\ &= \sum_{C \in \text{mc}} t_C, \end{aligned} \tag{4.9}$$

as claimed.

The states  $\psi_C$  in part (a) are  $\text{KMS}_{\ln \rho(A)}$  states with  $m^{\psi_C} = (1 + \|z^C\|_1)^{-1}(z^C, x^C, 0)$ , and hence (4.6) says that  $Am^{\psi_C} = \rho(A)m^{\psi_C}$ . We know from [6, Proposition 2.1(c)] that  $m^\phi$  is a probability measure with  $Am^\phi \leq \rho(A)m^\phi$ . Thus

$$\begin{aligned}
A\left(m^\phi - \sum_{C \in \text{mc}} t_C m^{\psi_C}\right) &= A m^\phi - \sum_{C \in \text{mc}} t_C A m^{\psi_C} = A m^\phi - \sum_{C \in \text{mc}} t_C \rho(A) m^{\psi_C} \\
&\leq \rho(A) m^\phi - \sum_{C \in \text{mc}} t_C \rho(A) m^{\psi_C} = \rho(A) \left(m^\phi - \sum_{C \in \text{mc}} t_C m^{\psi_C}\right).
\end{aligned} \tag{4.10}$$

For  $v \in E^0 \setminus H$ , we saw in (4.8) that

$$m_v^\phi = \phi(p_v) \geq \sum_{C \in \text{mc}} t_C (1 + \|z^C\|_1)^{-1} z_v^C = \sum_{C \in \text{mc}} t_C m_v^{\psi_C}.$$

For  $v \in K$ , say  $v \in C$ , we have from the definition of  $t_C$  that

$$m_v^\phi = t_C (1 + \|z^C\|_1)^{-1} x_v^C = t_C \psi_C(p_v) = t_C m_v^{\psi_C};$$

for  $v \in H \setminus K$ , Proposition 4.1 gives  $m_v^\phi = m_v^{\psi_C} = 0$  for all  $C$ . Thus the difference satisfies  $(m^\phi - \sum_{C \in \text{mc}} t_C m^{\psi_C})|_H = 0$ .

If  $\sum_{C \in \text{mc}} t_C = 1$ , then we have  $m^\phi = \sum_C t_C m^{\psi_C}$  because both are probability measures and  $m^\phi \geq \sum_C t_C m^{\psi_C}$ . Then, since  $\phi$  and  $\sum_C t_C \psi_C$  are  $\text{KMS}_{\ln \rho(A)}$  states of  $(\mathcal{TC}^*(E), \alpha)$  which agree on projections, [6, Proposition 2.1] implies that  $\phi = \sum_C t_C \psi_C$ .

If  $\sum_{C \in \text{mc}} t_C < 1$ , then

$$m := \left(1 - \sum_{C \in \text{mc}} t_C\right)^{-1} \left(m^\phi - \sum_{C \in \text{mc}} t_C m^{\psi_C}\right) \Big|_{E^0 \setminus H}$$

is a probability measure, and the calculation (4.10) implies that  $m$  is subinvariant for the graph  $E \setminus H$ . Since  $\rho(A_{E^0 \setminus H}) < \rho(A)$ , applying [6, Theorem 3.1] to the graph  $E \setminus H$ , with  $\beta = \ln \rho(A)$  and  $\epsilon = (1 - \rho(A)^{-1} A_{E^0 \setminus H})^{-1} m$ , gives a  $\text{KMS}_{\ln \rho(A)}$  state  $\phi_\epsilon$  on  $(\mathcal{TC}^*(E \setminus H), \alpha)$  such that  $\phi_\epsilon(p_v) = m_v$  for  $v \in E^0 \setminus H$ . Now

$$\left(1 - \sum_C t_C\right) (\phi_\epsilon \circ q_H) + \sum_C t_C \psi_C$$

is a  $\text{KMS}_{\ln \rho(A)}$  state on  $(\mathcal{TC}^*(E), \alpha)$  which agrees with  $\phi$  on vertex projections, and hence

$$\phi = \left(1 - \sum_{C \in \text{mc}} t_C\right) (\phi_\epsilon \circ q_H) + \sum_C t_C \psi_C. \quad \square$$

Since  $\rho(A_{E^0 \setminus H}) < \rho(A)$ , [6, Theorem 3.1] describes the  $\text{KMS}_{\ln \rho(A)}$  states on  $\mathcal{TC}^*(E \setminus H)$ . We write  $y^{E \setminus H}$  for the vector in  $[1, \infty)^{E^0 \setminus H}$  described in [6, Theorem 3.1(a)],  $\Delta_{\ln \rho(A)}^{E \setminus H}$  for the simplex  $\{\epsilon \mid \epsilon \cdot y^{E \setminus H} = 1\}$  in  $[0, \infty)^{E^0 \setminus H}$ , and  $\phi_\epsilon$  for the  $\text{KMS}_{\ln \rho(A)}$  state on  $\mathcal{TC}^*(E \setminus H)$  described in [6, Theorem 3.1(b)].

**COROLLARY 4.5.** *Every  $\text{KMS}_{\ln \rho(A)}$  state on  $\mathcal{TC}^*(E)$  has the form*

$$\phi_{r, \epsilon, t} := r(\phi_\epsilon \circ q_H) + (1 - r) \left( \sum_{C \in \text{mc}(E)} t_C \psi_C \right) \tag{4.11}$$

for some  $r \in [0, 1]$ ,  $\epsilon \in \Delta_{\ln \rho(A)}^{E \setminus H}$  and  $t \in S_{\text{mc}}$ . We have  $\phi_{r, \epsilon, t} = \phi_{r', \epsilon', t'}$  if and only if  $(r\epsilon, (1 - r)t) = (r'\epsilon', (1 - r')t')$ .

*Proof.* Theorem 4.3(b) shows that each  $\text{KMS}_{\ln \rho(A)}$  state has the form (4.11).

Suppose that  $(r\epsilon, (1-r)t) = (r'\epsilon', (1-r')t')$ . Then  $(1-r) \sum t_C \psi_C = (1-r') \sum t'_C \psi_C$ , and so  $\phi_{r,\epsilon,t} - \phi_{r',\epsilon',t'} = (r\phi_\epsilon - r'\phi_{\epsilon'}) \circ q_H$ . Since  $\sum t_C = \sum t'_C = 1$ , we also have  $1-r = 1-r'$  and hence  $r = r'$ . So either  $r = 0$  or  $\epsilon = \epsilon'$ , and in either case,  $r\phi_\epsilon = r'\phi_{\epsilon'}$ , giving  $\phi_{r,\epsilon,t} - \phi_{r',\epsilon',t'} = 0$ .

Now suppose that  $\phi_{r,\epsilon,t} = \phi_{r',\epsilon',t'}$ . Fix  $C \in \text{mc}(E)$  and  $v \in C$ . For  $C' \in \text{mc}(E)$ , formula (4.3) shows that  $\psi_{C'}(p_v) = \delta_{C,C'}(1 + \|z^C\|)^{-1} x_v^C$ . Since  $q_H(p_v) = 0$ ,

$$0 = \phi_{r,\epsilon,t}(p_v) - \phi_{r',\epsilon',t'}(p_v) = ((1-r)t_C - (1-r')t'_C)(1 + \|z^C\|)^{-1} x_v^C.$$

Parts (a) and (d) of [16, Theorem 1.5] imply that  $x_v^C > 0$ , and so  $(1-r)t_C = (1-r')t'_C$ . It remains to show that  $r\epsilon = r'\epsilon'$ . We have  $r = 1 - \|(1-r)t\|_1 = 1 - \|(1-r')t'\|_1 = r'$ , and so  $0 = \phi_{r,\epsilon,t} - \phi_{r',\epsilon',t'} = r(\phi_\epsilon \circ q_H - \phi_{\epsilon'} \circ q_H)$ . If  $r = 0$ , then we trivially have  $r\epsilon = r'\epsilon'$ . Suppose that  $r \neq 0$ . Then  $\phi_\epsilon \circ q_H = \phi_{\epsilon'} \circ q_H$ . Proposition 3.1 implies that  $q_H^*$  is injective, so  $\phi_\epsilon = \phi_{\epsilon'}$ ; since  $\epsilon \mapsto \phi_\epsilon$  is injective [6, Theorem 3.1(b)], we deduce that  $\epsilon = \epsilon'$ .  $\square$

### 5. The KMS simplices for a fixed inverse temperature

In this section, we consider a finite directed graph  $E$  and a real number  $\beta$ , and aim to describe the extreme points of the  $\text{KMS}_\beta$  simplices of  $\mathcal{TC}^*(E)$  and  $C^*(E)$ . The states described in Theorem 4.3 will be some of them. We generate some more candidates by applying [6, Theorem 3.1] to a graph of the form  $E \setminus H$ . We continue to use the recursive description of the saturation  $\Sigma H$  described in §3.

**PROPOSITION 5.1.** *Suppose that  $H$  is a hereditary subset of  $E^0$  and  $\beta > \ln \rho(A_{E^0 \setminus H})$ . For each  $v \in E^0 \setminus H$  the series  $\sum_{\mu \in (E \setminus H)^* v} e^{-\beta|\mu|}$  converges with sum  $y_v \geq 1$ ; let  $y$  be the vector  $(y_v)$  in  $[1, \infty)^{E^0 \setminus H}$ . Then for each  $v \in E^0 \setminus H$ , there is a  $\text{KMS}_\beta$  state  $\phi_v^H$  of  $\mathcal{TC}^*(E \setminus H)$  such that*

$$\phi_v^H(s_\mu s_v^*) = \delta_{\mu,v} e^{-\beta|\mu|} (1 - e^{-\beta} A_{E^0 \setminus H})^{-1}(s(\mu), v) y_v^{-1} \quad \text{for } \mu, v \in (E \setminus H)^*. \quad (5.1)$$

The states  $\{\phi_v^H \mid v \in E^0 \setminus H\}$  are the extremal  $\text{KMS}_\beta$  states of  $\mathcal{TC}^*(E \setminus H)$ .

*Proof.* Applying [6, Theorem 3.1(a)] to  $E \setminus H$  shows that the series defining  $y_v$  converges. We define  $\epsilon^v \in [0, \infty)^{E^0 \setminus H}$  by  $\epsilon_u^v = \delta_{u,v} y_v^{-1}$ . Then  $\epsilon^v \cdot y = 1$ , and the corresponding probability measure  $m^v = (1 - e^{-\beta} A_{E^0 \setminus H})^{-1} \epsilon^v$  in [6, Theorem 3.1(a)] has entries

$$m_w^v = (1 - e^{-\beta} A_{E^0 \setminus H})^{-1}(w, v) y_v^{-1} \quad \text{for } w \in E^0 \setminus H.$$

Thus by [6, Theorem 3.1(b)], there is a  $\text{KMS}_\beta$  state  $\phi_v^H$  of  $\mathcal{TC}^*(E \setminus H)$  satisfying (5.1). It follows from [6, Theorem 3.1(c)] that the  $\phi_v^H$  are the extreme points of the simplex of  $\text{KMS}_\beta$  states (as observed in [6, Remark 3.2]).  $\square$

**COROLLARY 5.2.** *Let  $v \in E^0$  and  $\beta > 0$ . Suppose that there is a hereditary subset  $H$  of  $E^0$  such that  $v \notin H$  and  $\ln \rho(A_{E^0 \setminus H}) < \beta$ . Then there is a  $\text{KMS}_\beta$  state  $\phi_{\beta,v}$  of  $(\mathcal{TC}^*(E), \alpha)$  such that for every pair  $\mu, v \in E^*$ , we have*

$$\phi_{\beta,v}(s_\mu s_v^*) = \begin{cases} 0 & \text{if } s(\mu)E^*v = \emptyset, \\ \delta_{\mu,v} \left( e^{-\beta|\mu|} \sum_{\lambda \in s(\mu)E^*v} e^{-\beta|\lambda|} \right) y_v^{-1} & \text{if } s(\mu)E^*v \neq \emptyset. \end{cases} \quad (5.2)$$

For every  $H$  satisfying these hypotheses, we have  $\phi_{\beta,v} = \phi_v^H \circ q_H$ .

Notice that (5.2) implies that the state  $\phi_{\beta,v}$  does not depend on the choice of the hereditary set  $H$  satisfying  $v \notin H$  and  $\ln \rho(A_{E^0 \setminus H}) < \beta$ .

*Proof.* Proposition 5.1 gives us a  $\text{KMS}_\beta$  state  $\phi_v^H$  of  $(\mathcal{TC}^*(E \setminus H), \alpha)$ . Because  $H$  is hereditary, every path  $\lambda$  in  $E^*v$  lies entirely in  $E \setminus H$ . Thus (5.1) implies that for every  $\mu, \nu \in (E \setminus H)^*$ , we have

$$\phi_v^H(s_\mu s_\nu^*) = \delta_{\mu,\nu} \left( e^{-\beta|\mu|} \sum_{\lambda \in s(\mu)(E \setminus H)^*v} e^{-\beta|\lambda|} \right) y_v^{-1} = \delta_{\mu,\nu} \left( e^{-\beta|\mu|} \sum_{\lambda \in s(\mu)E^*v} e^{-\beta|\lambda|} \right) y_v^{-1};$$

notice that (5.1) is zero if  $s(\mu)E^*v = \emptyset$ , and in that case we need to interpret the empty sum on the right-hand side as 0. For  $\mu, \nu \in E^*$  with  $s(\mu) = s(\nu) \in H$ , we have  $q_H(s_\mu s_\nu^*) = 0$ . Thus for arbitrary  $\mu, \nu \in E^*$  with  $s(\mu) = s(\nu)$ , we have

$$\phi_v^H \circ q_H(s_\mu s_\nu^*) = \begin{cases} 0 & \text{if } s(\mu) = s(\nu) \in H, \\ \delta_{\mu,\nu} \left( e^{-\beta|\mu|} \sum_{\lambda \in s(\mu)E^*v} e^{-\beta|\lambda|} \right) y_v^{-1} & \text{if } s(\mu) = s(\nu) \in E^0 \setminus H, \end{cases}$$

and  $\phi_{\beta,v} := \phi_v^H \circ q_H$  is a  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E), \alpha)$  satisfying (5.2).  $\square$

**THEOREM 5.3.** *Suppose that  $E$  is a finite directed graph and  $\beta$  is a real number, and denote by  $\alpha$  all the actions of  $\mathbb{R}$  obtained by lifting gauge actions on Toeplitz algebras and graph algebras. Let  $H_\beta$  be the hereditary closure in  $E^0$  of  $\{C \in E^0 / \sim \mid \ln \rho(A_C) > \beta\}$ .*

- (a) *If  $H_\beta = E^0$ , then  $(\mathcal{TC}^*(E), \alpha)$  has no  $\text{KMS}_\beta$  states.*
- (b) *Suppose that  $H_\beta \neq E^0$  and that  $\beta > \ln \rho(A_{E^0 \setminus H_\beta})$ . For  $v \in E^0 \setminus H_\beta$ , there is a  $\text{KMS}_\beta$  state  $\phi_{\beta,v}$  of  $(\mathcal{TC}^*(E), \alpha)$  satisfying (5.2). Then*

$$\{\phi_{\beta,v} \mid v \in E^0 \setminus H_\beta\}$$

*are the extreme points of the  $\text{KMS}_\beta$  simplex of  $(\mathcal{TC}^*(E), \alpha)$ . A  $\text{KMS}_\beta$  state factors through  $C^*(E)$  if and only if it belongs to the convex hull of*

$$\{\phi_{\beta,v} \mid v \text{ is a source in } E \setminus \Sigma H_\beta\}.$$

- (c) *Suppose that  $H_\beta \neq E^0$  and that  $\beta = \ln \rho(A_{E^0 \setminus H_\beta})$ . Let  $K_\beta$  be the hereditary closure in  $E^0$  of  $\{C \in E^0 / \sim \mid \ln \rho(A_C) \geq \beta\}$ . For  $v \in E^0 \setminus K_\beta$ , there is a  $\text{KMS}_\beta$  state  $\phi_{\beta,v}$  of  $(\mathcal{TC}^*(E), \alpha)$  satisfying (5.2). For  $C \in \text{mc}(E \setminus H_\beta)$ , let  $\psi_C^{H_\beta}$  be the  $\text{KMS}_\beta$  state of  $(\mathcal{TC}^*(E \setminus H_\beta), \alpha)$  obtained by applying Theorem 4.3(a) to the graph  $E \setminus H_\beta$ . Then the states*

$$\{\psi_C := \psi_C^{H_\beta} \circ q_{H_\beta} \mid C \in \text{mc}(E \setminus H_\beta)\} \cup \{\phi_{\beta,v} \mid v \in E^0 \setminus K_\beta\} \quad (5.3)$$

*are the extreme points of the  $\text{KMS}_\beta$  simplex of  $(\mathcal{TC}^*(E), \alpha)$ . A  $\text{KMS}_\beta$  state factors through  $C^*(E)$  if and only if it belongs to the convex hull of*

$$\{\psi_C \mid C \in \text{mc}(E \setminus H_\beta)\} \cup \{\phi_{\beta,v} \mid v \text{ is a source in } E \setminus \Sigma K_\beta\}. \quad (5.4)$$

Both  $H_\beta$  and  $K_\beta$  are hereditary subsets of  $E^0$ , and  $H_\beta \subset K_\beta$ . Obviously the proof of the theorem must exploit the specific nature of these two sets, but some of our arguments are more general, and we separate out some lemmas. Throughout this section,  $E$  is a finite directed graph.



LEMMA 5.4. *Suppose that  $I$  is an ideal in a  $C^*$ -algebra  $A$ , that  $\phi_1, \dots, \phi_n$  are states of  $A$ , and that  $\lambda_i \in (0, \infty)$  for  $1 \leq i \leq n$ . Then  $\sum_{j=1}^n \lambda_j \phi_j$  factors through  $A/I$  if and only if  $\phi_i$  factors through  $A/I$  for all  $i$ .*

*Proof.* If each  $\phi_i$  factors through  $A/I$ , then so does every linear combination. So suppose that  $\sum_j \lambda_j \phi_j$  factors through  $C^*(E)$ . For a positive element  $a$  in  $I$  and each  $i$ , we have

$$0 = \sum_{j=1}^n \lambda_j \phi_j(a) \geq \lambda_i \phi_i(a) \geq 0,$$

and since  $\lambda_i > 0$ , this forces  $\phi_i(a) = 0$ . Since  $I$  is spanned by its positive elements, we deduce that  $\phi_i$  vanishes on  $I$ , and hence  $\phi_i$  factors through  $A/I$ .  $\square$

LEMMA 5.5. *Suppose that  $H \subset E^0$  is hereditary and that  $\phi$  is a  $\text{KMS}_\beta$ -state of  $\mathcal{TC}^*(E \setminus H)$  which factors through  $C^*(E \setminus H)$ . If  $\phi(p_v) = 0$  for all  $v \in \Sigma H \setminus H$ , then the state  $\phi \circ q_H$  of  $\mathcal{TC}^*(E)$  factors through  $C^*(E)$ .*

*Proof.* The hypothesis says that there is a state  $\bar{\phi}$  of  $C^*(E \setminus H)$  such that  $\phi = \bar{\phi} \circ \pi_{E \setminus H}$ . Let  $J$  be the ideal of  $C^*(E \setminus H)$  generated by  $\{p_v \mid v \in \Sigma H \setminus H\}$ . Then [6, Lemma 2.2] implies that  $\bar{\phi}$  factors through  $C^*(E \setminus H)/J$ . Theorem 4.1(b) of [1] implies that there is an isomorphism of  $C^*(E \setminus \Sigma H)$  onto  $C^*(E \setminus H)/J$  which takes  $\bar{s}_e$  to  $s_e + J$ . So there is a  $\text{KMS}_\beta$  state  $\bar{\bar{\phi}}$  of  $C^*(E \setminus \Sigma H)$  such that  $\phi = \bar{\bar{\phi}} \circ \bar{q}_{\Sigma H \setminus H} \circ \pi_{E \setminus H}$ . By considering the images of generators of  $\mathcal{TC}^*(E)$ , one checks that the diagram

$$\begin{array}{ccccc} C^*(E \setminus H) & \xleftarrow{\pi_{E \setminus H}} & \mathcal{TC}^*(E \setminus H) & \xleftarrow{q_H} & \mathcal{TC}^*(E) \\ \bar{q}_{\Sigma H \setminus H} \downarrow & & & & \downarrow \pi_E \\ C^*(E \setminus \Sigma H) & \xleftarrow{\bar{q}_{\Sigma H}} & & & C^*(E) \end{array} \quad (5.5)$$

commutes. Thus  $\phi \circ q_H$  factors through the state  $\bar{\bar{\phi}} \circ \bar{q}_{\Sigma H}$  of  $C^*(E)$ .  $\square$

LEMMA 5.6. *Suppose that  $E$  is a finite directed graph with vertex matrix  $A$ , and that  $\beta > \ln \rho(A)$ . Suppose that  $G$  is a hereditary subset of  $E^0$ , and let  $y^E \in [1, \infty)^{E^0}$  and  $y^{E \setminus G}$  be the vectors of [6, Theorem 3.1] for the graphs  $E$  and  $E \setminus G$ . If  $\epsilon \in [0, 1]^{E^0}$  satisfies  $\epsilon \cdot y = 1$  and  $\epsilon|_G = 0$ , then  $\epsilon|_{E \setminus G}$  satisfies  $(\epsilon|_{E \setminus G}) \cdot y^{E \setminus G} = 1$ , and the corresponding  $\text{KMS}_\beta$  states on the Toeplitz algebras satisfy  $\phi_\epsilon = \phi_{\epsilon|_{E \setminus G}} \circ q_G$ .*

*Proof.* For  $w \in E^0 \setminus G$ , we have  $(E \setminus G)^* w = E^* w$ , and hence

$$y_w^{E \setminus G} = \sum_{\mu \in (E \setminus G)^* w} e^{-\beta|\mu|} = \sum_{\mu \in E^* w} e^{-\beta|\mu|} = y_w^E.$$

Thus  $y^{E \setminus G} = y|_{E \setminus G}$ , and  $1 = \epsilon \cdot y^E = (\epsilon|_{E \setminus G}) \cdot (y|_{E \setminus G}) = (\epsilon|_{E \setminus G}) \cdot y^{E \setminus G}$ .

Since  $G$  is hereditary, for  $v \in E^0$  we have

$$m_v = ((1 - e^{-\beta} A)^{-1} \epsilon)_v = \begin{cases} ((1 - e^{-\beta} A_{E^0 \setminus G})^{-1} \epsilon|_{E \setminus G})_v & \text{if } v \in E^0 \setminus G, \\ 0 & \text{if } v \in G, \end{cases}$$

and hence

$$\phi_\epsilon(p_v^E) = \begin{cases} \phi_{\epsilon|_{E \setminus G}}(p_v^{E \setminus G}) & \text{if } v \in E^0 \setminus G, \\ 0 & \text{if } v \in G. \end{cases}$$

Thus  $\phi_\epsilon$  and  $\phi_{\epsilon|_{E \setminus G}} \circ q_G$  agree on the vertex projections  $\{p_v\}$  in  $\mathcal{TC}^*(E)$ , and since both are  $\text{KMS}_\beta$  states, [6, Proposition 2.1(a)] implies that they are equal.  $\square$

**LEMMA 5.7.** *Suppose that  $H$  is a hereditary subset of  $E$  and  $\beta > \ln \rho(A_{E^0 \setminus H})$ . Let  $v \in E^0 \setminus H$ , and let  $\phi_v^H$  be the state of  $\mathcal{TC}^*(E \setminus H)$  described in Proposition 5.1. Then  $\phi_v^H \circ q_H$  factors through  $C^*(E)$  if and only if  $v$  is a source in  $E \setminus \Sigma H$ .*

*Proof.* Suppose that  $v$  is a source in  $E \setminus \Sigma H$ . Then  $v$  must be a source in  $E$ : otherwise, we have  $s(r^{-1}(v)) \subset \Sigma H$ , and saturation implies that  $v \in \Sigma H$ . In particular,  $v$  is a source in  $E \setminus H$ , and [6, Corollary 6.1(a)] implies that  $\phi_v^H$  factors through  $C^*(E \setminus H)$ . With a view to applying Lemma 5.5, we take  $w \in \Sigma H \setminus H$ . Since  $\Sigma H$  is hereditary,  $wE^n v = \emptyset$  for all  $n$ , and (4.4) implies that  $(1 - e^{-\beta} A_{E^0 \setminus H})^{-1}(w, v) = 0$ . Thus

$$\phi_v^H(p_w) = (1 - e^{-\beta} A_{E^0 \setminus H})^{-1}(w, v) y_v^{-1} = 0,$$

and Lemma 5.5 implies that  $\phi_v^H \circ q_H$  factors through  $C^*(E)$ .

Now suppose that  $\phi_v^H \circ q_H$  factors through  $C^*(E)$ . Since  $\phi_v^H \circ q_H(p_w) = 0$  for  $w \in H$ , Lemma 3.4(b) implies that  $\phi_v^H \circ q_H$  vanishes on  $\{p_w \mid w \in \Sigma H\}$ . Thus it follows from [6, Lemma 2.2] that  $\phi_v^H \circ q_H$  factors through  $C^*(E \setminus \Sigma H)$ . Since  $\phi_v^H(p_v) \neq 0$ , we deduce that  $v \in E^0 \setminus \Sigma H$ . Thus  $\epsilon_w^v = (y_v^{E \setminus H})^{-1} \delta_{v,w}$  vanishes for  $w$  in the hereditary set  $\Sigma H$ , and Lemma 5.6 implies that  $\phi_v^H \circ q_H = \phi_{\epsilon^v|_{E \setminus \Sigma H}} \circ q_{\Sigma H \setminus H}$ . Since  $\phi_v^H \circ q_H$  factors through  $C^*(E)$ , for  $w \in E^0 \setminus \Sigma H$  we have

$$\begin{aligned} \phi_{\epsilon^v|_{E \setminus \Sigma H}} \left( p_w - \sum_{e \in w(E \setminus \Sigma H)^1} s_e s_e^* \right) &= \phi_{\epsilon^v|_{E \setminus \Sigma H}} \circ q_{\Sigma H \setminus H} \left( p_w - \sum_{e \in w(E \setminus H)^1} s_e s_e^* \right) \\ &= \phi_v^H \circ q_H \left( p_w - \sum_{e \in w E^1} s_e s_e^* \right) = 0. \end{aligned}$$

Applying [6, Lemma 2.2] to  $E \setminus \Sigma H$  shows that  $\phi_{\epsilon^v|_{E \setminus \Sigma H}}$  factors through  $C^*(E \setminus \Sigma H)$ . We have  $\beta > \rho(A_{E^0 \setminus H})$ , so [6, Corollary 6.1(a)] implies that  $\epsilon^v|_{E \setminus \Sigma H}$  is supported on the sources of  $E \setminus \Sigma H$ , and hence  $v$  is a source in  $E \setminus \Sigma H$ .  $\square$

*Proof of Theorem 5.3.* (a) We suppose that  $\mathcal{TC}^*(E)$  has a  $\text{KMS}_\beta$  state  $\phi$ , and prove that  $H_\beta \neq E^0$ . The set  $\bigcup \{C \in E^0 / \sim \mid \ln \rho(A_C) > \beta\}$  generates  $H_\beta$  as a hereditary set, and so Proposition 3.1 implies that  $\phi(p_w) = 0$  for all  $w \in H_\beta$ . Hence  $1 = \phi(1) = \sum_{v \notin H_\beta} \phi(p_v)$ , and  $H_\beta$  cannot be all of  $E^0$ .

(b) Applying Corollary 5.2 with  $H = H_\beta$  gives the existence of the state  $\phi_{\beta,v}$ , and the last comment in Corollary 5.2 implies that  $\phi_{\beta,v} = \phi_v^{H_\beta} \circ q_{H_\beta}$ . We can apply Proposition 5.1 with  $H = H_\beta$ , and deduce that the states  $\phi_v^{H_\beta}$  for  $v \notin H_\beta$  are the extreme points of the  $\text{KMS}_\beta$  simplex of  $\mathcal{TC}^*(E \setminus H_\beta)$ . Since  $H_\beta$  is not all of  $E^0$ , the final statement of Proposition 3.1 implies that  $q_{H_\beta}^*$  is an isomorphism of the  $\text{KMS}_\beta$  simplex

of  $\mathcal{TC}^*(E \setminus H_\beta)$  onto that of  $\mathcal{TC}^*(E)$ . Hence the states  $\phi_{\beta,v} = \phi_v^{H_\beta} \circ q_{H_\beta}$  are the extreme points of the  $\text{KMS}_\beta$  simplex of  $\mathcal{TC}^*(E)$ .

Lemma 5.7 implies that  $\phi_{\beta,v} = \phi_v^{H_\beta} \circ q_{H_\beta}$  factors through  $C^*(E)$  if and only if  $v$  is a source in  $E \setminus \Sigma H_\beta$ . So Lemma 5.4 implies that a  $\text{KMS}_\beta$  state  $\phi$  factors through  $C^*(E)$  if and only if it belongs to the convex hull of  $\{\phi_{\beta,v} \mid v \text{ is a source in } E \setminus \Sigma H_\beta\}$ .

(c) We can apply Corollary 5.2 with  $H = K_\beta$  to get the state  $\phi_{\beta,v} = \phi^{K_\beta} \circ q_{K_\beta}$ . As in (b),  $q_{H_\beta}^*$  is an isomorphism of the  $\text{KMS}_\beta$  simplex of  $\mathcal{TC}^*(E \setminus H_\beta)$  onto that of  $\mathcal{TC}^*(E)$ . Since  $\beta = \ln \rho(A_{E^0 \setminus H_\beta})$  is real,  $\rho(A_{E^0 \setminus H_\beta})$  cannot be 0, and [6, Lemma A.1(b)] implies that  $E \setminus H_\beta$  has at least one cycle. The set  $K_\beta \setminus H_\beta$  is generated as a hereditary subset of  $E^0 \setminus H_\beta$  by the minimal critical components of  $E \setminus H_\beta$ , and hence is the set  $H$  in Theorem 4.3 for the graph  $E \setminus H_\beta$ . Thus Corollary 4.5 implies that the  $\text{KMS}_\beta$  states of  $\mathcal{TC}^*(E \setminus H_\beta)$  have the form  $\phi_{r,\epsilon,t}$ , and that the extreme points are the ones of the form  $\phi_{1,\epsilon^v,t} = \phi_v^{K_\beta \setminus H_\beta} \circ q_{K_\beta \setminus H_\beta} = \phi_{\beta,v}$  or  $\phi_{0,\epsilon,\delta_C} = \psi_C^{H_\beta}$ . Proposition 2.1 implies that  $q_{K_\beta \setminus H_\beta} \circ q_{H_\beta} = q_{K_\beta}$ . Thus the  $\text{KMS}_\beta$  simplex of  $\mathcal{TC}^*(E)$  is the convex hull of the set (5.3).

It remains to show that a convex combination of the states (5.3) factors through  $C^*(E)$  if and only if it belongs to the convex hull of the set (5.4). Lemma 5.7 implies that  $\phi_v^{K_\beta} \circ q_{K_\beta}$  factors through  $C^*(E)$  if and only if  $v$  is a source in  $E \setminus \Sigma K_\beta$ . We claim that the  $\psi_C^{H_\beta} \circ q_{H_\beta}$  all factor through  $C^*(E)$ . To see this, fix  $C \in \text{mc}(E \setminus H_\beta)$ . Theorem 4.3(a) implies that  $\psi_C^{H_\beta}$  factors through  $C^*(E \setminus H_\beta)$ . We have  $vE^n C \neq \emptyset$  for all  $v \in C$  and  $n \in \mathbb{N}$  because  $C$  is a non-trivial connected component. Since  $C \cap H_\beta = \emptyset$ , we deduce that  $C$  does not intersect any of the sets  $S_k H_\beta$  of (3.4), and hence  $C \cap \Sigma H_\beta = \emptyset$ . Then because  $\Sigma H_\beta$  is hereditary, we have  $wE^* C = \emptyset$  for all  $w \in \Sigma H_\beta$ . Hence (4.5) implies that  $z_w^C = 0$  for all  $w \in \Sigma H_\beta \setminus H_\beta$ , and so (4.3) implies that  $\psi_C^{H_\beta}(p_w) = 0$  for all  $w \in \Sigma H_\beta \setminus H_\beta$ . Now Lemma 5.5 implies that  $\psi_C = \psi_C^{H_\beta} \circ q_{H_\beta}$  factors through  $C^*(E)$ .  $\square$

Theorem 5.3 describes the  $\text{KMS}_\beta$  simplex for each fixed  $\beta$ . However, it also makes sense to fix a vertex  $v$ , and ask for which  $\beta$  there is a state  $\phi_{\beta,v}$  of  $(\mathcal{TC}^*(E), \alpha)$  as in Corollary 5.2.

**COROLLARY 5.8.** *Suppose that  $E$  is a finite directed graph and  $v \in E^0$ . Define*

$$\beta_v := \max\{\ln \rho(A_C) \mid C \leq v\}.$$


*Then there is a state  $\phi_{\beta,v}$  satisfying (5.2) if and only if  $\beta > \beta_v$ .*

*Proof.* First suppose that there exists such a state  $\phi_{\beta,v}$ . Then there is a hereditary set  $H$  such that  $v \notin H$  and  $\ln \rho(A_{E^0 \setminus H}) < \beta$ . But then any  $C$  with  $C \leq v$  lies in  $E^0 \setminus H$ , and  $\ln \rho(A_C) \leq \ln \rho(A_{E^0 \setminus H}) < \beta$ . Thus  $\beta_v = \max\{\ln \rho(A_C) \mid C \leq v\} < \beta$ . Conversely, suppose that  $\beta > \beta_v$ . Then the hereditary closure  $K_\beta$  of  $\{C \mid \ln \rho(A_C) \geq \beta\}$  does not contain  $v$ : for if so, then there exists  $C \in \{C \mid \ln \rho(A_C) \geq \beta\}$  with  $C \leq v$ , and we have  $\beta_v \geq \beta$ . Thus we can apply Corollary 5.2 with  $H = K_\beta$  to deduce the existence of  $\phi_{\beta,v}$ .  $\square$

## 6. Examples

We give some examples to show how we can use Theorem 5.3 to compute all the KMS states on  $\mathcal{TC}^*(E)$  and  $C^*(E)$ . Since we want to focus on how the different components of  $E$  interact, we consider graphs in which the components are small.


*Example 6.1.* The following graph  $E$



has two strongly connected components  $\{v\}$  and  $\{w\}$ . Both are non-trivial components, with  $A_{\{v\}} = (2)$ ,  $A_{\{w\}} = (3)$  and  $\rho(A) = 3$ .

- For  $\beta > \ln \rho(A) = \ln 3$ , the set  $H_\beta$  of Theorem 5.3 is empty, and Theorem 5.3(b) gives a one-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\beta,v}$  and  $\phi_{\beta,w}$ . None of these factor through  $C^*(E)$ .
- At  $\beta = \ln 3$ ,  $H_\beta$  is still empty, but  $K_{\ln 3}$  is the hereditary closure of  $\{w\}$ , which is all of  $E^0$ . The only critical component is  $\{w\}$ , and hence Theorem 5.3(c) gives a unique  $\text{KMS}_{\ln 3}$  state  $\psi_{\{w\}}$  which factors through  $C^*(E)$ .
- For  $\beta < \ln 3$ ,  $H_\beta = E^0$ , and  $(\mathcal{TC}^*(E), \alpha)$  has no  $\text{KMS}_\beta$  states.

*Example 6.2.* Reversing the horizontal arrow in the previous example makes a big difference. The graph  $E$  is now



The strongly connected components are still  $\{v\}$  and  $\{w\}$ , but now the minimal critical component  $\{w\}$  is hereditary.

- For  $\beta > \ln 3 = \ln \rho(A)$ ,  $H_\beta = \emptyset$ , and Theorem 5.3(b) gives a one-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\beta,v}$  and  $\phi_{\beta,w}$ . None of these factor through  $C^*(E)$ .
- For  $\beta = \ln 3$ , we have  $H_\beta = \emptyset$  and  $K_\beta = \{w\}$ . Theorem 5.3(c) gives a one-dimensional simplex of  $\text{KMS}_{\ln 3}$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\ln 3,v}$  and  $\psi_{\{w\}}$ , and only  $\psi_{\{w\}}$  factors through  $C^*(E)$ . (We work out a formula for  $\psi_{\{w\}}$  at the end of this example.)
- For  $\ln 2 < \beta < \ln 3$ ,  $H_\beta = \{w\}$ , and Theorem 5.3(b) gives a single  $\text{KMS}_\beta$  state  $\phi_{\beta,v}$  on  $(\mathcal{TC}^*(E), \alpha)$ , which does not factor through  $C^*(E)$ .
- For  $\beta = \ln 2$ ,  $H_\beta = \{w\}$  and  $K_\beta = \{v, w\} = E^0$ . The graph  $E \setminus H_\beta$  has a single critical component  $\{v\}$ , and Theorem 5.3(c) gives a unique  $\text{KMS}_{\ln 2}$  state  $\psi_{\{v\}}$  on  $(\mathcal{TC}^*(E), \alpha)$ . This state factors through  $C^*(E)$ .
- For  $\beta < \ln 2$ , there are no  $\text{KMS}_\beta$  states.

We can make the construction of these states quite explicit. We illustrate by working through the construction of the  $\text{KMS}_{\ln 3}$  state  $\psi_{\{w\}}$ . The unimodular Perron–Frobenius


eigenvector for the matrix  $A_{\{w\}} = (3)$  is the scalar  $x_w^{\{w\}} = 1$ , and the vector  $z^{\{w\}}$  in (4.2) is the scalar

$$z_v^{\{w\}} = \rho(A)^{-1}(1 - \rho(A)^{-1}A_{\{v\}})^{-1}A(v, w)x_w^{\{w\}} = 3^{-1}(1 - 3^{-1} \cdot 2)^{-1}1 \cdot 1 = 3^{-1} \cdot 3 = 1.$$

Thus  $\|1 + z^{\{w\}}\|_1 = 2$ ,  $\psi_{\{w\}}(p_v) = \psi_{\{w\}}(p_w) = 2^{-1}$ , and

$$\psi_{\{w\}}(s_\mu s_\nu^*) = \delta_{\mu, \nu} 3^{-|\mu|} 2^{-1} \quad \text{for } \mu, \nu \in E^*.$$

*Example 6.3.* We now replace the component  $\{w\}$  with a two-vertex component whose critical inverse temperature still exceeds that of the component  $\{v\}$ . This gives an example in which the  $\text{KMS}_\beta$  simplex changes dimension both as  $\beta$  decreases to  $\ln \rho(A)$ , and as  $\beta$  passes through  $\ln \rho(A)$ .



The strongly connected components are  $\{v\}$  and  $\{w, u\}$ . The block corresponding to the latter is  $A_{\{w, u\}} = \begin{pmatrix} 2 & 2 \\ 2 & 0 \end{pmatrix}$ , which has spectral radius  $\rho(A_{\{w, u\}}) = \gamma := 1 + \sqrt{5}$ . Since  $\rho(A) = \max\{\rho(A_{\{v\}}), \rho(A_{\{w, u\}})\} = \gamma$ ,  $\{w, u\}$  is a minimal critical component.

- For  $\beta > \ln \gamma$ ,  $H_\beta = \emptyset$ , and Theorem 5.3(b) gives a two-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\beta, v}$ ,  $\phi_{\beta, w}$  and  $\phi_{\beta, u}$ . None of these factor through  $C^*(E)$ .
- For  $\beta = \ln \gamma$ , we have  $H_\beta = \emptyset$  and  $K_\beta = \{w, u\}$ . Theorem 5.3(c) gives a one-dimensional simplex of  $\text{KMS}_{\ln \gamma}$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\ln \gamma, v}$  and  $\psi_{\{w, u\}}$ . Only  $\psi_{\{w, u\}}$  factors through  $C^*(E)$ .
- For  $0 < \beta < \ln \gamma$ , we have  $\{w, u\} \subseteq H_\beta$ , and so the  $\text{KMS}_\beta$  simplex is similar to that of Example 6.2. In particular, the dimension of the  $\text{KMS}_\beta$  simplex drops again to 0 as  $\beta$  drops below  $\ln \gamma$ , and the simplex disappears altogether for  $\beta < \ln 2$ .

*Example 6.4.* In the next graph  $E$ , we have added two trivial components, and now the subtleties involving saturations in Theorem 5.3 come into play.



- For  $\beta > \ln 3$ , we have  $H_\beta = \emptyset$ , and Theorem 5.3(b) gives us a three-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\beta, v}$ ,  $\phi_{\beta, w}$ ,  $\phi_{\beta, u_1}$  and  $\phi_{\beta, u_2}$ . The state  $\phi_{\beta, u_1}$  factors through  $C^*(E)$ .
- At  $\beta = \ln 3$ , we have  $H_\beta = \emptyset$  and  $K_\beta = \{w\}$ . Theorem 5.3(c) gives us a three-dimensional simplex of  $\text{KMS}_{\ln 3}$  states, with extreme points  $\phi_{\ln 3, v}$ ,  $\phi_{\ln 3, u_1}$  and  $\phi_{\ln 3, u_2}$  alongside the state  $\psi_{\{w\}}$  associated to the critical component  $\{w\}$  in  $K_\beta$ . Now  $\Sigma K_\beta = \{u_2, w\}$ , and the vertex  $u_1$  is a source in  $E \setminus \Sigma K_\beta$ . Thus both  $\psi_{\{w\}}$  and  $\phi_{\ln 3, u_1}$  factor through  $\text{KMS}_{\ln 3}$  states of  $(C^*(E), \alpha)$ .


- For  $\ln 2 < \beta < \ln 3$ , we have  $H_\beta = \{w\}$ , and Theorem 5.3(b) gives us a two-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\phi_{\beta,v}$ ,  $\phi_{\beta,u_1}$  and  $\phi_{\beta,u_2}$ . Since  $\Sigma H_\beta = \{u_2, w\}$ , only the state  $\phi_{\beta,u_1}$  factors through  $C^*(E)$ .
- For  $\beta = \ln 2$ , we have  $H_\beta = \{w\}$  and  $K_\beta = E^0$ . The only critical component in  $E \setminus H_\beta$  is  $\{v\}$ , and hence Theorem 5.3(c) implies that  $(\mathcal{TC}^*(E), \alpha)$  has a unique  $\text{KMS}_{\ln 2}$  state  $\psi_{\{v\}}$ , and that this state factors through  $C^*(E)$ .
- For  $\beta < \ln 2$ , the hereditary closure of  $H_\beta = \{v, w\}$  is all of  $E^0$ , and  $(\mathcal{TC}^*(E), \alpha)$  has no  $\text{KMS}_\beta$  states.

*Example 6.5.* Our next graph  $E$  is the one from Example 6.4 with the edge between  $u_1$  and  $v$  reversed.




- For  $\beta > \ln 3$  and  $\beta = \ln 3$ , we still have a three-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$ . However, for this graph  $u_1$  is not a source in  $E \setminus \Sigma K_{\ln 3} = E^0 \setminus \{u_2, w\}$ , and only the  $\text{KMS}_{\ln 3}$  state  $\psi_{\{w\}}$  factors through  $C^*(E)$ .
- For  $\ln 2 < \beta < \ln 3$ , we still have  $H_\beta = \{w\}$  and a two-dimensional simplex of  $\text{KMS}_\beta$  states. For this graph, none of these  $\text{KMS}_\beta$  states factors through  $C^*(E)$ .
- At  $\beta = \ln 2$ ,  $K_\beta = \{v, u_2, w\}$ , and we have a one-dimensional simplex of  $\text{KMS}_{\ln 2}$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\psi_{\{u_2, w\}}$  and  $\phi_{\ln 2, u_1}$ . The state  $\psi_{\{u_2, w\}}$  factors through  $C^*(E)$ .
- For  $\beta < \ln 2$ , we have  $H_\beta = \{v, u_2, w\}$ , and a single  $\text{KMS}_\beta$  state  $\phi_{\beta, u_1}$  on  $(\mathcal{TC}^*(E), \alpha)$ . Since  $\Sigma H_\beta$  is all of  $E^0$ , this state does not factor through  $C^*(E)$ .

*Example 6.6.* We now add a source  $u_3$  to the graph of Example 6.5.



The vertex  $u_3$  belongs to the complement of  $H_\beta$  and of  $K_\beta$  for all  $\beta$ . So at every  $\beta$ , the new vertex  $u_3$  gives an extreme point  $\phi_{\beta, u_3}$  of the  $\text{KMS}_\beta$  simplex of  $\mathcal{TC}^*(E)$ , and this state factors through  $C^*(E)$ .

*Example 6.7.* The following graph  $E$



has three components  $C$  with  $\rho(A) = \rho(AC)$ , but only  $\{v\}$  and  $\{w\}$  are minimal.

- For  $\beta > \ln 2$ , we have a three-dimensional simplex of  $\text{KMS}_\beta$  states on  $(\mathcal{TC}^*(E), \alpha)$ , and none of them factor through  $C^*(E)$ .
- At  $\beta = \ln 2$ , we have a two-dimensional simplex of  $\text{KMS}_{\ln 2}$  states on  $(\mathcal{TC}^*(E), \alpha)$  with extreme points  $\psi_{\{v\}}$ ,  $\psi_{\{w\}}$  and  $\phi_{\beta,u}$ . Of these, only  $\psi_{\{v\}}$  and  $\psi_{\{w\}}$  factor through  $C^*(E)$ .
- For  $0 \leq \beta < \ln 2$ , there is a unique  $\text{KMS}_\beta$  state on  $\mathcal{TC}^*(E)$ , which only factors through  $C^*(E)$  when  $\beta = 0$ . This  $\text{KMS}_0$  state is the invariant trace on  $C^*(E)$  that is obtained by lifting the trace on  $C^*(E \setminus \Sigma H) \cong C(\mathbb{T})$  given by integration against Haar measure on  $\mathbb{T}$ .

## 7. Concluding remarks

7.1. *Critical inverse temperatures.* We say that  $\beta$  is a *critical inverse temperature* if  $H_\beta \neq E^0$  and  $\beta = \ln \rho(A_{E^0 \setminus H_\beta})$ . Theorem 5.3(c) says that these are precisely the inverse temperatures at which we have states of the form  $\psi_C$ , and that these states factor through  $C^*(E)$ ; for all but the smallest critical  $\beta$ , we also have states of the form  $\phi_{\beta,v}$ .

Every critical inverse temperature  $\beta$  has the form  $\ln \rho(A_C)$  for some component  $C$ , but as our examples show, not every  $\ln \rho(A_C)$  need be critical (for example,  $\beta = \ln 2$  in Example 6.1). So to find the critical  $\beta$  for a given finite graph  $E$ , we compute the numbers  $\beta = \ln \rho(A_C)$ , identify the sets  $H_\beta$  by looking at the graph, and discard the numbers which are not critical. The set of critical inverse temperatures is always finite (with cardinality bounded by  $|E^0|$ ), but could in general be arbitrarily large.

Since there are finitely many critical values, we can list them in increasing order. Then for  $\beta$  between two consecutive critical values, say  $\beta \in (\beta_C, \beta_D)$ , Theorem 5.3(b) gives a simplex of  $\text{KMS}_\beta$  states with extreme points  $\{\phi_{\beta,v} \mid v \in E^0 \setminus H_\beta\}$ .

For the Toeplitz algebra, the range of possible inverse temperatures  $\beta$  is either  $\mathbb{R}$  (if  $E$  has a source which does not talk to any non-trivial component  $C$ ) or  $[\beta_l, \infty)$ , where  $\beta_l$  is the smallest critical inverse temperature. But for the graph algebra  $C^*(E)$  there are interesting number-theoretic restrictions on the possible values of critical  $\beta$  and thus on the range of possible inverse temperatures. We use results of Lind [12], and refer to the treatment in [13, §11.1].

Suppose that  $E$  is a directed graph without sources and suppose that  $C^*(E)$  has a  $\text{KMS}_\beta$  state. Then  $\beta = \ln \rho(A_C)$  for some strongly connected component  $C$  of  $E$ . Since  $\rho(A_C)$  is the Perron–Frobenius eigenvalue of  $A_C$ , it is a root of the characteristic polynomial  $\det(xI - A_C)$ , which is a monic polynomial of degree  $n$  with integer coefficients. Thus  $\rho(A_C)$  is an algebraic integer. For each algebraic integer  $\lambda$  there is a unique *minimal polynomial*  $q_\lambda(x) \in \mathbb{Q}[x]$  that is monic, irreducible and has  $q_\lambda(\lambda) = 0$  [8, Proposition 6.1.7]; the other roots of this polynomial are called the *conjugates* of  $\lambda$ . A *Perron number* is an algebraic integer  $\lambda \geq 1$  that is strictly larger than the absolute value of all its other conjugates.

**PROPOSITION 7.1.** *Suppose that  $\beta > 0$ . Then  $e^{p\beta}$  is a Perron number for some  $p \in \mathbb{N}$  if and only if there exists a graph  $E$  without sources such that the gauge dynamics on  $C^*(E)$  has a  $\text{KMS}_\beta$  state.*

*Proof.* Let  $E$  be a graph such that  $C^*(E)$  has a  $\text{KMS}_\beta$  state, and choose a component  $C$  such that  $\beta = \ln \rho(A_C)$ , as above. Let  $p$  be the period of the irreducible matrix  $A_C$ . Then  $(e^\beta)^p = e^{p\beta}$  is a Perron number by the implication (1)  $\implies$  (3) of [13, Theorem 11.1.5]. Conversely, if  $e^{p\beta}$  is a Perron number for some  $p \in \mathbb{N}$ , the implication (3)  $\implies$  (1) of the same theorem gives the existence of a non-negative integer matrix  $A$  with spectral radius  $e^\beta$ . Thus for the graph  $E$  with vertex matrix  $A$ ,  $C^*(E)$  has a  $\text{KMS}_\beta$  state.  $\square$

It is easy to produce Perron numbers, and also algebraic integers  $\lambda \geq 1$  that are not Perron numbers. For example,  $(5 - \sqrt{5})/2$  is an algebraic integer with minimal polynomial  $x^2 - 5x + 5$ , and hence the conjugates are  $(5 \pm \sqrt{5})/2$ . Thus Proposition 7.1 implies that there is no graph without sources such that  $C^*(E)$  has a KMS state with inverse temperature  $\ln((5 - \sqrt{5})/2)$ . Note that  $x^2 - 5x + 5$  is the characteristic polynomial of

$$A = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix},$$

which is the vertex matrix of a graph with two vertices.

**7.2. Connections with the results of Carlsen and Larsen.** In their recent preprint [2], Carlsen and Larsen study the KMS states of generalized gauge dynamics on the relative graph algebras of possibly infinite graphs using the partial action techniques developed by Exel and Laca in [4]. Their results apply in particular to finite graphs<sup>†</sup>, where taking their function  $N : E^1 \rightarrow \mathbb{R}$  to be identically  $e$  gives the action  $\alpha : \mathbb{R} \rightarrow \text{Aut } \mathcal{TC}^*(E)$  studied here.

To make the connection, we observe that the sum  $y_v := \sum_{\mu \in E^*v} e^{-\beta|\mu|}$  in [6, Theorem 3.1] is the same as that defining the ‘fixed-target partition function’  $Z_v(\beta)$  in [2, equation (5.8)] (see also [4, Definition 9.3]). Our results allow us to identify the intervals of convergence of these partition functions:

**LEMMA 7.2.** *Let  $E$  be a finite graph. For  $v \in E^0$ , take  $\beta_v = \max\{\ln \rho(A_C) \mid C \leq v\}$  as in Corollary 5.8. Then the series  $\sum_{\mu \in E^*v} e^{-\beta|\mu|}$  converges if and only if  $\beta > \beta_v$ .*

*Proof.* If  $\beta > \beta_v$ , then taking  $H = H_{\beta_v}$  in Proposition 5.1 shows that the series converges. On the other hand, suppose  $\beta \leq \beta_v$ , and let  $C$  be a component such that  $C \leq v$  and  $\beta \leq \ln \rho(A_C)$ . Choose a path  $\lambda$  in  $CE^*v$ . Then

$$\sum_{\mu \in E^*v} e^{-\beta|\mu|} \geq e^{-\beta|\lambda|} \sum_{\mu' \in CE^*r(\lambda)} e^{-\beta|\mu'|},$$

and the series  $\sum_{\mu' \in CE^*r(\lambda)} e^{-\beta|\mu'|}$  diverges because  $\rho(e^{-\beta}A_C) \geq 1$  and  $A_C$  is irreducible. Thus  $\sum_{\mu \in E^*v} e^{-\beta|\mu|}$  diverges too.  $\square$

The factors  $\delta_{\mu,v}$  in our formulas for the values of KMS states show that all the KMS states on  $\mathcal{TC}^*(E)$  and  $C^*(E)$  factor through the expectation onto the diagonal  $D := \overline{\text{span}}\{s_\lambda s_\lambda^* \mid \lambda \in E^*\}$ . The restriction to  $D$  is then given by a measure  $\nu$  on the

<sup>†</sup> Though in [2] they use the non-functorial convention for paths in directed graphs, so strictly speaking one would have to apply their results to the opposite graph  $E^{\text{opp}} = (E^0, E^1, s, r)$ .



spectrum of  $D$ , which is  $E^* \cup E^\infty$ . Exel and Laca say that a KMS state  $\psi$  is of *finite type* if this measure  $\nu$  is supported on the set  $E^*$  of finite paths, and of *infinite type* if  $\nu$  is supported on the set  $E^\infty$  of infinite paths [4]. (These are described as *infinite type (A)* in [2]; for finite  $E$ ,  $E^\infty$  has no wandering infinite paths, and hence there are no states which are of their infinite type (B).)

The states  $\phi_{\beta,v}$  have the form  $\phi_\epsilon \circ q_H$  with  $\epsilon$  a point mass supported at  $v$ . In [6, §6.4] we described measures on  $E^*$  for the states of the form  $\phi_\epsilon$ , so they and the  $\phi_{\beta,v}$  are of finite type. The states  $\psi_C$  factor through  $C^*(E)$ , and are of infinite type; to see this<sup>†</sup>, we use the fact that  $\psi_C$  factors through  $C^*(E)$  to compute

$$\nu(\{\lambda\}) = \nu(Z(\lambda)) - \sum_{r(e)=s(\lambda)} \nu(Z(\lambda e)) = \psi_C(s_\lambda s_\lambda^*) - \sum_{r(e)=s(\lambda)} \psi_C(s_\lambda e s_\lambda^*) = 0. \quad (7.1)$$

Thus for  $\beta$  between two critical values, say  $\beta \in (\beta_C, \beta_D)$ , the set  $E_{\beta\text{-reg}}^0$  in [2, Definition 5.5] is  $E \setminus H_{\beta_D}$  and for  $\beta$  critical it is  $E \setminus K_\beta$ . The set  $E_{\beta\text{-crit}}^0$  is empty unless  $\beta$  is critical, and it is the union of the critical components in  $E \setminus H_\beta$  if  $\beta$  is critical. If  $\beta_C$  is critical and there are sources in  $E \setminus \Sigma K_{\beta_C}$ , then  $(C^*(E), \alpha)$  has  $\text{KMS}_{\beta_C}$  states of both finite and infinite type.

*Acknowledgements.* This research has been supported by the Marsden Fund of the Royal Society of New Zealand, the Natural Sciences and Engineering Research Council of Canada, and the Australian Research Council. The authors are grateful for the opportunity to work on this project at the Banff International Research Station in November 2013.

## REFERENCES

- [1] T. Bates, D. Pask, I. Raeburn and W. Szymański. The  $C^*$ -algebras of row-finite graphs. *New York J. Math.* **6** (2000), 307–324.
- [2] T. M. Carlsen and N. S. Larsen. Partial actions and KMS states on relative graph  $C^*$ -algebras. *Preprint* arXiv:1311.0912.
- [3] M. Enomoto, M. Fujii and Y. Watatani. KMS states for gauge action on  $\mathcal{O}_A$ . *Math. Japon.* **29** (1984), 607–619.
- [4] R. Exel and M. Laca. Partial dynamical systems and the KMS condition. *Comm. Math. Phys.* **232** (2003), 223–277.
- [5] N. J. Fowler and I. Raeburn. The Toeplitz algebra of a Hilbert bimodule. *Indiana Univ. Math. J.* **48** (1999), 155–181.
- [6] A. an Huef, M. Laca, I. Raeburn and A. Sims. KMS states on the  $C^*$ -algebras of finite graphs. *J. Math. Anal. Appl.* **405** (2013), 388–399.
- [7] A. an Huef, M. Laca, I. Raeburn and A. Sims. KMS states on  $C^*$ -algebras associated to higher-rank graphs. *J. Funct. Anal.* **266** (2014), 265–283.
- [8] K. Ireland and M. Rosen. *A Classical Introduction to Modern Number Theory (Graduate Texts in Mathematics, 84)*, 2nd edn. Springer, New York, 1990.
- [9] T. Kajiwara and Y. Watatani. KMS states on finite-graph  $C^*$ -algebras. *Kyushu J. Math.* **67** (2013), 83–104.
- [10] A. Kumjian and D. Pask. Higher rank graph  $C^*$ -algebras. *New York J. Math.* **6** (2000), 1–20.
- [11] A. Kumjian, D. Pask, I. Raeburn and J. Renault. Graphs, groupoids and Cuntz–Krieger algebras. *J. Funct. Anal.* **144** (1997), 505–541.

<sup>†</sup> To see what goes wrong with this argument for states  $\phi_{\beta,v}$  which factor through  $C^*(E)$ , notice that then Theorem 5.3 implies that  $v$  is a source in some  $E \setminus \Sigma H_\beta$  or  $E \setminus \Sigma K_\beta$ , and hence  $v$  is also a source in  $E$ . But then no Cuntz–Krieger relation is imposed at  $v$  in  $C^*(E)$ , so (7.1) is not valid.

- [12] D. Lind. The entropies of topological Markov shifts and a related class of algebraic integers. *Ergod. Th. & Dynam. Sys.* **4** (1984), 283–300.
- [13] D. Lind and B. Marcus. *An Introduction to Symbolic Dynamics and Coding*. Cambridge University Press, Cambridge, 1995.
- [14] D. Pask, I. Raeburn and N. A. Weaver. A family of 2-graphs arising from two-dimensional subshifts. *Ergod. Th. & Dynam. Syst.* **29** (2009), 1613–1639.
- [15] I. Raeburn. *Graph Algebras (CBMS Regional Conference Series in Mathematics, 103)*. American Mathematical Society, Providence, RI, 2005.
- [16] E. Seneta. *Non-Negative Matrices and Markov Chains*, 2nd edn. Springer, New York, 1981.
- [17] S. P. Smith. Category equivalences involving graded modules over path algebras of quivers. *Adv. Math.* **230** (2012), 1780–1810.
- [18] S. P. Smith. Shift equivalence and a category equivalence involving graded modules over path algebras of quivers. *Preprint*, arXiv:1108.4994.
- [19] K. Thomsen. KMS weights on groupoid and graph  $C^*$ -algebras. *J. Funct. Anal.* **266** (2014), 2959–2988.
- [20] D. Yang. Type III von Neumann algebras associated with  $\mathcal{O}_\theta$ . *Bull. Lond. Math. Soc.* **44** (2012), 675–686.