

Octa-*O*-acetyl- α,α -trehalose ethanol disolvateMagnus Färnbäck,^{a,b} Lars Eriksson^{b*} and Göran Widmalm^a^aDepartment of Organic Chemistry, Arrhenius Laboratory, Stockholm University, S-106 91 Stockholm, Sweden, and ^bDivision of Structural Chemistry, Arrhenius Laboratory, Stockholm University, S-106 91 Stockholm, Sweden

Correspondence e-mail: lerik@struc.su.se

Key indicators

Single-crystal synchrotron study
 $T = 110$ K
Mean $\sigma(\text{C}-\text{C}) = 0.003$ Å
Disorder in solvent or counterion
 R factor = 0.043
 wR factor = 0.112
Data-to-parameter ratio = 19.1For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The glycosidic torsion angles φ_{H} and $\varphi_{\text{H}'}$ of the title compound, $\text{C}_{28}\text{H}_{38}\text{O}_{19} \cdot 2\text{C}_2\text{H}_5\text{OH}$, with values of -40.2 and -42.6° , respectively, are closely similar to those in the structure of the same compound crystallized from ethyl acetate. Of particular interest is that the solvent molecules in this structure take up a similar volume to those in the ethyl acetate solvate, thereby highlighting relationships between solvent and solute in these crystal structures.


Received 1 July 2004

Accepted 30 July 2004

Online 7 August 2004

Comment

Investigation of the conformation of carbohydrates in the solid state is usually performed using either NMR spectroscopy or X-ray crystallography. These techniques can also reveal insights about the effects of intramolecular stabilization factors, such as hydrogen bonds (Färnbäck, 2003). When an oligosaccharide crystallizes in the solid state, the extent to which conformational changes occur, relative to the major conformation in solution, may vary a good deal. Trehaloses are non-reducing disaccharides. The naturally occurring α,α -trehalose, α -D-Glcp-(1 \leftrightarrow 1)- α -D-Glcp, has antifreeze cell-protecting properties (Argülles, 2000). We describe here the structure of octa-*O*-acetyl- α,α -trehalose as an ethanol solvate, (I).



The non-derivatized α,α -trehalose does not show any internal hydrogen bonding in the global energy minimum conformation (Liu *et al.*, 1997) and the title compound crystallizes in a conformation similar to that determined by molecular mechanics calculations of α,α -trehalose. The glycosidic torsion angles φ_{H} ($\text{C}1'-\text{O}1-\text{C}1-\text{H}1$) and $\varphi_{\text{H}'}$ ($\text{H}1'-\text{C}1'-\text{C}1-\text{H}1$) have values of -40.2 and -42.6° , respectively (Fig. 1), similar to those of the ethyl acetate solvate (Baddeley *et al.*, 2001), *viz.* -41.8 and -43.2° . In both sugar residues, the hydroxymethyl torsion angle ω ($\text{O}5-\text{C}5-\text{C}6-\text{O}6$ and $\text{O}5'-\text{C}5'-\text{C}6'-\text{O}6'$) indicates the *gauche-trans* conformation, which is one of the two important conformations in solution. The carbonyl groups of the *O*-acetyl groups




Figure 1
The molecular structure of (I), showing 50% probability displacement ellipsoids. Disordered solvent molecules have been omitted for clarity.




Figure 2
Stereoview of an overlay plot, showing the similarity of the two rings in the trehalose molecule.

are present in synclinal arrangements compared with the C—H bonds at each substituted secondary C atom. The distance between the two anomeric H atoms is 2.47 Å (these atoms have been positioned geometrically in this study). This calculated distance may be of importance for comparison with distances derived from NMR experiments (Poveda *et al.*, 1997), as has been shown for other trehalose systems. Both pyranoside rings are in a 4C_1 chair conformation, as indicated by their Cremer & Pople (1975) parameters, $Q = 0.573$ (2) Å, $q_2 = 0.009$ (2) Å, $q_3 = 0.573$ (2) Å, $\theta = 1.0$ (2)° and $\psi = 6$ (13)° for the O5 → C5 direction, and $Q = 0.550$ (2) Å, $q_2 = 0.064$ (2) Å, $q_3 = 0.546$ (2) Å, $\theta = 6.8$ (2)° and $\psi = 0.5$ (17)° for the O5' → C5' direction. The average C—C bond length is 1.526 (7) Å, close to the average C—C bond distance of 1.52 Å found in carbohydrates in crystal structures (Pérez *et al.*,




Figure 3
The packing of slightly more than one unit-cell content, viewed along the *c* axis. Disordered solvent molecules are shown in blue. H atoms have been omitted for clarity.

2000). The most salient difference between the two sugar residues is the conformation of the *O*-acetyl groups attached to C6 and C6'. The two sugar rings, optimally fitted to each other (Norrestam, 2004) as shown in Fig. 2, have an average deviation of the atomic positions of only 0.15 Å.

Two ethanol molecules are present in the asymmetric unit. The solvent molecules are modelled with twofold disorder in each position and are arranged in columns along the *c* axis, as shown in Fig. 3. The ethanol molecules display high displacement parameters, but give a stable convergent refinement, although isotropic displacement parameters were used. Both of the solvent molecules show plausible hydrogen bonds to the octa-*O*-acetyl trehalose molecule. A comparison with the ethyl acetate solvate (Baddeley *et al.*, 2001) shows that the unit-cell dimensions of the current ethanol disolvate are essentially the same for the *c* axis but 3% larger for *b* and 3% shorter for *a*. The unit-cell volume is 0.4% smaller for the ethanol disolvate; this may be attributed both to the lower temperature of the present study (110 K) in comparison with 150 K used in the study of the ethyl acetate solvate, and to a denser packing due to the possibility of hydrogen bonding in the ethanol disolvate. The solvent molecules are packed in a closely similar position seen in projection along the *b* axis, with small changes occurring in the packing along the *c* axis but with slightly larger deviations in the *a*-axis direction.

Experimental

Octa-*O*-acetyl- α,α -trehalose was synthesized from α,α -trehalose, in which the glucosyl groups have the *D* absolute configuration, by *O*-acetylation using standard procedures (Wolrom & Thompson, 1963). It was crystallized by slow evaporation from ethanol at ambient temperature, yielding needles which were collected and mounted in capillaries with some mother liquor. The scattering power of the crystals was very weak, and it was necessary to collect data with synchrotron radiation (SR) at beamline I711 at the Swedish SR facility MAXLAB, Lund, Sweden.

Crystal data

$C_{28}H_{38}O_{19} \cdot 2C_2H_6O$	Synchrotron radiation, $\lambda = 0.891 \text{ \AA}$
$M_r = 770.72$	Cell parameters from 999 reflections
Orthorhombic, $P2_12_12_1$	$\theta = 3.0\text{--}29.0^\circ$
$a = 13.756 (4) \text{ \AA}$	$\mu = 0.11 \text{ mm}^{-1}$
$b = 15.667 (4) \text{ \AA}$	$T = 110 (2) \text{ K}$
$c = 17.934 (4) \text{ \AA}$	Needle, colourless
$V = 3865.0 (17) \text{ \AA}^3$	$0.20 \times 0.05 \times 0.03 \text{ mm}$
$Z = 4$	
$D_x = 1.324 \text{ Mg m}^{-3}$	

Data collection

Bruker SMART 1K area-detector diffractometer	9124 independent reflections
ω scans	4887 reflections with $I > 2\sigma(I)$
Absorption correction: numerical (<i>X-RED32</i> ; Stoe & Cie, 1997)	$R_{\text{int}} = 0.090$
$T_{\text{min}} = 0.97$, $T_{\text{max}} = 1.00$	$\theta_{\text{max}} = 46.0^\circ$
76 752 measured reflections	$h = -22 \rightarrow 22$
	$k = 0 \rightarrow 25$
	$l = 0 \rightarrow 28$

Refinement

Refinement on F^2	H-atom parameters constrained
$R[F^2 > 2\sigma(F^2)] = 0.044$	$w = 1/[\sigma^2(F_o^2) + (0.05P)^2]$
$wR(F^2) = 0.112$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.95$	$(\Delta/\sigma)_{\text{max}} = 0.002$
9124 reflections	$\Delta\rho_{\text{max}} = 0.39 \text{ e \AA}^{-3}$
478 parameters	$\Delta\rho_{\text{min}} = -0.36 \text{ e \AA}^{-3}$

Table 1

Selected geometric parameters (\AA , $^\circ$).

C1—O5	1.423 (2)	C5—O5	1.445 (2)
C1—O1	1.425 (2)	C1'—O5'	1.426 (2)
O1—C1'	1.428 (2)	C5'—O5'	1.447 (3)
C1—O1—C1'	112.65 (12)	C1'—O5'—C5'	113.65 (12)
C1—O5—C5	113.34 (14)		
H1—C1—O1—C1'	−40.2	H1'—C1'—O1—C1	−42.6

Table 2

Hydrogen-bonding geometry (\AA , $^\circ$).

$D\text{—}H \cdots A$	$D\text{—}H$	$H \cdots A$	$D \cdots A$	$D\text{—}H \cdots A$
O11—H11 \cdots CO3 ⁱ	0.82	2.15	2.948 (6)	168
O21—H21 \cdots O5'	0.82	2.13	2.746 (6)	132

Symmetry code: (i) $1 - x, \frac{1}{2} + y, \frac{3}{2} - z$.

A 'rigid-bond' restraint was applied to the atomic displacement parameters (Rollet, 1970). H atoms were positioned geometrically

and allowed to ride on their parent atoms, with C—H distances for CH, CH₂ and CH₃ groups set at 0.98, 0.97 and 0.96 \AA , respectively. The absolute configuration is known from the stereochemistry of the reactants in the synthesis. Due to the absence of heavy atoms with anomalous scattering effects, Friedel pairs were merged. The four disordered solvent molecules were modelled with isotropic displacement parameters due to correlation problems with anisotropic parameters. The relative occupancies of the disordered solvent molecules are 0.486 (5):0.514 (5) for one and 0.410 (6):0.590 (6) for the other.

Data collection: *SMART* (Siemens, 1998); cell refinement: *SMART*; data reduction: *SAINTE* (Siemens, 1998); program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *DIAMOND* (Bergerhoff, 1996); software used to prepare material for publication: *PLATON* (Spek, 2003).

This work was supported by a grant from the Swedish Research Council.

References

- Argüelles, J. C. (2000). *Arch. Microbiol.* **174**, 217–224.
- Baddeley, T. C., Clow, S. M., Cox, P. J., McLaughlin, A. M. & Wardell, J. (2001). *Acta Cryst.* **E57**, o456–o457.
- Bergerhoff, G. (1996). *DIAMOND*. Gerhard-Domagk-Straße 1, 53121 Bonn, Germany.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Färnäck, M. (2003). Doctoral thesis, Stockholm University, Sweden.
- Liu, Q., Schmidt, R. K., Teo, B., Karplus, P. A. & Brady, J. W. (1997). *J. Am. Chem. Soc.* **119**, 7851–7862.
- Norrestam, R. (2004). *ROTERA*. Stockholm University, Sweden.
- Peréz, S., Gautier, C. & Imbert, A. (2000). *Oligosaccharides in Chemistry and Biology*, edited by B. Ernst, G. W. Hart & P. Sinay, pp. 969–1001. Weinheim: Wiley-VCH.
- Poveda, A., Vicent, C. Penadés, S. & Jiménez-Barbero, J. (1997). *Carbohydr. Res.* **301**, 5–10.
- Rollet, J. S. (1970). *Crystallographic Computing*, edited by F. R. Ahmed, S. R. Hall & C. P. Huber, pp. 167–181. Copenhagen: Munksgaard.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.
- Siemens (1998). *SMART* and *SAINTE*. Siemens Analytical X-ray Instruments Inc., Madison, Wisconsin, USA.
- Stoe & Cie (1997). *X-RED32*. Version 1.09. Stoe & Cie GmbH, Hilpertstraße 10, D-64295 Darmstadt, Germany.
- Wolrom, M. L. & Thompson, A. (1963). *Methods Carbohydr. Chem.* **2**, 211–215.