

Crystallization and preliminary X-ray diffraction
analysis of the lectin from *Canavalia gladiata* seeds

Frederico B. M. B. Moreno,^a
Plínio Delatorre,^{a,b} Beatriz T.
Freitas,^a Bruno A. M. Rocha,^a
Emmanuel P. Souza,^a Felipe
Facó,^a Fernanda Canduri,^{c,d}
André L. H. Cardoso,^b Valder N.
Freire,^e José L. Lima Filho,^f
Alexandre H. Sampaio,^g Juan J.
Calvete,^h Walter F. De Azevedo
Jr^{c,d} and Benildo S. Cavada^{a*}

^aDepartamento de Bioquímica e Biologia Molecular, Universidade Federal do Ceará, Fortaleza, CE 60451-970, Brazil, ^bGrupo de Química Biológica, Departamento de Ciências Biológicas, Universidade Regional do Cariri, Crato, CE 63105-000, Brazil, ^cDepartamento de Física, UNESP, São José do Rio Preto, SP 15054-000, Brazil, ^dCenter for Applied Toxinology, CEPID, FAPESP, Av. Vital Brasil 1500, São Paulo, SP 05503-900, Brazil, ^eDepartamento de Física, Universidade Federal do Ceará, Fortaleza, CE 60451-970, Brazil, ^fLaboratório de Imunopatologia Keizo Asami, Univesidade Federal de Pernambuco, Recife, PE 50670-901, Brazil, ^gDepartamento de Engenharia de Pesca, Universidade Federal do Ceará, Fortaleza, CE 60451-970, Brazil, and ^hInstituto de Biomedicina de Valencia, CSIC, 46010 Valencia, Spain

Correspondence e-mail: bscavada@ufc.br

The seed lectin from *Canavalia gladiata* was purified and crystallized. Orthorhombic crystals belonging to space group $C22_1$ grew within three weeks at 293 K using the hanging-drop vapour-diffusion method. Using synchrotron X-ray radiation, a complete structural data set was collected at 2.3 Å resolution. The preliminary crystal structure of the lectin, determined by molecular replacement, had a correlation coefficient of 0.569 and an R factor of 0.412.

Received 10 May 2004

Accepted 14 June 2004

1. Introduction

The legume commonly known as the sword bean, *Canavalia gladiata*, belongs to the genus *Canavalia* (family Leguminosae, subfamily Papilionoideae, tribe Diocleinae). This bean represents a cheap source of protein and calories in several areas of Northeast Brazil. Sword beans show favourable agronomic features that make them suitable for cultivation in the tropics and have a high average yield compared with that of soybeans (Ekanayake *et al.*, 2000). Although *C. gladiata* seeds contain a large amount of protein (35%), the protein exhibits a low level of sulfur-containing amino acids (Siddhuraju & Becher, 2001). Anti-nutrient compounds such as phenolics, tannins, saponins, non-proteic amino acids, protease and α -amylase inhibitors and lectins are also present in sword-bean seeds. Lectins are proteins of non-immune origin whose structure includes at least a carbohydrate-binding domain (Peumans *et al.*, 2001). *C. gladiata* seed lectin (CGL) shows sugar-binding specificities similar to those of cona-*navalin* A (ConA), the well characterized lectin from *C. ensiformis* seeds (Kojima *et al.*, 1991). The lectin from *C. gladiata* has been sequenced (Yamauchi & Minamikawa, 1990) and when compared with ConA exhibits 95% amino-acid sequence identity. However, the Diocleinae lectins have been found to express distinct biological properties, as observed for the lectins from *C. brasiliensis* seeds (ConBr) and ConA, which share 99% amino-acid sequence (Sanz-Aparicio *et al.*, 1997). CGL binds to saccharides containing α -D-mannose or α -D-glucose residues and to oligosaccharides or polysaccharides containing these mono-saccharides.

Of the leguminous lectins isolated from the tribe Diocleinae, only those purified from the seeds of *C. ensiformis* (Hardman & Ainsworth, 1972), *C. brasiliensis* (Grangeiro *et al.*, 1997), *Dioclea grandiflora* (Rozwarski *et al.*, 1998),

D. guianensis (Gomes *et al.*, 1994) and *Cratylia mollis* (Souza *et al.*, 2003) have had their three-dimensional structures resolved by X-ray crystallography. These lectins have 99, 98, 83, 88 and 78% identity in their amino-acid sequences with CGL, respectively. They are dimeric or tetrameric structures built up of dome-shaped monomers consisting of two β -sheets of seven and six antiparallel β -strands interconnected by turns and loops (Calvete *et al.*, 1999). Although this group of lectins exhibits a high degree of primary structure similarity, a few amino-acid residue substitutions in key positions endow the homologous lectins with distinct biological effects. Thus, the seed lectins from *C. brasiliensis* (ConBr) and *C. ensiformis* (ConA) differ in just two amino-acid residues in their 237-residue monomer sequences. This minor difference results in lectins that exhibit distinct *in vitro* and *in vivo* biological activities, such as induction of rat paw oedema (Bento *et al.*, 1993), peritoneal macrophage spreading in mouse (Rodríguez *et al.*, 1992) and human lymphocyte stimulation (Barral-Neto *et al.*, 1992). The three-dimensional structures of the lectin monomers of ConA and ConBr can be superimposed with an r.m.s. deviation of 0.65 Å (Sanz-Aparicio *et al.*, 1997). However, structural differences are amplified in the lectin tetramers at 1.62 Å, suggesting that the distinct quaternary structures, which in turn determine the spatial arrangement of the four carbohydrate-binding sites of the lectin tetramers, may account for the differing biological properties of ConA and ConBr (Sanz-Aparicio *et al.*, 1997).

The quaternary structure variation exhibited by lectins that have nearly the same subunit structure precludes the use of a molecular-modelling approach to establish structure–function correlations. Thus, crystallographic studies have been carried out in order to establish the tertiary and quaternary structures of the lectins from the Diocleae subtribe as a first step towards their possible use in nano-

biotechnology and molecular-biology studies. Currently, 54 lectin structures from the Diocleae subtribe have been deposited in the Protein Data Bank (Berman *et al.*, 2000). However, 50 entries correspond to diverse forms of ConA. With the aim of establishing the crystal structure of the lectin from seeds of *C. gladiata*, we report here its purification, crystallization and preliminary X-ray diffraction analysis.

2. Materials and methods

2.1. Purification of *C. gladiata* seed lectin

Mature *C. gladiata* seeds were collected in Ceará state, Northeast Brazil. The seeds were grounded to a fine powder in a coffee mill and then defatted with *n*-hexane. Soluble proteins were extracted at 291 K for 4 h by continuous stirring with 0.15 M NaCl [1:10(*w*:*v*)], followed by centrifugation at 10 000g at 277 K for 20 min. The supernatant was applied onto a Sephadex G-50 column (5 × 25 cm), which had been equilibrated with 0.15 M NaCl containing 5 mM CaCl₂ and MnCl₂, as described by Ceccatto *et al.* (2002). CaCl₂ and MnCl₂ were added in order to retain carbohydrate-binding activity. The column was then washed with equilibration buffer at a flow rate of 45 ml h⁻¹ until the effluent absorbance at

280 nm was below 0.05. The bound lectin was eluted with 0.1 M glycine pH 2.6, dialyzed extensively against distilled water and lyophilized. The affinity chromatography fraction was further purified using an Äkta chromatographic system and a Mono-Q column (5 × 0.5 cm) equilibrated with 20 mM Tris-HCl pH 7.0 and developed with a linear gradient of 20 mM Tris-HCl pH 7.0 containing 1.0 M NaCl at a flow rate of 1 ml min⁻¹ and a slope of 5% NaCl min⁻¹. The *C. gladiata* lectin recovered in the unbound fraction was exhaustively dialyzed against distilled water and freeze-dried. The purity of CGL was assessed by SDS-PAGE.

2.2. Crystallization, data collection and processing

For crystallization trials, the purified lectin was dissolved at a concentration of 10 mg ml⁻¹ in 20 mM Tris-HCl pH 7.5 containing 5 mM CaCl₂ and MnCl₂. Crystals of CGL were grown in Linbro plates at 293 K by the vapour-diffusion method (Jancarik & Kim, 1991) in hanging drops, using Crystal Screen from Hampton Research. The drops were composed of equal volumes (3 µl) of protein solution and 0.1 M Tris-HCl pH 8.5 with 2.0 M ammonium sulfate and were equilibrated against 500 µl reservoir solution. Single crystals were transferred to a cryoprotectant solution consisting of 25% glycerol and 75% 0.1 M Tris-HCl pH 8.5 with 2.0 M ammonium sulfate and flash-frozen in a stream of nitrogen at 100 K. X-ray diffraction of cryoprotected crystals was performed at a wavelength of 1.4310 Å using Synchrotron Radiation Source (CPr station, Laboratório Nacional de Luz Síncrotron-LNLS, Campinas, Brazil) and a complete data set was collected using a CCD (MAR Research) in 120 frames with an oscillation range of 1°. The data were indexed, integrated and scaled using *MOSFLM* and *SCALA* (Collaborative Computational Project, Number 4, 1994).

3. Results and discussion

Small crystals of CGL appeared in 0.1 M Tris-HCl pH 8.5 containing 2.0 M ammonium sulfate. However, these crystals were not suitable for X-ray diffraction analysis. To optimize the condition of crystallization, changes in the pH range were tested. The best crystals grew in 0.1 M Tris-HCl pH 9.0, 2.0 M ammonium sulfate within 20 d (Fig. 1a) to maximum dimensions of approximately 0.4 × 0.2 × 0.3 mm and diffracted to a maximum resolution of 2.0 Å.

The crystals belong to the orthorhombic space group C22₁ (unit-cell parameters *a* = 101.81, *b* = 117.21, *c* = 243.61 Å). Data in the 51.99–2.31 Å resolution range were scaled using *SCALA* to an *R*_{merge} of 0.073 (0.354) with an (*I*/*σ*(*I*)) of 8.9 (2.4). The final data set to 2.31 Å resolution, with a total of 279 320 observed and 60 699 unique reflections, was 99.1% (99.8%) complete. The values in parentheses are for the highest resolution shell (2.42–2.31 Å). The refined mosaic spread of the crystal was 0.38°. Assuming the presence of four 25.5 kDa lectin monomers in the asymmetric unit, the calculated Matthews coefficient (*V*_M; Matthews, 1968) was 3.56 Å³ Da⁻¹, indicating a solvent content of 65.4%. Like ConA, CGL is a tetramer in the asymmetric unit.

The preliminary crystal structure of CGL was determined by standard molecular-replacement methods using the program *MOLREP* (Vagin & Teplyakov, 1997). The atomic coordinates of various lectin monomers were used in the search for a structural model. The best result was obtained with ConA (PDB code 1gkb; Kantardjieff *et al.*, 2002). The best solution had a final correlation coefficient of 0.569 and an *R* factor of 0.412. After placing the molecule in the unit cell, rigid-body refinement was performed using the *X-PLOR* program (Brünger, 1992). Refinement resulted in a model with an *R*_{free} of 0.375 and an *R* factor of 0.287. Refinement of the X-ray crystallographic structure of CGL is in progress.

This work was partly financed by Fundação Cearense de Apoio ao Desenvolvimento Científico e Tecnológico-FUNCAP, Conselho Nacional de Desenvolvimento Científico e Tecnológico-CNPq, FAPESP (SMOLBNet, 01/07532-0), Universidade Regional do Cariri-URCA, Coordenação de Aperfeiçoamento de Pessoal de Nível Superior-CAPES, National Synchrotron Light Laboratory-LNLS, Brazil and grant No. BMC2001-3337 from the Ministerio de Educación y Ciencia, Madrid, Spain. AHS, BSC, JLLF, WFA and VNF are senior investigators of CNPq.

References

- Barral-Neto, M., Santos, S. B., Barral, A., Moreira, L. I. M., Santos, C. F., Moreira, R. A., Oliveira, J. T. A. & Cavada, B. S. (1992). *Immunol. Invest.* **21**, 297–303.
- Bento, C. A. M., Cavada, B. S., Oliveira, J. T. A., Moreira, R. A. & Barja-Fidalgo, C. (1993). *Agents Actions*, **38**, 48–54.
- Berman, H. M., Westbrook, J., Feng, Z., Gilliland, G., Bhat, T. N., Weissig, H., Shindyalov, I. N. &




Figure 1
(a) Native crystal of the lectin from *C. gladiata*. (b) The crystal diffraction pattern at 2.0 Å.

- Bourne, P. E. (2000). *Nucleic Acids Res.* **28**, 235–242.
- Brünger, A. T. (1992). *X-PLOR Version 3.1. A System for Crystallography and NMR*. New Haven: Yale University Press.
- Calvete, J. J., Thole, H. H., Raida, M., Urbanke, C., Romero, A., Grangeiro, T. B., Ramos, M. V., Almeida da Rocha, I. M., Guimarães, F. N. & Cavada, B. S. (1999). *Biochem. Biophys. Acta*, **1430**, 367–375.
- Ceccatto, V. M., Cavada, B. S., Nunes, E. P., Nogueira, N. A., Grangeiro, M. B., Moreno, F. B., Teixeira, E. H., Sampaio, A. H., Alves, M. A., Ramos, M. V., Calvete, J. J. & Grangeiro, T. B. (2002). *Protein Pept. Lett.* **9**, 67–73.
- Collaborative Computational Project, Number 4 (1994). *Acta Cryst.* **D50**, 760–763.
- Ekanayake, S., Jansz, E. R. & Nair, B. M. (2000). *Plant Foods Hum. Nutr.* **55**, 305–321.
- Gomes, J. C., Ferreira, R. R., Cavada, B. S., Moreira, R. A. & Oliveira, J. T. (1994). *Agents Actions*, **41**, 132–135.
- Grangeiro, T. B., Schriefer, A., Calvete, J. J., Raida, M., Urbanke, C., Barral-Netto, M. & Cavada, B. S. (1997). *Eur. J. Biochem.* **248**, 43–48.
- Hardman, K. D. & Ainsworth, C. F. (1972). *Biochemistry*, **11**, 4910–4919.
- Jancarik, J. & Kim, S.-H. (1991). *J. Appl. Cryst.* **24**, 409–411.
- Kantardjieff, K., Hocht, P., Segelke, B., Tao, F. & Rupp, B. (2002). *Acta Cryst.* **D58**, 735–743.
- Kojima, K., Ogawa, H., Seno, N. & Matsumoto, I. (1991). *Carbohydr. Res.* **213**, 275–282.
- Matthews, B. W. (1968). *J. Mol. Biol.* **33**, 491–497.
- Peumans, W. J., Van Damme, E. J., Barre, A. & Rouge, P. (2001). *Adv. Exp. Med. Biol.* **491**, 27–54.
- Rodríguez, D., Cavada, B. S., Oliveira, J. T. A., Moreira, R. A. & Russo, M. (1992). *Braz. J. Med. Biol. Res.* **25**, 823–826.
- Rozwarski, D. A., Swami, B. M., Brewer, C. F. & Sacchettini, J. C. (1998). *J. Biol. Chem.* **273**, 32818–32825.
- Sanz-Aparicio, J., Hermoso, J., Grangeiro, T. B., Calvete, J. J. & Cavada, B. S. (1997). *FEBS Lett.* **405**, 114–118.
- Siddhuraju, P. & Becher, K. (2001). *Nahrung*, **45**, 224–233.
- Souza, G. A., Oliveira, P. S. L., Trapani, S., Santos, A. C. O., Rosa, J. C., Laure, H. J., Faça, V. M., Correia, M. T. S., Tavares, G. A., Oliva, G., Coelho, L. C. B. B. & Greene, L. J. (2003). *Glycobiology*, **13**, 961–972.
- Vagin, A. & Teplyakov, A. J. (1997). *J. Appl. Cryst.* **30**, 1022–1025.
- Yamauchi, D. & Minamikawa, T. (1990). *FEBS Lett.* **260**, 127–130.