

The Prevalence of Endemic Goiter and Its Relationship to Urine Iodine in Primary School Age Children in Khoramabad

Mahnaz Mardani¹, Fateme Gholaami², Pouria Rezapour³, Farzad Ebrahimzadeh⁴

¹Associate Professor of Nutrition, Lorestan University of Medical Sciences, Khoramabad, Iran;

²GP, Lorestan University of Medical Sciences, Khoramabad, Iran;

³Ms Student of English literature, Kharazmi University, Tehran, Iran;

⁴Instructor of Statistics, Lorestan University of Medical Sciences, Khoramabad, Iran

Correspondence:

Mahnaz Mardani,
Associate professor of Nutrition,
Lorestan University of Medical Sciences,
Khoramabad, Iran

Tel: +98 66 33204878

Fax: +98 66 33120172

Email: msmardani@yahoo.com

Received: 6 December 2014

Revised: 20 January 2015

Accepted: 9 March 2015

Abstract

Backgrounds: Iodine deficiency disorder is a major public health problem in areas where there is a lack of iodine in drinking water, soil, and food. This study was carried out to assess the prevalence of goiter among school children in the age group of 6-10 years in Khorramabad city. Also, the level of urine iodine was evaluated.

Methods: The study was conducted from January 2012 to January 2013 in 1125 school children of 6-10 years of age, attending all the schools of Khorramabad city. The subjects were selected through stratified, random, and cluster sampling methods. Their Goiter degrees were evaluated according to WHO classification. Excretory urine iodine was measured based on micrograms per deciliter. Furthermore, the digestion method was used for experimentation.

Results: Out of the 1125 subjects, 866 (77%) had Goiter. 24.2% were diagnosed with Goiter 1a, 45.3% with goiter 1b, and 7.2% with type 2 Goiter. Regarding Goiter, there was no difference between the two sexes. The prevalence of goiter increased with age, but it decreased with the increase of parental educational level. Also, goiter prevalence was lower in families with a higher socioeconomic status. The median urinary iodine was 17.1 micrograms per deciliter, which is desirable in terms of the WHO criterion.

Conclusion: The results of the study indicated that using iodizing edible salt to fight with iodine deficiency was effective and helped Khorramabad reach the International indexes of iodine control. Despite this, Khorramabad is one of the hyper- endemic regions in the world.

Please cite this article as: Mardani M, Gholaami F, Rezapour P, Ebrahimzadeh F. The Prevalence of Endemic Goiter and Its Relationship to Urine Iodine in Primary School Age Children in Khoramabad. *J Health Sci Surveillance Sys.* 2015;3(2):71-75.

Keywords: Endemic goiter, Prevalence, Students, Urine iodine

Introduction

Today, it has been confirmed that the lack of iodine in food and drinking water can affect the function of the thyroid gland.^{1,2} Iodine deficiency can cause problems called Iodine Deficiency Disorders (IDD). The most serious problems are endemic Goiter, Hypothyroidism, Stillbirth, Congenital Malformations,³ minor brain damage, irreversible nerve-brain-psychotic regeneration

problems, and infertility.⁴ Goiter is an enlarged thyroid gland, the most common symptom of Iodine deficiency.^{5,6} Endemic Goiter is common in mountainous regions and areas with deficiency of iodine in drinking water, soil, and food.^{7,8} Based on global researches, around 2 billion people in the world have IDD, especially in Southern Asia and the African Desert.⁹

WHO highlighted the importance of IDD prevalence in 1986. It also confirmed programs to

eradicate IDD, such as public health programs to decrease the level of goiter to 10% among the students carried out by 2000.^{10,11} In studies, until 1990, around 40 million infants per year were prone to congenital malformations of the brain due to their mothers' lack of dietary iodine. This rate reached 28 million children by 1997, a clear reduction, but it is still regarded as high.¹²

According to previously carried out studies, iodine deficiency and endemic Goiter used to be epidemic in all of the provinces of the Islamic Republic of Iran.^{13,14} In reaction to the high prevalence of endemic Goiter in some regions of Iran (up to 89.5% in Semirum), a national program was started in 1989 to control IDD.¹⁴ The mandatory use of iodizing salt was legislated in 1994.¹⁵

The legislation of the prevention of IDD was followed by a high level of success in Iran^{13,16} but, despite sufficient iodine intake, there was goiter prevalence in some enriched regions with endemic iodine. And there also was a mild to moderate degree of Goiter. Thus, the prevalence of goiter cannot be justified only by the lack of iodine. Among the suggested factors, there were also unexpected factors, such as unknown *Goitrogens*, no use of iodized salt, no standard of Iodized salt, thyroid self-defending diseases, or the lack of some micronutrients such as Iron and selenium.¹⁷

Due to different reports on Goiter in our country, the lack of current information on Goiter prevalence in KhorramAbad city, the importance of children groups, and occasional controlling programs held only every five year in the country, and omission of disorders caused by lack of iodine, this study (2012-2013) was carried out to determine the prevalence of endemic goiter, to measure the level of urine iodine and test the diet iodine level among school children aged between 6 and 10 in Khorramabad.

Materials and Methods

This study was carried out in 2012-2013 as a cross-sectional study. 1125 Khorramabad city school children, aged between 6 and 10, participated in the study. They were chosen by cluster sampling method among girls and boys at elementary schools. Biographic and background information of each student was registered through documentation, the parents were interviewed, and the results were registered in each student's questionnaire. Then each student was visited by a trained medical doctor for clinical examination of the thyroid gland and the Goiter rates determined by the WHO protocol were resolved. Based on World Health Organization standards,¹⁸ when the thyroid gland cannot be seen and is untouchable, it is in zero level; when it is not visible but it is touchable, it is leveled in 1A. When the thyroid gland is seen by head extension, it is 1b; also, if it is observable

in the normal form of the head, it is type 2, and if it is observable from 6 meters it is Goiter type 3.

Analysis of urine iodine is a basic tool for recognizing iodine levels; more than 90% of the received iodine is visible in urine. Urine samples of the student were collected to determine their iodine level. Iodine testing was done by acidic digestion method, in which urine details are digested by Chloride *acid* in high temperatures resulting in the release of iodine. According to Global Health Organization's recommendations, the level of urine excretion of iodine lower than 2µg/dl means severe lack of iodine, between 2 and 4.9 µg/dl means medium lack of iodine, between 5 and 9.9 µg per dl indicates a low level of lack of iodine, between 10 and 19.9 shows a situation without lack of iodine, between 20 and 29.9µg/ dl means the reception of more than necessary iodine, and more than 30 means much more reception of iodine than necessary. In this study, data analysis was done using SPSS software; χ^2 independence method, Spearman correlation coefficient, and statistical descriptive methods were used to analyze the data of the experiments.

Results

1125 students aged between 6-10 years participated in this study; they were visited by doctors and urine samples were taken. 77% of the students had Goiter, 24.4% had type 1a, 45.3 % 1b, and 7.2 % Goiter type 2. No case of Goiter type 3 was seen (Table1).

The median urinary iodine concentration was 17.1 µ/dl, the mean urinary iodine was determined to be 18.98 and the standard deviation of urinary iodine was 10.3. According to urinary iodine concentration, 6.8% of the students had moderate to severe iodine deficiency, which is considered to be within the range allowed (up to 20%) and 11.3% had mild iodine deficiency. 40.4% had normal urinary iodine, 25.9% had more than needed iodine, and 15.6% had very high urinary iodine levels. Of the 1125 students surveyed, 50.8% were male and 49.2% female.

According to the results of Chi-square test of independence, there was no significant relationship between gender and Goiter prevalence (P=0.81). Goiter prevalence in boys was 76.7% and in girls 77.2; the difference was not statistically significant (Table1).

Of the 1125 students surveyed, 18.7% were 6-years-old, 19.8% were 7-years-old, 18.8% were 8-years-old, 20.8% were 9-years-old, and 22% were 10-years-old.

The prevalence of goiter in 6 year old students was 73.3% and it was 80.6% in 10 year old ones. Based on Spearman correlation coefficient, there was no significant relationship between urine iodine level and

Table 1: Adaptive table of students categorized according to sex and goiter prevalence

Grade	Goiter	Sex		Total
		Boy	Girl	
0	Frequency	133	126	259
	percent	23.3	22.8	23
a ¹	Frequency	140	135	275
	percent	24.5	24.4	24.5
b ¹	Frequency	254	256	510
	percent	44.4	46.3	45.3
2	Frequency	45	36	81
	percent	7.9	6.5	7.2
Total	Frequency	572	553	1125
	percent	100	100	100

family dimension. Medium poverty to high poverty levels of students' families with >6 and < 3 members were 12% and 21.4% respectively, and in students with 3-4 and 5-6 members these percentages were 6.3% and 6.5% respectively; the difference was not significant.

Discussions

Lack of iodine and its related disorders still exist worldwide; in most parts of Iran, such as Khoarmabad, it seems to be endemic. Many studies have been done in different provinces to determine goiter prevalence and the present study is the most recent one.

As prevalence of Goiter was more than 50% in this region, Khorramabad is considered one of the hyper-endemic regions of the world. The present prevalence of goiter compared to a 1989 study which was 100%¹⁹ shows reduction and it has increased compared to 66% high prevalence in 1994.²⁰ Furthermore, the results indicated that the median urine iodine is 17.1 µg per dl which is within the range of the Global Health standard.¹⁸ Researchers around the world are still trying to find the reason for high existence of Goiter regardless of the sufficient intake of iodine.

Kapil and colleagues²¹ reported that the total Goiter rate is 16.8% and median Urinary Iodine concentration level was 115 µg/L amongst school age children in district Pauri, Uttarakhand of India. In the Girma and colleagues²² study, the total goiter prevalence of 750 school children aged between 6 and 18 years in Metekel Zone, northwest Ethiopia was 39.5% and their median urinary iodine concentration range was 39.9 µg/L. Uzun and colleagues have shown that the prevalence of hypothyroidism in children aged 6-12 and 13-19 years of the West Black Sea region of Turkey was 10.4% and 18.9%, respectively, and the median urinary iodine concentration levels in children aged 6-12 and 13-19 years were 83 µg/l and 78 µg/l, respectively.²³ Although Kocak and colleagues have reported that the overall goiter prevalence was 26.5% in an adult population in a formerly iodine-deficient area of Turkey,²⁴ Kapil and colleagues have found that the total goiter prevalence was 15.8% in the children

aged of 6-12 years in district Kangra, Himachal Pradesh after 60 years of salt iodization.²⁵

The thyroid gland changes slowly; thus, the intake of iodine salt is not capable of controlling goiter prevalence. This may be one of the reasons for the high level of goiter regardless of iodine intake. Zimmerman's study on Iodine deficiency,²⁶ a study in Turkey on Zinc deficiency,²⁷ and a study in Sudan on the lack of vitamin A and malnutrition²⁸ were introduced as reasons of the slow decreasing rate of goiter endemic. The effect of anti-thyroid antibodies is one of the reasons of goiter prevalence regardless of the sufficient iodine consumption. Boukis' study, carried out in 1983 in a region diagnosed with iodine deficiency in Greece, showed that anti-thyroid antibodies of goiter patients could not be measured before treatment by oily iodine but 3 -6 months after treatment with oily iodine these antibodies were positive in 42.8% of the patients.²⁹ In a Ashrafi and colleagues' study in Isfahan, due to the prevalence of anti-thyroid antibodies in patients with goiter in comparison to healthy people, self defense was determined as a reason for simple goiter after public use of iodized salt.³⁰ The same was the result for a study in Semirum, carried out in 2007, which indicated that anti-thyroid antibodies were recognized as a reason for the remaining of goiter regardless of the consumption of iodized salt.¹⁷

Havach's study in 1985 showed that adding iodine to food to treat goiter due to the lack of iodine resulted in the prevalence of self defended thyroiditis and an increase in anti-thyroid antibodies.³¹ Furthermore, Zhang and colleagues³² concluded that the excessive intake of iodine might serve as a risk factor for toxic nodular goiter, thyroid papillary carcinoma and Hashimoto's thyroiditis.

In this study, there was no significant relationship between goiter and sex; this confirms the results of studies in 2001 in Tehran,³³ Semirum in 2007¹⁷ Kocak in 2014,²³ Marwaha in 2012³⁴ and Markazi in 2001³⁵ and many other studies.³⁶⁻³⁸ But, in studies such as those carried out in Tehran in 2007³⁹ and in Kashan in 2008⁴⁰ and some other studies, the prevalence of

goiter was determined higher among girls.

In a 1997 study carried out in Kashan, the prevalence of goiter was determined higher among boys compared to girls.⁴¹ In this study, there was a significant and linear relationship between age and prevalence of goiter; aging increases the prevalence of goiter. This shows the increase in public knowledge and the effect of the consumption of iodized salt.

In similar studies in Bandar Abas in 2000⁴² and in Tehran in 2007, aging showed a higher prevalence of goiter.³⁹ In a study in 2002 in Yazd, there was no relationship between age and goiter.⁴³

In this study, there was no significant relationship between goiter prevalence and family dimensions, although goiter among single parent families and families with more than 6 members was higher compared to families with 3 to 6 members.

Conclusion

This study demonstrated that iodine deficiency is still a severe public health problem in Khorramabad city, about 25 years after the introduction of salt iodine supplementation program. This study suggests that autoimmunity may be one of the mechanisms responsible for goiter persistence, after iodine replenishment, in this iodine deficient region, but the role of other factors should also be considered. We suggest the further confirmation of the existing monitoring system for the quality of iodized salt. Also, there is a need for further studies in all Iranian provinces to achieve proper elimination of IDD in the community.

Conflict of Interest: None declared.

References

- Shikholeslam K, Sadvandian S. Disorders of iodine deficiency in water. *Yearly Teaching Journal of Health Ministry* 1996; 28: 1-45.
- Azizi F. Iodine deficiency disorders. *Journal Of Shahid Beheshti Medical Faculty*. 1995; 2: 24-40.
- Hetz B. *The prevention and control of iodine deficiency disorders*. Amsterdam, Elsevier. 1993; 27: 1-16.
- Delange F. Iodine deficiency in Europe and its consequences. An update, *Eur J Nucl Med Mol Imaging* 2002; 29: 404-16.
- Delange F, Dunn J. Iodine deficiency in: Braverman LE, Utiger RD, editors. *The Thyroid: A fundamental and clinical text* 9th ed. Philadelphia : Lippincott Williams & Wilkins 2005; 26: 264-88.
- Stone T. Endemic goiter, nutrition and landholding in Bangladesh. *Int J Epidemiol* 1984; 13: 58-64.
- Koutras D, Christakis G, Trichopoulos D, Dakovoutetaki A, Kyriakopoulos V, Fontanares P, et al. Endemic goiter in Greece. Nutritional status, growth, and skeletal development of goitrous and nongoitrous population. *AMJ Clin Nutr* 1973; 26: 1360-8.
- Koutras D, Papapetrou P, Yataganas X, Malamos B. Dietary sources of iodine in area with and without iodine deficiency goiter. *AMJ Clin Nutr* 1970; 23: 870-4.
- Zimmermann M. Iodine deficiency. Human Nutrition Laboratory, Swiss Federal Institute of Technology Zurich, CH-8092 Zurich, Switzerland and Division of Human Nutrition, Wageningen University, 6708 Wageningen, The Netherlands. *Endocrine Reviews* 2009; 30: 376-408.
- Azizi F, Sheikholeslam R, Hedayati M. Goiter survey and urinary iodine concentration in 8 to 10 years old school-children from Fars province in 1996. *Shahid Beheshti University (MC) & Health Service Endocrine & Metabolism Research Center* 2001; 3: 37-42.
- Azizi F, Delshad H, Amouzegar A. Marked reduction in goiter prevalence and eventual normalization of urinary iodine concentration in Iranian school-children 10 years after salt iodization (Third National survey of Iodine Deficiency Disorders 2000). *Shahid Beheshti University (MC) & Health Services Endocrine & Metabolism Research Center* 2008; 10: 191-203.
- Fumee C, West C, Vander Haar F, Hautvast J. Efficacy of oral iodized oil is associated with anthropometric status in severely iodine deficient school children in rural Malawi. *Br J Nutr* 2008; 84(3): 345-52.
- Azizi F, Navai L, Fattahi F. Goiter prevalence, urinary iodine excretion, thyroid function and anti-thyroid antibodies after 12 years of salt iodization in shahriar, Iran. *Int J Vitam Nutr Res* 2002; 72(5): 291-5.
- Emami A, Amini M, Azizzadeh A. Goiter prevalence from iodine deficiency after lipidol injection in Hana school children. *Faculty Med J* 1994; 38: 47-53.
- Azizi F, Sheikholeslam R, Hedayati M, Mimran P, Malekafzali H, Kimiagar M, et al. Sustainable control of iodine deficiency in Iran: Beneficial result of the implementation of mandatory law on salt iodization. *J Endocrinol Invest* 2002; 25(5): 409-13.
- Hashemipour M, Amini M, Gheisari A, Sharifei S, Iranpour R, Aminorroaya A. Comparison of urinary iodine excretion in neonates and their mothers in Isfahan, Iran. *Endocrine Pract* 2002; 8(5): 347-50.
- Hashemipour M, Amini M, Aminorroaya A, Dastjerdi M, Rezvanian H, Kachoei A, et al. High prevalence of goiter in an iodine replete area: Do thyroid auto-antibodies play a role?. *Asia Pac J Clin Nutr* 2007; 16(3): 403-10.
- World Health Organization, United Nations Childrens Fund, and International council for control of Iodine Deficiency Disorders Indicators for assessing iodine deficiency disorders and their control through salt iodization. Geneva: World Health Organization, 1994. (WHO/NUT 94.6).
- The committee of fighting with the complications of iodine deficiency, Quick assessment of goiter, Lorestan state's findings, 1989.

- 20 Sotudemaram A, Papi A, Ravanshad S, Sadeghi Hasanabadi A. Assessment of goiter prevalence in 6-10 years old school children of Khorramabad city, 1994. *Journal of Kerman University of Medical Sciences* 1998; 5(3): 142-7.
- 21 Kapil U, Pandey RM, Prakash S, Sareen N, Bhadoria AS. Assessment of iodine deficiency in school age children in Nainital District, Uttarakhand State. *Asia Pac J Clin Nutr* 2014; 23(2): 278-81.
- 22 Girma K, Nibret E, Gedefaw M. The status of iodine nutrition and iodine deficiency disorders among school children in Metekel Zone, northwest Ethiopia. *Ethiop J Health Sci* 2014 Apr; 24(2): 109-16.
- 23 Uzun H, Gozkaya S, Yesildal N, Okur M, Arslanoglu İ, Kocabay K, et al. The prevalence of goiter and hypothyroidism among school children 6 years after introduction of a mandatory salt iodination program in a severely iodine-deficient area of the West Black Sea region of Turkey. *J Trop Pediatr* 2014 Aug; 60(4): 318-21.
- 24 Kocak M, Erem C, Deger O, Topbas M, Ersoz HO, Can E. Current prevalence of goiter determined by ultrasonography and associated risk factors in a formerly iodine-deficient area of Turkey. *Endocrine* 2014 Sep; 47(1): 290-8.
- 25 Kapil U, Pandey RM, Kabra M, Jain V, Sareen N, Bhadoria AS, et al. Status of iodine deficiency in district Kangra, Himachal Pradesh after 60 years of salt iodization. *Eur J Clin Nutr* 2013 Aug; 67(8): 827-8.
- 26 Zimmerman M, Adou P, Torresani T. Persistence of goiter despite oral iodine supplementation in goiterous children with iron deficiency anemia in Cote de Ivoir. *Am J Clin Nutr* 2000; 71: 81-93.
- 27 Ozata M, Aydin A. Iodine and zinc, but not selenium and copper def exists in a male Turkish population with endemic goiter. *Biol Trace Elem Res* 1999; 69: 211-16.
- 28 Rezvanian H, Hashemipour M, Aminoraaya A, Kachui A, Moaddab M, Siavash M. Is iodine deficiency the cause of endemic goiter in Semiröm region?. *Iran Journal of Endocrinology & Metabolism* 2006; 8: 351-6.
- 29 Boukis M, Koutras D, Sauvatzoglu A. Thyroid hormone and immunological studies in endemic goiter. *J Clin Endocrinol Metab* 1983; 57: 859-65.
- 30 Ashrafi F, Momen-Zadeh M, Amini M. Study of Anti Thyroidal Antibodies in Simple Goiter Patients and Comparison with a Control Group . *JSSU* 2004; 12: 12-16.
- 31 Havach H, Escalante D, Onativia A. Thyroid carcinoma and thyroiditis in an endemic goiter region before and after iodine prophylaxis. *Acta Endocrinol (Copenh)* 1985; 68: 55-60.
- 32 Zhang JY, Li SM, Leng JL, Chen YJ, Pu J, Li JM, et al. Changes of the spectrum on thyroid disease after the ten-year implementation of universal salt iodization in Guangxi Zhuang Autonomous Region. *Zhonghua Liu Xing Bing Pang FXXue Za Zhi* 2013 Oct; 34(10): 970-4. (abstract).
- 33 Delshad H, Mehran L, Sheikholeslam R, Mirmiran P, Azizi F. Assessment of goiter prevalence and urine iodine level in 7-10 years old school children of Tehran state in the year 2001. *The Journal of Research in Medicine* 2006; 30: 177-81.
- 34 Marwaha RK, Tandon N, Ganie MA, Kanwar R, Sastry A, Garg MK, et al. Status of thyroid function in Indian adults: two decades after universal salt iodization. *J Assoc Physicians India* 2012 Apr; 60: 32-6.
- 35 Naghavi M, Rezai Ashtiani A, Sheikholeslam R, Tashakkori N, Haiforouh S, Azizi F. Assessment of iodine efficacy situation in 7-10 years old school children of Markazi state in the year 2001. *Arak Medical University Journal of Science & Research* 2005; 4: 43-9.
- 36 Mehran L, Sheikholeslam R, Samadpour K, Hajipour R, Soleimani B, Khadivi R, et al. Assessment of goiter prevalence & urine iodine level in 7-10 years old school children of Chahar mahal bakhtiari state in the year 2001. *Iran Journal of Endocrinology & metabolism* 2006; 8: 9-15.
- 37 Hedayati M, Sheikholeslam R, Chaman R, Soleimani B, Padyab M, Azizi F. Assessment of goiter prevalence & urine iodine level in 7-10 years old school children of Kakhiluye va buyer ahmad state in the year 2001. *The Journal of Research in Medicine* 2005; 2: 113-17.
- 38 Hedayati M, Mirmiran P, Hatamizade P, Jafarzade H, Hajipour R, Azizi F. Goiter prevalence and urine iodine level in 7-10 years old school children of Ardebil state in the year 2001, 17 years after salt iodization. *Ardebil Medical University Journal of Research & Science* 2005; 6: 300-5.
- 39 Mirmiran P, Amouzgar A, Azizi F, Vasei M, Hedayati M, Mehrabi Y. Assessment of goiter prevalence and urine iodine level in 8-10 years old school children of Tehran state in the year 2007, 17 years after salt iodization. *The Journal of Research in Medicine* 2005; 40: 153-8.
- 40 Honarpisheh A, Taghavi A, Akbari H. Association between goiter prevalence and urine iodine levels in 6-12-year-old children in central Iran. *Iran J Med Sci* 2009; 34: 242-5.
- 41 Honarpishe A, Razavizade R. Goiter prevalence in the school children of mountainous region of kashan in the year 1997. *The Science & Research Journal of Feiz* 1999; 11: 94-7.
- 42 Soleimanizade L, Salehi L. The prevalence & etiology of goiter in school children of Bandar Abbas city in the year 2000. *The Journal of Medical University of Shahre kord city* 2002; 4: 40-7.
- 43 Mozaffari Khosravi H, Dehghani A, Afkhani M. Goiter prevalence and urine iodine in 6-11 years old school children of Yazd city in the year 2002. *The Scientific Journal of Medical University of Rafsanjan* 2004; 2: 96-103.