ON BOUND STATES FOR SYSTEMS OF WEAKLY COUPLED SCHRÖDINGER EQUATIONS IN ONE SPACE DIMENSION

MICHAEL MELGAARD

ABSTRACT. We establish the Birman-Schwinger relation for a class of Schrödinger operators $-d^2/dx^2 \otimes 1_{\mathcal{H}} + V$ on $L^2(\mathbb{R}, \mathcal{H})$, where $\mathcal{H}$ is an auxiliary Hilbert space and V is an operator-valued potential. As an application we give an asymptotic formula for the bound states which may arise for a weakly coupled Schrödinger operator with a matrix potential (having one or more thresholds). In addition, for a two-channel system with eigenvalues embedded in the continuous spectrum we show that, under a small perturbation, such eigenvalues turn into resonances.

1. INTRODUCTION

In a recent paper [22] (see also [21]) we studied spectral and scattering theory for the two-channel Schrödinger operator

$$\mathbf{H} = \tilde{\mathbf{H}}_0 + \mathbf{V} = \begin{pmatrix} -\frac{d^2}{dx^2} & 0 \\ 0 & -\frac{d^2}{dx^2} + 1 \end{pmatrix} + \begin{pmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{pmatrix} \tag{1.1}$$

on the Hilbert space $L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$. In the low-energy limit, where the spectral parameter tends to the boundary point of the continuous spectrum of $\mathbf{H}$, viz. the point zero, we deduced asymptotic expansions for the resolvent of $\mathbf{H}$ and, as an application, we obtained asymptotic expansions for the scattering matrix associated with the pair $(\mathbf{H}, \tilde{\mathbf{H}}_0)$ as the energy parameter tends to zero. Besides being interesting from the mathematical point of view, the study of spectral and scattering theory for $\mathbf{H}$, having thresholds at 0 and 1, also works as a useful exercise towards analogous investigations for various multichannel quantum system with more than one threshold (see, e.g., [23]) because it describes many actual physical phenomena to a good approximation.

If we replace $\tilde{\mathbf{H}}_0$ in (1.1) by $\mathbf{H}_0 = -d^2/dx^2 \otimes 1_{\mathbb{C}^N}$ and $\mathbf{V}$ by an $N \times N$ matrix potential, we obtain the (usual) matrix Schrödinger operator on $L^2(\mathbb{R}, \mathbb{C}^N)$ having a single threshold at 0. The latter, of course, has attracted a lot of attention during the years. Among recent results we mention low-energy asymptotics for the corresponding scattering

To appear in Journal of Mathematical Physics **43**, no. 11 (2002).

The author is a Marie Curie Post-Doc Fellow, supported by the European Union under grant no. HPMF-CT-2000-00973.

matrix [2, 3], Levinson's theorem [14], Lieb-Thirring inequalities [20, 4] and quantum design [7].

A natural question, which seems not to have been addressed in the literature, concerns how negative energy levels may arise in a system of weakly coupled Schrödinger equations. In the scalar-valued setting, weakly coupled bound states for Schrödinger operators have been investigated in various dimensions (see [19, Chapter VI] and [31, 15, 16]).

In this work we generalize the scalar-valued result obtained by Simon in dimension one [31] to the analogous matrix-valued setting.

We begin in the more abstract framework of Schrödinger operators with operator-valued potentials given formally by $H = -d^2/dx^2 \otimes 1_{\mathcal{H}} + V$ on $L^2(\mathbb{R}, \mathcal{H})$, where $\mathcal{H}$ is an auxiliary Hilbert space and the potential V is a $\mathcal{B}(\mathcal{H})$-valued, measurable function on $\mathbb{R}$ such that $V(x)$ is symmetric for almost all x. In Section 3 we define the Hamiltonian H by means of quadratic forms (Proposition 3.1) and in Section 4 we establish the celebrated Birman-Schwinger relation (Proposition 4.2), which transforms the eigenvalue problem for H into an eigenvalue problem for a compact operator; the so-called Birman-Schwinger operator.

Equipped with the Birman-Schwinger relation we study weakly coupled bound states in Section 5. We restrict our attention to Schrödinger operators with matrix-valued potentials. In Section 5.1 we consider two-channel Hamiltonians with one and two thresholds, resp. First we consider $\mathbf{H}(g) = -d^2/dx^2 \otimes 1_{\mathbb{C}^2} + g\mathbf{V}(x)$, where $\mathbf{V}$ is a 2×2 matrix potential. Theorem 5.2 reveals how non-positive eigenvalues of an auxiliary matrix $\mathbf{S}$, defined in (5.2), give rise to negative eigenvalues E_{ij} of $\mathbf{H}(g)$ provided g is small enough. The eigenvalues E_{ij} satisfy an asymptotic perturbation formula in which we derive the first few coefficients explicitly (see (5.3)). Second, we consider the above-mentioned Hamiltonian (1.1), henceforth denoted $\tilde{\mathbf{H}}(g)$, having thresholds at 0 and 1. In Theorem 5.6 we show how a negative eigenvalue of an auxiliary matrix $\tilde{\mathbf{S}}$, defined in (5.9), generates a negative eigenvalue of $\tilde{\mathbf{H}}(g)$. However, if one compares the proofs of Theorems 5.2 and 5.6 (in particular, the expressions for the matrices $\mathbf{T}_0$ and $\tilde{\mathbf{T}}_0$), it seems that the argument used in the proof of Theorem 5.2(ii), cannot be modified in order to treat the situation where zero is an eigenvalue of $\tilde{\mathbf{S}}$. Thus, it remains an attractive open problem to show that the zero eigenvalue of $\tilde{\mathbf{S}}$ (may) gives rise to a negative eigenvalue of $\tilde{\mathbf{H}}(g)$. In Section 5.2 we state the generalization of Theorem 5.2 to the N-channel Hamiltonian $-d^2/dx^2 \otimes 1_{\mathbb{C}^N} + \mathbf{V}(x)$, where $\mathbf{V}$ is an $N \times N$ matrix potential.

Having studied how negative eigenvalues arise for multichannel Hamiltonians under weak coupling, it is natural to address the problem of perturbation of embedded eigenvalues for a multichannel Schrödinger operator with a matrix-valued potential. In Section 6 we consider a two-channel Hamiltonian having eigenvalues embedded in its continuous spectrum. When perturbed by a "short range" potential, we show

that such eigenvalues move into the complex plane and become resonances. In particular, we verify Fermi's golden rule (see, e.g., [27, 32]).

There is a vast literature on 2×2 operator-valued matrices, e.g. in system theory (see e.g. [8]) and in semigroup theory (see e.g. [11]). Most notably in this context is the substantial number of questions of a general nature which have been answered on spectral theory recently, see e.g. the survey [33]. However, the methods therein are not related to ours although some of the questions addressed clearly are, e.g. the appearance of resonances discussed by Mennicken and Motovilov [24].

2. Preliminaries

1) *Vector-valued functions.* Let $\mathcal{H}$ be a separable Hilbert space with scalar product and norm denoted by $\langle \cdot, \cdot \rangle_{\mathcal{H}}$ and $\| \cdot \|_{\mathcal{H}}$. Then a function $\psi(x)$ from $\mathbb{R}$ to $\mathcal{H}$ is measurable if the scalar-valued functions $\langle \psi(x), \phi \rangle_{\mathcal{H}}$ are measurable, where ϕ denotes an arbitrary vector of $\mathcal{H}$. If $\psi(x)$ is such a measurable function, then $\|\psi(x)\|_{\mathcal{H}}$ is also measurable (as a function with non-negative values). Thus $L^p(\mathbb{R}, \mathcal{H})$ is defined as the set of equivalence classes of measurable functions $\psi(x)$ from $\mathbb{R}$ to $\mathcal{H}$, which satisfy that $\int_{\mathbb{R}} \|\psi(x)\|_{\mathcal{H}}^p dx$ is finite if $p < \infty$ and $\|\psi\|_\infty = \text{ess sup } \|\psi(x)\|_{\mathcal{H}} < \infty$ if $p = \infty$. The measure dx is the Lebesgue measure. For any p the $L^p(\mathbb{R}, \mathcal{H})$ space is a Banach space with norm $\| \cdot \|_p = (\int_{\mathbb{R}} \| \cdot \|_{\mathcal{H}}^p dx)^{1/p}$. In the case $p = 2$, $L^2(\mathbb{R}, \mathcal{H})$ is a complex and separable Hilbert space with scalar product $\langle \phi, \psi \rangle_2 = \int_{\mathbb{R}} \langle \phi, \psi \rangle_{\mathcal{H}} dx$ and corresponding norm $\|\psi\|_2 = \langle \psi, \psi \rangle_2^{1/2}$. For $n \in \mathbb{N}$, $1 \leq p < \infty$, the Sobolev space $W^{n,p}(\mathbb{R}, \mathcal{H})$ is defined as the space of those $\psi \in L^p(\mathbb{R}, \mathcal{H})$, for which all derivatives (weak sense) up to order n are in $L^p(\mathbb{R}, \mathcal{H})$. If $p = 2$, $W^{n,2}(\mathbb{R}, \mathcal{H})$ is a separable Hilbert space denoted by $H^n(\mathbb{R}, \mathcal{H})$ with scalar product $\langle \phi, \psi \rangle_{H^n(\mathbb{R}, \mathcal{H})} = \int_{\mathbb{R}} \sum_{\alpha=0}^n \langle (d/dx)^\alpha \phi, (d/dx)^\alpha \psi \rangle_{\mathcal{H}}$ and norm denoted by $\|\psi\|_{H^n(\mathbb{R}, \mathcal{H})}$.

2) *Operators.* Below $\mathcal{H}$, $\mathcal{H}_1$, $\mathcal{H}_2$ are separable Hilbert spaces. For a linear operator T, the notations $\mathcal{D}(T)$, $\text{Ran}\,(T)$, $\text{Ker}\,(T)$, T^*, $\overline{T}$, $\sigma(T)$, $\rho(T)$ are standard, see for example [25]. By I we denote the identity operator. The resolvent of a self-adjoint operator T is denoted by $R(T, z) = (T - zI)^{-1}$. By $\mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ and $\mathbf{S}_\infty(\mathcal{H}_1, \mathcal{H}_2)$ we denote respectively the sets of bounded and compact operators acting from $\mathcal{H}_1$ into $\mathcal{H}_2$. With the usual operator norm $\mathcal{B}(\mathcal{H}_1, \mathcal{H}_2)$ is a Banach space. We set $\mathcal{B}(\mathcal{H}) := \mathcal{B}(\mathcal{H}, \mathcal{H})$ and $\mathbf{S}_\infty(\mathcal{H}) := \mathbf{S}_\infty(\mathcal{H}, \mathcal{H})$.

3) *Trace classes of compact operators.* If $T \in \mathbf{S}_\infty(\mathcal{H})$ then the non-zero eigenvalues of $|T| = \sqrt{T^*T}$ are called the singular numbers or s-numbers of T. Let $\{s_j(T)\}$ denote the (possibly finite) non-increasing sequence of the singular numbers of T; every number counted according to its multiplicity as an eigenvalue of $|T|$. For $0 < p < \infty$ the von Neumann-Schatten class $\mathbf{S}_p(\mathcal{H}_1, \mathcal{H}_2)$ is the set of $T \in \mathbf{S}_\infty(\mathcal{H}_1, \mathcal{H}_2)$ for

which the functional

$$\|T\|^p_{\mathbf{S}_p(\mathcal{H}_1,\mathcal{H}_2)} := \sum_j [s_j(T)]^p$$

is finite. The functional $\|\cdot\|_{\mathbf{S}_p(\mathcal{H}_1,\mathcal{H}_2)}$ is a norm for $p \geq 1$ and the normed space $\mathbf{S}_p(\mathcal{H}_1,\mathcal{H}_2)$ is a Banach space. For $p < 1$ the functional is a quasinorm. For additional properties of the spaces $\mathbf{S}_p$ of compact operators we refer [5, Chapter 11]. The sets $\mathbf{S}_1(\mathcal{H}_1,\mathcal{H}_2)$ and $\mathbf{S}_2(\mathcal{H}_1,\mathcal{H}_2)$ are called the trace class and Hilbert-Schmidt class, respectively.

4) *Operator-valued functions.* Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be two separable Hilbert spaces. From above, a function $\mathbb{R} \ni x \to \psi(x) \in \mathcal{H}$ is measurable if and only if all the functions $\mathbb{R} \ni x \to \langle \psi(x), \phi\rangle_{\mathcal{H}} \in \mathbb{C}$ are measurable. As a result of Pettis Measurability Theorem (see, e.g.,[10, Theorem II.1.2]) the following properties are equivalent for a $\mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)$-valued function $\mathbb{R} \ni x \mapsto T(x)$:
(i) $\forall \phi \in \mathcal{H}_2$, $\forall \psi \in \mathcal{H}_1$, $\mathbb{R} \ni x \to \langle \phi, T(x)\psi\rangle_{\mathcal{H}_2} \in \mathbb{C}$ is measurable,
(ii) $\forall \psi \in \mathcal{H}_1$, $\mathbb{R} \ni x \to T(x)\psi \in \mathcal{H}_2$ is measurable.
We say that a function $\mathbb{R} \ni x \mapsto T(x) \in \mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)$ is measurable if it satisfies any one of the above properties (i)-(ii). In the affirmative case, $\|T(x)\|_{\mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)}$ is also measurable because

$$\|T(x)\|_{\mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)} = \sup_{\psi\in\mathcal{D}_1} \left(\|T(x)\psi\|_{\mathcal{H}_2}/\|\psi\|_{\mathcal{H}_1}\right),$$

where $\mathcal{D}_1$ is a countable dense subset of $\mathcal{H}_1$. Moreover, we can define $L^p(\mathbb{R}, \mathcal{B}(\mathcal{H}_1,\mathcal{H}_2))$ as the linear space of (equivalence classes of) measurable functions $T : \mathbb{R} \to \mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)$ such that $\|T(\cdot)\|_{\mathcal{B}(\mathcal{H}_1,\mathcal{H}_2)} \in L^p(\mathbb{R})$.

For the functional calculus for self-adjoint operators we recall the following result which can be found in, e.g., [6, Proposition V.1.2].

Proposition 2.1. *If for each $x \in \mathbb{R}$, $T(x)$ is a self-adjoint operator on $\mathcal{H}$ and $\{E_{T(x)}(A); A$ Borel set of $\mathbb{R}\}$ denotes its resolution of the identity, the following three properties are equivalent:*
(i) $\mathbb{R} \ni x \to E_{T(x)}(A) \in \mathcal{B}(\mathcal{H})$ is measurable for all Borel sets A,
(ii) $\mathbb{R} \ni x \to e^{-itT(x)} \in \mathcal{B}(\mathcal{H})$ is measurable for all $t \in \mathbb{R}$,
(iii) $\mathbb{R} \ni x \to (T(x) - \zeta)^{-1} \in \mathcal{B}(\mathcal{H})$ is measurable for all $\zeta \in \mathbb{C}\backslash\mathbb{R}$.

5) *Fourier transform.* Suppose $\psi \in L^1(\mathbb{R}^d, \mathcal{H})$. Then we define its Fourier transform $(\mathcal{F}\psi)(\xi) = \widehat{\psi}(\xi) := (2\pi)^{-1/2} \int_{\mathbb{R}} e^{ix\xi}\psi(x)\, dx$ which is an element of $L^\infty(\mathbb{R}, \mathcal{H})$. If $\psi \in L^1(\mathbb{R}, \mathcal{H}) \cap L^2(\mathbb{R}, \mathcal{H})$, then $\widehat{\psi} \in L^2(\mathbb{R}, \mathcal{H})$ with $\|\widehat{\psi}\|_{L^2} = \|\psi\|_{L^2}$. The Fourier transform can then be extended by continuity to a unitary mapping of the Hilbert space $L^2(\mathbb{R}, \mathcal{H})$ into itself.

We have the following criterion.

Lemma 2.2. *Let T be an operator on $L^2(\mathbb{R}, \mathcal{H})$ defined by*

$$(T\phi)(x) = \int_{\mathbb{R}} t(x,\xi)\phi(\xi)\, d\xi, \tag{2.1}$$

where $t(x,\xi) \in \mathcal{B}(\mathcal{H})$ for each (x,ξ). Then T is a Hilbert-Schmidt operator on $L^2(\mathbb{R},\mathcal{H})$ if and only if

$$\int_{\mathbb{R}_x}\int_{\mathbb{R}_\xi} \mathrm{tr}_{\mathcal{H}}[t(x,\xi)^* t(x,\xi)]\, d\xi dx < \infty.$$

In this case,

$$\|T\|^2_{\mathbf{S}_2(L^2(\mathbb{R},\mathcal{H}))} = \int_{\mathbb{R}_x}\int_{\mathbb{R}_\xi} \mathrm{tr}_{\mathcal{H}}[t(x,\xi)^* t(x,\xi)]\, d\xi dx.$$

Proof. The Hilbert space $\mathcal{H}$ is isomorphic to some $L^2(Y)$ space and therefore it suffices to establish the statement for an operator T on $L^2(\mathbb{R},L^2(Y))$ defined by (2.1) for some $t(x,\xi) \in \mathcal{B}(L^2(Y))$. Since $L^2(\mathbb{R},L^2(Y))$ is isomorphic to $L^2(\mathbb{R}\times Y)$, the rephrased assertion follows immediately from [25, Theorem VI.23]. □

3. The Hamiltonian $H = H_0 + V$

As in the scalar-valued case the quadratic form

$$h_0[\psi,\psi] := \int_{\mathbb{R}} \|(d/dx)\psi(x)\|^2_{\mathcal{H}}\, dx \tag{3.1}$$

is closed in $L^2(\mathbb{R},\mathcal{H})$ on the domain $H^1(\mathbb{R},\mathcal{H})$. Thus, this form generates a self-adjoint operator H_0 on $L^2(\mathbb{R},\mathcal{H})$. The free Hamiltonian H_0 corresponds to the "Laplacian" $-d^2/dx^2 \otimes 1_{\mathcal{H}}$ on $L^2(\mathbb{R},\mathcal{H})$.

A potential V is a $\mathcal{B}(\mathcal{H})$-valued, measurable function on $\mathbb{R}$. Assume that $V(x)$ is symmetric for almost all x, i.e. $V(x)^* = V(x)$ for almost all x. The operator $V(x) \in \mathcal{B}(\mathcal{H})$ has a unique representation[1] in the form $V(x) = U(x)|V(x)|$, where $|V(x)|$ is the modulus of $V(x)$ defined by $|V(x)| = (V(x)^*V(x))^{1/2} = (V(x)V(x))^{1/2}$. We have that $|V(x)|$ is a non-negative, selfadjoint operator belonging to $\mathcal{B}(\mathcal{H})$ and, moreover, $\||V(x)|\|_{\mathcal{B}(\mathcal{H})} = \|V(x)\|_{\mathcal{B}(\mathcal{H})}$. The operator $U(x)$ is a partial isometry with initial domain $\overline{\mathrm{Ran}\,|V(x)|}$, final domain $\overline{\mathrm{Ran}\,V(x)}$ and $\mathrm{Ker}\,U(x) = \mathrm{Ker}\,V(x)$. Observe that $U(x)^*U(x) = P_{\overline{\mathrm{Ran}\,|V(x)|}}$ and $U(x)U(x)^* = P_{\overline{\mathrm{Ran}\,V(x)}}$, where P_M denotes the orthogonal projection onto a closed subspace M. The modulus $|V(x)|$ possesses exactly one non-negative, self-adjoint square-root $|V(x)|^{1/2} \in \mathcal{B}(\mathcal{H})$. The square-root $|V(x)|^{1/2}$ commutes with every bounded operator which commutes with $|V(x)|$. We may define $V(x)^{1/2} = U(x)|V(x)|^{1/2}$ such that $V(x) = V(x)^{1/2}|V(x)|^{1/2}$. Moreover, $V(x)^{1/2} \in \mathcal{B}(\mathcal{H})$ with $\|V(x)^{1/2}\|_{\mathcal{B}(\mathcal{H})} = [\|V(x)\|_{\mathcal{B}(\mathcal{H})}]^{1/2}$ and adjoint $(V(x)^{1/2})^* = |V(x)|^{1/2}U(x)^*$. From Proposition 2.1 it follows that $|V|$, $|V|^{1/2}$ and $V^{1/2}$ are $\mathcal{B}(\mathcal{H})$-valued measurable functions on $\mathbb{R}$.

We want to establish the following result.

[1]The representation is not unique if the potential vanishes on a set of positive measure

Proposition 3.1.
(i) If $V \in L^1(\mathbb{R}, \mathcal{B}(\mathcal{H}))$ then the real-valued quadratic form

$$v[\psi, \psi] := \int_{\mathbb{R}} \langle V(x)^{1/2}\psi(x), |V(x)|^{1/2}\psi(x)\rangle_{\mathcal{H}}\, dx$$

is H_0 form-bounded with relative bound zero.
(ii) If $V \in L^1(\mathbb{R}, \mathbf{S}_2(\mathcal{H}))$ then v is H_0 form-compact.

It follows from Proposition 3.1(i) and the KLMN theorem [26, Theorem X.17] that the form sum

$$h[\psi, \psi] := h_0[\psi, \psi] + v[\psi, \psi]$$

is closed and semi-bounded from below on $H^1(\mathbb{R}^d, \mathcal{H})$ and thus generates a self-adjoint operator $H = H_0 + V$ on $L^2(\mathbb{R}, \mathcal{H})$. From Proposition 3.1(ii) and Weyl's essential spectrum theorem it follows that $\sigma_{ess}(H) = \sigma_{ess}(H_0) = [0, \infty)$.

Proof of Proposition 3.1. The "kernel" of the resolvent of H_0 is given by (see, e.g., [28, Theorem 9.5.2])

$$Q(x - y; \sqrt{|E|}) = \frac{e^{-\sqrt{|E|}|x-y|}}{2\sqrt{|E|}}, \ E < 0. \tag{3.2}$$

(i) To show that the form v is infinitesimally H_0 form-bounded, it suffices to show that the form

$$w[\phi] = \langle |V|^{1/2}(H_0 - E)^{-1/2}\phi, V^{1/2}(H_0 - E)^{-1/2}\phi\rangle_{L^2(\mathbb{R},\mathcal{H})}$$

is bounded on $L^2(\mathbb{R}, \mathcal{H})$ and that its norm

$$\|w\| := \inf_{\phi \in L^2(\mathbb{R},\mathcal{H})} \frac{|\langle |V|^{1/2}(H_0 - E)^{-1/2}\phi, V^{1/2}(H_0 - E)^{-1/2}\phi\rangle|}{\|\phi\|^2}$$

tends to zero as $E \to -\infty$. By the definition of $\|w\|$, and since U in $V^{1/2} = U|V|^{1/2}$ is a partial isometry, we have that

$$\|w\| \leq \left\| |V|^{1/2}(H_0 - E)^{-1/2} \right\|^2_{\mathcal{B}(L^2(\mathbb{R},\mathcal{H}))}. \tag{3.3}$$

Therefore, it is enough to show that the right-hand side of the latter tends to zero as $E \to -\infty$.

We consider first $V \in L^\infty(\mathbb{R}, \mathcal{B}(\mathcal{H}))$. For such V we have that

$$\left\| |V|^{1/2}(H_0 - E)^{-1/2} \right\|^2_{\mathcal{B}(L^2)} = \left\| |V|^{1/2}(H_0 - E)^{-1}|V|^{1/2} \right\|_{\mathcal{B}(L^2)} \tag{3.4}$$

Let $\alpha = \sqrt{|E|}$ and $\phi \in L^2(\mathbb{R}, \mathcal{H})$. Then Hölder's inequality yields that

$$\begin{aligned} &\left\| [|V|^{1/2}(H_0 + \alpha^2)^{-1}|V|^{1/2}\phi](x) \right\|_{\mathcal{H}} \\ &\quad \leq \frac{1}{2\alpha} \left\| V(x) \right\|^{1/2}_{\mathcal{B}(\mathcal{H})} \left\| V \right\|_{L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))} \left\| \phi \right\|_{L^2(\mathbb{R},\mathcal{H})}. \end{aligned}$$

The latter implies that

$$\begin{aligned}
&\left\| |V|^{1/2}(H_0+\alpha^2)^{-1}|V|^{1/2}\phi\right\|^2_{L^2(\mathbb{R},\mathcal{H})} \\
&\quad\le \int_{\mathbb{R}} \frac{1}{4\alpha^2}\,\|V(x)\|_{\mathcal{B}(\mathcal{H})}\,\|V\|^2_{L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))}\,\|\phi\|^2_{L^2(\mathbb{R},\mathcal{H})}\,dx \\
&\quad\le \frac{1}{4\alpha^2}\|V\|^3_{L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))}\|\phi\|^2_{L^2(\mathbb{R},\mathcal{H})}.
\end{aligned}$$

In conjunction with (3.4), the latter shows that the right-hand side of (3.3) tends to zero as $E\to-\infty$, which establishes assertion (i) for $V\in L^\infty(\mathbb{R},\mathcal{H})$. A standard approximation argument yields the assertion for general V.

(ii). It suffices to show that the form

$$w[\phi]=\langle |V|^{1/2}(H_0-E)^{-1/2}\phi, V^{1/2}(H_0-E)^{-1/2}\phi\rangle$$

defines a compact operator in $L^2(\mathbb{R},\mathcal{H})$. Under the assumption in (i) we already know that w generates a bounded, self-adjoint operator W in $L^2(\mathbb{R},\mathcal{H})$. Let us show that W is a Hilbert-Schmidt operator. From

$$\mathrm{tr}_{L^2}(W^*W)=\mathrm{tr}\left(V^{1/2}(H_0-E)^{-1}(V^{1/2})^*|V|^{1/2}(H_0-E)^{-1}(|V|^{1/2})^*\right),$$

we see that it is enough to show that $W_1=|V|^{1/2}(H_0-E)^{-1}(|V|^{1/2})^*$ and $W_2=V^{1/2}(H_0-E)^{-1}(V^{1/2})^*$ are Hilbert-Schmidt operators on $L^2(\mathbb{R},\mathcal{H})$. It is enough to show it for W_2; the proof for W_1 is similar. The operator W_2 has integral "kernel"

$$K_{W_2}(x-y;\alpha)=V(x)^{1/2}(2\alpha)^{-1}e^{-\alpha|x-y|}(V(y)^{1/2})^*,\quad \alpha=\sqrt{-E}>0.$$

Using the criterion in Lemma 2.2 and the assumption in (ii), we estimate as follows:

$$\begin{aligned}
&\int_{\mathbb{R}}\int_{\mathbb{R}}\mathrm{tr}_{\mathcal{H}}[K_{W_2}(x-y;\alpha)^*K_{W_2}(x-y;\alpha)]\,dxdy=\int_{\mathbb{R}}\int_{\mathbb{R}}\left(\frac{e^{-\alpha|x-y|}}{2\alpha}\right)^2 \\
&\qquad\times\mathrm{tr}_{\mathcal{H}}[(V(y)^{1/2})^{**}(V(x)^{1/2})^*V(x)^{1/2}(V(y)^{1/2})^*]\,dxdy \\
&=\int_{\mathbb{R}}\int_{\mathbb{R}}\left(\frac{e^{-\alpha|x-y|}}{2\alpha}\right)^2 \\
&\times\mathrm{tr}_{\mathcal{H}}[V(y)^{1/2}|V(x)|^{1/2}U(x)^*U(x)|V(x)|^{1/2}(V(y)^{1/2})^*]\,dxdy \\
&=\int_{\mathbb{R}}\int_{\mathbb{R}}\left(\frac{e^{-\alpha|x-y|}}{2\alpha}\right)^2\mathrm{tr}_{\mathcal{H}}[|V(x)||V(y)|]\,dxdy \\
&\le\frac{1}{4\alpha^2}\int_{\mathbb{R}}\|V(x)\|_{\mathbf{S}_2(\mathcal{H})}\,dx\int_{\mathbb{R}}\|V(y)\|_{\mathbf{S}_2(\mathcal{H})}\,dy.
\end{aligned}$$

This shows that W_2 is a Hilbert-Schmidt operator in $L^2(\mathbb{R},\mathcal{H})$. □

We note that $V\in L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))$ implies that $|V|\in L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))$ and, in view of Proposition 3.1(i), $|V|$ is infinitesimally H_0 form-bounded.

Consequently, the following mappings are bounded:

$$V,\ |V| \ : \ H^1(\mathbb{R},\mathcal{H}) \to H^{-1}(\mathbb{R},\mathcal{H}) \tag{3.5}$$
$$|V|^{1/2}, V^{1/2} \ : \ H^1(\mathbb{R},\mathcal{H}) \to L^2(\mathbb{R},\mathcal{H}) \tag{3.6}$$
$$|V|^{1/2}, V^{1/2} \ : \ L^2(\mathbb{R},\mathcal{H}) \to H^{-1}(\mathbb{R},\mathcal{H}). \tag{3.7}$$

The qualitative behaviour of any possible negative eigenvalues of $H_0 + gV$ as $g \to 0$ is described by the following simple result.

Proposition 3.2. *If $V \in L^1(\mathbb{R}, \mathcal{B}(\mathcal{H}))$ then any negative eigenvalues of $H_0 + gV$ approach zero as g tends to zero.*

Proof. Following [31] it suffices to show that there are positive constants g_0 and C such that $H_0 + gV \geq -Cg$ for all $g_0 > g > 0$.

Let $\mathcal{F}$ denote the Fourier transform of vector-valued functions in $L^2(\mathbb{R},\mathcal{H})$ (see Part 5 in Section 2). We observe that, as for scalar-valued functions, a function ϕ whose Fourier transform is integrable is bounded and continuous with the usual estimate

$$\|\phi(x)\|_{\mathcal{H}} \leq \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} \|\mathcal{F}\phi(\xi)\|_{\mathcal{H}}\, d\xi. \tag{3.8}$$

For an arbitrary $\gamma > 0$, Hölder's inequality yields that

$$\begin{aligned}
&\left(\int_{\mathbb{R}} \|\mathcal{F}\phi(\xi)\|_{\mathcal{H}}\, d\xi\right)^2 \\
&\qquad \leq \left(\int_{\mathbb{R}} (\xi^2+\gamma^2)^{-1}\, d\xi\right)\left(\int_{\mathbb{R}} (\xi^2+\gamma^2)\|\mathcal{F}\phi(\xi)\|_{\mathcal{H}}^2\, d\xi\right) \\
&\qquad = \frac{\pi}{\gamma}\|(-i\xi+\gamma)\mathcal{F}\phi\|^2_{L^2(\mathbb{R},\mathcal{H})} = \frac{\pi}{\gamma}\|(H_0^{1/2}+\gamma)\phi\|^2_{L^2(\mathbb{R},\mathcal{H})} \\
&\qquad \leq \frac{2\pi}{\gamma}\left\{\|H_0^{1/2}\phi\|^2_{L^2(\mathbb{R},\mathcal{H})} + \|\gamma\phi\|^2_{L^2(\mathbb{R},\mathcal{H})}\right\}.
\end{aligned} \tag{3.9}$$

Let $d = \max\{1/\gamma, 1\}$. Then (3.8) and (3.9) imply that

$$\|\phi(x)\|_{\mathcal{H}} \leq d\left\{\|H_0^{1/2}\phi\|^2_{L^2(\mathbb{R},\mathcal{H})} + \|\phi\|^2_{L^2(\mathbb{R},\mathcal{H})}\right\} \tag{3.10}$$

for any $\phi \in \mathcal{D}(H_0^{1/2}) = H^1(\mathbb{R},\mathcal{H})$. The Sobolev type inequality (3.10) implies that

$$\begin{aligned}
h_g[\phi] \ &\geq \ \|H_0^{1/2}\|^2_{L^2(\mathbb{R},\mathcal{H})} - g\int_{\mathbb{R}} \|V(x)\|_{\mathcal{B}(\mathcal{H})}\|\phi(x)\|^2_{\mathcal{H}}\, dx \\
&\geq \ (1-gd_1)\|H_0^{1/2}\|^2_{L^2(\mathbb{R},\mathcal{H})} - gd_1\|\phi\|^2_{L^2(\mathbb{R},\mathcal{H})}
\end{aligned}$$

where $d_1 = d\|V\|^2_{L^1(\mathbb{R},\mathcal{B}(\mathcal{H}))}$. When we take $C = d_1$, $g_0 = d_1^{-1}$ and $0 < g < g_0$, we arrive at $h_g[\phi] \geq -Cg$ as desired. □

4. The Birman-Schwinger relation

The Birman-Schwinger relation has been established rigorously for various classes of operators in the scalar-valued setting (see, e.g., [30, 31, 17]. It asserts that E is a negative eigenvalue of $H = -d^2/dx^2 + V$ if and only if -1 is an eigenvalue of the operator $V^{1/2}(-d^2/dx^2 - E)^{-1}|V|^{1/2}$. Formally this is obvious since $\phi = V^{1/2}\psi$ is a solution to $V^{1/2}(-d^2/dx^2 - E)^{-1}|V|^{1/2}\phi = -\phi$.

Here we provide a simple proof of the Birman-Schwinger relation in our concrete operator-valued setting. For this purpose we introduce the Birman-Schwinger operator $K_E(V) = V^{1/2}(H_0 - E)^{-1}|V|^{1/2}$, $E < 0$, where H_0 is the nonnegative, self-adjoint operator associated with the quadratic form h_0 in (3.1). Setting $\alpha^2 = -E$, its integral "kernel" is given by

$$K_\alpha(x, y) = V(x)^{1/2}(2\alpha)^{-1}e^{-\alpha|x-y|}|V(y)|^{1/2}, \quad \alpha > 0.$$

We have the following result.

Lemma 4.1. *If $V \in L^1(\mathbb{R}, \mathbf{S}_2(\mathcal{H}))$ then the Birman-Schwinger operator $K_E(V)$, $E < 0$, is a Hilbert-Schmidt operator on $L^2(\mathbb{R}, \mathcal{H})$; in particular $K_E(V)$ is a compact operator. Moreover, $\|K_E(V)\|_{\mathcal{B}(L^2(\mathbb{R},\mathcal{H}))} \to 0$ as $E \to -\infty$.*

Proof. We argue as for the operator W_2 in the proof of Proposition 3.1(ii). We omit the details. □

Having introduced the compact Birman-Schwinger operator we may formulate the Birman-Schwinger relation.

Proposition 4.2. *Let $V \in L^1(\mathbb{R}, \mathbf{S}_2(\mathcal{H}))$. Then $E < 0$ is an eigenvalue of $H = H_0 + V$ (defined by a quadratic form) having multiplicity $\varkappa$ if and only if -1 is an eigenvalue of $K_E(V)$ having geometric multiplicity $\varkappa$.*

To establish Proposition 4.2 we need the following two results.

Lemma 4.3. *If $V \in L^1(\mathbb{R}, \mathcal{B}(\mathcal{H}))$ then for $E < 0$ the operators $|V|^{1/2} \times (H_0 - E)^{-1/2}$ and $V^{1/2}(H_0 - E)^{-1/2}$ are bounded on $L^2(\mathbb{R}, \mathcal{H})$.*

Proof. Since V is H_0 form-bounded, it follows immediately from (3.6) and (3.7) in conjunction with the fact that the operator $(H_0 - E)^{-1/2}$ is a bounded map from the domain $L^2(\mathbb{R}, \mathcal{H})$ to the range $H^1(\mathbb{R}, \mathcal{H})$. □

Lemma 4.4. *Let S and T be bounded operators on the Hilbert space $\mathcal{K}$. Then $\sigma(ST)\backslash\{0\} = \sigma(TS)\backslash\{0\}$. Moreover, $\lambda \neq 0$ is an eigenvalue of ST having geometric multiplicity m if and only if λ is an eigenvalue of TS having geometric multiplicity m.*

Proof. This is a simplified version of Theorem 2(i) in [9]. □

Proof of Proposition 4.2. Let h_0 be the form of H_0, let v be the form of V and let $h = h_0 + v$ be their form sum. According to Lemma 4.3

the operators $|V|^{1/2}(H_0-E)^{-1/2}$ and $V^{1/2}(H_0-E)^{-1/2}$ are bounded on $L^2(\mathbb{R},\mathcal{H})$ and, consequently, the operator $I+[|V|^{1/2}(H_0-E)^{-1/2}]^*V^{1/2}$ $(H_0-E)^{-1/2}$ is bounded on $L^2(\mathbb{R},\mathcal{H})$. Moreover, the operator $A^{-1}=(H_0-E)^{1/2}$ has domain $H^1(\mathbb{R},\mathcal{H})$ and range $L^2(\mathbb{R},\mathcal{H})$. Thus, we may introduce an auxiliary sesquilinear form a defined on the form domain $H^1(\mathbb{R},\mathcal{H})\times H^1(\mathbb{R},\mathcal{H})$ by

$$a[\phi,\psi]=\left\langle\left[I+(|V|^{1/2}A)^*V^{1/2}A\right]A^{-1}\phi,A^{-1}\psi\right\rangle. \tag{4.1}$$

We re-write a and find that

$$a[\phi,\psi]=\underbrace{\langle A^{-1}\phi,A^{-1}\psi\rangle}_{a_1[\phi,\psi]}+\underbrace{\langle(|V|^{1/2}A)^*V^{1/2}AA^{-1}\phi,A^{-1}\psi\rangle}_{a_2[\phi,\psi]}. \tag{4.2}$$

Clearly,

$$a_1[\phi,\psi]=h_0[\phi,\psi]-E\langle\phi,\psi\rangle \tag{4.3}$$

and, since $|V|^{1/2}A$ is bounded on $L^2(\mathbb{R},\mathcal{H})$,

$$\begin{aligned}a_2[\phi,\psi] &= \langle V^{1/2}AA^{-1}\phi,[|V|^{1/2}A]^{**}A^{-1}\psi\rangle\\ &= \langle V^{1/2}\phi,|V|^{1/2}AA^{-1}\psi\rangle=v[\phi,\psi].\end{aligned} \tag{4.4}$$

Hence, (4.2)-(4.4) shows that the forms a and $h-E$ are identical.

Suppose that $E<0$ is an eigenvalue of $H=H_0+V$, i.e. there exists an eigenfunction $\psi\in\mathcal{D}(H)$, $\psi\neq 0$, such that $(H-E)\psi=0$. This is equivalent to $(h-E)[\psi,\phi]=0$ for all $\phi\in H^1(\mathbb{R},\mathcal{H})$. Since the forms h and a are identical, we may introduce $u=A^{-1}\phi$ and deduce that

$$\begin{aligned}0=a[\psi,\phi] &= \left\langle\left[I+(|V|^{1/2}A)^*V^{1/2}A\right]A^{-1}\psi,A^{-1}\phi\right\rangle,\ \forall\phi\in H^1(\mathbb{R},\mathcal{H}),\\ &= \left\langle\left[I+(|V|^{1/2}A)^*V^{1/2}A\right]A^{-1}\psi,u\right\rangle,\ \forall\phi\in L^2(\mathbb{R},\mathcal{H}),\end{aligned}$$

because u runs through $L^2(\mathbb{R},\mathcal{H})$ as ϕ runs through $H^1(\mathbb{R},\mathcal{H})$. Consequently, $\left(I+[|V|^{1/2}A]^*V^{1/2}A\right)v=0$, where $v=A^{-1}\psi=(H_0-E)^{1/2}\psi$, so $-1\in\sigma_p([|V|^{1/2}A]^*V^{1/2}A)$. By reversing the arguments leading to the latter conclusion, we infer that

$$E\in\sigma_p(H_0+V)\text{ if and only if }-1\in\sigma_p([|V|^{1/2}A]^*V^{1/2}A) \tag{4.5}$$

Since $(H_0-E)^{1/2}$ is injective from the domain $L^2(\mathbb{R},\mathcal{H})$ to the range $H^1(\mathbb{R},\mathcal{H})$, the arguments above also show that the multiplicities of the eigenvalues E and -1 must be equal. In view of Lemma 4.4 and the definition of A, (4.5) implies that

$$\begin{aligned}&E\in\sigma_p(H_0+V)\text{ if and only if}\\ &\quad -1\in\sigma_p(V^{1/2}(H_0-E)^{-1/2}[|V|^{1/2}(H_0-E)^{-1/2}]^*)\end{aligned} \tag{4.6}$$

and the multiplicities of E and -1 are equal. But $[|V|^{1/2}(H_0-E)^{-1/2}]^*=(H_0-E)^{-1/2}|V|^{1/2})$ and therefore, in view of the definition of $K_E(V)$, (4.6) yields that $E\in\sigma_p(H_0+V)$ if and only if $-1\in\sigma_p(K_E(V))$. $\square$

If g is fixed and $\lambda(\alpha)$ is an eigenvalue of $K_\alpha(V)$ then the Birman-Schwinger relation asserts that any solution $\alpha_g > 0$ of

$$g\lambda(\alpha_g) = -1 \tag{4.7}$$

is associated to the eigenvalue $E(g) = -\alpha_g^2$ of $H(g)$. The latter equation plays a crucial role in Section 5.

Define the operators L_α and M_α by their "kernels":

$$L_\alpha(x,y) = \frac{1}{2\alpha} V(x)^{1/2} |V(y)|^{1/2}, \tag{4.8}$$

$$M_\alpha(x,y) = \frac{1}{2\alpha} V(x)^{1/2} [[e^{-\alpha|x-y|} - 1]|V(y)|^{1/2} \tag{4.9}$$

Moreover, we introduce the operator M_0 with "kernel"

$$M_0(x,y) = -\frac{1}{2} V(x)^{1/2} |x-y| |V(y)|^{1/2}. \tag{4.10}$$

Imitating [31] we obtain the following result.

Lemma 4.5. *If $\int_{\mathbb{R}} (1+|x|^2) \left\|V(x)\right\|_{\mathbf{S}_2(\mathcal{H})} dx < \infty$ then the following assertions are valid:*
(i) The operator M_0 is Hilbert-Schmidt on $L^2(\mathbb{R}, \mathcal{H})$.
(ii) As $\alpha \downarrow 0$, the operator M_α converges to M_0 in the Hilbert-Schmidt norm on $L^2(\mathbb{R}, \mathcal{H})$.
(iii) The Birman-Schwinger operator $gK_\alpha(V)$ has eigenvalue -1 if and only if the same is true for $g(1+gM_\alpha)^{-1}L_\alpha$.

Proof.
(i) It follows from the estimate

$$\int_{\mathbb{R}} \int_{\mathbb{R}} \mathrm{tr}_{\mathcal{H}} [M_0(x,y)^* M_0(x,y)] \, dxdy$$
$$\leq \frac{1}{2} \int \int (|x|^2 + |y|^2) \left\| |V(x)| \right\|_{\mathbf{S}_2(\mathcal{H})} \left\| |V(y)| \right\|_{\mathbf{S}_2(\mathcal{H})} dxdy < \infty.$$

(ii) We want to show that

$$\int \int \mathrm{tr}_{\mathcal{H}} \left[(M_\alpha - M_0)(x,y)^* (M_\alpha - M_0)(x,y) \right] dxdy \longrightarrow 0 \tag{4.11}$$

as $\alpha \downarrow 0$. Now,

$$\mathrm{tr}_{\mathcal{H}} \left[(M_\alpha - M_0)(x,y)^* (M_\alpha - M_0)(x,y) \right]$$
$$= \left| \frac{1}{2\alpha} (e^{-\alpha|x-y|} - 1) + \frac{1}{2}|x-y| \right|^2 \mathrm{tr}_{\mathcal{H}} \left[|V(x)||V(y)| \right],$$

and since $\left| \frac{1}{2\alpha} (e^{-\alpha|x-y|} - 1) + \frac{1}{2}|x-y| \right| \to 0$ as $\alpha \downarrow 0$, we have the pointwise convergence

$$\mathrm{tr}_{\mathcal{H}} \left[(M_\alpha - M_0)(x,y)^* (M_\alpha - M_0)(x,y) \right] \longrightarrow 0 \text{ as } \alpha \downarrow 0. \tag{4.12}$$

Moreover,

$$\operatorname{tr}_{\mathcal{H}}[M_\alpha(x,y)^* M_\alpha(x,y)] = \left|\frac{1}{2\alpha}(e^{-\alpha|x-y|}-1)\right|^2 \operatorname{tr}_{\mathcal{H}}\left[|V(x)||V(y)|\right]$$
$$\leq \left|\frac{1}{2}|x-y|\right|^2 \operatorname{tr}_{\mathcal{H}}\left[|V(x)||V(y)|\right] = \operatorname{tr}_{\mathcal{H}}[M_0(x,y)^* M_0(x,y)]. \quad (4.13)$$

It follows from (i) and (4.12)-(4.13) in conjunction with Lebesgue's dominated convergence theorem that (4.11) holds.
(iii) It follows from (4.13) that $\|M_\alpha\|_{\mathcal{B}(L^2(\mathbb{R},\mathcal{H}))} \leq \|M_\alpha\|_{HS} \leq \|M_0\|_{HS}$. Hence, $\|M_\alpha\|_{\mathcal{B}(L^2(\mathbb{R},\mathcal{H}))}$ is bounded *independently* of $\alpha \in (0,\alpha_0]$ for some $\alpha_0 > 0$. Therefore, for g small enough, $\|gM_\alpha\|_{\mathcal{B}(L^2(\mathbb{R},\mathcal{H}))} < 1$ and, consequently, $(1+gM_\alpha)^{-1}$ exists and is bounded for these g and α. In particular, we may write $1+gK_\alpha(V) = (1+gM_\alpha)[1+g(1+gM_\alpha)^{-1}L_\alpha]$ from which the assertion follows. □

5. Weakly coupled bound states

Throughout this section operators (resp. vectors) are denoted by boldface capital (resp. small) letters to emphasize their matrix (resp. vector) structure.

5.1. **Two-channel Hamiltonians with matrix-valued potentials.**

We consider the case where the potential is a 2×2 matrix-valued potential $\mathbf{V}(x)$ with measurable functions V_{ij} on $\mathbb{R}$ as entries. The Euclidean inner product and norm in $\mathbb{C}^2$ are denoted by $\langle\cdot,\cdot\rangle_{\mathbb{C}^2}$ and $\|\cdot\|_{\mathbb{C}^2}$, respectively.

Assumption 5.1.
(a) $\mathbf{V}(x)$ is symmetric, i.e. $\overline{V_{ji}} = V_{ij}$.
(b)
$$\int_{\mathbb{R}} (1+|x|^2)\,\|\mathbf{V}(x)\|_{\mathcal{B}(\mathbb{C}^2)}\,dx < \infty.$$
(c) The functions V_{ij} are real-valued.

5.1.1. *Two-channel Hamiltonian with a single threshold.* First we consider the Hamiltonian $\mathbf{H}(g) = \mathbf{H}_0 + g\mathbf{V}(x)$ in $L^2(\mathbb{R},\mathbb{C}^2)$, defined in Proposition 3.1 by means of forms. Formally, we may write the Hamiltonian as

$$\mathbf{H}(g) = \mathbf{H}_0 + g\mathbf{V} = \begin{pmatrix} -\frac{d^2}{dx^2} & 0 \\ 0 & -\frac{d^2}{dx^2}\end{pmatrix} + g\begin{pmatrix} V_{11} & V_{12} \\ V_{21} & V_{22}\end{pmatrix} \quad (5.1)$$

in $L^2(\mathbb{R},\mathbb{C}^2) = L^2(\mathbb{R})\oplus L^2(\mathbb{R})$. Under Assumption 5.1 we know that its essential spectrum equals the half-axis starting at the (threshold) point zero.

Define the matrices $\mathbf{S}$ and $\mathbf{T}$ by

$$\mathbf{S} = \int_{\mathbb{R}} \mathbf{V}(x)\,dx, \quad \mathbf{T}_0 = -\frac{1}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)|x-y|\mathbf{V}(y)\,dydx. \quad (5.2)$$

We establish the following result.

Theorem 5.2. *Let* $\mathbf{V}$ *obey Assumption 5.1(a)-(b) and let* $\mathbf{H}(g) = \mathbf{H}_0 + g\mathbf{V}(x)$ *be the self-adjoint Hamiltonian on* $L^2(\mathbb{R}, \mathbb{C}^2)$ *defined in Proposition 3.1 by means of forms.*
(i) Assume that the matrix $\mathbf{S}$*, defined in (5.2), has* n (≤ 2) *negative eigenvalues, denoted by* s_i*, with multiplicities* $\varkappa_i$*. Then, for a small enough* g*, the two-channel Hamiltonian* $\mathbf{H}(g)$ *has precisely* $\sum_{i=1}^{n} \varkappa_i$ *negative eigenvalues (taking into account multiplicity)* E_{ij} *satisfying the formulas*

$$(-E_{ij}(g))^{1/2} = -\frac{g}{2}s_i + \frac{g^2}{2}\langle \mathbf{v}_{ij}, \mathbf{T}_0\mathbf{v}_{ij}\rangle_{\mathbb{C}^2} + O(g^3), \qquad (5.3)$$
$$i = 1, \dots, n, \quad j = 1, \dots, \varkappa_i,$$

where $\mathbf{T}_0$ *is defined in (5.2) and* $\mathbf{v}_{ij}$ *are the eigenvectors corresponding to the eigenvalue* s_i *of* $\mathbf{S}$*.*
(ii) Suppose that $\mathbf{V}$ *obey Assumption 5.1(c) and that the matrix* $\mathbf{S}$ *has* n *non-positive eigenvalues, denoted by* s_i*, with multiplicities* $\varkappa_i$*. If the eigenvectors* $\mathbf{v}_{0j}$*,* $j = 1, \dots, \varkappa_i$*, associated with the eigenvalue zero of* $\mathbf{S}$ *satisfy* $\langle \mathbf{v}_{0j}, \mathbf{T}_0\mathbf{v}_{0j}\rangle_{\mathbb{C}^2} \neq 0$ *then the conclusion of part (i) remains valid.*

Proof. According to the Birman-Schwinger relation formulated in Proposition 4.2, $E(g) < 0$ is an eigenvalue of $\mathbf{H}(g)$ if and only if -1 is an eigenvalue of $g\mathbf{K}_\alpha(\mathbf{V})$ with $\alpha^2 = -E(g)$. Furthermore, in view of Lemma 4.5(iii), the operator $g\mathbf{K}_\alpha(\mathbf{V})$ has eigenvalue -1 if and only if the same is true for $g(1 + g\mathbf{M}_\alpha)^{-1}\mathbf{L}_\alpha$. Now let us denote the (unknown) eigenvalues and eigenfunctions of $(1 + g\mathbf{M}_\alpha)^{-1}\mathbf{L}_\alpha$ by $\mu_k(g, \alpha)$ and $\mathbf{\Psi}_k(x; g, \alpha)$, respectively, viz.

$$(1 + g\mathbf{M}_\alpha)^{-1}\mathbf{L}_\alpha\mathbf{\Psi}_k(x; g, \alpha) = \mu_k(g, \alpha)\mathbf{\Psi}_k(x; g, \alpha). \qquad (5.4)$$

Let $\mathbf{u}_k \in \mathbb{C}^2$ be a constant vector. We insert

$$\mathbf{\Psi}_k(x; g, \alpha) = \frac{1}{2\alpha}(1 + g\mathbf{M}_\alpha)^{-1}|\mathbf{V}(x)|^{1/2}\mathbf{u}_k$$

into (5.4) and obtain

$$\mathbf{R}_g\mathbf{u}_k = \mu_k(g, \alpha)\mathbf{u}_k, \qquad (5.5)$$

where $\mathbf{R}_g$ is the matrix

$$\mathbf{R}_g = \frac{1}{2\alpha}\int_{\mathbb{R}} \mathbf{V}^{1/2}(x)[(1 + g\mathbf{M}_\alpha)^{-1}|\mathbf{V}|^{1/2}](x)\, dx.$$

Define $\mathbf{S}$ as in (5.2) and, moreover, define

$$\mathbf{T}(\alpha) = \int_{\mathbb{R}} \mathbf{V}^{1/2}(x)[\mathbf{M}_\alpha|\mathbf{V}|^{1/2}](x)\, dx.$$

Then we have that

$$\mathbf{R}_g = \frac{1}{2\alpha}\mathbf{S} - \frac{g}{2\alpha}\mathbf{T}(\alpha) + \mathrm{O}(g^2)$$

for small g.
(i) By assumption the matrix $\mathbf{S}$ has n negative eigenvalues, denoted by s_k. For simplicity we assume that the eigenvalues s_k are simple. The corresponding eigenvectors are denoted by $\mathbf{v}_k$. We apply the regular perturbation theory to the eigenvalue problem (5.5) and we find that

$$\mu_k(g,\alpha) = \frac{1}{2\alpha}s_k - \frac{g}{2\alpha}\langle \mathbf{v}_k, \mathbf{T}(\alpha)\mathbf{v}_k\rangle_{\mathbb{C}^2} + \mathrm{O}(g^2).$$

Define the matrix $\mathbf{T}_0$ as in (5.2) and

$$\mathbf{T}_1 = \frac{1}{4}\int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)|x-y|^2\mathbf{V}(y)\,dxdy.$$

Then we have that $\mathbf{T}(\alpha) = \mathbf{T}_0 + \alpha\mathbf{T}_1 + \mathrm{O}(\alpha^2)$. In this way we find that the eigenvalues associated with the eigenvalue problem (5.5) are

$$\mu_k(g,\alpha) = \frac{1}{2\alpha}s_k - \frac{g}{2\alpha}\langle \mathbf{v}_k, \mathbf{T}_0\mathbf{v}_k\rangle_{\mathbb{C}^2} + \mathrm{O}(g^2).$$

Together with the comments following the proof of Proposition 4.2, the latter implies that the solution to (4.7) is

$$\alpha_g = -\frac{g}{2}s_k + \frac{g^2}{2}\langle \mathbf{v}_k, \mathbf{T}_0\mathbf{v}_k\rangle_{\mathbb{C}^2} + \mathrm{O}(g^2). \tag{5.6}$$

Clearly, (5.6) implies that each negative eigenvalue s_k of $\mathbf{S}$ gives rise to precisely one negative eigenvalue $E_k(g)$ of $\mathbf{H}(g)$ obeying the asymptotic formula

$$(-E_k(g))^{1/2} = -\frac{g}{2}s_k + \frac{g^2}{2}\langle \mathbf{v}_k, \mathbf{T}_0\mathbf{v}_k\rangle_{\mathbb{C}^2} + \mathrm{O}(g^3).$$

(ii) We investigate the situation where zero is an eigenvalue of $\mathbf{S}$ (as above we restrict ourselves to the case where zero is simple). Let $\mathbf{S}\mathbf{v}_0 = 0$ for some $\mathbf{v}_0 \neq 0$. Taylor's formula yields

$$\begin{aligned}
&\left\langle \mathbf{v}_0, \int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)\frac{e^{-\alpha|x-y|}}{2\alpha}\mathbf{V}(y)\,dx\,dy\mathbf{v}_0\right\rangle_{\mathbb{C}^2} \\
&\quad = \frac{1}{2\alpha}\langle \mathbf{v}_0, \mathbf{S}^2\mathbf{v}_0\rangle_{\mathbb{C}^2} - \frac{1}{2}\left\langle \mathbf{v}_0, \int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)|x-y|\mathbf{V}(y)\,dx\,dy\mathbf{v}_0\right\rangle_{\mathbb{C}^2} \\
&\quad +\alpha\left\langle \mathbf{v}_0, \int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)O(|x-y|^2)\mathbf{V}(y)\,dx\,dy\mathbf{v}_0\right\rangle_{\mathbb{C}^2}.
\end{aligned}$$

Since $\mathbf{S}\mathbf{v}_0 = 0$ by assumption, the first term equals zero. As $\alpha \downarrow 0$, we obtain that

$$\begin{aligned}
\langle \mathbf{v}_0, \mathbf{T}_0\mathbf{v}_0\rangle_{\mathbb{C}^2} &= \lim_{\alpha\downarrow 0}\left\langle \mathbf{v}_0, \int\int \mathbf{V}(x)\frac{e^{-\alpha|x-y|}}{2\alpha}\mathbf{V}(y)\,dxdy\,\mathbf{v}_0\right\rangle_{\mathbb{C}^2} \\
&= \lim_{\alpha\downarrow 0}\sum_{i,j,k}\int\int \frac{e^{-\alpha|x-y|}}{2\alpha}V_{ik}(x)V_{kj}(y)(\mathbf{v}_0)_i\overline{(\mathbf{v}_0)_j}\,dxdy.
\end{aligned} \tag{5.7}$$

Let $\mathcal{F}$ denote the one-dimensional Fourier transform and let $(\mathcal{F}\mathbf{V})(\xi)$ denote the matrix with elements $(\mathcal{F}\mathbf{V})_{ij}(\xi) = (\mathcal{F}V_{ij})(\xi)$ satisfying $\mathcal{F}V_{ij} = \mathcal{F}V_{ji}$ because V is symmetric and V_{ij} are real-valued. Using the latter in conjunction with the Fourier transform of $(1/2\alpha)e^{-\alpha|x|}$, which equals $1/(\xi^2+\alpha^2)$, we find that

$$\begin{aligned}\text{r.h.s. of (5.7)} &= \lim_{\alpha\downarrow 0}\sum_{i,j,k}\int \frac{1}{\xi^2+\alpha^2}(\mathcal{F}V_{ik})(\xi)\overline{(\mathcal{F}V_{kj})(\xi)}(\mathbf{v}_0)_i\overline{(\mathbf{v}_0)_j}\,d\xi \\ &= \int \frac{1}{\xi^2}\langle \mathbf{v}_0, (\mathcal{F}\mathbf{V})^*(\xi)(\mathcal{F}\mathbf{V})(\xi)\mathbf{v}_0\rangle_{\mathbb{C}^2}\,d\xi \\ &= \int \frac{1}{\xi^2}\,\|(\mathcal{F}\mathbf{V})(\xi)\mathbf{v}_0\|^2_{\mathbb{C}^2}\;d\xi \geq 0, \qquad (5.8)\end{aligned}$$

By assumption, $\langle \mathbf{v}_0, \mathbf{T}_0\mathbf{v}_0\rangle_{\mathbb{C}^2} \neq 0$ and therefore (5.8) implies that there is also a negative eigenvalue of $\mathbf{H}(g)$ associated with the eigenvalue zero of $\mathbf{S}$. □

Remark 5.3. The reasoning in the proof of Theorem 5.2(ii) requires that the entries V_{ij} in the potential $\mathbf{V}$ are real-valued. A substantial improvement would be to establish the result for complex-valued entries.

Example 5.4 (Square-well potentials). Let $\chi_{[0,1]}$ denote the characteristic function associated with the interval $[0,1]$. Choose the following entries of $\mathbf{V}$:

$$\begin{aligned}V_{11}(x) &= -5\chi_{[0,1]}(x), \quad V_{22}(x) = -3\chi_{[0,1]}(x), \\ V_{12}(x) &= V_{21}(x) = -3a\chi_{[0,1]}(x), \;\; a>0.\end{aligned}$$

Then the matrix $\mathbf{S}$ equals

$$\mathbf{S} = \begin{pmatrix} -5 & -3a \\ -3a & -3 \end{pmatrix}$$

and it has two real eigenvalues given by $-4 \pm \sqrt{1+9a^2}$. Thus the following cases are possible: 1) If $a > \sqrt{5/3}$ there is exactly one negative eigenvalue of $\mathbf{S}$, namely $-4-\sqrt{1+9a^2}$. 2) If $a < \sqrt{5/3}$ there are two negative eigenvalues of $\mathbf{S}$, namely $-4\pm\sqrt{1+9a^2}$. 3) If $a = \sqrt{5/3}$ there are two nonpositive eigenvalues of $\mathbf{S}$, namely $-4-\sqrt{1+9a^2}$ and 0.

5.1.2. *Two-channel Hamiltonian with two thresholds.* As an example of a Hamiltonian with more than one threshold, we consider the one in (1.1), having thresholds at 0 and 1. Henceforth its free Hamiltonian is denoted by $\tilde{\mathbf{H}}_0$. The essential spectrum of $\tilde{\mathbf{H}}_0$ is the union of the half-axes starting at the thresholds, i.e. $\sigma_{ess}(\tilde{\mathbf{H}}_0) = [0,\infty)$. The resolvent of $\tilde{\mathbf{H}}_0$ is given by

$$(\tilde{\mathbf{H}}_0+\alpha^2)^{-1} = \begin{pmatrix} (-d^2/dx^2+\alpha^2)^{-1} & 0 \\ 0 & (-d^2/dx^2+\alpha^2+1)^{-1} \end{pmatrix}, \;\; \alpha>0,$$

where the entries have the integral kernels

$$\frac{1}{2\alpha}e^{-\alpha|x-y|} \quad \text{and} \quad \frac{1}{2\sqrt{\alpha^2+1}}e^{-\sqrt{\alpha^2+1}|x-y|}.$$

It is easy to show that the assertions of Proposition 3.1 are valid if one replaces $\mathbf{H}_0$ by $\tilde{\mathbf{H}}_0$. In this way we obtain a self-adjoint realization of the formal Hamiltonian $\tilde{\mathbf{H}}_0 + g\mathbf{V}$ in $L^2(\mathbb{R}, \mathbb{C}^2)$. Moreover, the Birman-Schwinger relation in Proposition 4.2 holds for $\tilde{\mathbf{H}}(g) = \tilde{\mathbf{H}}_0 + g\mathbf{V}$.

Define the operators $\tilde{\mathbf{L}}_\alpha$ and $\tilde{\mathbf{M}}_\alpha$ by their "kernels"

$$\begin{aligned}
\tilde{\mathbf{L}}_\alpha(x,y) &= \frac{1}{2\alpha}\mathbf{V}(x)^{1/2}\begin{pmatrix}1 & 0\\ 0 & 0\end{pmatrix}|\mathbf{V}(y)|^{1/2},\\
\tilde{\mathbf{M}}_\alpha(x,y) &= \mathbf{V}(x)^{1/2}\begin{pmatrix}\frac{1}{2\alpha}[e^{-\alpha|x-y|}-1] & 0\\ 0 & \frac{e^{-\sqrt{\alpha^2+1}|x-y|}}{2\sqrt{\alpha^2+1}}\end{pmatrix}|\mathbf{V}(y)|^{1/2}.
\end{aligned}$$

Moreover, we introduce the operator $\tilde{\mathbf{M}}_0$ by its kernel

$$\tilde{\mathbf{M}}_0(x,y) = \mathbf{V}(x)^{1/2}\begin{pmatrix}-\frac{1}{2}|x-y| & 0\\ 0 & \frac{1}{2}e^{-|x-y|}\end{pmatrix}|\mathbf{V}(y)|^{1/2}.$$

By making a few obvious changes to the proof of Lemma 4.5 we obtain the following result.

Lemma 5.5. *Assume that* $\int_{\mathbb{R}}(1+|x|^2)\,\|\mathbf{V}(x)\|_{\mathcal{B}(\mathbb{C}^2)}\,dx < \infty$. *If* K_α, L_α, M_α *and* M_0 *in Lemma 4.5 are replaced by* $\tilde{\mathbf{K}}_\alpha$, $\tilde{\mathbf{L}}_\alpha$, $\tilde{\mathbf{M}}_\alpha$ *and* $\tilde{\mathbf{M}}_0$ *then the assertions (i)-(iii) of Lemma 4.5 are still valid.*

Define the matrices

$$\tilde{\mathbf{S}} = \begin{pmatrix}\int_{\mathbb{R}} V_{11}(x)\,dx & \int_{\mathbb{R}} V_{12}(x)\,dx\\ 0 & 0\end{pmatrix}, \tag{5.9}$$

$$\tilde{\mathbf{T}}_0 = \int_{\mathbb{R}}\int_{\mathbb{R}}\begin{pmatrix}1 & 0\\ 0 & 0\end{pmatrix}\mathbf{V}(x)\begin{pmatrix}-\frac{1}{2}|x-y| & 0\\ 0 & \frac{e^{-|x-y|}}{2}\end{pmatrix}\mathbf{V}(y)\,dxdy. \tag{5.10}$$

For the Hamiltonian $\tilde{\mathbf{H}}(g)$ we are able to derive an analogue of part (i) in Theorem 5.2.

Theorem 5.6. *Let* $\mathbf{V}$ *obey Assumption 5.1(a)-(c) and let* $\tilde{\mathbf{H}}(g) = \tilde{\mathbf{H}}_0 + g\mathbf{V}(x)$ *be the self-adjoint Hamiltonian on* $L^2(\mathbb{R}, \mathbb{C}^2)$ *defined in Proposition 3.1 by means of forms.*
Assume that the matrix $\tilde{\mathbf{S}}$, *defined in (5.9), has a negative eigenvalue* $\tilde{s}$ *(such an eigenvalue is simple if it exists). Then, for a small enough coupling constant* g, *the eigenvalue* $\tilde{s}$ *of* $\tilde{\mathbf{S}}$ *gives rise to exactly one negative eigenvalue* $\tilde{E}$ *of the two-channel Hamiltonian* $\tilde{\mathbf{H}}(g)$. *The negative eigenvalue* $\tilde{E}$ *satisfies the formula*

$$(-\tilde{E}(g))^{1/2} = -\frac{g}{2}\tilde{s} + \frac{g^2}{2}\langle\tilde{\mathbf{v}}, \tilde{\mathbf{T}}_0\tilde{\mathbf{v}}\rangle_{\mathbb{C}^2} + O(g^3), \tag{5.11}$$

where $\tilde{\mathbf{T}}_0$ *is defined in (5.10) and* $\tilde{\mathbf{v}}$ *is the eigenvector corresponding to the eigenvalue* $\tilde{s}$ *of* $\tilde{\mathbf{S}}$.

Proof. Imitating the proof of Theorem 5.2 we arrive at the eigenvalue problem

$$\tilde{\mathbf{R}}_g \mathbf{u}_k = \mu_k(g,\alpha)\mathbf{u}_k, \tag{5.12}$$

where $\tilde{\mathbf{R}}_g$ is the matrix

$$\tilde{\mathbf{R}}_g = \frac{1}{2\alpha}\int_{\mathbb{R}} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{V}^{1/2}(x)[(1+g\tilde{\mathbf{M}}_\alpha)^{-1}|\mathbf{V}|^{1/2}](x)\,dx.$$

Define the matrix $\tilde{\mathbf{S}}$ as in (5.9) and, moreover, define

$$\tilde{\mathbf{T}}(\alpha) = \int_{\mathbb{R}} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \mathbf{V}^{1/2}(x)[\tilde{\mathbf{M}}_\alpha|\mathbf{V}|^{1/2}](x)\,dx.$$

Then we may write

$$\tilde{\mathbf{R}}_g = \frac{1}{2\alpha}\tilde{\mathbf{S}} - \frac{g}{2\alpha}\tilde{\mathbf{T}}(\alpha) + \mathrm{O}(g^2)$$

for small g. From here on everything depends on the possible eigenvalues of $\tilde{\mathbf{S}}$. Let

$$a = \int_{\mathbb{R}} V_{11}(x)\,dx.$$

The following cases may occur: I. If $a \neq 0$ then there are two subcases. I.1. If $a > 0$ then $\tilde{\mathbf{S}}$ has the eigenvalue zero and the positive eigenvalue a, each of multiplicity one. I.2. If $a < 0$ then $\tilde{\mathbf{S}}$ has the eigenvalue zero and the negative eigenvalue a, each having multiplicity one. II. If $a = 0$ then $\tilde{\mathbf{S}}$ has the eigenvalue zero with multiplicity one.

Repeating the reasoning in the first part of the proof of Theorem 5.2 we show that a negative eigenvalue of $\tilde{\mathbf{S}}$ (from I.2 it has multiplicity one) generates exactly one negative eigenvalue of $\tilde{\mathbf{H}}(g)$ provided g is small enough. □

Remark 5.7. The matrix $\tilde{\mathbf{S}}$ always has the eigenvalue zero. It remains an open problem to settle whether or not the latter gives rise to a negative eigenvalue of $\tilde{\mathbf{H}}(g)$ for a sufficiently small g.

5.2. **N-channel Hamiltonian with matrix-valued potentials.** In this section we consider the case where the potential is a $N \times N$ matrix-valued potential $\mathbf{V}(x)$ with measurable functions V_{ij} on $\mathbb{R}$ as entries.

Assumption 5.8.
(a) $\mathbf{V}(x)$ is symmetric, i.e. $\overline{V_{ji}} = V_{ij}$.
(b)

$$\int_{\mathbb{R}} (1+|x|^2)\,\|V(x)\|_{\mathcal{B}(\mathbb{C}^N)}\,dx < \infty.$$

(c) The functions V_{ij} are real-valued.

Define the matrices $\mathbf{S}$ and $\mathbf{T}_0$ by

$$\mathbf{S} = \int_{\mathbb{R}} \mathbf{V}(x)\,dx, \quad \mathbf{T}_0 = -\frac{1}{2}\int_{\mathbb{R}}\int_{\mathbb{R}} \mathbf{V}(x)|x-y|\mathbf{V}(y)\,dydx. \tag{5.13}$$

We have the following result.

Theorem 5.9. *Let* $\mathbf{V}$ *obey Assumption 5.8(a)-(b) and let* $\mathbf{H}(g) = \mathbf{H}_0 + g\mathbf{V}(x)$ *be the self-adjoint Hamiltonian on* $L^2(\mathbb{R}, \mathbb{C}^2)$ *defined in Proposition 3.1 by means of forms.*
(i) Assume that the matrix $\mathbf{S}$*, defined in (5.13), has* n *negative eigenvalues, denoted by* s_i*, with multiplicities* $\varkappa_i$*. Then, for a small enough* g*, the* N*-channel Hamiltonian* $\mathbf{H}(g)$ *has precisely* $\sum_{i=1}^{n} \varkappa_i$ *negative eigenvalues (taking into account multiplicity)* E_{ij} *satisfying the formulas*

$$(-E_{ij}(g))^{1/2} = -\frac{g}{2}s_i + \frac{g^2}{2}\langle \mathbf{v}_{ij}, \mathbf{T}_0\mathbf{v}_{ij}\rangle_{\mathbb{C}^N} + O(g^3), \qquad (5.14)$$
$$i = 1, \dots, n, \quad j = 1, \dots, \varkappa_i,$$

where $\mathbf{T}_0$ *is defined in (5.13) and* $\mathbf{v}_{ij}$ *are the eigenvectors corresponding to the eigenvalue* s_i *of* $\mathbf{S}$*.*
(ii) Suppose that $\mathbf{V}$ *obey Assumption 5.8(c) and that the matrix* $\mathbf{S}$ *has* n *non-positive eigenvalues, denoted by* s_i*, with multiplicities* $\varkappa_i$*. If the eigenvectors* $\mathbf{v}_{0j}$*,* $j = 1, \dots, \varkappa_i$*, associated with the eigenvalue zero of* $\mathbf{S}$ *satisfy* $\langle \mathbf{v}_{0j}, \mathbf{T}_0\mathbf{v}_{0j}\rangle_{\mathbb{C}^N} \neq 0$ *then the conclusion of part (i) remains valid.*

Proof. The proof is a straightforward generalization of the proof of Theorem 5.2. □

Remark 5.10. One of the referees pointed out that Theorem 5.9 was proven in Šeba [29]. Therein, however, Theorem 3 is incorrect because the quantity $\int_{\mathbb{R}} (1/p^2)(\mathbf{a}_0, \mathcal{F}(V)^2(p)\mathbf{a}_0)_n \, dp$ (in Šeba's notation) is not necessarily different from zero. Moreover, the Birman-Schwinger relation (in the matrix-valued setting) is stated without proof.

6. Perturbation of embedded eigenvalues

For the sake of completeness we consider perturbation of two-channel diagonal Hamiltonians with one-dimensional Schrödinger operators as component Hamiltonians, having eigenvalues embedded in its continuous spectrum.

6.1. **Two-channel Hamiltonians.**

Consider the formal expression

$$\mathbf{H}(g) = \mathbf{H}(0) + g\mathbf{V} = \begin{pmatrix} H_{11} & 0 \\ 0 & H_{22} \end{pmatrix} + g \begin{pmatrix} V_{11}(x) & V_{12}(x) \\ V_{21}(x) & V_{22}(x) \end{pmatrix} \qquad (6.1)$$

in $\mathcal{H} = L^2(\mathbb{R}) \oplus L^2(\mathbb{R})$, where

$$H_{11} = -\frac{d^2}{dx^2} + W_{11}(x) \text{ and } H_{22} = -\frac{d^2}{dx^2} + 1 + W_{22}(x). \qquad (6.2)$$

We impose the following assumptions on the potentials W_{jj}, $j = 1, 2$.

Assumption 6.1. Suppose that the real-valued, measurable functions W_{jj}, $j = 1, 2$, satisfy:
(a) $W_{jj} \neq 0$.
(b) The bound

$$|W_{jj}(x)| \leq C\,(1 + |x|^2)^{-1-\delta} \tag{6.3}$$

holds for some $C, \delta > 0$ and all x.
(c) $\int_{\mathbb{R}} W_{jj}(x)\,dx \leq 0$.
(d) W_{jj} extends to a function analytic in the sector

$$\mathcal{A}_{\alpha_0} = \{\, z \in \mathbb{C} \,:\, |\arg z| \leq \alpha_0 \,\}$$

for some $\alpha_0 > 0$. Moreover, the bound (6.3) holds in this sector.

Under Assumption 6.1(a)-(c) the operator $H_{11} = -\frac{d^2}{dx^2} + W_{11}(x)$ is self-adjoint in $L^2(\mathbb{R})$ and $\sigma(H_{11}) = \sigma_d(H_{11}) \cup \sigma_{ess}(H_{11}) = \sigma_d(H_{11}) \cup [0, \infty)$ with a non-empty discrete spectrum $\mu_1 < \mu_2 < \cdots < \mu_N < 0$, which is simple and finite [31]. The corresponding normalized eigenfunctions ϕ_n , $n = 1, 2, \ldots, N$, are exponentially decaying. The analyticity requirement in Assumption 6.1(d) is convenient to adopt for analyzing the resonance behaviour. Similarly, the operator $H_{22} = -\frac{d^2}{dx^2} + 1 + W_{22}(x)$ is self-adjoint in $L^2(\mathbb{R})$ and $\sigma(H_{22}) = \sigma_d(H_{22}) \cup \sigma_{ess}(H_{22}) = \sigma_d(H_{22}) \cup [1, \infty)$ with a non-empty discrete spectrum $\nu_1 < \nu_2 < \cdots < \nu_M < 1$ which is simple and finite. The corresponding normalized eigenfunctions χ_m , $m = 1, 2, \ldots, M$, are exponentially decaying.

Consider the unperturbed Hamiltonian $\mathbf{H}(0) = \mathrm{diag}\,(H_{11}, H_{22})$. Assumption 6.1 ensures that

$$\begin{aligned} \sigma_c(\mathbf{H}(0)) &= \sigma_{ess}(\mathbf{H}(0)) = \sigma_{ess}(H_{11}) \cup \sigma_{ess}(H_{22}) \\ &= [0, \infty) \cup [1, \infty) = [0, \infty). \end{aligned}$$

Thus, the continuous spectrum of $\mathbf{H}(0)$ is the union of the two half-lines starting at 0 and 1. This motivates the definition of the threshold set $\mathsf{T} = \{\, 0, 1 \,\}$. Furthermore, $\sigma_p(\mathbf{H}(0)) = \sigma_p(H_{11}) \cup \sigma_p(H_{22})$. Among this (finite) set of eigenvalues, a (finite) subset is isolated or situated at the threshold 0, while the rest satisfying the condition $0 < \nu_m < 1$ is embedded in the continuous spectrum of $\mathbf{H}(0)$. For the sake of simplicity we make the following assumption.

Assumption 6.2. Suppose that none of the embedded eigenvalues ν_m of $\mathbf{H}(0)$ coincide with the threshold 0.

We impose the following conditions on the components of the perturbation $\mathbf{V}$.

Assumption 6.3. Suppose that the real-valued, measurable functions V_{ij}, $i, j = 1, 2$, satisfy:
(a) The bound

$$|V_{ij}(x)| \leq C\,(1 + |x|^2)^{-1-\delta} \tag{6.4}$$

holds for some $C, \delta > 0$ and all x.
(b) V_{ij} extends to a function analytic in the sector $\mathcal{A}_{\alpha_0}$ (see Assumption 6.1(d)) for some $\alpha_0 > 0$. Moreover, the bound (6.4) holds in this sector.

6.2. **Complex dilation.** We use a complex deformation. For θ real define S_θ on $L^2(\mathbb{R})$ by the unitary operator

$$(S_\theta \psi) = e^{\theta/2}\psi(e^\theta x), \ \ \psi \in L^2(\mathbb{R}). \tag{6.5}$$

S_θ is a one-parameter unitary group on $L^2(\mathbb{R})$. It is easy to see that S_θ leave $\mathcal{D}(-d^2/dx^2) = H^2(\mathbb{R})$ invariant and that

$$H_{11,\theta} := S_\theta H_{11} S_\theta^{-1} = -e^{-2\theta}\frac{d^2}{dx^2} + W_{11,\theta}(x) = -e^{-2\theta}\frac{d^2}{dx^2} + W_{11}(e^\theta x),$$

Let $\mathcal{A}_0 = \{\, \theta \,:\, |\operatorname{Im}\theta| \leq \min\{\alpha_0, \pi/4\}\,\}$ (cf. Assumption 6.1(d)). Under Assumption 6.1, $H_{11,\theta}$ obviously has a continuation to a type (A) family of m-sectorial operators analytic in the sense of Kato [13] for $\theta \in \mathcal{A}_0$. Likewise,

$$H_{22,\theta} := S_\theta H_{22} S_\theta^{-1} = -e^{-2\theta}\frac{d^2}{dx^2} + 1 + W_{22,\theta}(x) = -e^{-2\theta}\frac{d^2}{dx^2} + 1 + W_{22}(e^\theta x)$$

has a continuation to a type (A) analytic family of operators on $\mathcal{A}_0$. From standard Aguilar-Combes theory [1] we determine the spectra of $H_{11,\theta}$ and $H_{22,\theta}$:

$$\begin{aligned}\sigma(H_{11,\theta}) &= \{\mu_1, \mu_2, \ldots, \mu_N\} \cup \{e^{-2\theta}\lambda \,:\, \lambda \in [0,\infty)\,\},\\ \sigma(H_{22,\theta}) &= \{\nu_1, \nu_2, \ldots, \nu_M\} \cup \{e^{-2\theta}\lambda + 1 \,:\, \lambda \in [0,\infty)\,\}.\end{aligned}$$

Having extended S_θ in (6.5) analytically to $\mathcal{A}_0$ we may define

$$\mathbf{S}_\theta \Psi = \begin{pmatrix} S_\theta & 0 \\ 0 & S_\theta \end{pmatrix}\begin{pmatrix}\psi_1 \\ \psi_2\end{pmatrix}, \quad \Psi \in \mathcal{H}.$$

Due to its diagonal structure, the Hamiltonian

$$\mathbf{H}_\theta(0) := \mathbf{S}_\theta \mathbf{H}(0) \mathbf{S}_\theta^{-1} = \begin{pmatrix} H_{11,\theta} & 0 \\ 0 & H_{22,\theta}\end{pmatrix}$$

has a continuation to a type (A) analytic family of operators in the sector $\mathcal{A}_0$. Furthermore,

$$\begin{aligned}\sigma(\mathbf{H}_\theta(0)) &= \sigma(H_{11,\theta}) \cup \sigma(H_{22,\theta})\\ &= \{\mu_1, \mu_2, \ldots, \mu_N\} \cup \{\nu_1, \nu_2, \ldots, \nu_M\}\\ &\quad \cup \{e^{-2\theta}\lambda \,:\, \lambda \in [0,\infty)\,\} \cup \{e^{-2\theta}\lambda + 1 \,:\, \lambda \in [0,\infty)\,\}.\end{aligned}$$

In particular, the eigenvalues embedded in $\sigma(\mathbf{H}(0))$ are discrete eigenvalues of $\mathbf{H}_\theta(0)$ for θ nonreal.

Henceforth E_0 denotes any of the embedded eigenvalues ν_m of $\mathbf{H}(0)$. Let $\mathbf{R}_0(\theta;\zeta)$ denote the resolvent of $\mathbf{H}_\theta(0)$. Since E_0 is an isolated eigenvalue of $\mathbf{H}_\theta(0)$, we may choose a contour Γ around E_0 such that Γ belongs to the resolvent set of $\mathbf{H}_\theta(0)$ and E_0 is the only eigenvalue

of $\mathbf{H}_\theta(0)$ contained inside of Γ. Moreover, let $\mathbf{P}_\theta$ denote the eigenprojection associated with the eigenvalue E_0 and put

$$\mathbf{S}_\theta^{(p)} := \frac{1}{2\pi i}\int_\Gamma \frac{\mathbf{R}_0(\theta;\zeta)}{(E_0-\zeta)^p}\,d\zeta, \quad p \geq 1. \tag{6.6}$$

Then $\mathbf{P}_\theta = -\mathbf{S}_\theta^{(0)}$ and $\widehat{\mathbf{R}}_0(\theta;\zeta) := \mathbf{S}_\theta^{(1)}$ is the reduced resolvent of $\mathbf{H}_\theta(0)$ at the point ζ. Define

$$\mathbf{V}_\theta = \mathbf{S}_\theta \mathbf{V} \mathbf{S}_\theta^{-1} = \begin{pmatrix} V_{11,\theta} & V_{12,\theta} \\ V_{21,\theta} & V_{22,\theta} \end{pmatrix} \text{ with } V_{ij,\theta}(x) = V_{ij}(e^\theta x).$$

Then we have the following result.

Lemma 6.4. *Let Assumption 6.1 and Assumption 6.3 hold. Let Γ be the contour described above and let $\mathbf{S}_\theta^{(p)}$ be defined in (6.6).*
(i) If $\operatorname{Im}\theta \in (0,\alpha_0)$ then there exists a constant $C_\theta > 0$ such that

$$\max_{\zeta\in\Gamma} \|g\mathbf{V}_\theta \mathbf{R}_0(\theta;\zeta)\| \leq C_\theta |g|. \tag{6.7}$$

If ζ is replaced by $\zeta_0 = \min\{\mu_1,\nu_1\} - 1$ then the constant in (6.7) is independent of θ and the estimate holds for all $|\operatorname{Im}\theta| < \alpha_0$.
(ii) For $p \geq 0$ there exists a constant $C_\theta > 0$ such that

$$\|g\mathbf{V}_\theta \mathbf{S}_\theta^{(p)}\| \leq C_\theta \frac{|\Gamma|}{2\pi}\frac{|g|}{[\operatorname{dist}(E_0,\Gamma)]^p}.$$

Proof. The contour Γ is by assumption contained in the resolvent of $\mathbf{H}_\theta(0)$. Since $\mathbf{R}_0(\theta;\cdot)$ is bounded and continuous and Γ is compact, there exists a constant $\tilde{C}_\theta$ such that $\max_{\zeta\in\Gamma}\|\mathbf{R}_0(\theta;\zeta)\| \leq \tilde{C}_\theta$.

Thus, $\max_{\zeta\in\Gamma}\|\mathbf{V}_\theta\mathbf{R}_0(\theta;\zeta)\| \leq \tilde{C}_\theta C_{\mathbf{V}}$, where $C_{\mathbf{V}}$ denotes a bound on the norm of $\mathbf{V}_\theta$, which is independent of θ by Assumption 6.3(b). This shows the first claim. Moreover, ζ_0 is to the left of the numerical range $\Theta(\mathbf{H}_\theta(0))$ of $\mathbf{H}_\theta(0)$ at the unit distance. Hence $\|\mathbf{R}_0(\theta;\zeta_0)\| = 1/[\operatorname{dist}(\zeta_0,\Theta(\mathbf{H}_\theta(0)))] = 1$. Therefore the constant $\tilde{C}_\theta$ in the above estimate may be replaced by 1. This verifies (i). The assertion (ii) follows immediately. □

Hence, provided g is small enough, it follows from Lemma 6.4(i) that $g\mathbf{V}_\theta$ is $\mathbf{H}_\theta(0)$-compact. The latter, in conjunction with [27, Lemma 1, page 16], implies that the perturbed operators $\mathbf{H}_\theta(g) = \mathbf{H}_\theta(0) + g\mathbf{V}_\theta$ are a type (A) analytic family of operators for $\theta \in \mathcal{A}_0$ and suitable small g. Since E_0 is an isolated, simple eigenvalue of $\mathbf{H}_\theta(0)$, the analyticity of $\mathbf{H}_\theta(g)$ allows us to apply regular perturbation theory. The next section is devoted to this task.

6.3. **Perturbation series and Fermi's golden rule.** Following Kato [13, Sections II.2 and VII.1] and using Lemma 6.4 we infer that $\mathbf{H}_\theta(g)$ has an eigenvalue near E_0 given by a convergent power series in g. The convergent series is given by

$$E(g) = E_0 + \sum_{j=1}^{\infty} E_j(g), \tag{6.8}$$

where

$$E_j(g) = \sum_{p_1+\cdots+p_j=j-1} \frac{(-1)^j}{j} \mathrm{tr} \prod_{i=1}^{j} g\mathbf{V}_\theta \mathbf{S}_\theta^{(p_i)} \tag{6.9}$$

In view of Lemma 6.4(ii) we see that $E_j(g) = O(g^j)$.

Let us compute the lowest-order terms of the series (6.8). Since $\mathrm{Rank}\,\mathbf{P}_\theta = 1$, $\mathbf{P}_\theta$ can be represented as

$$\mathbf{P}_\theta = \left\langle \cdot, \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix} \right\rangle \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix}$$

with $\chi_m^\theta := S_\theta \chi_m$, where χ_m is the eigenfunction associated with the eigenvalue E_0 of H_{22}. Indeed, $H_{22,\theta} = \chi_m^\theta = S_\theta H_{22} S_\theta^{-1} S_\theta \chi_m = E_0 S_\theta \chi_m = E_0 \chi_m^\theta$ and, consequently,

$$\mathbf{H}_\theta(0) \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix} = \begin{pmatrix} H_{11,\theta} & 0 \\ 0 & H_{22,\theta} \end{pmatrix} \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix} = E_0 \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix}.$$

We compute $E_1(g)$:

$$\begin{aligned} E_1(g) &= \mathrm{tr}\,(g\mathbf{V}_\theta \mathbf{P}_\theta) = g \left\langle \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix}, \mathbf{V}_\theta \begin{pmatrix} 0 \\ \chi_m^\theta \end{pmatrix} \right\rangle_{\mathcal{H}} \\ &= g \left\langle \begin{pmatrix} 0 \\ \chi_m \end{pmatrix}, \mathbf{V} \begin{pmatrix} 0 \\ \chi_m \end{pmatrix} \right\rangle_{\mathcal{H}} = g\langle \chi_m, V_{22}\chi_m \rangle_{L^2(\mathbb{R})}. \end{aligned} \tag{6.10}$$

We see that the first-order term is real and does not contribute to the resonance width. Next, we consider $E_2(g)$. According to (6.9),

$$E_2(g) = -g^2 \mathrm{tr}\,(\mathbf{P}_0 \mathbf{V}_\theta \widehat{\mathbf{R}}_0(\theta; E_0 - i0)\mathbf{V}_\theta \mathbf{P}_0).$$

Due to the standard constancy-in-θ argument (see e.g. [27, pages 55-56]), we may take the limit $\mathrm{Im}\,\theta \to 0$ and in this way we arrive at

$$\begin{aligned} E_2(g) &= -g^2 \mathrm{tr}\,(\mathbf{P}_0 \mathbf{V} \widehat{\mathbf{R}}_0(0; E_0 - i0)\mathbf{V}\mathbf{P}_0) \\ &= -g^2 \left\langle \begin{pmatrix} 0 \\ \chi_m \end{pmatrix}, \mathbf{V}\widehat{\mathbf{R}}_0(0; E_0 - i0)\mathbf{V} \begin{pmatrix} 0 \\ \chi_m \end{pmatrix} \right\rangle_{\mathcal{H}} \\ &= -g^2 \langle \chi_m, V_{21}[(H_{11} - E_0 + i0)^{-1}]\widehat{\ } V_{12}\chi_m \rangle_{L^2} \\ &\quad -g^2 \langle \chi_m, V_{22}[(H_{22} - E_0 + i0)^{-1}]\widehat{\ } V_{22}\chi_m \rangle_{L^2} \\ &= -g^2 \langle V_{12}\chi_m, [(H_{11} - E_0 + i0)^{-1}]\widehat{\ } V_{12}\chi_m \rangle_{L^2} \\ &\quad -g^2 \langle V_{22}\chi_m, [(H_{22} - E_0 + i0)^{-1}]\widehat{\ } V_{22}\chi_m \rangle_{L^2}, \end{aligned} \tag{6.11}$$

where the notation $[(H_{jj} - E_0 + i0)^{-1}]\widehat{\ }$ refers to the reduced resolvent of H_{jj}, $j = 1, 2$.

We restrict our focus to the imaginary part of $E_2(g)$, which determines the resonance width to leading order. For this purpose we introduce

$$R_k = \left((-d^2/dx^2 + W_{kk}(x) - E_0 + \mathfrak{t}_k - i0)^{-1}\right)\widehat{\ }, \quad k = 1, 2,$$

where $\mathfrak{t}_1 = 0$ and $\mathfrak{t}_2 = 1$ are the thresholds. Clearly,

$$\operatorname{Im} E_2(g) = -g^2 \sum_{k=1}^{2} \langle V_{k2}\chi_m (\operatorname{Im} R_k) V_{k2}\chi_m \rangle_{L^2(\mathbb{R})}. \tag{6.12}$$

Now, for $E > 0$, the resolvent equation yields that

$$\begin{aligned} &\operatorname{Im}(-d^2/dx^2 + W_{kk}(x) - E - i0)^{-1} = \\ &\quad = t_k(E + i0)^* \operatorname{Im}(-d^2/dx^2 - E - i0)^{-1} t_k(E + i0), \end{aligned} \tag{6.13}$$

where

$$t_k(\zeta) = \left[I + |W_{kk}|^{1/2}(-\partial_x^2 - \zeta)^{-1}|W_{kk}|^{1/2}\operatorname{Sgn}(W_{kk})\right]^{-1}.$$

The quantities $t_k(E + i0)$ are well-defined in view of Assumption 6.1. Furthermore, again for $E > 0$,

$$\operatorname{Im}(-d^2/dx^2 - E - i0)^{-1} = \frac{\pi}{2\sqrt{E}} \sum_{\sigma=\pm} (\gamma_E^\sigma)^* \gamma_E^\sigma, \tag{6.14}$$

where $\gamma_E^\sigma : H^1 \to \mathbb{C}$ is the trace operator which acts on the first Sobolev space $H^1(\mathbb{R})$ as follows (see, e.g., [18, Section IV.1]),

$$\gamma_E^\sigma \phi := \widehat{\phi}(\sigma\sqrt{E}), \ \ \sigma = \pm, \ \ E > 0.$$

Here, as usual, $\widehat{\phi}$ denotes the Fourier transform of ϕ. Using (6.13) and (6.14) we can rewrite the expression (6.12) in the following way,

$$\begin{aligned} \operatorname{Im} E_2(g) &= -g^2 \sum_{k=1}^{2} \langle V_{k2}\chi_m, (\operatorname{Im} R_k) V_{k2}\chi_m \rangle_{L^2} \\ &= -g^2 \sum_{k=1}^{2} \langle V_{k2}\chi_m, t_k(E_0 - \mathfrak{t}_k + i0)^* \\ &\quad \times \operatorname{Im}(-d^2/dx^2 - E_0 + \mathfrak{t}_k - i0)^{-1} t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m \rangle_{L^2} \end{aligned}$$

$$= -g^2 \sum_{k=1}^{2} \big\langle t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m, \mathrm{Im}\,(-d^2/dx^2 - E_0 + \mathfrak{t}_k - i0)^{-1}$$
$$\times t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m \big\rangle_{L^2}$$
$$= -g^2 \sum_{k=1}^{2} \sum_{\sigma=\pm} \frac{\pi}{2\sqrt{E_0 - \mathfrak{t}_k}} \big\langle \gamma^{\sigma}_{E_0-\mathfrak{t}_k} t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m, \gamma^{\sigma}_{E_0-\mathfrak{t}_k}$$
$$\times t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m \big\rangle_{\mathbb{C}}$$
$$= -g^2 \sum_{k=1}^{2} \sum_{\sigma=\pm} \frac{\pi}{2\sqrt{E_0 - \mathfrak{t}_k}} |\gamma^{\sigma}_{E_0-\mathfrak{t}_k} t_k(E_0 - \mathfrak{t}_k + i0) V_{k2}\chi_m|^2. \qquad (6.15)$$

In this way we have established the following result.

Theorem 6.5. *Let Assumption 6.1 and Assumption 6.3 hold. Let ν_m be a simple eigenvalue of the operator H_{22} defined in (6.2) giving rise to the eigenvalue $E_0 = \nu_m$ embedded in the continuous spectrum of $\mathbf{H}(0)$. Let E_0 satisfy Assumption 6.2. For a small enough coupling constant g, the eigenvalue E_0 of $\mathbf{H}(0)$ turns into a resonance, i.e. $E_0 \notin \sigma(\mathbf{H}(g))$. The coordinates of its corresponding pole is given by (6.9)-(6.11). In particular, Fermi's golden rule takes the explicit form (6.15).*

Remark 6.6. If Assumption 6.2 is not fulfilled, i.e. we have an eigenvalue of H_{22} at the threshold point 0 of the continuous spectrum of $\mathbf{H}(0)$, complex dilation breaks down. An insight into this problem was established in [12]. For abstract Hamiltonians $\mathbf{H}(g)$ having the structure found in (6.1), it was shown that under small off-diagonal perturbations this eigenvalue never moves into the continuous spectrum.

7. Acknowledgments

The author thanks J. Brasche and G. Rozenblum for several useful discussions. Moreover, he is grateful that one of the referees pointed out an error in Part 4 of Section 2 and the other referee made him aware of the paper [29].

References

[1] J. Aguilar, J. M. Combes, "Applications of a commutation formula," Comm. Math. Phys. **22** (1971), 269-310.

[2] T. Aktosun, M. Klaus, "Small-energy asymptotics for the Schrödinger equation on the line," Inverse Problems **17** (2001), 619-632.

[3] T. Aktosun, M. Klaus, C. van der Mee, "Small-energy asymptotics of the scattering matrix for the matrix Schrödinger equation on the line," J. Math. Phys. **42** (2001), 4627-4652.

[4] R. Benguria, M. Loss, "A simple proof of a theorem of Laptev and Weidl," Math. Res. Lett. **7** (2000), 195-203.

[5] M. Sh. Birman, M. Z. Solomjak, *Spectral theory of selfadjoint operators in Hilbert space* (D. Reidel Publishing Co., Dordrecht, 1987).

[6] R. Carmona, J. Lacroix, *Spectral theory of random Schrödinger operators* (Birkhäuser Boston, Inc., Boston, 1990).

[7] V. M. Chabanov, B. N. Zakhariev, I. V. Amirkhanov, "Toward the quantum design of multichannel systems. The inverse problem approach," Ann. Physics **285** (2000), 1-24.
[8] R. F. Curtain, H. J. Zwart, *An Introduction to infinite-dimensional Linear Systems Theory* (Springer, Berlin, New York, 1995).
[9] P. Deift, "Applications of a commutation formula," Duke Math. J. **45** (1978), 267-310.
[10] J. Diestel, and J. J. Uhl Jr., *Vector Measures* (AMS, Providence, 1977).
[11] K.-J. Engel, *Operator Matrices and Systems of Evolution Equations*, Manuscript (Tübingen, 1996).
[12] A. Jensen, M. Melgaard, "Perturbation of eigenvalues embedded at a threshold," Proc. Roy. Soc. Edinburgh **132A** (2002), 163-179.
[13] T. Kato, *Perturbation Theory for Linear Operators*, 2nd ed. (Springer, New York, 1976).
[14] K. A. Kiers, W. van Dijk, W., "Scattering in one dimension: The coupled Schrödinger equation, threshold behaviour and Levinson's theorem," J. Math. Phys. **37** (1996), 6033-6059.
[15] M. Klaus, "On the bound state of Schrödinger operators in one dimension," Ann. Physics **108** (1977), 288-300.
[16] M. Klaus, B. Simon, "Coupling constant thresholds in nonrelativistic quantum mechanics. I. Short-range two-body case," Ann. Physics **130** (1980), 251-281.
[17] M. Klaus, "Some applications of the Birman-Schwinger principle," Helv. Phys. Acta **55** (1982/83), 49-68.
[18] S. T. Kuroda, *An introduction to scattering theory.* Lecture Notes Series 51 (Aarhus Universitet, Matematisk Institut, Aarhus, 1978).
[19] L. D. Landau, E. M. Lifshitz, *Quantum mechanics: Non-relativistic theory. Course of Theoretical Physics, Vol. 3* (Pergamon Press Ltd., London-Paris, 1958).
[20] A. Laptev, T. Weidl, "Sharp Lieb-Thirring inequalities in high dimensions," Acta Math. **184** (2000), 87-111.
[21] M. Melgaard, "Spectral properties at a threshold for two-channel Hamiltonians. I. Abstract theory," J. Math. Anal. Appl. **256** (2001), 281-303.
[22] M. Melgaard, "Spectral properties at a threshold for two-channel Hamiltonians. II. Applications to scattering theory," J. Math. Anal. Appl. **256** (2001), 568-586.
[23] M. Melgaard, "New approach to quantum scattering near the lowest Landau threshold for a Schrödinger operator with a constant magnetic field," Few-Body Systems **32** (2002), 1-22.
[24] R. Mennicken, A. K. Motovilov, "Operator interpretation of resonances arising in spectral problems for 2×2 matrix Hamiltonians," in *Mathematical results in quantum mechanics (Prague, 1998)*, Oper. Theory Adv. Appl. **108**, edited by J. Dittrich, P. Exner and M. Tater (Birkhäuser, Basel, 1999), pp. 315–322.
[25] M. Reed, B. Simon, *Methods of modern mathematical physics. I: Functional analysis* (Academic Press, New York, 1980).
[26] M. Reed, M., B. Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness* (Academic Press, New York, 1975).
[27] M. Reed, B. Simon, *Methods of modern mathematical physics. IV: Analysis of operators* (Academic Press Inc., London, 1978).
[28] M. Schechter, *Operator methods in quantum mechanics* (North-Holland Publishing Co., New York, Amsterdam, 1981).
[29] P. Šeba, "Spectral properties of Schrödinger operators with matrix potentials. II.," J. Phys. A: Math. Gen. **19** (1986), 2573-2581.

[30] B. Simon, *Quantum mechanics for Hamiltonians defined as quadratic forms* (Princeton University Press, Princeton, 1971).
[31] B. Simon, "The bound state of weakly coupled Schrödinger operators in one and two dimensions," Ann. Phys. **97** (1976), 279-288.
[32] W. Thirring, *A Course in Mathematical Physics. Vol. 3. Quantum Mechanics of Atoms and Molecules.* Lecture Notes in Physics **141**. (Springer-Verlag, New York-Vienna, 1981).
[33] C. Tretter, "Spectral issues for block matrices," in *Differential Equations and Mathematical Physics (Birmingham, AL, 1999)*, AMS/IP Stud Adv. Math. **16**, edited by R. Weikard, G. Weinstein (Amer. Math. Soc., Providence, RI, 2000), pp. 407-423.

Department of Mathematics, Chalmers University of Technology, and Göteborg University, Eklandagatan 86, S-412 96 Göteborg, Sweden

E-mail address: melgaard@math.chalmers.se