

# A reverse entropy power inequality for log-concave random vectors

Keith Ball, Piotr Nayar\* and Tomasz Tkocz

18/09/2015

## Abstract

We prove that the exponent of the entropy of one dimensional projections of a log-concave random vector defines a  $1/5$ -seminorm. We make two conjectures concerning reverse entropy power inequalities in the log-concave setting and discuss some examples.

**2010 Mathematics Subject Classification.** Primary 94A17; Secondary 52A40, 60E15.

**Key words.** entropy, log-concave, reverse entropy power inequality.

## 1 Introduction

One of the most significant and mathematically intriguing quantities studied in information theory is the *entropy*. For a random variable  $X$  with density  $f$  its entropy is defined as

$$\mathcal{S}(X) = \mathcal{S}(f) = - \int_{\mathbb{R}} f \ln f \quad (1)$$

provided this integral exists (in the Lebesgue sense). Note that the entropy is translation invariant and  $\mathcal{S}(bX) = \mathcal{S}(X) + \ln |b|$  for any nonzero  $b$ . If  $f$  belongs to  $L_p(\mathbb{R})$  for some  $p > 1$ , then by the concavity of the logarithm and Jensen's inequality  $\mathcal{S}(f) > -\infty$ . If  $\mathbb{E}X^2 < \infty$ , then comparison with the standard Gaussian density and again Jensen's inequality yields  $\mathcal{S}(X) < \infty$ . Particularly, the entropy of a log-concave random variable is well defined and finite. Recall that a random vector in  $\mathbb{R}^n$  is called log-concave if it has a density of the form  $e^{-\psi}$  with  $\psi : \mathbb{R}^n \rightarrow (-\infty, +\infty]$  being a convex function.

---

\*Supported in part by the Institute for Mathematics and its Applications with funds provided by the National Science Foundation; supported in part by NCN grant DEC-2012/05/B/ST1/00412

The entropy power inequality (EPI) says that

$$e^{\frac{2}{n}\mathcal{S}(X+Y)} \geq e^{\frac{2}{n}\mathcal{S}(X)} + e^{\frac{2}{n}\mathcal{S}(Y)}, \quad (2)$$

for independent random vectors  $X$  and  $Y$  in  $\mathbb{R}^n$  provided that all the entropies exist. Stated first by Shannon in his seminal paper [22] and first rigorously proved by Stam in [23] (see also [6]), it is often referred to as the Shannon-Stam inequality and plays a crucial role in information theory and elsewhere (see the survey [16]). Using the AM-GM inequality, the EPI can be *linearised*: for every  $\lambda \in [0, 1]$  and independent random vectors  $X, Y$  we have

$$\mathcal{S}(\sqrt{\lambda}X + \sqrt{1-\lambda}Y) \geq \lambda\mathcal{S}(X) + (1-\lambda)\mathcal{S}(Y) \quad (3)$$

provided that all the entropies exist. This formulation is in fact equivalent to (2) as first observed by Lieb in [20], where he also shows how to derive (3) from Young's inequality with sharp constants. Several other proofs of (3) are available, including refinements [13], [15], [26], versions for the Fisher information [11] and recent techniques of the minimum mean-square error [25].

If  $X$  and  $Y$  are independent and identically distributed random variables (or vectors), inequality (3) says that the entropy of the normalised sum

$$X_\lambda = \sqrt{\lambda}X + \sqrt{1-\lambda}Y \quad (4)$$

is at least as big as the entropy of the summands  $X$  and  $Y$ ,  $\mathcal{S}(X_\lambda) \geq \mathcal{S}(X)$ . It is worth mentioning that this phenomenon has been quantified, first in [12], which has deep consequences in probability (see the pioneering work [4] and its sequels [1, 2] which establish the rate of convergence in the entropic central limit theorem and the “second law of probability” of the entropy growth, as well as the independent work [18], with somewhat different methods). In the context of log-concave vectors, Ball and Nguyen in [5] establish dimension free lower bounds on  $\mathcal{S}(X_{1/2}) - \mathcal{S}(X)$  and discuss connections between the entropy and major conjectures in convex geometry; for the latter see also [10].

In general, the EPI cannot be reversed. In [7], Proposition V.8, Bobkov and Christyakov find a random vector  $X$  with a finite entropy such that  $\mathcal{S}(X+Y) = \infty$  for every independent of  $X$  random vector  $Y$  with finite entropy. However, for log-concave vectors and, more generally, convex measures, Bobkov and Madiman have recently addressed the question of reversing the EPI (see [8, 9]). They show that for any pair  $X, Y$  of independent log-concave random vectors in  $\mathbb{R}^n$ , there are linear volume preserving maps  $T_1, T_2 : \mathbb{R}^n \rightarrow \mathbb{R}^n$  such that

$$e^{\frac{2}{n}\mathcal{S}(T_1(X)+T_2(Y))} \leq C(e^{\frac{2}{n}\mathcal{S}(X)} + e^{\frac{2}{n}\mathcal{S}(Y)}),$$

where  $C$  is some universal constant.

The goal of this note is to further investigate in the log-concave setting some new forms of what could be called a reverse EPI. In the next section we present our results. The last section is devoted to their proofs.

## Acknowledgements

The authors would like to thank Assaf Naor for pointing out the Aoki-Rolewicz theorem as well as for fruitful discussions without which Theorem 1 would not have been discovered. They are also indebted to Mokshay Madiman for his help with tracking down several references.

## 2 Main results and conjectures

Suppose  $X$  is a symmetric log-concave random vector in  $\mathbb{R}^n$ . Then any projection of  $X$  on a certain direction  $v \in \mathbb{R}^n$ , that is the random variable  $\langle X, v \rangle$  is also log-concave. Here  $\langle \cdot, \cdot \rangle$  denotes the standard scalar product in  $\mathbb{R}^n$ . If we know the entropies of projections in, say two different directions, can we say anything about the entropy of projections in related directions? We make the following conjecture.

**Conjecture 1.** Let  $X$  be a symmetric log-concave random vector in  $\mathbb{R}^n$ . Then the function

$$N_X(v) = \begin{cases} e^{\mathcal{S}(\langle v, X \rangle)} & v \neq 0, \\ 0 & v = 0 \end{cases}$$

defines a norm on  $\mathbb{R}^n$ .

The homogeneity of  $N_X$  is clear. To check the triangle inequality, we have to answer really a two-dimensional question: *is it true that for a symmetric log-concave random vector  $(X, Y)$  in  $\mathbb{R}^2$  we have*

$$e^{\mathcal{S}(X+Y)} \leq e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)}? \tag{5}$$

Indeed, this applied to the vector  $(\langle u, X \rangle, \langle v, X \rangle)$  which is also log-concave yields  $N_X(u+v) \leq N_X(u) + N_X(v)$ . Inequality (5) can be seen as a reverse EPI, cf. (2). It is not too difficult to show that this inequality holds up to a multiplicative constant.

**Proposition 1.** *Let  $(X, Y)$  be a symmetric log-concave random vector on  $\mathbb{R}^2$ . Then*

$$e^{\mathcal{S}(X+Y)} \leq e(e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)}).$$

*Proof.* The argument relies on the well-known observation that for a log-concave density  $f: \mathbb{R} \rightarrow [0, +\infty)$  its maximum and entropy are related (see for example [5] or [10]),

$$-\ln \|f\|_\infty \leq \mathcal{S}(f) \leq 1 - \ln \|f\|_\infty. \quad (6)$$

Suppose that  $w$  is an even log-concave density of  $(X, Y)$ . The densities of  $X, Y$  and  $X + Y$  equal respectively

$$f(x) = \int w(x, t) dt, \quad g(x) = \int w(t, x) dt, \quad h(x) = \int w(x - t, t) dt. \quad (7)$$

They are even and log-concave, hence attain their maximum at zero. By the result of Ball (Busemann's theorem for symmetric log-concave measures, see [3]), the function  $\|x\|_w = (\int w(tx) dt)^{-1}$  is a norm on  $\mathbb{R}^2$ . Particularly,

$$\begin{aligned} \frac{1}{\|h\|_\infty} &= \frac{1}{h(0)} = \frac{1}{\int w(-t, t) dt} = \|e_2 - e_1\|_w \leq \|e_1\|_w + \|e_2\|_w \\ &= \frac{1}{\int w(t, 0) dt} + \frac{1}{\int w(0, t) dt} = \frac{1}{f(0)} + \frac{1}{g(0)} = \frac{1}{\|f\|_\infty} + \frac{1}{\|g\|_\infty}. \end{aligned}$$

Using (6) twice we obtain

$$e^{\mathcal{S}(X+Y)} \leq \frac{e}{\|h\|_\infty} \leq e \cdot \left( \frac{1}{\|f\|_\infty} + \frac{1}{\|g\|_\infty} \right) \leq e \cdot (e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)}).$$

□

Recall that the classical result of Aoki and Rolewicz says that a  $C$ -quasi-norm (1-homogeneous function satisfying the triangle inequality up to a multiplicative constant  $C$ ) is equivalent to some  $\kappa$ -semi-norm ( $\kappa$ -homogeneous function satisfying the triangle inequality) for some  $\kappa$  depending only on  $C$  (to be precise, it is enough to take  $\kappa = \ln 2 / \ln(2C)$ ). See for instance Lemma 1.1 and Theorem 1.2 in [19]. In view of Proposition 1, for every symmetric log-concave random vector  $X$  in  $\mathbb{R}^n$  the function  $N_X(v)^\kappa = e^{\kappa \mathcal{S}(\langle X, v \rangle)}$  with  $\kappa = \frac{\ln 2}{1 + \ln 2}$  is equivalent to some nonnegative  $\kappa$ -semi-norm. Therefore, it is natural to relax Conjecture 1 and ask whether there is a positive universal constant  $\kappa$  such that the function  $N_X^\kappa$  itself satisfies the triangle inequality for every symmetric log-concave random vector  $X$  in  $\mathbb{R}^n$ . Our main result answers this question positively.

**Theorem 1.** *There exists a universal constant  $\kappa > 0$  such that for a symmetric log-concave random vector  $X$  in  $\mathbb{R}^n$  and two vectors  $u, v \in \mathbb{R}^n$  we have*

$$e^{\kappa \mathcal{S}(\langle u+v, X \rangle)} \leq e^{\kappa \mathcal{S}(\langle u, X \rangle)} + e^{\kappa \mathcal{S}(\langle v, X \rangle)}. \quad (8)$$

*Equivalently, for a symmetric log-concave random vector  $(X, Y)$  in  $\mathbb{R}^2$  we have*

$$e^{\kappa \mathcal{S}(X+Y)} \leq e^{\kappa \mathcal{S}(X)} + e^{\kappa \mathcal{S}(Y)}. \quad (9)$$

*In fact, we can take  $\kappa = 1/5$ .*

*Remark 1.* If we take  $X$  and  $Y$  to be independent random variables uniformly distributed on the intervals  $[0, t]$  and  $[0, 1]$  with  $t < 1$ , then (9) becomes  $e^{\kappa t/2} \leq 1+t^\kappa$ . Letting  $t \rightarrow 0$  shows that necessarily  $\kappa \leq 1$ . We believe that this is the extreme case and the optimal value of  $\kappa$  equals 1.

*Remark 2.* Inequality (9) with  $\kappa = 1$  can be easily shown for log-concave random vectors  $(X, Y)$  in  $\mathbb{R}^2$  for which one marginal has the same law as the other one rescaled, say  $Y \sim tX$  for some  $t > 0$ . Note that the symmetry of  $(X, Y)$  is not needed here. This fact in the essential case of  $t = 1$  was first observed in [14]. We recall the argument in the next section. Moreover, in that paper the converse was shown as well: given a density  $f$ , the equality

$$\max\{\mathcal{S}(X + Y), X \sim f, Y \sim f\} = \mathcal{S}(2X)$$

holds if and only if  $f$  is log-concave, thus characterizing log-concavity. For some bounds on  $\mathcal{S}(X \pm Y)$  in higher dimensions see [21] and [9].

It will be much more convenient to prove Theorem 1 in an equivalent form, obtained by linearising inequality (9).

**Theorem 2.** *Let  $(X, Y)$  be a symmetric log-concave vector in  $\mathbb{R}^2$  and assume that  $\mathcal{S}(X) = \mathcal{S}(Y)$ . Then for every  $\theta \in [0, 1]$  we have*

$$\mathcal{S}(\theta X + (1 - \theta)Y) \leq \mathcal{S}(X) + \frac{1}{\kappa} \ln(\theta^\kappa + (1 - \theta)^\kappa), \quad (10)$$

where  $\kappa > 0$  is a universal constant. We can take  $\kappa = 1/5$ .

*Remark 3.* Proving Conjecture 1 is equivalent to showing the above theorem with  $\kappa = 1$ .

Notice that in the above reverse EPI we estimate the entropy of linear combinations of summands whose joint distribution is log-concave. This is different from what would be the straightforward reverse form of the EPI (3) for independent summands with weights  $\sqrt{\lambda}$  and  $\sqrt{1 - \lambda}$  preserving variance. Suppose that the summands  $X, Y$  are independent and identically distributed, say with finite variance and recall (4). Then, as we mentioned in the introduction, the EPI says that the function  $[0, 1] \ni \lambda \rightarrow \mathcal{S}(X_\lambda)$  is minimal at  $\lambda = 0$  and  $\lambda = 1$ . Following this logic, reversing the EPI could amount to determining the  $\lambda$  for which the maximum of this function occurs. Our next result shows that the somewhat natural guess of  $\lambda = 1/2$  is false in general.

**Proposition 2.** *For each positive  $\lambda_0 < \frac{1}{2(2+\sqrt{2})}$  there is a symmetric continuous random variable  $X$  of finite variance for which  $\mathcal{S}(X_{\lambda_0}) > \mathcal{S}(X_{1/2})$ .*

Nevertheless, we believe that in the log-concave setting the function  $\lambda \mapsto \mathcal{S}(X_\lambda)$  should behave nicely.

**Conjecture 2.** Let  $X$  and  $Y$  be independent copies of a log-concave random variable. Then the function

$$\lambda \mapsto \mathcal{S}(\sqrt{\lambda}X + \sqrt{1-\lambda}Y)$$

is concave on  $[0, 1]$ .

## 3 Proofs

### 3.1 Theorems 1 and 2 are equivalent

To see that Theorem 2 implies Theorem 1 let us take a symmetric log-concave random vector  $(X, Y)$  in  $\mathbb{R}^2$  and take  $\theta$  such that  $\mathcal{S}(X/\theta) = \mathcal{S}(Y/(1-\theta))$ , that is,  $\theta = e^{\mathcal{S}(X)}/(e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)}) \in [0, 1]$ . Applying Theorem 2 with the vector  $(X/\theta, Y/(1-\theta))$  and using the identity  $\mathcal{S}(X/\theta) = \mathcal{S}(X) - \ln \theta = -\ln(e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)})$  gives

$$\mathcal{S}(X + Y) \leq \mathcal{S}(X/\theta) + \frac{1}{\kappa} \ln \left( \frac{e^{\kappa \mathcal{S}(X)} + e^{\kappa \mathcal{S}(Y)}}{(e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)})^\kappa} \right) = \frac{1}{\kappa} \ln (e^{\kappa \mathcal{S}(X)} + e^{\kappa \mathcal{S}(Y)}),$$

so (9) follows.

To see that Theorem 1 implies Theorem 2, take a log-concave vector  $(X, Y)$  with  $\mathcal{S}(X) = \mathcal{S}(Y)$  and apply (9) to the vector  $(\theta X, (1-\theta)Y)$ , which yields

$$\begin{aligned} \mathcal{S}(\theta X + (1-\theta)Y) &\leq \frac{1}{\kappa} \ln (\theta^\kappa e^{\kappa \mathcal{S}(X)} + (1-\theta)^\kappa e^{\kappa \mathcal{S}(Y)}) \\ &= \mathcal{S}(X) + \frac{1}{\kappa} \ln (\theta^\kappa + (1-\theta)^\kappa). \end{aligned}$$

### 3.2 Proof of Remark 2

Let  $w: \mathbb{R}^2 \rightarrow [0, +\infty)$  be the density of such a vector and let  $f, g, h$  be the densities of  $X, Y, X + Y$  as in (7). The assumption means that  $f(x) = tg(tx)$ . By convexity,

$$\mathcal{S}(X + Y) = \inf \left\{ - \int h \ln p, p \text{ is a probability density on } \mathbb{R} \right\}.$$

Using Fubini's theorem and changing variables yields

$$\begin{aligned} - \int h \ln p &= - \iint w(x, y) \ln p(x + y) \, dx dy \\ &= -\theta(1-\theta) \iint w(\theta x, (1-\theta)y) \ln p(\theta x + (1-\theta)y) \, dx dy \end{aligned}$$

for every  $\theta \in (0, 1)$  and a probability density  $p$ . If  $p$  is log-concave we get

$$\begin{aligned} \mathcal{S}(X + Y) &\leq -\theta^2(1 - \theta) \iint w(\theta x, (1 - \theta)y) \ln p(x) \, dx dy \\ &\quad - \theta(1 - \theta)^2 \iint w(\theta x, (1 - \theta)y) \ln p(y) \, dx dy \\ &= -\theta^2 \int f(\theta x) \ln p(x) dx - (1 - \theta)^2 \int g((1 - \theta)y) \ln p(y) dy. \end{aligned}$$

Set

$$p(x) = \theta f(\theta x) = t\theta g(t\theta x)$$

with  $\theta$  such that  $t\theta = 1 - \theta$ . Then the last expression becomes

$$\theta \mathcal{S}(X) + (1 - \theta) \mathcal{S}(Y) - \theta \ln \theta - (1 - \theta) \ln(1 - \theta).$$

Since  $\mathcal{S}(Y) = \mathcal{S}(X) + \ln t = \mathcal{S}(X) + \ln \frac{1-\theta}{\theta}$ , we thus obtain

$$\mathcal{S}(X + Y) \leq \mathcal{S}(X) - \ln \theta = \mathcal{S}(X) + \ln(1 + t) = \ln(e^{\mathcal{S}(X)} + e^{\mathcal{S}(Y)}).$$

### 3.3 Proof of Theorem 2

The idea of our proof of Theorem 2 is very simple. For small  $\theta$  we bound the quantity  $\mathcal{S}(\theta X + (1 - \theta)Y)$  by estimating its derivative. To bound it for large  $\theta$ , we shall crudely apply Proposition 1. The exact bound based on estimating the derivative reads as follows.

**Proposition 3.** *Let  $(X, Y)$  be a symmetric log-concave random vector on  $\mathbb{R}^2$ . Assume that  $\mathcal{S}(X) = \mathcal{S}(Y)$  and let  $0 \leq \theta \leq \frac{1}{2(1+e)}$ . Then*

$$\mathcal{S}(\theta X + (1 - \theta)Y) \leq \mathcal{S}(X) + 60(1 + e)\theta. \quad (11)$$

The main ingredient of the proof of the above proposition is the following lemma. We postpone its proof until the next subsection.

**Lemma 1.** *Let  $w : \mathbb{R}^2 \rightarrow \mathbb{R}_+$  be an even log-concave function. Define  $f(x) = \int w(x, y) dy$  and  $\gamma = \int w(0, y) dy / \int w(x, 0) dx$ . Then we have*

$$\iint \frac{-f'(x)}{f(x)} y w(x, y) dx dy \leq 30\gamma \int w.$$

*Proof of Proposition 3.* For  $\theta = 0$  both sides of inequality (11) are equal. It is therefore enough to prove that  $\frac{d}{d\theta} \mathcal{S}(\theta X + (1 - \theta)Y) \leq 60(1 + e)$  for  $0 \leq \theta \leq \frac{1}{2(1+e)}$ . Let  $f_\theta$  be the density of  $X_\theta = \theta X + (1 - \theta)Y$ . Note that  $f_\theta = e^{-\varphi_\theta}$ , where  $\varphi_\theta$  is

convex. Let  $\frac{d\varphi_\theta}{d\theta} = \Phi_\theta$  and  $\frac{df_\theta}{d\theta} = F_\theta$ . Then  $\Phi_\theta = -F_\theta/f_\theta$ . Using the chain rule we get

$$\begin{aligned}\frac{d}{d\theta}S(\theta X + (1 - \theta)Y) &= -\frac{d}{d\theta}\mathbb{E}\ln f_\theta = \frac{d}{d\theta}\mathbb{E}\varphi_\theta(X_\theta) \\ &= \mathbb{E}\Phi_\theta(X_\theta) + \mathbb{E}\varphi'_\theta(X_\theta)(X - Y).\end{aligned}$$

Moreover,

$$\begin{aligned}\mathbb{E}\Phi_\theta(X_\theta) &= -\mathbb{E}F_\theta(X_\theta)/f_\theta(X_\theta) = -\int F_\theta(x)dx \\ &= -\frac{d}{d\theta}\int f_\theta(x)dx = 0.\end{aligned}$$

Let  $Z_\theta = (X_\theta, X - Y)$  and let  $w_\theta$  be the density of  $Z_\theta$ . Using Lemma 1 with  $w = w_\theta$  gives

$$\begin{aligned}\frac{d}{d\theta}S(\theta X + (1 - \theta)Y) &= -\mathbb{E}\left(\frac{f'_\theta(X_\theta)}{f_\theta(X_\theta)}(X - Y)\right) \\ &= -\int \frac{f_\theta(x)}{f_\theta(x)}yw_\theta(x, y)dx dy \leq 30\gamma_\theta,\end{aligned}$$

where  $\gamma_\theta = \int w_\theta(0, y)dy / \int w_\theta(x, 0)dx$ . It suffices to show that  $\gamma_\theta \leq 2(1 + e)$  for  $0 \leq \theta \leq \frac{1}{2(1+e)}$ . Let  $w$  be the density of  $(X, Y)$ . Then  $w_\theta(x, y) = w(x + (1 - \theta)y, x - \theta y)$ . To finish the proof we again use the fact that  $\|v\|_w = (\int w(tv)dt)^{-1}$  is a norm. Note that

$$\gamma_\theta = \frac{\int w_\theta(0, y)dy}{\int w_\theta(x, 0)dx} = \frac{\int w((1 - \theta)y, -\theta y)dy}{\int w(x, x)dx} = \frac{\|e_1 + e_2\|_w}{\|(1 - \theta)e_1 - \theta e_2\|_w}.$$

Let  $f(x) = \int w(x, y)dy$  and  $g(x) = \int w(y, x)dy$  be the densities of real log-concave random variables  $X$  and  $Y$ , respectively. Observe that by (6) we have

$$\|f\|_\infty^{-1} \leq e^{\mathcal{S}(X)} \leq e\|f\|_\infty^{-1}, \quad \|g\|_\infty^{-1} \leq e^{\mathcal{S}(Y)} \leq e\|g\|_\infty^{-1}.$$

Since  $\|f\|_\infty^{-1} = f(0)^{-1} = \|e_1\|_w$ ,  $\|g\|_\infty^{-1} = g(0)^{-1} = \|e_2\|_w$  and  $\mathcal{S}(X) = \mathcal{S}(Y)$ , this gives  $e^{-1} \leq \|e_1\|_w / \|e_2\|_w \leq e$ . Thus, by the triangle inequality

$$\begin{aligned}\gamma_\theta &\leq \frac{\|e_1\|_w + \|e_2\|_w}{(1 - \theta)\|e_1\|_w - \theta\|e_2\|_w} \\ &\leq \frac{(1 + e)\|e_1\|_w}{(1 - \theta)\|e_1\|_w - \theta e\|e_1\|_w} = \frac{1 + e}{1 - \theta(1 + e)} \\ &\leq 2(1 + e).\end{aligned}$$

□

*Proof of Theorem 2.* We can assume that  $\theta \in [0, 1/2]$ . Using Proposition 1 with the vector  $(\theta X, (1 - \theta)Y)$  and the fact that  $\mathcal{S}(X) = \mathcal{S}(Y)$  we get  $\mathcal{S}(\theta X + (1 - \theta)Y) \leq$

$\mathcal{S}(X) + 1$ . Thus, from Proposition 3 we deduce that it is enough to find  $\kappa > 0$  such that

$$\min\{1, 60(1+e)\theta\} \leq \kappa^{-1} \ln(\theta^\kappa + (1-\theta)^\kappa), \quad \theta \in [0, 1/2]$$

(if  $60(1+e)\theta < 1$  then  $\theta < \frac{1}{2(1+e)}$  and therefore Proposition 3 indeed can be used in this case). By the concavity and monotonicity of the right hand side it is enough to check this inequality at  $\theta_0 = (60(1+e))^{-1}$ , that is, we have to verify the inequality  $e^\kappa \leq \theta_0^\kappa + (1-\theta_0)^\kappa$ . We check that this is true for  $\kappa = 1/5$ .  $\square$

### 3.4 Proof of Lemma 1

We start off by establishing two simple and standard lemmas. The second one is a limiting case of the so-called Grünbaum theorem, see [17] and [24].

**Lemma 2.** *Let  $f : \mathbb{R} \rightarrow \mathbb{R}_+$  be an even log-concave function. For  $\beta > 0$  define  $a_\beta$  by*

$$a_\beta = \sup\{x > 0, f(x) \geq e^{-\beta} f(0)\}.$$

*Then we have*

$$2e^{-\beta} a_\beta \leq \frac{1}{f(0)} \int f \leq 2(1 + \beta^{-1} e^{-\beta}) a_\beta.$$

*Proof.* Since  $f$  is even and log-concave, it is maximal at zero and nonincreasing on  $[0, \infty)$ . Consequently, the left hand inequality immediately follows from the definition of  $a_\beta$ . By comparing  $\ln f$  with an appropriate linear function, log-concavity also guarantees that  $f(x) \leq f(0)e^{-\beta \frac{x}{a_\beta}}$  for  $|x| > a_\beta$ , hence

$$\int f \leq 2a_\beta f(0) + 2 \int_{a_\beta}^{\infty} f(0) e^{-\beta \frac{x}{a_\beta}} dx = 2a_\beta f(0) + 2f(0) \frac{a_\beta}{\beta} e^{-\beta}$$

which gives the right hand inequality.  $\square$

**Lemma 3.** *Let  $X$  be a log-concave random variable. Let  $a$  satisfy  $\mathbb{P}(X > a) \leq e^{-1}$ . Then  $\mathbb{E}X \leq a$ .*

*Proof.* Without loss of generality assume that  $X$  is a continuous random variable and that  $\mathbb{P}(X > a) = e^{-1}$ . Moreover, the statement is translation invariant, so we can assume that  $a = 0$ . Let  $e^{-\varphi}$  be the density of  $X$ , where  $\varphi$  is convex. There exists a function  $\psi$  of the form

$$\psi(x) = \begin{cases} ax + b, & x \geq L \\ +\infty, & x < L \end{cases}$$

such that  $\psi(0) = \varphi(0)$  and  $e^{-\psi}$  is the probability density of a random variable  $Y$  with  $\mathbb{P}(Y > a) = e^{-1}$ . One can check, using convexity of  $\varphi$ , that  $\mathbb{E}X \leq \mathbb{E}Y$ . We have  $1 = \int e^{-\psi} = \frac{1}{a} e^{-(b+aL)}$  and  $e^{-1} = \int_0^\infty e^{-\psi} = \frac{1}{a} e^{-b}$ . It follows that  $aL = -1$  and we have  $\mathbb{E}X \leq \mathbb{E}Y = \frac{1}{a} (L + \frac{1}{a}) e^{-(b+aL)} = 0$ .  $\square$

We are ready to prove Lemma 1.

*Proof of Lemma 1.* Without loss of generality let us assume that  $w$  is strictly log-concave and  $w(0) = 1$ . First we derive a pointwise estimate on  $w$  which will enable us to obtain good pointwise bounds on the quantity  $\int yw(x, y)dy$ , relative to  $f(x)$ . To this end, set unique positive parameters  $a$  and  $b$  to be such that  $w(a, 0) = e^{-1} = w(0, b)$ . Consider  $l \in (0, a)$ . We have

$$w(-l, 0) = w(l, 0) \geq w(a, 0)^{l/a} w(0, 0)^{1-l/a} = e^{-l/a}.$$

Fix  $x > 0$  and let  $y > \frac{b}{a}x + b$ . Let  $l$  be such that the line passing through the points  $(0, b)$  and  $(x, y)$  intersect the  $x$ -axis at  $(-l, 0)$ , that is  $l = \frac{bx}{y-b}$ . Note that  $l \in (0, a)$ . Then

$$\begin{aligned} e^{-1} = w(0, b) &\geq w(x, y)^{b/y} w(-l, 0)^{1-b/y} \geq w(x, y)^{b/y} e^{-\frac{l}{a}(1-b/y)} \\ &= \left[ w(x, y) e^{-\frac{l}{a} \frac{y-b}{y}} \right]^{b/y}, \end{aligned}$$

hence

$$w(x, y) \leq e^{x/a - y/b}, \quad \text{for } x > 0 \text{ and } y > \frac{b}{a}x + b.$$

Let  $X$  be a random variable with log-concave density  $y \mapsto w(x, y)/f(x)$ . Let us take  $\beta = b + b \ln(\max\{f(0), b\})$  and

$$\alpha = \frac{b}{a}x - b \ln f(x) + \beta.$$

Since  $f$  is maximal at zero (as it is an even log-concave function), we check that

$$\alpha \geq \frac{b}{a}x - b \ln f(0) + \beta \geq \frac{b}{a}x + b,$$

so we can use the pointwise estimate on  $w$  and get

$$\int_{\alpha}^{\infty} w(x, y) dy \leq e^{x/a} \int_{\alpha}^{\infty} e^{-y/b} dy = b e^{x/a - \alpha/b} = \frac{b}{\max\{f(0), b\}} e^{-1} f(x) \leq e^{-1} f(x).$$

This means that  $\mathbb{P}(X > \alpha) \leq e^{-1}$ , which in view of Lemma 3 yields

$$\frac{1}{f(x)} \int yw(x, y) dy = \mathbb{E}X \leq \alpha = \frac{b}{a}x - b \ln f(x) + \beta, \quad \text{for } x > 0.$$

Having obtained this bound, we can easily estimate the quantity stated in the lemma. By the symmetry of  $w$  we have

$$\iint \frac{-f'(x)}{f(x)} yw(x, y) dx dy = 2 \iint_{x>0} \frac{-f'(x)}{f(x)} yw(x, y) dx dy.$$

Since  $f$  decreases on  $[0, \infty)$ , the factor  $-f'(x)$  is nonnegative for  $x > 0$ , thus we can further write

$$\begin{aligned} \iint \frac{-f'(x)}{f(x)} y w(x, y) dx dy &\leq 2 \int_0^\infty -f'(x) \left( \frac{b}{a} x - b \ln f(x) + \beta \right) dx \\ &= 2f(0)(-b \ln f(0) + \beta) + 2 \int_0^\infty f(x) \left( \frac{b}{a} - b \frac{f'(x)}{f(x)} \right) dx \\ &= 2f(0)b \left( 1 + \ln \frac{\max\{f(0), b\}}{f(0)} \right) + \frac{b}{a} \int w + 2f(0)b. \end{aligned}$$

Now we only need to put the finishing touches to this expression. By Lemma 2 applied to the functions  $x \mapsto w(x, 0)$  and  $y \mapsto w(0, y)$  we obtain

$$\frac{b}{a} \leq \frac{e}{2} 2(1 + e^{-1}) \frac{\int w(0, y) dy}{\int w(x, 0) dx} = (e + 1)\gamma$$

and  $b/f(0) \leq e/2$ . Estimating the logarithm yields

$$1 + \ln \frac{\max\{f(0), b\}}{f(0)} \leq \frac{\max\{f(0), b\}}{f(0)} \leq \frac{e}{2}.$$

Finally, by log-concavity,

$$\int w(x, y) dx dy \geq \int \sqrt{w(2x, 0)w(0, 2y)} dx dy = \frac{1}{4} \int \sqrt{w(x, 0)} dx \int \sqrt{w(0, y)} dy$$

and

$$\int w(x, 0) dx \leq \sqrt{w(0, 0)} \int \sqrt{w(x, 0)} dx = \int \sqrt{w(x, 0)} dx.$$

Combining these two estimates we get

$$f(0) = \int w(0, y) dy \leq \int \sqrt{w(0, y)} dy \leq \frac{4 \int w}{\int w(x, 0) dx}$$

and consequently,

$$f(0)b \leq \frac{e}{2} f(0)f(0) \leq 2ef(0) \frac{\int w}{\int w(x, 0) dx} = 2e\gamma \int w.$$

Finally,

$$\iint \frac{-f'(x)}{f(x)} y w(x, y) dx dy \leq (2e^2 + 5e + 1)\gamma \int w$$

and the assertion follows.  $\square$

### 3.5 Proof of Proposition 2

For a real number  $s$  and nonnegative numbers  $\alpha \leq \beta$  we define the following trapezoidal function

$$T_{\alpha, \beta}^s(x) = \begin{cases} 0 & \text{if } x < s \text{ or } x > s + \alpha + \beta, \\ x - s & \text{if } s \leq x \leq s + \alpha, \\ \alpha & \text{if } s + \alpha \leq x \leq s + \beta, \\ s + \alpha + \beta - x & \text{if } s + \beta \leq x \leq s + \alpha + \beta. \end{cases}$$

The motivation is the following convolution identity: for real numbers  $a, a'$  and nonnegative numbers  $h, h'$  such that  $h \leq h'$  we have

$$\mathbf{1}_{[a, a+h]} \star \mathbf{1}_{[a', a'+h']} = T_{h, h'}^{a+a'}. \quad (12)$$

It is also easy to check that

$$\int_{\mathbb{R}} T_{\alpha, \beta}^s = \alpha\beta. \quad (13)$$

We shall need one more formula: for any real number  $s$  and nonnegative numbers  $A, \alpha, \beta$  with  $\alpha \leq \beta$  we have

$$I(A, \alpha, \beta) = \int_{\mathbb{R}} AT_{\alpha, \beta}^s \ln(AT_{\alpha, \beta}^s) = A\alpha\beta \ln(A\alpha) - \frac{1}{2}A\alpha^2. \quad (14)$$

Fix  $0 < a < b = a + h$ . Let  $X$  be a random variable with the density

$$f(x) = \frac{1}{2h} (\mathbf{1}_{[-b, -a]}(x) + \mathbf{1}_{[a, b]}(x)).$$

We shall compute the density  $f_\lambda$  of  $X_\lambda$ . Denote  $u = \sqrt{\lambda}$ ,  $v = \sqrt{1-\lambda}$  and without loss of generality assume that  $\lambda \leq 1/2$ . Clearly,  $f_\lambda(x) = \frac{1}{u}f(\frac{\cdot}{u}) \star \frac{1}{v}f(\frac{\cdot}{v})(x)$ , so by (12) we have

$$\begin{aligned} f_\lambda(x) &= \left( \mathbf{1}_{u[-b, -a]} \star \mathbf{1}_{v[-b, -a]} + \mathbf{1}_{u[a, b]} \star \mathbf{1}_{v[-b, -a]} \right. \\ &\quad \left. + \mathbf{1}_{u[-b, -a]} \star \mathbf{1}_{v[a, b]} + \mathbf{1}_{u[a, b]} \star \mathbf{1}_{v[a, b]} \right)(x) \cdot \frac{1}{(2h)^2 uv} \\ &= \left( \underbrace{T_{uh, vh}^{-(u+v)b}}_{T_1}(x) + \underbrace{T_{uh, vh}^{ua-vb}}_{T_2}(x) + \underbrace{T_{uh, vh}^{-ub+va}}_{T_3}(x) + \underbrace{T_{uh, vh}^{(u+v)a}}_{T_4}(x) \right) \cdot \frac{1}{(2h)^2 uv}. \end{aligned}$$

This symmetric density is superposition of 4 trapezoid functions  $T_1, T_2, T_3, T_4$  which are certain shifts of the same trapezoid function  $T_0 = T_{uh, vh}^0$ . The shifts may overlap depending on the value of  $\lambda$ . Now we shall consider two particular values of  $\lambda$ .

*Case 1:*  $\lambda = 1/2$ . Then  $u = v = 1/\sqrt{2}$ . Notice that  $T_0$  becomes a triangle looking function and  $T_2 = T_3$ , so we obtain

$$f_{1/2}(x) = \frac{1}{2h^2} \left( T_{h/\sqrt{2}, h/\sqrt{2}}^{-b\sqrt{2}} + 2T_{h/\sqrt{2}, h/\sqrt{2}}^{-h/\sqrt{2}} + T_{h/\sqrt{2}, h/\sqrt{2}}^{a\sqrt{2}} \right)(x).$$

If  $h/\sqrt{2} < a\sqrt{2}$  then the supports of the summands are disjoint and with the aid of identity (14) we obtain

$$\mathcal{S}(X_{1/2}) = -2I\left(\frac{1}{2h^2}, \frac{h}{\sqrt{2}}, \frac{h}{\sqrt{2}}\right) - I\left(\frac{1}{h^2}, \frac{h}{\sqrt{2}}, \frac{h}{\sqrt{2}}\right) = \ln(2h) + \frac{1}{2}.$$

*Case 2: small  $\lambda$ .* Now we choose  $\lambda = \lambda_0$  so that the supports of  $T_1$  and  $T_2$  intersect in such a way that the down-slope of  $T_1$  adds up to the up-slope of  $T_2$  giving a flat piece. This happens when  $-b(u+v) + vh = ua - bv$ , that is,

$$\sqrt{\frac{1-\lambda_0}{\lambda_0}} = \frac{v}{u} = \frac{a+b}{h} = 2\frac{a}{h} + 1. \quad (15)$$

The earlier condition  $a/h > 1/2$  implies that  $\lambda_0 < 1/5$ . With the above choice for  $\lambda$  we have  $T_1 + T_2 = T_{uh,2vh}^{-b(u+v)}$ , hence by symmetry

$$f_\lambda = \left( T_{uh,2vh}^{-b(u+v)} + T_{uh,2vh}^{-ub+va} \right) \cdot \frac{1}{(2h)^2 uv}.$$

As long as  $-ub+va > 0$ , the supports of these two trapezoid functions are disjoint. Given our choice for  $\lambda$ , this is equivalent to  $v/u > b/a = 1 + h/a = 1 + 2/(v/u - 1)$ , or putting  $v/u = \sqrt{1/\lambda_0 - 1}$ , to  $\lambda_0 < \frac{1}{2(2+\sqrt{2})}$ . Then also  $\lambda_0 < 1/5$  and we get

$$\mathcal{S}(X_\lambda) = -2I \left( \frac{1}{(2h)^2 uv}, uh, 2vh \right) = \ln(4vh) + \frac{u}{4v} = \ln(4h\sqrt{1-\lambda_0}) + \frac{1}{4} \sqrt{\frac{\lambda_0}{1-\lambda_0}}.$$

We have

$$\mathcal{S}(X_{\lambda_0}) - \mathcal{S}(X_{1/2}) = \ln 2 - \frac{1}{2} + \ln \sqrt{1-\lambda_0} + \frac{1}{4} \sqrt{\frac{\lambda_0}{1-\lambda_0}}.$$

We check that the right hand side is positive for  $\lambda_0 < \frac{1}{2(2+\sqrt{2})}$ . Therefore, we have shown that for each such  $\lambda_0$  there is a choice for the parameters  $a$  and  $h$  (given by (15)), and hence a random variable  $X$ , for which  $\mathcal{S}(X_{\lambda_0}) > \mathcal{S}(X_{1/2})$ .

## References

- [1] Artstein, S., Ball, K. M., Barthe, F., Naor, A., On the rate of convergence in the entropic central limit theorem, *Probab. Theory Related Fields* 129 (2004), no. 3, 381–390.
- [2] Artstein, S., Ball, K. M., Barthe, F., Naor, A., Solution of Shannon’s problem on the monotonicity of entropy, *J. Amer. Math. Soc.* 17 (2004), no. 4, 975–982.
- [3] Ball, K., Logarithmically concave functions and sections of convex sets in  $\mathbb{R}^n$ , *Studia Math.* 88 (1988), 69–84.
- [4] Ball, K., Barthe, F., Naor, A., Entropy jumps in the presence of a spectral gap, *Duke Math. J.* 119 (2003), no. 1, 41–63.
- [5] Ball, K., Nguyen, V. H., Entropy jumps for isotropic log-concave random vectors and spectral gap, *Studia Math.* 213 (2012), no. 1, 81–96.

- [6] Blachman, N. M., The convolution inequality for entropy powers, *IEEE Trans. Information Theory* IT-11 (1965) 267–271.
- [7] Bobkov, S. G., Chistyakov, G. P., Entropy power inequality for the Rényi entropy, *IEEE Transactions on Information Theory*, vol. 61 (2015), no. 2, pp. 708–714.
- [8] Bobkov, S. G., Madiman, M., Reverse Brunn-Minkowski and reverse entropy power inequalities for convex measures, *J. Funct. Anal.* 262 (2012), no. 7, 3309–3339.
- [9] Bobkov, S. G., Madiman, M., On the problem of reversibility of the entropy power inequality. *Limit theorems in probability, statistics and number theory*, 61–74, Springer Proc. Math. Stat., 42, Springer, Heidelberg, 2013.
- [10] Bobkov, S. G., Madiman, M., The entropy per coordinate of a random vector is highly constrained under convexity conditions, *IEEE Trans. Inform. Theory*, 57 (2011), no. 8, 4940–4954.
- [11] Carlen, E. A., Superadditivity of Fisher’s information and logarithmic Sobolev inequalities, *J. Funct. Anal.* 101 (1991), no. 1, 194–211.
- [12] Carlen, E. A., Soffer, A., Entropy production by block variable summation and central limit theorems, *Comm. Math. Phys.* 140 (1991), no. 2, 339–371.
- [13] Costa, M. H. M., A new entropy power inequality, *IEEE Trans. Inform. Theory* 31 (1985), no. 6, 751–760.
- [14] Cover, T. M., Zhang, Z., On the maximum entropy of the sum of two dependent random variables. *IEEE Trans. Inform. Theory* 40 (1994), no. 4, 1244–1246.
- [15] Dembo, A., Simple proof of the concavity of the entropy power with respect to added Gaussian noise, *IEEE Trans. Inform. Theory* 35 (1989), no. 4, 887–888.
- [16] Dembo, A., Cover, T. M., Thomas, J. A., Information-theoretic inequalities, *IEEE Trans. Inform. Theory* 37 (1991), no. 6, 1501–1518.
- [17] Grünbaum, B., Partitions of mass-distributions and of convex bodies by hyperplanes, *Pacific J. Math.* 10 (1960), 1257–1261.
- [18] Johnson, O., Barron, A., Fisher information inequalities and the central limit theorem, *Probab. Theory Related Fields* 129 (2004), no. 3, 391–409.

- [19] Kalton, N. J., Peck, N. T., Roberts, James W., An F-space sampler. London Mathematical Society Lecture Note Series, 89. *Cambridge University Press, Cambridge*, 1984.
- [20] Lieb, E. H., Proof of an entropy conjecture of Wehrl, *Comm. Math. Phys.* 62 (1978), no. 1, 35–41.
- [21] Mokshay Madiman, M., Kontoyiannis, I., The Ruzsa divergence for random elements in locally compact abelian groups, arXiv:1508.04089.
- [22] Shannon, C. E., A mathematical theory of communication, *Bell System Tech. J.* 27, (1948). 379–423, 623–656.
- [23] Stam, A. J., Some inequalities satisfied by the quantities of information of Fisher and Shannon, *Information and Control* 2 1959, 101–112.
- [24] Szarek, S., On measures of symmetry and floating bodies, arXiv:1302.2076
- [25] Verdú, S., Guo, D., A simple proof of the entropy-power inequality, *IEEE Trans. Inform. Theory* 52 (2006), no. 5, 2165–2166.
- [26] Villani, C., A short proof of the "concavity of entropy power", *IEEE Trans. Inform. Theory* 46 (2000), no. 4, 1695–1696.

Keith Ball\*, k.m.ball@warwick.ac.uk

Piotr Nayar†, nayar@mimuw.edu.pl

Tomasz Tkocz\*, t.tkocz@warwick.ac.uk

\*Mathematics Institute, University of Warwick,  
Coventry CV4 7AL,  
UK

†Institute of Mathematics & Applications,  
Minneapolis MN 55455  
United States