

Prevalence and Correlates of Physical Spousal Violence Against Women in Slum and Nonslum Areas of Urban Bangladesh

Journal of Interpersonal Violence
26(13) 2592–2618
© The Author(s) 2011
Reprints and permission: <http://www.sagepub.com/journalsPermissions.nav>
DOI: 10.1177/0886260510388282
<http://jiv.sagepub.com>



William Sambisa¹, Gustavo Angeles¹, Peter M. Lance¹, Ruchira T. Naved², and Juliana Thornton¹

Abstract

This study explores the prevalence and correlates of past-year physical violence against women in slum and nonslum areas of urban Bangladesh. The authors use multivariate logistic regression to analyze data from the 2006 Urban Health Survey, a population-based survey of 9,122 currently married women aged between 15 and 49 who were selected using a multistage cluster sampling design. The prevalence of reported past-year physical spousal violence is 31%. Prevalence of past-year physical spousal violence is higher in slums (35%) than in non-slums (20%). Slapping/arm-twisting and pushing/shaking/throwing something at the women are the most commonly reported acts of physical abuse. Multivariate analysis shows that the risk of physical spousal abuse is lower among older women, women with post-primary education, and those belonging to rich households and women whose husbands considered their opinion in decision making. Women are at higher risk of abuse if they had many children, believe that married woman should work if the husband is not making enough money, and approve wife-beating norms. This

¹University of North Carolina at Chapel Hill

²ICDDR, B: Center for Health and Population Research, Dhaka, Bangladesh

Corresponding Author:

William Sambisa, Carolina Population Center of the University of North Carolina, Chapel Hill
Email: william.sambisa@gmail.com

study serves to confirm the commonness of physical spousal abuse in urban Bangladesh, demonstrating the seriousness of this multifaceted phenomenon as a social and public health issue. The present findings suggest the need for comprehensive prevention and intervention strategies that capitalize on the interplay of individual and sociocultural factors that cause physical spousal violence. Our study adds to a growing literature documenting domestic violence against women in urban areas of developing south Asian nations.

Keywords

physical spousal violence; wife-beating norms; urban Bangladesh

Introduction

Intimate partner violence (IPV) is a major public health problem engendering serious physical health effects for women, including deaths and disabilities due to injuries (Ellsberg et al., 2008; Plichta & Falik, 2001). In addition, IPV acts as a common cause of mental health problems (Garcia-Moreno, Jansen, Ellsberg, Heise, & Watts, 2005; Krug, Mercy, Dahlberg, & Zwi, 2002; Naved & Akhtar, 2008) and has adverse sexual and reproductive health outcomes (Campbell et al., 2002; Dunkle et al., 2004). During the recent decades, efforts to identify correlates and etiology of IPV have increased significantly in Bangladesh, dominated by studies conducted in rural areas (Bates, Schuler, Islam, & Islam, 2004; Bhuiya, Sharmin, & Hanifi, 2003; Decker et al., 2008; Koenig, Ahmed, Hossain, & Mozumder, 2003; Schuler, Hashemi, Riley, & Akhter, 1996; Silverman, Gupta, Decker, Kapur, & Raj, 2007). These studies suggest that IPV is widespread within the Bangladesh population and that Bangladeshi women are threatened with multiple forms of IPV, including physical, sexual, and emotional abuse (Naved, 2003; Wahed & Bhuiya, 2007). Population-based surveys place lifetime rate of physical IPV at between 42% and 76% and annual rate at between 16% and 67% (Bates et al., 2004; Decker et al., 2008; Garcia-Moreno et al., 2005; Naved, Azim, Bhuiya, & Persson, 2006; Schuler et al., 1996).

Existing research on determinants of IPV in Bangladesh suggests that specific types of risk factors—demographic, socioeconomic, relational and cultural variables—influence the prevalence of domestic violence against women (Johnston & Naved, 2008; Wahed & Bhuiya, 2007). This cross-sectional research indicates the importance of women's relative younger age within marriage as a factor associated with IPV, with risk of experiencing abuse declining with age (Ahmed, van Ginneken, Razzaque, & Alam, 2004; Naved & Persson, 2005). The leading explanatory hypothesis is the lack of

experience among younger women to avoid situations that trigger abuse or lack of kin support and integration among younger couples within the husband's extended family compared to older women (Counts, Brown, & Campbell, 1992). In addition, some studies have shown education as a consistent marker for IPV (Hadi, 2000). According to Naved and Persson, and Koenig et al., women with some level of primary education are less likely to be abused than those without education. The limited existing literature does not report a consistent story on the association between income-generating activities and IPV (Johnston & Naved, 2008). Some studies have shown a negative relationship between IPV and membership in microcredit groups, whereas others showed positive relationship, and still others showed no significant relationship (Chowdhury & Bhuiya, 2001; Koenig et al., 2003; Schuler et al., 1996). In terms of income-generating activities, Bates et al. (2004) and Naved and Persson found that rural women who earn independent incomes experience more IPV than do women who do not earn independent incomes.

Previous research indicates that IPV is associated with poverty (Amin, Khuda, Islam, & Levin, 1997) and having many children (Aklimunnessa, Khan, Kabir, & Mori, 2007). The frustrations and stresses imposed by poverty or financial problems upon family life can lead to abuse (MacMillan & Gartner, 1999), and having many children can limit space, demand more time and attention, and also increase stress levels, leading to greater IPV (Panda & Agarwal, 2005).

International studies with considerable geographic and cultural variation show that women living within societies dominated by patriarchal institutions, unequal power relationships between the genders, and male's control of resources have increased vulnerability to IPV (Dobash & Dobash, 1979). Research in Bangladesh has shown that women are more likely to be abused in relationships where men make more of the key decisions (Ahmed, 2005; Islam et al., 2004). However, some studies in Bangladesh have shown that a woman's increased status as evinced by greater autonomy and control over resources is more protective against IPV (Schuler et al., 1996). There are, however, inconsistencies on the finding on the relationship between women's autonomy and IPV. Koenig et al. (2003) have found that husband's lower decision-making power is positively related to greater IPV. Some research suggests that normative tolerance of wife abuse rooted in patriarchal ideology is strongly associated with IPV (Schuler, Bates, Maseljo, & Islam, 2006).

As predominately rural-focused studies have documented the prevalence and associated factors for partner abuse in Bangladesh, very few studies have examined the nature of IPV against women in urban areas of this country (Naved et al., 2006; Naved & Persson, 2005; Salam, Alim, & Noguchi, 2006).

In addition, even rarer are studies that examine differences in IPV in slum and non-slum areas within Bangladesh cities. There is also a scarcity of such studies in many developing countries (Jeyaseelan et al., 2007). Our study aims to fill this gap. As rapid urbanization is occurring in South Asia and elsewhere, urban areas have become the new frontier for public health concerns (Freudenberg, Galea, & Vlahov, 2005). Campbell and Campbell (2007) argued that current and future patterns of urban growth, combined with advances in the treatment of traditional scourges of communicable diseases, is causing a shift in the burden of diseases toward what is classified as Category 2 (noncommunicable) and Category 3 (injury/violence) conditions. The shift in the burden of public health concerns is argued to have a clear link to features of city growth—poverty and slum formation. Like all health-related behaviors, domestic violence should be understood in relation to the broad ecological and socioeconomic context in which it occurs (Heise, 1998). A better understanding on the nature of IPV in urban areas is needed to identify appropriate targets for prevention and interventions programs aimed at addressing the problem of domestic violence and promoting behavior change where necessary.

In this study, we investigate the nature of past-year physical spousal violence using the ecological model that outlines IPV as a multifaceted phenomenon grounded in an interplay of a multitude of factors operating at different levels of the victim's social ecology (Heise, 1998). These factors are best visualized as concentric circles. The innermost circle contains characteristics of the individual; the second circle refers to the relational context in which abuse takes place. The outermost circle refers to the dominant cultural views and attitudes within the society. Our main assumption is that multiple factors, rather than any one individual factor, are most likely to be responsible for IPV in Bangladesh. Specifically guided by previous findings (e.g., Kishor & Johnson, 2004; Koenig et al., 2003; Naved & Persson, 2005), we hypothesized that slum women would more likely experience physical violence than non-slum women. Adult women rather than teenage women, those with post-primary education and women employed for wages, would less likely experience physical violence. In addition, the risk of physical abuse would be greater for women with higher parity and those who believed that married women should work outside the home if the husband is not making enough money. We also expected that women who reported that it was justified for a husband to beat his wife under various scenarios would more likely be physically abused by their husbands. Furthermore, we hypothesized that those women whose husbands considered their opinion in decision making would less likely experience physical abuse. In sum, our study describes the

prevalence and identifies associated factors of past-year physical spousal violence among reproductive-aged women in slum and non-slum areas of urban Bangladesh using population-based data.

Method

Study Setting and Study Design

This study used data from the 2006 Urban Health Survey (UHS), a population-based cross-sectional household survey covering a target population of Bangladeshi women aged between 10 and 59 and residing in district municipalities and six city corporations: Dhaka, Chittagong, Kulna, Rajshahi, Barisal, and Sylhet. The study protocol, questionnaires, and interview guide were approved by the Institutional Review Boards of the University of North Carolina at Chapel Hill and the Bangladesh Medical Research Council, Dhaka, Bangladesh.

The UHS survey used a multistage cluster sample design with each selected city stratified into slum and non-slum areas for which these urban communities served as the primary sampling unit (PSU). District municipalities were included in the UHS sampling frame as PSUs (without reference to slum/non-slum partition) based on the sample frame from the 2004 Bangladesh Demographic and Health Survey (National Institute of Population Research and Training, Mitra Associates & ORC Macro, 2005). Using the probability proportional to size (PPS) selection method (Levy & Lemeshow, 1965), 510 PSUs were selected (254 from slums, 192 from non-slums, and 64 from district municipalities). In PPS sampling, a PSU's selection probability is proportional to its size measure. PPS sampling is often used in cluster sampling, whereby in our study we selected clusters (i.e., groups of sampling units: slum, non-slums, and district municipalities) of varying sizes in the first stage of selection. A list of households in each PSU was created to form the sampling frame for systematic selection of households. Within each cluster, 25 households were randomly selected. Demographic and socioeconomic information of each selected household and its members were recorded on a household form. Of 15,277 eligible women, 14,191 (93% response rate) completed the survey.

After obtaining informed verbal consent, female research assistants administered face-to-face interviews to all selected women. The interviews were conducted in privacy and in a nonjudgmental manner. For the survey, all questions were translated from English into Bengali, the national language of Bangladesh. Survey items included assessments of demographic

and socioeconomic status, birth history, maternal health, current physical health, mental health using the Self-Reported Questionnaire, HIV/AIDS, and smoking, alcohol, and drug use. The survey also included items about husband's perpetration of physical violence against his wife, which is the focus of our study. The direct and behaviorally explicit questions on physical violence and their time period ("ever" or "past year") were derived from the Revised Conflict Tactics Scale (CTS2; Straus, Hamby, Boney-McCoy, & Sugarman, 1996). The domestic violence questions were adapted for local relevance in consultation with experts in Bangladesh. The study followed the WHO guidelines on ethics and safety in studies on violence against women (World Health Organization, 2001). Assessments on gender role beliefs were included in the domestic violence module as was household decision making and normative acceptance of violence against women.

Study Participants

For the current study, we applied the following inclusion criteria: currently married women of reproductive age (aged 15-49 years) for whom both household decision making and physical violence data were collected. Of 14,191 interviewed women, 1,529 (11%) were dropped from the analysis because they were never married, and importantly, they were not asked questions on domestic violence and household decision making. An additional 1,110 (8%) divorced, widowed, or separated women and 1,074 (8%) women aged ≤ 14 and ≥ 50 years were dropped from the analysis. As the purpose of our analysis is to compare slum and non-slum violence, we also excluded 1,356 (10%) women from district municipalities because the sample could not be partitioned into the residential categories of interest (slum vs. non-slum). The final analytical sample was 9,122 currently married women (5,128 from slums and 3,994 from non-slums), thus retaining 64% of the full survey sample.

Measures

Dependent Variable

Our outcome variable is occurrence of physical violence against a woman by her husband in the past year. Five "yes-no" violence tactic items assessed physical spousal violence: (1) push, shake, or throw something that could hurt you, (2) slap or twist your arms, (3) punch with a fist or with something that could hurt you, (4) kick or drag you, and (5) strangulate, burn, or attempt to kill you. These items were combined into a dichotomous variable: no

physical violence (score = 0) and any physical violence (score ≥ 1). We restrict our analysis to cases of physical violence that occurred in the past year (rather than “ever”) so as to establish a reasonable temporal ordering between physical spousal violence and the selected independent variables

Independent Variables

Independent variables for this analysis fall into four domains: sociodemographic characteristics, gender role beliefs, attitudes toward partner violence, and relational factors. Sociodemographic characteristics included place of residence, age, education, income-earning activity, participation in microcredit programs, religion, household wealth, age at first marriage, and number of children. Household wealth was assessed using a combination of measures of household asset and amenities ownership, and housing tenure, an approach commonly used to assess wealth in this setting (Rutstein & Johnson, 2004).

Respondent’s view on gender role beliefs was derived from two questions that assessed whether it was acceptable for married women to work outside the home. First, women were asked, “If for some reason the husband cannot make enough money for the family, do you believe that it is acceptable for married women to work outside the home to earn an income,” and the response was yes (coded = 1) or no (coded = 0). Second, women were also asked, “If the husband is making money, do you believe that it is acceptable for married women to work outside the home to earn an income.” The response was yes (coded = 1) or no (coded = 0).

Our study adopted a composite measure on acceptance of partner violence under various scenarios. A binary variable, the wife-beating norm, was developed by inquiring of the woman whether or not it is justified for a husband to beat his wife if she “neglects the children,” “argues with her husband,” “fails to provide food on time,” “visits her family without her husband’s permission,” and “visits a friend without her husband’s permission.” The response was no = 0 and yes = 1. Respondents answering “no” to all of the attitudes items were coded 0, and those who responded in the affirmation on one or several of the attitudes items were coded 1.

Finally, the study included the following relational factors: Husband considers wife’s opinion in decision making and household decision-making power. The variable, husband considers wife’s opinion in decision making, assessed whether women were consulted by their husbands when making decisions on “large household purchases,” “minor daily household expenses,” and “when he wants to visit family or friends.” The “yes” (code = 1) or “no”

(code = 0) responses were summed (Cronbach's $\alpha = .85$) and then dichotomized: no consideration (score = 0) and some consideration (score ≥ 1). A categorical variable of household decision-making power was constructed based on inquiring of the women who in their opinion bore the responsibility of making final decisions on "the women's own health care," "children's health"; "large household purchases"; "everyday household purchases"; "visits to family, friends, or relatives"; and "what food to be cooked each day." The responses to these items were coded into one of three relationship power types: egalitarian, husband dominated, and wife dominated.

Statistical Analysis

Statistical analyses were performed using STATA Release 10 (Stata Corporation, 2007). Basic descriptive statistics (means) were computed to describe the characteristics of the study sample as well as the prevalence of past-year physical violence within marriage and for the specific acts of abuse, together with 95% confidence limits. Next, we used chi-square analyses to assess bivariate association between past-year physical spousal violence and the selected background characteristics of women. Given that the dependent variable is dichotomous, multivariate logistic regression was used to examine factors associated with past-year physical spousal violence. We also tested whether selected explanatory variables had moderating effects on slum residence by creating interaction terms between these variables. None of the interaction terms achieved significance; hence, we excluded them in the analysis. STATA's survey commands were used to account for the cluster sampling design. In addition, in order to present results that are representative to the survey population in slums and non-slums, we apply weights to our statistical analyses. Following the logit model, we defined P_i , the probability that the i th respondent was abused by her husband in the past year:

$$P_i = 1/(1+e^{-Z_i})$$

where $Z_i = b_0 + b_1 X_{1i} + b_2 X_{2i} + b_3 X_{3i} + \dots + b_{24} X_{24i}$ and where the subscript i refer to the i th respondent. The X s represents the sociodemographics, gender role beliefs, attitudes toward partner violence, and relational characteristics of the respondent that determine the P_i . To enhance interpretability, coefficients from the logit regression were transformed to marginal effects using 5,000 bootstrap samples (Efron & Tibshirani, 1993). The marginal effects of the X s, evaluated at their sample means, were shown to be represented by $\partial P_i / \partial X_i = b_j / (1 - P_i) P_i$. The marginal effects approximate an increased or

decreased probability that a woman was abused by her husband in the past year, given a one-unit increase in the covariate. For dummy variables (i.e., variables recoded as 0 or 1) the marginal effects represent a predicted increase or decrease for a discrete change from 0 to 1.

Results

Sample Characteristics

Table 1 presents summary statistics of selected background characteristics of married women by stratum. More than half (56%) of slum women were aged 29 years or less, in contrast to 48% of nonslum women. Higher rates of non-literacy were found among slum women (47%), whereas higher educational attainment was more common among nonslum women (30%). Approximately 29% of slum women were employed for wages, in contrast to 17% from the nonslums. As expected, a higher proportion of slum women (32%) reported participating in credit programs, in contrast to 27% from the nonslums. The sample was predominantly Muslim. Sixty-five percent of slum women belonged to poor households, in contrast to one-in-five of nonslum women (23%), indicating that slum women live in greater poverty. Although the legal minimum age at marriage for women in Bangladesh is 18 years, a majority of women got married before they reached 18 years: 79% of slum women got married before age 18, in contrast to 60% of nonslum women.

About one third of slum women (32%) reported that it was tolerable for wives to work outside the home if the husband is making enough money, in contrast to 35% of nonslum women. About half of slum women (50%) believed that husbands should beat their wives under various circumstances, in contrast to 36% of nonslum women. A majority of women in both urban stratum reported that their husbands considered their opinions when making decisions.

Past-year prevalence of physical spousal violence. The results in Table 2 reveal that in the past year, about a third of women surveyed (31%) had been physically assaulted by their husbands. We also found significant variations in prevalence of past-year physical spousal violence by stratum, both in terms of overall prevalence and specific acts of violence ($p < .001$). Past-year rates of physical spousal violence were higher among slum women (34.8%) than among nonslum women (19.8%). Overall, some 28.3% of the women reported experiencing slapping or arm-twisting, followed by 18.2% experiencing pushing, shaking, or throwing something, and the least commonly cited acts of physical violence being strangulation, burning, or attempting to kill (3.5%).

Table 1. Sample Profile of Currently Married Bangladeshi Women Aged Between 15 and 49 Years by Urban Stratum, UHS, 2006

Characteristic	Slum		Nonslum	
	M	SE	M	SE
Sociodemographic characteristics				
Age (years)				
15-19	0.132	0.011	0.068	0.005
20-29	0.424	0.015	0.407	0.012
30-39	0.300	0.008	0.350	0.011
40-49	0.144	0.008	0.175	0.010
Education				
No education	0.468	0.013	0.197	0.011
Some primary	0.298	0.008	0.230	0.012
Some secondary	0.234	0.011	0.573	0.020
Income-earning activity				
Not employed	0.667	0.016	0.803	0.014
Self-employed	0.047	0.005	0.024	0.004
Employed for wages	0.286	0.017	0.173	0.014
Participation in microcredit programs				
Nonparticipation	0.682	0.015	0.725	0.021
Participation	0.318	0.015	0.275	0.021
Religion				
Non-Muslim	0.044	0.008	0.127	0.030
Muslim	0.956	0.008	0.873	0.030
Household wealth				
Poor	0.646	0.019	0.231	0.027
Middle	0.201	0.011	0.173	0.013
Rich	0.152	0.016	0.516	0.033
Age at first marriage				
≤ 17 years	0.786	0.008	0.589	0.018
≥ 18 years	0.214	0.008	0.410	0.018
Number of children				
None	0.118	0.006	0.123	0.010
1-2	0.417	0.010	0.511	0.017
3-4	0.296	0.010	0.284	0.016
≥ 5	0.169	0.007	0.082	0.009

(continued)

Table 1. (continued)

Characteristic	Slum		Nonslum	
	M	SE	M	SE
Gender role beliefs				
Acceptable for wife to work, if husband is not making enough money				
No	0.113	0.007	0.142	0.013
Yes	0.887	0.007	0.858	0.013
Acceptable for wife to work, if husband is making enough money				
No	0.678	0.012	0.650	0.019
Yes	0.322	0.012	0.350	0.019
Attitudes toward partner violence				
Wife-beating norms				
No wife-beating norm (score = 0)	0.505	0.019	0.639	0.024
Some wife-beating norms (score ≥ 1)	0.495	0.019	0.361	0.024
Relational factors				
Husband considers wife's opinion in decision making				
No opinion considered	0.119	0.007	0.099	0.013
Some opinion considered	0.881	0.007	0.901	0.013
Household decision-making power				
Egalitarian	0.110	0.008	0.103	0.013
Husband dominated	0.071	0.007	0.060	0.009
Wife dominated	0.829	0.009	0.837	0.012
Number of women	5,128		3,994	

Table 2 also shows the marked disparities of magnitude of specific acts of physical violence experienced by women living in slum and nonslum areas. Overall, the pattern of past-year physical violence was similar across the residential categories. However, slum women reported higher exposure of past-year physical spousal violence across all five acts of abuse, in contrast to nonslum women (also see Figure 1). The most commonly reported act of abuse was slapping or arm-twisting (10% in nonslums vs. 21% in slums). Pushing, shaking, or throwing something at the women was the second most common act of abuse. The least commonly cited act of abuse was

Table 2. Prevalence of Past-Year Physical Spousal Violence by Urban Stratum, UHS 2006

Prevalence of Physical Spousal Violence	Slum		Nonslum		Total	
	%	95% CI	%	95% CI	%	95% CI
Any past-year physical violence ^a	34.8	(32.56-37.11)	19.76	(16.94-22.91)	30.96	(29.11-32.87)
Specific acts of physical abuse						
Pushing, shaking or throwing something at her	20.73	(18.75-22.87)	10.67	(9.31-12.21)	18.16	(16.62-19.81)
Slapping or twisting arms	31.87	(29.18-34.96)	17.72	(15.14-20.64)	28.26	(26.17-30.45)
Punching,	15.50	(14.07-17.06)	8.70	(6.88-10.94)	13.77	(12.62-15.00)
Kicking or dragging	11.78	(10.30-13.42)	6.39	(5.08-8.00)	10.40	(9.30-11.62)
Strangulation, burning, or attempting to kill	4.02	(3.41-4.75)	2.03	(1.37-3.00)	3.52	(3.01-4.10)

Note: Currently married women, aged between 15 and 49 years, urban slum (n = 5,128), urban nonslum (n = 3,994), and total (n = 9,122). CI = confidence interval

a. Any one or more specific acts of physical abuse are classified as physical spousal violence.




Figure 1. Prevalence of past-year physical spousal violence by urban strata.

strangulation, burning, or attempting to kill, with a rate of prevalence of 2% in non-slums versus 4% in slums.

Bivariate Analysis of Factors Associated With Past-Year Physical Spousal Violence

Chi-square analyses of bivariate associations between past-year physical spousal violence and selected characteristics of women are displayed in Table 3 by stratum. Overall, this table suggests that similar factors operate in both slum and non-slum settings. For both urban strata, physical violence was less common with increasing age of women and post-primary educational attainment ($p < .001$ for all). For non-slum women, physical violence was higher among those who participated in microcredit programs ($p < .05$). For slum women, the proportion experiencing abuse was nonsignificantly higher for those participating in microcredit programs than among those not participating in microcredit programs. Women who worked for wages reported the highest rate of abuse ($p < .005$), for non-slum area only. Physical violence was higher among Muslim women residing in slum areas ($p < .001$), but not among non-slum Muslim women. Physical violence was less common with increasing household wealth ($p < .001$ for both urban strata). Similarly for both settings, physical violence was lower among women reporting age at first marriage as ≥ 18 years ($p < .05$). For slum women, physical abuse somewhat declined with increasing parity ($p < .001$), and those who indicated it is acceptable for married women to work if the husband is not making enough money reported higher physical abuse ($p < .01$). For non-slum women, physical abuse was lowest among those reporting that it is acceptable for a wife to work if the husband is making enough money ($p < .05$). For both urban strata, physical abuse was higher among women reporting that it is acceptable for husbands to beat their wives under various scenarios. Also, χ^2 test for husband's consideration of wife's opinion in decision making were significant ($p < .05$ for all). For slum women only, physical violence was significantly associated with household decision-making power ($p < .05$).

Multivariate Analysis of Factors Associated With Past-Year Physical Spousal Violence

Table 4 presents the regression coefficients, marginal effects, and robust standard errors. Marginal effects are calculated at the mean values for all other regressors from the logistic regression model predicting women's risk of experiencing physical spousal violence in the past year. In analyses stratified by place of residence (not shown) there were no consistent differences in the

Table 3. Past-Year Physical Spousal Violence Among Married Bangladeshi Women by Selected Background Characteristics According to Urban Stratum, UHS 2006

Characteristic	% physical abuse and 95% confidence interval (CI)			
	Slum (%)		Nonslum (%)	
Sociodemographic characteristics				
Age (years)		***		***
15-19	38.55	(31.80-45.78)	23.85	(16.78-32.73)
20-29	40.76	(37.75-43.84)	25.65	(21.09-30.80)
30-39	31.52	(28.75-34.44)	16.99	(13.50-21.17)
40-49	20.59	(17.14-24.53)	10.00	(6.70-14.68)
Education		***		***
No education	38.00	(35.03-41.05)	32.42	(25.27-40.49)
Some primary	38.71	(35.46-42.06)	26.03	(20.33-32.66)
Some secondary	23.45	(20.12-27.16)	12.90	(11.17-14.85)
Participation in microcredit programs				*
Nonparticipation	34.21	(31.70-36.81)	18.70	(15.97-20.59)
Participation	36.05	(31.77-40.56)	23.94	(17.95-31.15)
Employment status				**
Not employed	33.64	(30.97-36.42)	18.12	(15.17-21.51)
Self-employed	35.34	(28.78-42.49)	19.19	(11.48-32.47)
Employed for wages	37.40	(33.86-41.07)	27.32	(21.74-33.70)
Religion		**		
Non-Muslim	20.53	(13.72-29.55)	18.51	(10.05-31.58)
Muslim	35.45	(33.18-37.78)	19.94	(17.65-22.45)
Household wealth		***		***
Poor	41.16	(38.82-43.55)	36.28	(31.26-41.62)
Middle	26.52	(22.56-30.88)	26.54	(20.57-33.50)
Rich	18.75	(15.49-22.52)	11.40	(9.55-13.55)
Age at first marriage		*		**
≤17 years	35.90	(33.57-38.29)	22.58	(18.58-27.15)
≥18 years	30.77	(26.70-35.16)	15.71	(12.85-19.06)
Number of children		***		
None	31.48	(26.69-36.70)	19.11	(13.92-25.66)
1-2	37.99	(34.42-41.69)	19.93	(16.06-24.45)
3-4	34.19	(30.35-38.25)	18.18	(14.97-21.90)
≥ 5	30.28	(26.64-34.19)	25.14	(15.24-38.55)

(continued)

Table 3. (continued)

Characteristic	% physical abuse and 95% confidence interval (CI)			
	Slum (%)		Nonslum (%)	
Gender role beliefs				
Acceptable for wife to work, if husband is not making enough money		**		
No	29.19	(24.99-33.78)	14.88	(10.43-20.79)
Yes	35.51	(33.19-37.90)	20.56	(17.38-24.16)
Acceptable for wife to work, if husband is making enough money				*
No	34.12	(31.70-36.62)	21.64	(17.86-25.96)
Yes	36.22	(33.17-39.40)	16.26	(13.43-19.56)
Attitudes toward partner violence				
Wife-beating norms		***		***
No wife-beating norms	29.65	(26.33-33.20)	15.75	(13.07-18.85)
Some wife-beating norms	40.04	(37.41-42.73)	26.85	(22.42-31.79)
Relational factors				
Husband considers wife opinion in decision making		***		***
No opinion considered	47.12	(41.69-52.63)	40.57	(32.06-49.70)
Some opinion considered	33.13	(30.93-35.41)	17.47	(14.95-20.31)
Household decision-making power				
Egalitarian	39.80	(34.83-44.98)	19.34	(12.51-28.68)
Husband dominated	29.75	(23.35-37.05)	26.02	(19.05-34.46)
Wife dominated	34.80	(32.55-37.11)	19.36	(16.47-22.61)
Number of women	5,128		3,994	

Note: The test statistics are designed based on F-ratio produced from Pearson's χ^2 statistics.
 * $p < .05$; ** $p < .01$; *** $p < .001$.

final models predicting past-year physical spousal violence; thus, we analyzed the two residences together. In our analysis, marginal effects have been multiplied by 100 for ease of interpretation.

As regards sociodemographic characteristics, slum residence was not significantly associated with physical spousal violence, whereas age was

Table 4. Regression Coefficients, Marginal Effects, and Standard Errors for Past-Year Physical Spousal Violence by Selected Background Characteristics, UHS 2006

Characteristic	β	Marginal effects	SE
Sociodemographic characteristics			
Place of residence (ref: nonslum)			
Slum	0.131	0.025	0.021
Age (years) (ref: 15-19)			
20-29	-0.094	-0.020	0.032
30-39	-0.675	-0.133***	0.038
40-49	-1.250	-0.222***	0.039
Education (ref: no education)			
Some primary	-0.073	-0.015	0.018
Some secondary	-0.582	0.108***	0.021
Income-earning activity (ref: not employed)			
Self-employed	0.041	0.008	0.030
Employed for wages	0.001	0.001	0.017
Participation in microcredit programs (ref: nonparticipation)			
Participation	0.131	0.025	0.018
Religion (ref: non-Muslim)			
Muslim	0.293	0.053	0.036
Household wealth (ref: poor)			
Middle	-0.466	-0.093***	0.016
Rich	-0.840	-0.156***	0.020
Age at first marriage (ref: ≤ 17 years)			
≥ 18 years	-0.011	-0.002	0.019
Number of children (ref: zero)			
1-2	0.405	0.072**	0.023
3-4	0.464	0.083**	0.027
≥ 5	0.579	0.105***	0.033
Gender role beliefs			
Acceptable for wife to work, if husband not making money (ref: no)			
Yes	0.287	0.052**	0.019
Acceptable for wife to work, if husband making money (ref: no)			
Yes	0.070	0.013	0.013
Attitudes toward partner violence			
Wife-beating norms (ref: no wife-beating norm)			
Some wife-beating norms	0.451	0.086***	0.019

(continued)

Table 4. (continued)

Characteristic	β	Marginal effects	SE
Relational factors			
Husband considers wife opinion in decision making (ref: no opinion considered)			
Some opinion considered	-0.778	-0.159***	0.021
Household decision-making power (ref: egalitarian)			
Husband dominated	-0.436	-0.082**	0.033
Wife dominated	-0.195	-0.038*	0.022
Number of women		9,122	
Number of clusters		445	
F statistics		$F(22, 424) = 36.75$	

Note: ref. = reference category. Standard errors (SE) were adjusted for clustering at the strata level (i.e., the primary sampling unit in the UHS sample design)

* $p < .05$; ** $p < .01$; *** $p < .001$.

negatively related to spousal violence. Specifically, women aged 30 years or more, in contrast to those aged 19 years or less, had a lower probability of being physically abused by their husbands (13.3%-22.2%). Not surprisingly, having some post-primary education was negatively associated with physical spousal violence after controlling for other factors; the likelihood of being physically abused was 10.8% lower for women with some secondary education, as opposed to those with no education.

We also found evidence that household wealth was a negative predictor of physical spousal violence. More specifically, belonging to any of the wealth categories, as opposed to the poor category, significantly decreased the probability of being physically abused; the likelihood of being abused decreased by 9.3% for women in households with middle wealth and 15.6% for women in rich households. Number of children was a significant positive predictor of physical spousal violence; the likelihood of being physically abused increased by 7.2% for women having 1 to 2 children to about 10.5% for women having 3 or more children. The coefficients of income earning activity, participation in microcredit programs, religion, and age at first marriage were not statistically significant as predictors of past-year physical spousal violence.

With respect to gender role beliefs, only one factor within this domain emerged as a significant predictor of physical spousal violence. The belief that

a married woman should work outside the home if the husband is not making enough money significantly increased the probability of being physically abused by 5.2%, whereas the belief that a married woman should work outside the home if the husband is making enough money did not predict physical abuse.

Concerning attitudes toward partner violence, our results indicate that wife-beating norms were positively related to physical spousal violence, all else being equal; having some wife-beating norms significantly increased the probability of being physically abused by a husband by 8.6%.

The results of the multivariate analyses also provided support for the link between selected relational factors and physical spousal violence. Specifically, husband's consideration of wife's opinion in decision making was a negative predictor of physical spousal violence; the likelihood of being abused increased by 15.9%. Interestingly, the probability of being physically abused decreased by 8.2% for women whose husbands' dominated household decision making, whereas wife dominance in household decision making had a marginal effect on physical violence.

Discussion

To our knowledge, this is the first large population-based study to provide valuable data on the prevalence and associated factors of past-year physical spousal violence in slum and nonslum areas of urban Bangladesh. Furthermore, this study allows us to examine simultaneously the multiple risk factors that have been linked to IPV against women in Bangladesh and elsewhere. Our study reported an overall past-year rate of physical spousal violence of in this urban sample (31%). Previous studies that have explored the prevalence of physical spousal violence in Bangladesh found rates ranging from 19% to as high as 35% to 38% (Bates et al., 2004). Our prevalence rate is within the upper range of estimates documented in these previous reports. The past-year rate of physical spousal violence (31%) in our sample was higher than the past-year rate of physical spousal violence found in India (22%; Chaudhary, Girdhar, & Soni, 2009) but is consistent with the findings in Turkey (34%; Tokuç, Ekuklu, & Avcioglu, 2010). We also found that past-year physical spousal abuse varied by stratum: Abuse was higher among the poorest strata, namely, the slums (34.8%), with lower rates in nonslums (19.8%). The pattern of abuse was the same in both slum and nonslum areas, with slapping or arm-twisting more common than strangulation, burning or attempting to kill that may be related to impulsiveness in families or due to cultural attitudes toward killing (Haj-Yahia, 2003; Jewkes, Levin, &

Penn-Kekana, 2002). Our results show that physical spousal abuse is common event in slums and non-slums of urban Bangladesh. Given the potentially immediate and long-term deleterious consequences of physical abuse, it is realistic to state that domestic violence is a serious social and public health issue for women in urban Bangladesh.

Cross- and intra-national comparison of domestic violence is complicated due to differences in sampling methods, measures of violence, and willingness to disclose abuse in interview settings (Krug et al., 2002). Nevertheless, the prevalence of past-year physical spousal abuse found in our study is greater than those reported in six cities included in the Garcia-Moreno et al. (2005) multicountry study on domestic violence against women: 3.2% in Serbia and Montenegro, 7.9% in Thailand, 8.3% in Brazil, 14.8% in Tanzania, and 16.9% in Peru. The prevalence reported in our study, particularly for slums, is somewhat similar to rates found in other surveys in Tirana, Albania, and Cuernavaca Morelos, Mexico, giving rates of 37% and 35.8%, respectively (Burazeri et al., 2005; Rivera-Rivera et al., 2004). Our prevalence for non-slums (20%) is similar to that reported in previous Bangladesh studies (Naved et al., 2006).

The results of the current study confirmed our hypotheses that many sociodemographic characteristics affect the likelihood of women's risk of being physical abused even when other factors were statistically controlled. Whereas slum residence showed no significant effect on past-year physical spousal violence, our study identified higher age as an important protective factor against physical spousal violence. These results are consistent with studies in Bangladesh (Ahmed, 2005) and Iran (Ghazizadeh, 2003). One possible explanation of our finding is that women tend to accrue social status with rising age. As a consequence they are subjected to less abuse within marriage. In addition, having some post-primary education was also found to be a protective factor against spousal violence. Results were consistent with previous associational studies of the role of education on IPV (Ahmed, 2005; Vung, Ostergren, & Krantz, 2008). Such results highlight the critical role women's education play in improving their status within their households or improve spousal communication skills, thus reducing their vulnerability to abuse. Yet the protective effect of education can be slowed by the cultural context of rigid gender norms deeply entrenched in the society (Baden, Green, Goetz, & Guhathakurta, 1994). Nonetheless, investing in girl-child education and female literacy programs still remains a critical pathway to reducing abuse and improving women's status (Jewkes et al., 2002).

Our findings suggest that household wealth has a protective effect against physical spousal violence. These results corroborate studies from a wide range of setting such as Ukraine, India, and Cambodia (Dude, 2007; Kishor & Johnson, 2004), including Bangladesh (Bates et al., 2004; Koenig et al., 2003), stating the association between household wealth and IPV. Thus, policies and interventions aimed at reducing poverty may contribute to decreasing the risk of physical violence among women in urban Bangladesh, particularly for those living in slum areas. All measures of income-generating activities were not associated with spousal abuse; echoing earlier studies from Bangladesh (e.g., Ahmed, 2005), we found no association between physical spousal violence and income earning activity and participation in micro-credit programs.

Consistent with our hypotheses, the findings shows that there is an association between physical spousal violence and conservative ideas regarding gender role beliefs (Sugarman, 1996). Notably, the belief that married women should work outside the home if the husband is not making enough money was associated with elevated risk of abuse. Thus, the findings seem to support the idea often repeated in the literature that men in patriarchal societies subscribe to a notion of masculinity involving distinctly hierarchical gender positions and definitions of male success in terms of controlling women (Dobash & Dobash, 1979). In this context, it is not surprising to find that perceived challenges to male dominance are associated with wife abuse.

Similar to other studies documenting association between normative support of wife beating and IPV (Choi & Ting, 2008; Faramarzi, Esmailzadeh, & Mosavi, 2005), our study found that wife-beating norms are a strong risk factor for physical spousal violence. One explanation is that entrenched patriarchal attitudes defining IPV as a normal practice transcend economic and social boundaries in urban Bangladesh. Given the pervasiveness of such attitudes, there is a need to transform gender relationships in the whole society. This might be achieved through developing sociocultural interventions aimed at preventing IPV (Heise, 1998). Such approaches should aim to change the societal norms about IPV and provide the future generations with knowledge and skills needed for healthy relationships so as to shift from a society where IPV is widely condoned.

In contrast to results reported in the Philippines (Hindin & Adair, 2002), our study shows that husband-dominated household decision making was negatively related to physical spousal abuse. Similarly, our results suggest that when a man considers his wife's opinions in his decision making, a woman's exposure to IPV is reduced. Thus, men may be less abusive if they have control over their or partner's decision making. In Bangladeshi culture, men are

decision makers and women having elevated decision-making power may lead men to face some kind of masculinity identity crisis that results in them adopting aggressive behaviors in order to claim their domination (Goode, 1971).

This study has several limitations. First, this study examined associations between physical spousal violence and selected variables; thus, causality cannot be inferred. Nevertheless, the associations provide insights about the role of the selected variables in women's vulnerability to violence. Second, the study relied on self-reported experiences of IPV obtained through face-to-face interview, which are subject to response biases due to sensitivities and stigma associated with IPV and social desirability (Arias & Beach, 1987). However, the instrument used for data collection assessed behaviorally specific acts of physical abuse based on the revised Conflict Tactics Scale (CTS2; Straus et al., 1996). The CTS2 has been used in several studies in developing (e.g., India, Kenya, Malawi) and developed (e.g., North America) nations revealing consistent reliability and validity of the scale and its subscales (Hindin, Kishor, & Ansara, 2008; Straus, 2004). Third, the study excluded data that might allow greater precision in assessing the risk of being abused in adulthood and considered as potential important information in the context of designing interventions to reduce violence. For example, our survey excluded data on witnessing abuse between parents, experience of childhood abuse, and husband's risk factors such as age, education, risky sexual behaviors, drug, and alcohol use. This also limits the contextual understanding of abuse. Finally, by limiting the sample to married women the study precluded a complete profile of abused women in urban Bangladesh.

With these caveats, this study enhances our understanding of physical spousal violence, its magnitude, pattern, and associated factors in urban Bangladesh. The study points to the importance of studying IPV against women using a multifactor framework such as the ecological model (Heise, 1998). Similar to research worldwide, our results reveal that the occurrence of IPV against Bangladeshi women is related to an interplay of risk factors at the individual, relational, and societal level (DeMaris, Benson, Fox, Hill, & Van Wyk, 2003). Hence, a combination of strategies and several risk reduction options with strong individual and community involvement over an extended period are necessary for the reduction of IPV against Bangladeshi women.

The study's sample was unique in that it included important urban strata, namely, slums and non-slums. The sample represented a significantly larger population of individuals in urban Bangladesh than did previous studies that were based on small samples with limited geographic coverage (Naved et al., 2006). Our study serves to confirm the commonness of spousal abuse in Bangladesh—not limited to one stratum—therefore, interventions

oriented toward the whole society rather than risk groups are warranted. Furthermore, our study points out that spousal physical abuse is a complex phenomenon related with individual characteristics and deeply rooted in power relationships related to gender norms and normative use of abuse in conflict situations. The high rates of spousal physical abuse underscores the need for policy makers to increase their recognition of spousal abuse as a critical target in public health concerns and that comprehensive and culturally acceptable approaches to tackling domestic violence against women are needed.

Declaration of Conflicting Interests

The authors declared no potential conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

References

- Ahmed, S. M. (2005). Intimate partner violence against women: Experiences from a woman-focused development programme in Matlab, Bangladesh. *Journal of Health, Population, and Nutrition*, 23(1), 95-101.
- Ahmed, M. K., van Ginneken, J., Razzaque, A., & Alam, N. (2004). Violent deaths among women of reproductive age in rural Bangladesh. *Social Science & Medicine*, 59(2), 311-319.
- Aklimunnessa, K., Khan, M. M. H., Kabir, M., & Mori, M. (2007). Prevalence and correlates of domestic violence by husbands against wives in Bangladesh: Evidence from a national survey. *Journal of Men's Health and Gender*, 4(1), 52-63.
- Amin, M., Khuda, B., Islam, A., & Levin, A. (1997). *Do female education and mobility lead to less intra-family abuse?* Dhaka: Bangladesh.
- Arias, I., & Beach, S. R. H. (1987). Validity of self-reports of marital violence. *Journal of Family Violence*, 2(2), 139-149.
- Baden, S., Green, C., Goetz, A. M., & Guhathakurta, M. (1994). *Background report on gender issues in Bangladesh*, No. 26. Brighton, UK: Institute of Development Studies.
- Bates, L. M., Schuler, S. R., Islam, F., & Islam, K. (2004). Socioeconomic factors and processes associated with domestic violence in rural Bangladesh. *International Family Planning Perspectives*, 30(4), 190-199.
- Bhuiya, A., Sharmin, T., & Hanifi, S. M. (2003). Nature of domestic violence against women in a rural area of Bangladesh: Implication for preventive interventions. *Journal of Health, Population, and Nutrition*, 21(1), 48-54.

- Burazeri, G., Roshi, E., Jewkes, R., Jordan, S., Bjegovic, V., & Laaser, U. (2005). Factors associated with spousal physical violence in Albania: Cross-sectional study. *British Medical Journal*, *331*(7510), 197-201.
- Campbell, T., & Campbell, A. (2007). Emerging disease burdens and the poor in cities of the developing world. *Journal of Urban Health*, *84*(Suppl. 3), 54-64.
- Campbell, J., Jones, A. S., Dienemann, J., Kub, J., Schollenberger, J., O'Campo, P., et al. (2002). Intimate partner violence and physical health consequences. *Archives of Internal Medicine*, *162*, 1157-1163.
- Chaudhary, A., Girdhar, S., & Soni, R. K. (2009). Epidemiological correlates of domestic violence in married women in urban area of Ludhiana, Punjab, India. *The Internet Journal of Health*, *9*(1). Retrieved from http://www.ispub.com/journal/the_internet_journal_of_health/volume_9_number_1_12/article/epidemiological-correlates-of-domestic-violence-in-married-women-in-urban-area-of-ludhiana-punjab-india.html
- Choi, S. Y., & Ting, K. F. (2008). Wife beating in South Africa: An imbalance theory of resources and power. *Journal of Interpersonal Violence*, *23*, 834-852.
- Chowdhury, A., & Bhuiya, A. (2001). Do poverty alleviation programmes reduce inequities in health? The Bangladesh experience. In D. Leon & G. Walt (Eds.), *Poverty, inequality and health: An international perspective* (pp. 312-332). Oxford, UK: Oxford University Press.
- Counts, D. A., Brown, J. K., & Campbell, J. (1992). *Sanctions and sanctuary: Cultural perspectives on the beating of wives*. Boulder, CO: Westview.
- Decker, M. R., Miller, E., Kapur, N. A., Gupta, J., Raj, A., & Silverman, J. G. (2008). Intimate partner violence and sexually transmitted disease symptoms in a national sample of married Bangladeshi women. *International Journal of Gynecology and Obstetrics*, *100*(1), 18-23.
- DeMaris, A., Benson, M. L., Fox, G. L., Hill, T., & Van Wyk, J. (2003). Distal and proximal factors in domestic violence: A test of an integrated model. *Journal of Marriage and Family*, *65*, 652-667.
- Dobash, R. E., & Dobash, R. (1979). *Violence against wives: A case against the patriarchy*. New York: Free Press.
- Dude, A. (2007). Intimate partner violence and increased lifetime risk of sexually transmitted infection among women in Ukraine. *Studies in Family Planning*, *38*(2), 89-100.
- Dunkle, K. L., Jewkes, R. K., Brown, H. C., Gray, G. E., McIntyre, J. A., & Harlow, S. D. (2004). Gender-based violence, relationship power, and risk of HIV infection in women attending antenatal clinics in South Africa. *Lancet*, *363*, 1415-1421.
- Efron, B., & Tibshirani, R. (1993). *An introduction to the bootstrap*. London: Chapman and Hall.
- Ellsberg, M., Jansen, H. A., Heise, L., Watts, C. H., Garcia-Moreno, C., & WHO Multi-country Study on Women's Health and Domestic Violence Against Women Study

- Team. (2008). Intimate partner violence and women's physical and mental health in the WHO multi-country study on women's health and domestic violence: An observational study. *Lancet*, *371*, 1165-1172.
- Faramarzi, M., Esmailzadeh, S., & Mosavi, S. (2005). A comparison of abused and non-abused women's definitions of domestic violence and attitudes to acceptance of male dominance. *European Journal of Obstetrics, Gynecology, and Reproductive Biology*, *122*, 225-231.
- Freudenberg, N., Galea, S., & Vlahov, D. (2005). *Cities and the health of the public*. Nashville, TN: Vanderbilt University Press.
- Garcia-Moreno, C., Jansen, H. A., Ellsberg, M., Heise, L., & Watts, C. (2005). WHO multi-country study on women's health and domestic violence against women: Initial results on prevalence, health outcomes and women's responses. Geneva, Switzerland: World Health Organization.
- Ghazizadeh, A. (2003). Wife battering: A prospective study in Sanandaj city Iran. *Iranian Journal of Public Health*, *32*(2), 24-27.
- Goode, W. J. (1971). Force and violence in the family. *Journal of Marriage and the Family*, *33*, 624-636.
- Hadi, A. (2000). Prevalence and correlates of the risk of marital sexual violence in Bangladesh. *Journal of Interpersonal Violence*, *15*, 787-805.
- Haj-Yahia, M. M. (2003). Beliefs about wife beating among Arab men from Israel: The influence of their patriarchal ideology. *Journal of Family Violence*, *18*, 193-206.
- Heise, L. L. (1998). Violence against women: An integrated, ecological framework. *Violence Against Women*, *4*, 262-290.
- Hindin, M. J., & Adair, L. S. (2002). Who's at risk? Factors associated with intimate partner violence in the Philippines. *Social Science & Medicine*, *55*, 1385-1399.
- Hindin, M. J., Kishor, S., & Ansara, D. L. (2008). *Intimate partner violence among couples in 10 DHS countries: Predictors and Health Outcomes* (DHS Analytical Studies No. 18). Calverton, MD: Macro International Inc.
- Islam, F., Mailman, N., Acharya, K., Higgins, J., Islam, M. K., & Bates, L. M. (2004). *Violence against women within marriage in Bangladesh*. Washington, DC: Academy for Educational Development.
- Jeyaseelan, L., Kumar, S., Neelakantan, N., Peedicayil, A., Pillai, R., & Duvvury, N. (2007). Physical spousal violence against women in India: Some risk factors. *Journal of Biosocial Science*, *39*, 657-670.
- Jewkes, R., Levin, J., & Penn-Kekana, L. (2002). Risk factors for domestic violence: Findings from a South African cross-sectional study. *Social Science & Medicine*, *55*, 1603-1617.
- Johnston, H. B., & Naved, R. T. (2008). Spousal violence in Bangladesh: A call for a public-health response. *Journal of Health, Population, and Nutrition*, *26*, 366-377.
- Kishor, S., & Johnson, K. (2004). *Profiling domestic violence: A multicountry study*. Calverton, MD: ORC Macro, MEASURE DHS+.

- Koenig, M. A., Ahmed, S., Hossain, M. B., & Mozumder, A. B. M. K. A. (2003). Women's status and domestic violence in rural Bangladesh: Individual- and community-level effects. *Demography, 40*, 269-288.
- Krug, E. G., Mercy, J. A., Dahlberg, L. L., & Zwi, A. B. (2002). World report on violence and health. Geneva, Switzerland: World Health Organization.
- Levy, P. S., & Lemeshow, S. (1965). *Sampling*. New York: John Wiley.
- MacMillan, R., & Gartner, R. (1999). When she brings home the bacon: Labor-force participation and the risk of spousal violence against women. *Journal of Marriage and the Family, 61*, 947-958.
- National Institute of Population Research and Training, Mitra Associates, and ORC Macro. (2005). *Bangladesh demographic and health survey 2004*. Dhaka, Bangladesh: NIPORT, Mitra Associates; Calverton, MD: ORC Macro.
- Naved, R. T. (2003). *A situation analysis of violence against women in South Asia*. Kathmandu, Nepal: UNFPA.
- Naved, R. T., & Akhtar, N. (2008). Spousal violence against women and suicidal ideation in Bangladesh. *Women's Health Issues, 18*, 442-452.
- Naved, R. T., Azim, S., Bhuiya, A., & Persson, L. A. (2006). Physical violence by husbands: Magnitude, disclosure and help-seeking behavior of women in Bangladesh. *Social Science & Medicine, 62*, 2917-2929.
- Naved, R. T., & Persson, L. A. (2005). Factors associated with spousal physical violence against women in Bangladesh. *Studies in Family Planning, 36*, 289-300.
- Panda, P., & Agarwal, B. (2005). Marital violence, human development and women's property status in India. *World Development, 33*, 823-850.
- Plichta, S. B., & Falik, M. (2001). Prevalence of violence and its implications for women's health. *Women's Health Issues, 11*, 244-258.
- Rivera-Rivera, L., Lozano-Ponce, E., Salerom-Castro, J., Salazar-Martinez, E., Castro, R., & Hernandez-Avila, M. (2004). Prevalence and determinants of male partner violence against Mexican women: A population-based study. *Salud Publica De Mexico, 46*, 113-122.
- Rutstein, S., & Johnson, K. (2004). *The DHS wealth index*. Calverton, MD: ORC Macro.
- Salam, M. A., Alim, M. A., & Noguchi, T. (2006). Spousal abuse against women and its consequences on reproductive health: A study in the urban slums in Bangladesh. *Maternal and Child Health Journal, 10*(1), 83-92.
- Schuler, S. R. (1998). Men's violence against women in rural Bangladesh: Undermined or exacerbated by microcredit programmes? *Development in Practice, 8*(2), 148.
- Schuler, S. R., Bates, L. M., Maseljo, J., & Islam, F. (2006). *Women's rights and gender-based violence and recourse-seeking in rural Bangladesh*. Report prepared

- for the Bangladesh Human Rights Advocacy Program, Center for Civil Society and Governance, Academy for Educational Development. Dhaka: March 2006.
- Schuler, S. R., Hashemi, S. M., Riley, A. P., & Akhter, S. (1996). Credit programs, patriarchy and men's violence against women in rural Bangladesh. *Social Science & Medicine*, *43*, 1729-1742.
- Silverman, J. G., Gupta, J., Decker, M. R., Kapur, N., & Raj, A. (2007). Intimate partner violence and unwanted pregnancy, miscarriage, induced abortion, and stillbirth among a national sample of Bangladeshi women. *International Journal of Obstetrics and Gynecology*, *114*, 1246-1252.
- Stata Corporation. (2007). *Stata survey data reference manual (release 10)*. College Station, TX: Stata Press.
- Straus, M. A. (2004). Cross-cultural reliability and validity of the Revised Conflict Tactics Scales: A study of university student dating couples in 17 nations. *Cross-cultural Research*, *38*, 407-432.
- Straus, M. A., Hamby, S. L., Boney-McCoy, S., & Sugarman, D. B. (1996). The revised conflict tactics scales (CTS2): Development and preliminary psychometric data. *Journal of Family Issues*, *17*, 283-316.
- Sugarman, D. D. B. (1996). Patriarchal ideology and wife-assault: A meta-analytic review. *Journal of Family Violence*, *11*, 13-40.
- Tokuç, B., Ekuklu, G., & Avcioğlu, S. (2010). Domestic violence against married women in Edirne. *Journal of Interpersonal Violence*, *25*, 832-847.
- Vung, N. D., Ostergren, P. O., & Krantz, G. (2008). Intimate partner violence against women in rural Vietnam—different socio-demographic factors are associated with different forms of violence: Need for new intervention guidelines? *BMC Public Health*, *8*, 55.
- Wahed, T., & Bhuiya, A. (2007). Battered bodies & shattered minds: Violence against women in Bangladesh. *The Indian Journal of Medical Research*, *126*, 341-354.
- World Health Organization. (2001). Putting women first: Ethical and safety recommendations for research on domestic violence against women. Geneva, Switzerland: Author.
- World Health Organization. (2005). *WHO multi-country study on women's health and domestic violence against women: Summary report of initial results on prevalence, health outcomes and women's responses*. Geneva, Switzerland: Author.

Bios

William Sambisa, PhD, is a research associate with the Carolina Population Center of the University of North Carolina at Chapel Hill. He has worked in the family planning and HIV/AIDS sector in sub-Saharan Africa for more than 8 years as a monitoring and evaluation specialist. His research has focused on ethnicity, gender, and risky sexual

behavior. He has experience with analyses of household data sets and conduct of quantitative surveys in Bangladesh, Kenya, Zambia, and Zimbabwe.

Gustavo Angeles, PhD, is research assistant professor in the Department of Maternal and Child Health at the University of North Carolina at Chapel Hill, fellow at the Carolina Population Center, and deputy director of the MEASURE Evaluation Project. A health economist, his technical areas of expertise include monitoring and evaluation of population and health programs, household and health facility survey research, measurement of socioeconomic status, statistical modeling of health-related behaviors, and methods for evaluating the impact of interventions on human capital outcomes. He has extensive field experience and has provided technical assistance on evaluation and household survey data collection and analysis in Tanzania, Kenya, Paraguay, Ecuador, Nicaragua, India, and Bangladesh. He has organized and conducted workshops on monitoring and program impact evaluation in the United States, Vietnam, Thailand, Costa Rica, and Mexico.

Peter M. Lance, PhD, is an economist at the MEASURE Evaluation Project at the University of North Carolina at Chapel Hill. He has been conducting monitoring and evaluation and health interview surveys in Bangladesh for 5 years. His research focuses on program impact evaluation and the distribution of health and health-related behaviors in lower-income nations.

Ruchira Tabassum Naved, PhD, is a scientist with the public health sciences division of International Diarrheal Diseases Research, Bangladesh (ICDDR, B). She has more than 15 years of research experience in the field of gender. She has also worked with Population Council and International Food Policy Research Institute. Her recent research has particularly focused on violence against women and its effect on women and children. She combines both quantitative and qualitative research methods in her work.

Juliana Thornton, is a masters student in the Department of Maternal and Child Health at the University of North Carolina at Chapel Hill. Prior to this she worked as the director of research at an NGO called NOAH working with community groups to care for orphans and vulnerable children in South Africa. She holds a bachelor's degree cum laude in anthropology from the University of Chicago and has conducted survey research in South Africa and Tanzania. Her current research interests include HIV prevention, the impact of HIV/AIDS on women and children, women's rights, and child protection.