

**Molecular cloning and expression in  
Saccharomyces cerevisiae of two Aspergillus  
nidulans xylanase genes.**

J A Pérez-Gonzalez, L H De Graaff, J Visser and D Ramón  
*Appl. Environ. Microbiol.* 1996, 62(6):2179.

---

Updated information and services can be found at:  
<http://aem.asm.org/content/62/6/2179>

---

**CONTENT ALERTS**

*These include:*

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

---

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>  
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

## Molecular Cloning and Expression in *Saccharomyces cerevisiae* of Two *Aspergillus nidulans* Xylanase Genes

JOSÉ A. PÉREZ-GONZÁLEZ,<sup>1\*</sup> LEO H. DE GRAAFF,<sup>2</sup> JAAP VISSER,<sup>2</sup> AND DANIEL RAMÓN<sup>1</sup>

*Departamento de Biotecnología de los Alimentos, Instituto de Agroquímica y Tecnología de los Alimentos, Consejo Superior de Investigaciones Científicas, 46100 Burjassot, Valencia, Spain,<sup>1</sup> and Molecular Genetics of Industrial Microorganisms, Wageningen Agricultural University, NL-6703 HA Wageningen, The Netherlands<sup>2</sup>*

Received 8 November 1995/Accepted 6 March 1996

**Two *Aspergillus nidulans* genes, *xlnA* and *xlnB*, encoding the X<sub>22</sub> and X<sub>24</sub> xylanases from this fungus, respectively, have been cloned and sequenced. Their cDNAs have been expressed in a laboratory *Saccharomyces cerevisiae* strain under the control of a constitutive yeast promoter, resulting in the construction of recombinant xylanolytic yeast strains.**

The breakdown of xylan, a complex polymer of plant cell walls, requires the action of several enzymes, including xylanases,  $\beta$ -xylosidases, acetyl xylan esterases, and several debranching enzymes (2). Xylanases, which are responsible for depolymerization of the xylan backbone, have potentially broad biotechnological applications (2, 3, 32). At present, most commercial xylanolytic preparations are obtained from filamentous-fungus species. As with other xylanolytic microorganisms, filamentous fungi produce multiple xylanases (33) whose genes have been cloned and sequenced from the following: *Aureobasidium pullulans* (22), *Cochliobolus carbonum* (1), *Penicillium chrysogenum* (17), *Trichoderma reesei* (25, 30), *Aspergillus awamori* (18), *Aspergillus kawachi* (20, 21), and *Aspergillus tubingensis* (4). Little is known about the mechanisms controlling xylanase gene expression. Only in the case of the *A. tubingensis xlnA* gene has a regulatory element involved in xylan-specific induction been clearly identified in the gene promoter (4).

*Aspergillus nidulans* is genetically well characterized and hence an ideal system for the study of gene regulation. *A. nidulans* produces three major xylanases, X<sub>22</sub>, X<sub>24</sub>, and X<sub>34</sub> (10), so called by virtue of their molecular weights, which have been recently purified and characterized (7–9). This report presents the cloning, sequencing, and yeast heterologous expression of the *xlnA* and *xlnB* genes, which encode the X<sub>22</sub> and X<sub>24</sub> xylanases, respectively.

**Cloning of the *xlnA* and *xlnB* genes from *A. nidulans*.** An *A. nidulans* genomic library constructed in  $\lambda$  Charon 4A (34) was screened by heterologous hybridization using a DNA fragment containing the *A. tubingensis xlnB* gene (5) as a probe; 10 positive plaques were detected and purified. Two different patterns of hybridization signals were observed in Southern analyses of restriction endonuclease digestions of the recombinant phages (results not shown). These patterns correlated well with those seen on blots of *A. nidulans* genomic digests in which at least two specific signals were detected for each restriction-enzyme-treated DNA sample. From the phage DNAs, two fragments hybridizing to the *A. tubingensis xlnB* probe, a 6.2-kb *Bam*HI fragment containing the putative xylanase gene *xlnA* (see below) and a 5-kb *Eco*RI fragment containing the

putative xylanase gene *xlnB* (see below), were isolated and subcloned in pUC18.

**Nucleotide sequences of the *xlnA* and *xlnB* genes.** The DNA fragments hybridizing to the *A. tubingensis xlnB* probe were partially sequenced. In both cases, open reading frames interrupted by introns were identified by homology to the *A. tubingensis xlnA* (4) and *xlnB* (5) genes. The positions of the introns and their processing were confirmed by partial sequencing of the *xlnA* and *xlnB* cDNAs (see below). The intron position is conserved between the *xlnA* genes from both *A. nidulans* and *A. tubingensis*, though the intron is larger in the *A. nidulans* gene (68 versus 49 bp). The N termini of the translated open reading frames of *xlnA* and *xlnB* correspond to the N-terminal amino acid sequences (STPSSTGWSNGYYYSF and STPSSTGTSG) determined for *A. nidulans* X<sub>22</sub> and X<sub>24</sub>, respectively.

A signal peptide cleavage site (31) is predicted to be present between residues 19 and 20 after the initiation codon in the *xlnA* product and between residues 18 and 19 in the *xlnB* product (Fig. 1). In addition, 18- and 15-amino-acid putative propeptides preceding the N-terminal amino acids of the X<sub>22</sub> and X<sub>24</sub> mature proteins, respectively, would be removed by a protease at the monobasic cleavage sites (26). Such two-step proteolytic processing has been suggested for many other secreted fungal proteins (27, 28, 30) and shown for the *Aspergillus niger* glucoamylase (19). The *xlnA*-encoded mature protein is 187 amino acids long and has a calculated molecular weight of 20,247 and an isoelectric point of 6.33, in good agreement with the values determined for the purified X<sub>22</sub> enzyme (8). The *xlnB*-encoded mature protein is 182 amino acids long and has a calculated molecular weight of 20,088 and an isoelectric point of 4.22, values which differ from those determined for the purified X<sub>24</sub> protein (24,000 and <3.6, respectively) (9). The differences between the predicted and determined values may be due to glycosylation or to a particular configuration of this protein. There are no putative N glycosylation sites in the sequence of the *xlnB* product (nor were sites detected in the *xlnA* product sequence), and no sugar modifications were observed for the purified protein (6); hence, the second possibility is more likely.

Both *A. nidulans xlnA* and *xlnB* gene products showed homologies to those low-molecular-weight xylanases of bacterial and fungal (Fig. 1) origins classified in the G family of glycosyl hydrolases (14, 29). However, no clear homology was found with the F family xylanases. The identity values at the nucleotide and amino acid levels between the *xlnA* and *xlnB* genes

\* Corresponding author. Mailing address: Departamento de Biotecnología de los Alimentos, Instituto de Agroquímica y Tecnología de los Alimentos, Apartado de Correos 73, 46100 Burjassot, Valencia, Spain. Phone: 34-6-3900022. Fax: 34-6-3636301. Electronic mail address: jantonio@iata.csic.es.






FIG. 2. Growth (filled symbols) and xylanase activities (open symbols) in culture media inoculated with *S. cerevisiae* OL1::Yeplac181 (triangles), *S. cerevisiae* OL1::pYLA1 (circles), and *S. cerevisiae* OL1::pYLB1 (squares).

lanase activities was also tested in liquid cultures. Very low xylanase activities, measured as described above, were detected in uracil-deficient minimal medium culture filtrates for both recombinant yeast strains. However, activity was easily detectable in supernatants of yeast extract-peptone-dextrose (YPD)-rich medium cultures (Fig. 2). Although this is not a selective medium, the stabilities of both 2 $\mu$ m-based plasmids were found to be high by analysis of Ura<sup>+</sup> and Ura<sup>-</sup> phenotypes (80% of cells retained the Ura<sup>+</sup> phenotype after 60-h culture). Xylanase activity was observed from the beginning of growth for both recombinant strains (Fig. 2), which is consistent with constitutive expression directed by the actin gene promoter. The xylanase activity of the *xlnA*-transformed strain was higher than that of the *xlnB*-transformed strain, but neither of the recombinant strains reached the level of activity detected in *A. nidulans* transformants or even in the *A. nidulans* G191 host strain (Table 1).

Zymograms of culture filtrates were made in order to detect xylanase activities after sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE). Equal amounts of each sample (16  $\mu$ l from original culture filtrates) were loaded onto gels. As can be seen in Fig. 3A, two clear bands corresponding to X<sub>22</sub> and X<sub>24</sub> xylanase activities were detected in the *A. nidulans* sample. A band detected in the *S. cerevisiae* OL1::pYLA1 culture filtrate corresponded to that produced by X<sub>22</sub> in *A. nidulans*. A second band, corresponding to that produced by X<sub>24</sub> in *A. nidulans* but more intense, was detected in the culture filtrate from *S. cerevisiae* OL1::pYLB1. No bands were detected in the *S. cerevisiae* OL1::Yeplac181 control sample. For immunoassay, 16- $\mu$ l samples of culture filtrates were run on an SDS-PAGE gel, blotted to nitrocellulose, and exposed to antibody prepared against *A. tubingensis* xylanase A (Fig. 3B). A faint band corresponding to X<sub>22</sub> was detected in the *A. nidulans* sample. As expected, no band corresponding to X<sub>24</sub> xylanase was detected, as it is a very minor protein in *A. nidulans* culture filtrates (the zymogram was more sensitive, as it detected this activity). A faint protein band with the same mobility as that of X<sub>22</sub> was detected in the culture filtrate from *S. cerevisiae* OL1::pYLA1. Two closely spaced and more intense protein bands were observed in *S. cerevisiae* OL1::pYLB1. From their mobilities, these proteins could correspond to X<sub>24</sub>, with the larger one possibly being a glycosylated form of this protein. No bands were detected in the *S. cerevisiae* OL1::Yeplac181 control. The results from the zymogram and immunoassay analyses confirm that both the *xlnA* and *xlnB* genes are expressed in the corresponding *S. cerevisiae* recombinant strains.




FIG. 3. SDS-PAGE followed by zymogram analysis (A) or immunostaining (B) of extracellular xylanases produced by *A. nidulans* and *S. cerevisiae*. Culture filtrates (16  $\mu$ l) were applied from *A. nidulans* grown for 40 h in *Aspergillus* minimal medium containing 0.5% oat spelt xylan (lanes 1), *S. cerevisiae* OL1::Yeplac181 grown for 42 h in YPD (lanes 2), *S. cerevisiae* OL1::pYLA1 grown for 42 h in YPD (lanes 3), and *S. cerevisiae* OL1::pYLB1 grown for 42 h in YPD (lanes 4). The positions of molecular weight markers (in thousands) are indicated on the right.

The previous activity assays performed with liquid (Fig. 2) and solid media showed that *S. cerevisiae* OL1::pYLB1 produced less xylanase activity than did *S. cerevisiae* OL1::pYLA1. However, the activity detected on the zymogram and the amount of protein detected in the immunoassay were much higher in the former than they were in the latter. The lower level of X<sub>22</sub> activity detected on the zymogram could be explained by our observation that X<sub>22</sub> protein renaturation after SDS-PAGE is more difficult than X<sub>24</sub> protein renaturation (23a). The specific activity of X<sub>22</sub> on birchwood xylan and oat spelt xylan is three times that of X<sub>24</sub> (6) and may explain why *S. cerevisiae* OL1::pYLA1 showed more activity in liquid and solid media than did *S. cerevisiae* OL1::pYLB1, which secretes more protein, as detected by immunoassay. In addition, it is also possible that some modification of X<sub>24</sub> by yeast cells inactivates the enzyme.

**Nucleotide sequence accession numbers.** The EMBL accession numbers of the *A. nidulans* *xlnA* and *xlnB* gene sequences are Z49892 and Z49893, respectively.

This work was supported by grants from D.G. XII of the European Commission (BIOTECH BIO2-CT93-0174) and the Spanish Government (CICYT AL193-0809). J. A. Pérez-González was the recipient of a short-term fellowship from the European Molecular Biology Organization.

Thanks are due to A. P. MacCabe for critically reading the manuscript and to P. Sanz for gifts of *S. cerevisiae* OL1 and the Yeplac181 plasmid.

#### REFERENCES

1. Apel, P. C., D. G. Panaccione, F. R. Holden, and J. D. Walton. 1993. Cloning and targeted gene disruption of *XYL1*, a  $\beta$ 1,4-xylanase gene from the maize pathogen *Cochliobolus carbonum*. *Mol. Plant-Microbe Interact.* **6**:467-473.
2. Biely, P. 1985. Microbial xylanolytic systems. *Trends Biotechnol.* **3**:286-289.
3. Coughlan, M. P., and G. P. Hazlewood. 1993.  $\beta$ -1,4-D-Xylan-degrading enzyme systems: biochemistry, molecular biology and applications. *Biotechnol. Appl. Biochem.* **17**:259-289.
4. de Graaff, L., H. C. van den Broeck, A. J. J. van Ooijen, and J. Visser. 1994. Regulation of the xylanase-encoding *xlnA* gene of *Aspergillus tubingensis*. *Mol. Microbiol.* **12**:479-490.
5. de Graaff, L. H. Personal communication.
6. Fernández-Espinar, M. T. 1994. Ph.D. thesis. University of Valencia, Valencia, Spain.
7. Fernández-Espinar, M. T., F. Piñaga, L. de Graaff, J. Visser, D. Ramón, and S. Vallés. 1994. Purification, characterization and regulation of the synthesis of an *Aspergillus nidulans* acidic xylanase. *Appl. Microbiol. Biotechnol.* **42**: 555-562.
8. Fernández-Espinar, M. T., F. Piñaga, P. Sanz, D. Ramón, and S. Vallés.

1993. Purification and characterization of a neutral endoxylanase from *Aspergillus nidulans*. FEMS Microbiol. Lett. **113**:223–228.
9. Fernández-Espinar, M. T., F. Piñaga, S. Vallés, J. A. Pérez-González, and D. Ramón. Construction of an *Aspergillus nidulans* multicopy transformant for the *xlnB* gene and its use to purify the minor X<sub>24</sub> xylanase. Appl. Microbiol. Biotechnol., in press.
  10. Fernández-Espinar, M. T., D. Ramón, F. Piñaga, and S. Vallés. 1992. Xylanase production by *Aspergillus nidulans*. FEMS Microbiol. Lett. **91**:91–96.
  11. Frohman, M. A., M. K. Dush, and G. R. Martin. 1988. Rapid production of full-length cDNAs from rare transcripts: amplification using a single gene-specific oligonucleotide primer. Proc. Natl. Acad. Sci. USA **85**:8998–9002.
  12. Gietz, D., A. St. Jean, R. A. Woods, and R. H. Schiestl. 1992. Improved method for high efficiency transformation of intact yeast cells. Nucleic Acids Res. **20**:1425.
  13. Gietz, R. D., and A. Sugino. 1988. New yeast-*Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. Gene **74**:527–534.
  14. Gilkes, N. R., B. Henrissat, D. G. Kilburn, R. C. Miller, Jr., and R. A. J. Warren. 1991. Domains in microbial  $\beta$ -1,4-glycanases: sequence conservation, function, and enzyme families. Microbiol. Rev. **55**:303–315.
  15. González, R., D. Ramón, and J. A. Pérez-González. 1992. Cloning, sequence analysis and yeast expression of the *egl1* gene from *Trichoderma longibrachiatum*. Appl. Microbiol. Biotechnol. **38**:370–375.
  16. Goosen, T., F. van Engelenburg, F. Debets, K. Swart, K. Bos, and H. van der Broek. 1989. Tryptophan auxotrophic mutants in *Aspergillus niger*: inactivation of the *trpC* gene by co-transformation mutagenesis. Mol. Gen. Genet. **219**:282–288.
  17. Haas, H., E. Friedlin, G. Stöffler, and B. Redl. 1993. Cloning and structural organization of a xylanase-encoding gene from *Penicillium chrysogenum*. Gene **126**:237–242.
  18. Hensing, J. G. M., C. von Rotterdam, J. M. A. Verbakel, M. Roza, J. Maat, R. F. M. van Gorcom, and C. A. M. J. van den Hondel. 1994. Isolation and characterization of a 1,4- $\beta$ -endoxylanase gene of *A. awamori*. Curr. Genet. **26**:228–232.
  19. Innis, M. A., M. J. Holland, P. C. McCabe, G. E. Cole, V. P. Wittman, R. Tal, K. W. K. Watt, D. H. Gelfand, J. P. Holland, and J. H. Meade. 1985. Expression, glycosylation and secretion of an *Aspergillus* glucoamylase by *Saccharomyces cerevisiae*. Science **228**:21–26.
  20. Ito, K., T. Ikemasu, and T. Ishikawa. 1992. Cloning and sequencing of the *xynA* gene encoding xylanase A of *Aspergillus kawachi*. Biosci. Biotechnol. Biochem. **56**:906–912.
  21. Ito, K., K. Iwashita, and K. Iwano. 1992. Cloning and sequencing of the *xynC* gene encoding acid xylanase of *Aspergillus kawachi*. Biosci. Biotechnol. Biochem. **56**:1338–1340.
  22. Li, X.-L., and L. G. Ljungdahl. 1994. Cloning, sequencing, and regulation of a xylanase gene from the fungus *Aureobasidium pullulans* Y-2311-1. Appl. Environ. Microbiol. **60**:3160–3166.
  23. Ng, R., and J. Abelson. 1980. Isolation and sequence of the gene for actin in *S. cerevisiae*. Proc. Natl. Acad. Sci. USA **77**:3912–3916.
  - 23a. Pérez-González, J. A., L. H. De Graaff, J. Visser, and D. Ramón. Unpublished observation.
  24. Pérez-González, J. A., R. González, A. Querol, J. Sendra, and D. Ramón. 1993. Construction of a recombinant wine yeast strain expressing  $\beta$ -(1,4)-endoglucanase and its use in microvinification processes. Appl. Environ. Microbiol. **59**:2801–2806.
  25. Saarelainen, R., M. Paloheimo, R. Fagerström, P. L. Suominen, and K. M. H. Nevalainen. 1993. Cloning, sequencing and enhanced expression of the *Trichoderma reesei* endoxylanase II (pI 9) gene *xln2*. Mol. Gen. Genet. **241**:497–503.
  26. Schwartz, T. W. 1986. The processing of peptide precursors, 'proline-directed arginyl cleavage' and other monobasic processing mechanisms. FEBS Lett. **200**:1–9.
  27. Stålbrand, H., A. Saloheimo, J. Vehmaanperä, B. Henrissat, and M. Penttilä. 1995. Cloning and expression in *Saccharomyces cerevisiae* of a *Trichoderma reesei*  $\beta$ -mannanase gene containing a cellulose binding domain. Appl. Environ. Microbiol. **61**:1090–1097.
  28. Teeri, T. T., P. Lehtovaara, S. Kauppinen, I. Salovouri, and J. Knowles. 1987. Homologous domains in *Trichoderma reesei* cellulolytic enzymes: gene sequence and expression of cellobiohydrolase II. Gene **51**:43–52.
  29. Törrönen, A., C. P. Kubicek, and B. Henrissat. 1993. Amino acid sequence similarities between low molecular weight endo-1,4-xylanases and family H cellulases revealed by clustering analysis. FEBS Lett. **321**:135–139.
  30. Törrönen, A., R. L. Mach, R. Messner, R. González, N. Kalkkinen, A. Harkki, and C. P. Kubicek. 1992. The two major xylanases from *Trichoderma reesei*: characterization of both enzymes and genes. Bio/Technology **10**:1461–1465.
  31. von Heijne, G. 1986. A new method for predicting signal sequence cleavage sites. Nucleic Acids Res. **14**:4683–4690.
  32. Wong, K. K. Y., and J. N. Saddler. 1993. Applications of hemicellulases in the food, feed and pulp and paper industries, p. 127–143. In M. P. Coughlan and G. P. Hazlewood (ed.), Hemicellulose and hemicellulases. Portland Press, London.
  33. Wong, K. K. Y., L. U. L. Tan, and J. N. Saddler. 1988. Multiplicity of  $\beta$ -1,4-xylanase in microorganisms: functions and applications. Microbiol. Rev. **52**:305–317.
  34. Zimmermann, C. R., W. C. Orr, R. F. Leclerc, E. C. Barnard, and W. E. Timberlake. 1980. Molecular cloning and selection of genes regulated in *Aspergillus* development. Cell **23**:709–715.