

Short lists with short programs for functions

Nikolay Vereshchagin*

Moscow State University, Yandex and Higher School of Economics

Abstract

Let $\{\varphi_p\}$ be an optimal Gödel numbering of the family of computable functions (in Schnorr's sense), where p ranges over binary strings. Assume that a list of strings $L(p)$ is computable from p and for all p contains a φ -program for φ_p whose length is at most ε bits larger than the length of the shortest φ -program for φ_p . We show that for infinitely many p the list $L(p)$ must have $2^{|p|-\varepsilon-O(1)}$ strings. Here ε is an arbitrary function of p .

1 Results

A numbering of a family of computable functions of m variables is a computable partial function $\varphi : (\{0, 1\}^*)^{m+1} \rightarrow \{0, 1\}^*$. We call p a φ -index or a φ -program for the function $\langle x_1, \dots, x_m \rangle \mapsto \varphi(p, x_1, \dots, x_m)$, which is denoted as φ_p . A numbering φ is *universal* if for all computable partial functions f from $(\{0, 1\}^*)^m$ to $\{0, 1\}^*$ there is p with $\varphi_p = f$.

By $C_\varphi(f)$ we denote the minimal length of a φ -program for f (*Kolmogorov complexity of f with respect to φ*). A numbering φ has *Kolmogorov property*, if for every other numbering ψ there is a constant c such that $C_\varphi(f) \leq C_\psi(f) + c$ for all functions f .

A numbering φ is called a *Gödel numbering* if for every other numbering ψ there is a total computable function t (called a *translator* from ψ to φ) such that $\psi_p = \varphi_{t(p)}$ for all p . A Gödel numbering φ is called an *optimal Gödel numbering* if for all numberings ψ there is a translator t from ψ to φ that has additional property $|t(p)| \leq |p| + O(1)$ (the translator is *linearly*

*The work was done while visiting IMS (University of Singapore), the program "Algorithmic Randomness", 2-30 June 2014. The work was in part supported by the RFBR grant 12-01-00864.

bounded).¹ Here and further $|p|$ denotes the length of p . Every optimal Gödel numbering has Kolmogorov property but not the other way around.

Example 1. Here is an example of an optimal Gödel numbering φ of the family of computable functions of m variables. Let Φ denote a universal numbering of the family of computable functions of $m + 1$ variables. Let $p \mapsto \hat{p}$ denote a computable prefix encoding, for instance, $\hat{p} = 0^{|p|}1p$. Then $\varphi(\hat{p}q, x_1, \dots, x_m) = \Phi(p, q, x_1, \dots, x_m)$ is an optimal Gödel numbering of the family of computable functions of m variables. Indeed, the mapping $t(q) = \hat{p}q$ is a linearly bounded translator from the numbering Φ_p to φ .

The above definitions make sense also for $m = 0$. In this case φ_p is understood as $\varphi(p)$ if defined and as a special symbol \perp otherwise. Optimal Gödel numberings for $m = 0$ were called *standard machines* in [1] and we will use the same terminology. Kolmogorov complexity $C_U(x)$ of a string x with respect to a standard machine U is the usual Kolmogorov complexity (the minimal length of a U -program for x).

The paper [1] shows that for every standard machine U , given a string x we can find a short list of strings with a short program for x : the size (=cardinality) of the list is $O(|x|^2)$ and it contains a U -program for x of length at most $C_U(x) + O(1)$.

Is there a total algorithm that computes a short list with a short program for x from any U -program for x ? This question was asked recently by Alexander Shen [5]. Notice that there is no total algorithm that maps any program for x to x (otherwise the positive answer to the question would immediately follow from the cited result of [1]). We show that for every standard machine U and for every function ε of p there are infinitely pairs $(x, \text{its } U\text{-program } p)$ such that the size of $L(p)$ is exponential in both $|x| - \varepsilon$ and $|p| - \varepsilon$ provided $L(p)$ has a program for x of length at most $C_U(x) + \varepsilon$.

Let $C_{U,L}(x)$ denote the minimal length of a U -program $p \in L$ for x .

Theorem 1. *Let U be a standard machine and L a total computable function mapping (binary) strings to finite sets of strings. Then for some c for all k the following holds. There is a string x and its U -program p of length between k and $k + c$ such that $\#L(p) \geq 2^{|x|} - 2$ and $C_{U,L(p)}(x) \geq k$.*

Corollary 2 (A negative answer to Shen's question). *Let U, L be as in the theorem. Let $\varepsilon(p)$ denote $C_{U,L(p)}(U(p)) - |U(p)|$. Then for infinitely many*

¹ The term "optimal Gödel numbering" was introduced by Schnorr [4]. Teutsch and Zimand [2] call optimal Gödel numberings *Kolmogorov numberings*. However Kolmogorov has neither introduced nor studied them.

p the size of $L(p)$ is at least $2^{|p|-\varepsilon(p)-O(1)} - 2$. Moreover, for those p 's the size of $L(p)$ is at least $2^{|U(p)|} - 2$ and $|U(p)| > |p| - \varepsilon(p) - O(1)$.

Notice, that Kolmogorov complexity is less than the length (up to an additive constant) and hence the corollary holds for $\varepsilon(p) = C_{U,L(p)}(U(p)) - C_U(U(p))$ as well.

Proof of the corollary. Let p, x be the pairs existing by Theorem 1. The last inequality of the theorem implies $|x| + \varepsilon(p) = C_{U,L(p)}(x) \geq k = |p| + O(1)$ and hence $|x| \geq |p| - \varepsilon(p) - O(1)$. \square

Let us stress that L is assumed to be a total function. If we allowed L to be defined only on those strings p for which $U(p)$ halts, then there would be a computable list $L(p)$ of quadratic size (in the length of $x = U(p)$) with a program for x of length at most $C_U(x) + O(1)$, which follows from the result of [1].


Example 2. This example provides a family of computable lists for which the lower bounds for the size of L and for $C_{U,L}(x)$ established in Theorem 1 are tight.

The lower bound $k = |p| + O(1)$ for $C_{U,L}(x)$ is tight (up to an additive constant) for any list $L(p)$ containing p , for instance, for $L(p) = \{p\}$. For this list the lower bound for the size is tight too, however, this is not very impressive, as the list is too small.

There is much larger computable list $L(p) = \{p\} \cup \{0, 1\}^{<|p|}$ for which both lower bounds are tight. Indeed, the length of the string x in the theorem is $|p| + O(1)$, as $C_{U,L(p)}(x) = C_U(x) \leq |x| + O(1)$ and on the other hand $C_{U,L(p)}(x) \geq k = |p| + O(1)$.

Moreover, there are similar lists of any log-cardinality between 0 and $|p|$. Fix any computable function $p \mapsto i \leq |p|$ and consider the computable list $L_i(p) = \{p\} \cup \{0, 1\}^{<i}$. For this list we have $\#L(p) = 2^i$ and $C_{U,L(p)}(x) = C_U(x)$ if $i > C_U(x)$ and $C_{U,L(p)}(x) = |p|$ otherwise.

The parameters $(\log \#L_i(p), C_{U,L_i(p)}(x))$ for these lists are shown in the following picture (where we drop the subscript U):



More specifically, they lie on the horizontal straight line segments on the border of the gray area P .

Let us show that the lower bound of the size in the theorem is tight for all computable lists of the form $L_i(p)$. That is, we will show that the length of the string x existing by the theorem is $i + O(1)$. As $2^{|x|} - 2 \leq \#L_i(p) = 2^i$, we have $|x| - 1 \leq i$ and hence $|x| - 1 \leq i \leq |p| = C_{U, L_i(p)}(x) + O(1)$. If $i \leq C_U(x)$ then we have $|x| - 1 \leq i \leq C_U(x)$ and hence these inequalities are equalities up to an additive constant. Otherwise $i > C_U(x)$ and hence $C_{U, L_i(p)}(x) = C_U(x)$. In this case $|x| - 1 \leq i \leq C_{U, L_i(p)}(x) + O(1) = C_U(x) + O(1)$ and again these inequalities are equalities up to an additive constant.

Theorem 1 easily translates to optimal Gödel numberings of functions of arbitrary number of variables. For general case the statement is the following. Let Singl_x denote the function defined only on the tuple $\langle x, \dots, x \rangle$ with value x . Let $C_{\varphi, L}(f)$ denote the minimal length of a φ -program $p \in L$ for f .

Theorem 3. *Let φ be an optimal Gödel function of $m > 0$ variables and L a total computable function mapping strings to finite sets of strings. Then for some c for all k the following holds. There is a string x and a φ -program p of length between k and $k + c$ for the function Singl_x such that $\#L(p) \geq 2^{|x|} - 2$ and $C_{\varphi, L(p)}(\text{Singl}_x) \geq k$.*

Remark. Theorem 3 holds for numberings of enumerable sets with the singleton set $\{x\}$ in place of the function Singl_x . The proof is entirely similar.

For the string p from Theorem 3 we have

$$\log \#L(p) + C_{U, L(p)}(\varphi_p) \geq C_U(\varphi_p) + |p| - O(1). \quad (1)$$

Indeed, $\log \#L(p) \geq |x| - O(1) \geq C_U(\varphi_p) - O(1)$ and $C_{U, L(p)}(\varphi_p) \geq k \geq |p| - O(1)$. Summing these inequalities we get (1).

Theorem 3 answers a question asked recently by Teutsch and Zimand. For a numbering φ of computable functions of one variable, Teutsch and Zimand [2] considered the set of minimal programs for φ , where p is called *minimal*, if for all $q < p$ we have $\varphi_q \neq \varphi_p$. Here $<$ denotes the lexicographical ordering on binary strings (more precisely, $p < q$ iff $|p| < |q|$ or $|p| = |q|$ and p is lexicographically less than q). The minimal φ -program for a function φ_q is denoted by $\min_\varphi(q)$. Teutsch and Zimand showed the following.

- If φ is a Gödel numbering and a computable function L on input p returns a list $L(p)$ containing $\min_\varphi(p)$, then the size of that list cannot be constant.
- For every numbering φ with Kolmogorov property, if a computable function L on input p returns a list containing $\min_\varphi(p)$, then the size of the list must be $\Omega(|p|^2)$.
- There exists an optimal Gödel numbering φ such that if a computable function on input p returns a list containing $\min_\varphi(p)$, then the size of that list must be $\Omega(2^{|p|})$.

In summary, their results show that a computable list that contains the minimal φ -program cannot be too small.

Along the lines of the second result Teutsch and Zimand asked the following question: is there an optimal Gödel numbering φ with a computable list $L(p)$ that contains $\min_\varphi(p)$ and has size $O(|p|^2)$?

Theorem 3 implies the negative answer to this question. Indeed, if $\min_\varphi(p) \in L(p)$ for all p then $C_{\varphi, L(p)}(\varphi_p) = C_\varphi(\varphi_p)$ for all p . By (1) for the pair p, x existing by the theorem the size of $L(p)$ must be at least $2^{|p| - O(1)}$. In other words, the third result of Teutsch and Zimand holds for *all* optimal Gödel numberings φ .

So far we were constructing for a given computable function L inputs p such that the list $L(p)$ has large parameters $\#L(p)$ and $C_{U, L(p)}(U(p))$. Let us consider the “short list with short programs” problem from the other end. Are there p ’s such that *every* short list L computable from p by a total algorithm has high parameters $\#L(p)$ and $C_{U, L}(U(p))$? In this form the question is trivial: we can hard-wire the shortest U -program q for $U(p)$ into a total algorithm which will return the list $\{q\}$, which has optimal parameters. The question becomes reasonable if we restrict the complexity, say by $O(\log |p|)$, of the total algorithm producing the list from p .

To make this question precise consider the *total complexity* $CT_\Phi(a|b)$ defined as the minimal length of a Φ -program of a total function that maps

b to a . Here Φ is an optimal Gödel numbering of computable functions of one variable.

Fix a natural δ (the upper bound for total complexity). Then for each p consider the set

$$S_p^\delta = \{(i, j) \mid \exists L, CT(L|p) \leq \delta, \#L \leq 2^i, C_{U,L}(x) \leq j\},$$

where x stands for $U(p)$. The larger this set is the better parameters may have lists L with small $CT(L|p)$. If $\delta \geq \log |p| + O(1)$ then the list $\{0, 1\}^i$ for $i = C_U(U(p))$ and the list $\{p\}$ witness that the set S_p^δ includes the entire gray set P on the picture from Example 1.

The set S_p^δ may be much larger than the gray set P . For instance, this happens when p is a shortest program for $x = U(p)$. In this case the set S_p^δ coincides with the set of all points above the dashed line. Are there infinitely many p such that the set S_p^δ is close to the gray set P in the picture? In other words, are there infinitely many p such that for every list L with $CT_\Phi(L|p) \leq \delta$ either $\log_2 \#L > C_U(x)$, or $C_{U,L}(x) \geq |p|$ (with certain accuracy)? A positive answer is provided by the following

Theorem 4. *Let U be a standard machine. For all n and all $k > n$ there is a string x with $C_U(x) = n + O(1)$ and its U -program p of length at most $k + O(1)$ such that for all $\delta < k - \log k - O(1)$ and all L with $CT_\Phi(L|p) = \delta$ either $\#L \geq 2^{n - \delta - \log k - O(1)}$ or $C_{U,L}(x) \geq k - 1$.*

Notice that the inequality $C_{U,L}(x) \geq k - 1$ for $L = \{p\}$ implies that $|p| \geq k - O(1)$ and hence $|p| = k - O(1)$.

2 The proofs

We first drop in Theorems 1 and 3 the requirement $|p| \geq k$. As a reward, the lower bound for the list size will be a little bit stronger: $2^{|x|} - 1$ in place of $2^{|x|} - 2$.

Proof of Theorem 1. Let us first show that the statement of Theorem 1 is invariant: if it holds for some standard machine U then it holds for any other standard machine U' . Indeed, assume that Theorem 1 holds for a standard machine U . To show Theorem 1 for another standard machine U' and a list $L'(p')$, choose a linearly bounded translator t from U' to U and a linearly bounded translator s from U to U' . Let c' be a constant with $|t(p')| \leq |p'| + c'$.

Apply Theorem 1 to the machine U and the list $L(p) = t(L'(s(p)))$. By Theorem 1 for all k there is a string x and its U -program p of length at most $k + c' + c$ such that $\#L(p) > 2^{|x|} - 1$ and the list $L(p)$ does not contain any U -program for x of length less than $k + c'$.

Let $p' = s(p)$. By construction,

$$|p'| \leq |p| + O(1) \leq k + c' + c + O(1).$$

We also have

$$\#L'(p') \geq \#t(L'(p')) \geq 2^{|x|} - 1.$$

Finally the list $t(L(p'))$ does not contain any U -program of length less than $k + c'$ for x . Hence the list $L(p')$ does not contain any U' -program of length less than k for x .

Thus it suffices to prove Theorem 1 for the standard machine U from Example 1, that is for $U(\hat{p}q) = \Phi(p, q)$ where Φ is a Gödel numbering of the family of computable functions of one variable.

We will let $p = \hat{r}q$ where q is a string of length k and r does not depend on k . The statement of the theorem will follow from the following properties of r, q and the function $V \doteq \Phi_r$ (of one variable):

- q is a V -program of a string x such that
- $\#L(\hat{r}q) \geq 2^{|x|} - 1$ and
- the list $L(\hat{r}q)$ contains no U -program for x of length less than k .

Notice that the string $p = \hat{r}q$ has all the required properties.

It remains to find such V, r and q . The computable function V and its Φ -program r will be defined using the Kleene fixed point theorem [3]. By that theorem we may assume that computing V we have access to a Φ -program r for V . We construct an algorithm that enumerates the graph of V .

The algorithm enumerating the graph of V . We maintain for all k a string q_k of length k and a string x_k . At the start let q_k be any string of length k and let x_k be the empty string. Enumerate all the pairs $\langle q_k, x_k \rangle$ into the graph of V thus letting $V(q_k) = x_k$.

Then we start an enumeration of the graph of U . Each time a new pair appears in that enumeration, we look if the current situation is good or not. We consider the current situation *good for* k if the pair $\langle q_k, x_k \rangle$ has been enumerated into the graph of V , $\#L(\hat{r}q_k) \geq 2^{|x_k|} - 1$ and the list $L(\hat{r}q_k)$ has no \bar{U} -program for x_k of length less than k , where \bar{U} denotes the sub-function of U consisting of all pairs enumerated so far.

At the start $\bar{U} = \emptyset$ and thus the situation is good for all k . Each time a new pair appears in the enumeration of the graph of U , we look whether the situation has become bad for some k . Obviously that may happen only if a pair $\langle s, x_k \rangle$ with $|s| < k$ and $s \in L(\hat{r}q_k)$ is enumerated. In that case pick a new string q of length k (“new” means that q has not been used as q_k earlier). Let n be the integer with $2^{n+1} - 1 > \#L(\hat{r}q) \geq 2^n - 1$. For all strings x of length at most n consider the set $S(x) = \{s \mid \bar{U}(s) = x, |s| < k\}$ of \bar{U} -programs for x of length less than k . Pick any string x of length at most n such that $S(x)$ does not intersect the list $L(\hat{r}q)$. As $\#L(\hat{r}q) < 2^{n+1} - 1$ and the number of x 's is $2^{n+1} - 1$, there is such x . Then let $q_k = q$, $x_k = x$ and enumerate the pair $\langle q, x \rangle$ into the graph of V . The situation has become good for k . **End of Algorithm.**

By Kleene's theorem for some r this algorithm enumerates the graph of the function Φ_r . Let us show that for each k , starting from some moment the situation is good for k . Indeed, for any k the situation may become bad less than 2^k times for k , as that may happen only after a new pair of the form $\langle s, x_k \rangle$ with $|s| < k$ has appeared. On the other hand, the number of strings q of length k is 2^k and hence we indeed are able to repair the situation $2^k - 1$ times. \square

Proof of Theorem 3. Let Singl_\perp stand for the nowhere defined function. There is a linearly bounded total computable translator t mapping any U -program for $x \in \{0, 1\}^* \cup \{\perp\}$ (for a standard machine U) to a φ -program for the function Singl_x . There is also a linearly bounded total computable translator s mapping any φ -program for Singl_x back to a U -program for x . Given a list L we just apply Theorem 1 to the list $L'(p') = s(L(t(p')))$ and $k + c'$, where c' is a constant with $|s(p)| \leq |p| + c'$. \square

It remains to prove Theorems 1 and 3 as they are stated, that is, with the requirement $|p| \geq k$ and with the lower bound $2^{|x|} - 2$ for the list size. Given any computable list $L(p)$ we add p into the list and apply Theorem 1 in the proven form to the resulting list $L'(p)$. The list $L'(p)$ does have a U -program for x of length $|p|$ and has no U -program for x of length less than k . This implies that $|p| \geq k$. The program p fulfills Theorem 1 in the original form.

The proof of Theorem 3 is entirely similar.

Remark. As Jason Teutsch observed, one can prove Theorem 1 without using the fixed point theorem. To this end we modify the construction of V so that V becomes a standard machine. Specifically, we first let $V_{0q} = U_q$ for

all strings $0q$ starting with zero, where U is any standard machine. Then we define V_{1q} so that for all k there is a string $1q$ of length $k + 1$ such that

- $1q$ is a V -program of a string x such that
- $\#L(1q) \geq 2^{|x|} - 1$ and
- the list $L(1q)$ contains no V -program for x of length less than k .

This can be done by the same technique. The function V defined in this way satisfies the theorem. As we already observed, this implies that the theorem holds for all standard machines.

Proof of Theorem 4. The proof is very similar to that of Theorem 1. We construct a computable function V such that for all $k > n$ there are strings q, x of lengths k, n , respectively, with

- $V(q) = x$,
- $C_U(x) > n - 1$,
- for all $\delta < k - \log k - 2$ and all L with $CT_\Phi(L|q) = \delta$ and $\#L < 2^{n-\delta-\log k-1}$ we have $C_{U,L}(x) \geq k - 1$.

The algorithm enumerating the graph of V . This time we maintain for all k a bunch of pairs $\{(q_{kn}, x_{kn}) \mid n = 0, 1, \dots, k - 1\}$. The length of q_{kn} is k and the length of x_{kn} is n . At the start let q_{kn} be the n th string of length k and let x_{kn} be the first string of length n (independent of k). Enumerate all the pairs $\langle q_{kn}, x_{kn} \rangle$ into the graph of V thus letting $V(q_{kn}) = x_{kn}$.

Then we start an enumeration of the graph of U and an enumeration of the graph of Φ . We denote by \bar{U} and $\bar{\Phi}$ the sub-functions of U and Φ consisting of all pairs (triples) enumerated so far. For each k we look if the situation is *good for k* . This means that for all $n < k$ the pair $\langle q_{kn}, x_{kn} \rangle$ has been enumerated into the graph of V , $C_{\bar{U}}(x_{kn}) \geq n - 1$ and for all $\delta < k - \log k - 2$ and all L with $CT_{\bar{\Phi}}(L|q_{kn}) = \delta$ and $\#L < 2^{n-\delta-\log k-1}$ we have $C_{\bar{U},L}(x_{kn}) \geq k - 1$. Here $CT_{\bar{\Phi}}(L|q)$ means the minimal length of p such that $\bar{\Phi}_p$ is defined on all q' of length k and $\bar{\Phi}_p(q) = L$.

At the start \bar{U} and $\bar{\Phi}$ are empty and thus the situation is good for all k . Each time a new pair (triple) appears in the enumeration of the graphs of U or Φ , we look whether the situation has become bad for some k . This may happen only if $C_{\bar{U}}(x_{kn})$ has become less than $n - 1$ (for some $n < k$) or a new list L with $CT_{\bar{\Phi}}(L|q_{kn}) < k - \log k - 2$ appeared or for an old list L the value $C_{\bar{U},L}(x_{kn})$ has become less than $k - 1$ (for some $n < k$). In

all the cases we first change q_{kn} and then we change x_{kn} . The string q_{kn} is replaced by any a new string q of length k (“new” means that q has not been used as q_{k^*} earlier). The string x_{kn} is replaced by any string x of length n such that $C_{\bar{U}}(x) \geq n - 1$ and the set $S(x) = \{s \mid \bar{U}(s) = x, |s| < k - 1\}$ does not intersect the union (over all δ) of all lists L of cardinality less than $2^{n-\delta-\log k-1}$ with $CT_{\bar{\Phi}}(L|q) = \delta$. Notice that for every p there is only one list L with $\bar{\Phi}_p(q) = L$ thus the total number of strings in all these lists is less than $\sum_{\delta < k} 2^\delta \cdot 2^{n-\delta-\log k-1} = 2^{n-1}$. On the other hand, the number of strings x of length n with $C_{\bar{U}}(x) \geq n - 1$ is more than 2^{n-1} . Thus there is such x .

Then let $q_{kn} = q$, $x_{kn} = x$ and enumerate the pair $\langle q, x \rangle$ into the graph of V . The situation has become good for k . **End of Algorithm.**

We have to show that we are able to choose a new string of length k each time we need one. Any replacement of a string of the form q_{k^*} is caused by

- discovering a new p of length less than $k - 2 \log k - 2$ such that Φ_p is defined on all strings of length k (this may cause replacement of the whole bunch of q_{kn} 's, for all $n < k$), or
- discovering a new U -program r of length less than $k - 1$, which may cause the replacement of q_{kn} only if $U(r) = x_{kn}$ thus for a single n , or
- discovering a new halting U -program of length less than $n - 1$ for x_{kn} , which again may cause the replacement of q_{kn} only for a single n .

Thus the total number of strings q_{k^*} we need is less than

$$k + \sum_{\delta < k - \log k - 2} k 2^\delta + 2^{k-1} < 2^k.$$

To prove the theorem let p be the U -program of x obtained from q by translation from V to U . Then $|p| \leq k + O(1)$. Notice that $CT_{\bar{\Phi}}(L|q) \leq CT_{\bar{\Phi}}(L|p) + O(1)$. Indeed, let s be a linearly bounded translator from V to U . Then $\bar{\Phi}(r, s(q))$ is a computable function hence there is a total computable function t with $\bar{\Phi}_{t(r)}(q) = \bar{\Phi}_r(s(q))$. If $\bar{\Phi}_r$ is total then so is $\bar{\Phi}_{t(r)}$. Hence $CT_{\bar{\Phi}}(L|q) \leq CT_{\bar{\Phi}}(L|s(q)) + O(1)$. \square

Acknowledgments. The author is sincerely grateful to Alexander Shen for asking the question and hearing the preliminary version of the proof of the result. The author is grateful to Jason Teutsch for the idea of how to omit the use of the fixed point theorem. The author is grateful to Alexander Rubtsov for discovering “intermediate” lists. The author is also grateful to the hospitality of the IMS of University of Singapore.

References

- [1] Bruno Bauwens, Anton Makhlin, Nikolay Vereshchagin, Marius Zimand. Short lists with short programs in short time. In Proceedings 28-th IEEE Conference on Computational Complexity (CCC), Stanford, CA, pages 98–108, June 2013. ECCC report TR13-007.
- [2] Jason Teutsch and Marius Zimand. On approximate decidability of minimal programs. 2014. Available from <http://arxiv.org/abs/1409.0496> and <http://people.cs.uchicago.edu/~teutsch/papers/teutschpubs.html>.
- [3] Hartley Rogers, Jr., The Theory of Recursive Functions and Effective Computability, MIT Press, 1987.
- [4] C.P. Schnorr. Optimal enumerations and optimal Gödel numberings. *Mathematical Systems Theory*, 8(2):182–191, 1975.
- [5] Alexander Shen. A talk on a meeting during the IMS program “Algorithmic Randomness” (IMS, University of Singapore, 2–30 June 2014).