

Sufficient Conditions of Existence of Fix-Free Codes

Yekhanin Sergey¹
 Department of Cybernetics and
 Computer Science
 Moscow State University
 e-mail: yekhanin@cityline.ru

Abstract — Code is fix-free if no codeword is a prefix or a suffix of any other. In this paper we improve the best-known sufficient conditions on existence of fix-free codes by a new explicit construction. We also discuss the well-known Kraft-type conjecture on the existence of fix-free codes basing on the results obtained by computer checking.

I. INTRODUCTION

Let $C(k_1, k_2, \dots, k_n)$ denote a binary variable-length code C with k_1 codewords of length 1, k_2 codewords of length 2, ... and k_n codewords of length n .

Let k_1, \dots, k_n be arbitrary positive integers. To simplify further presentation we denote $\sum_{i=1}^n \frac{k_i}{2^i}$ by $\chi(k_1, \dots, k_n)$.

Recall that a code is called prefix-free {resp. suffix-free}, if no codeword is beginning {resp. ending} of another one. Code C , which is simultaneously prefix-free and suffix-free is called fix-free.

Our goal is to develop sufficient conditions on the existence of fix-free codes. The well-known Kraft-type conjecture [1] by R. Ahlswede is that for any values of parameters k_1, k_2, \dots, k_n , such that

$$\chi(k_1, k_2, \dots, k_n) \leq \frac{3}{4}, \quad (1)$$

there exists a fix-free code $C(k_1, k_2, \dots, k_n)$

The next section gives a short overview of particular cases in which the conjecture is proved.

The following lemma [1] shows that, if true, bound (1) is the best possible.

Lemma 1: For any $\varepsilon > 0$ there exist parameters k_1, k_2, \dots, k_n such that $\chi(k_1, k_2, \dots, k_n) \leq \frac{3}{4} + \varepsilon$ and there exists no fix-free code $C(k_1, k_2, \dots, k_n)$.

II. STATEMENT OF RESULTS

In this section we formulate the known sufficient conditions on existence of fix-free codes. Proofs of the next two theorems can be found in [1].

Theorem 1: If $\chi(k_1, \dots, k_n) \leq \frac{1}{2}$ then there exists a fix-free $C(k_1, \dots, k_n)$.

Theorem 2: Suppose that if $i < j$ and $k_i > 0$ and $k_j > 0$, then $i < 2j$. Then $\chi(k_1, \dots, k_n) \leq \frac{3}{4}$ implies the existence of a fix-free code $C(k_1, \dots, k_n)$.

The main result presented in current paper is formulated by the next theorem. The sketch of proof is given in section three.

Theorem 3: Let k_1, k_2, \dots, k_n be arbitrary nonnegative integers. Suppose that the following statements are true:

$$\chi(k_1, \dots, k_n) \leq \frac{3}{4},$$

$$\exists p : k_1 = \dots = k_{p-1} = 0, \text{ and } \frac{k_p}{2^p} + \frac{k_{p+1}}{2^{p+1}} \geq \frac{1}{2}.$$

This implies the existence of fix-free code $C(k_1, k_2, \dots, k_n)$.

Corollary 1: If $\chi(k_1, \dots, k_n) \leq \frac{3}{4}$ and $k_1 = 1$ then there exists a fix-free code $C(k_1, k_2, \dots, k_n)$.

Note, that all the theorems formulated above are particular cases of conjecture (1). We have applied computer programming to check the conjecture for the small values of n . Thus the following theorem was obtained.

Theorem 4: Let k_1, \dots, k_n be arbitrary nonnegative integers such that $\chi(k_1, \dots, k_n) \leq \frac{3}{4}$ and $n \leq 8$ then there exists a fix-free code $C(k_1, \dots, k_n)$.

One more sufficient condition of existence of fix-free codes is given in [2].

III. SKETCH OF THE PROOF

We say that set $D \subseteq \{0, 1\}^n$ is *left regular* {*right regular*} if all $(n-1)$ -prefixes {suffixes} of words from D are pairwise distinct.

Let $C(k_1, \dots, k_n)$ be a fix-free code. We say that a vector $w \in \{0, 1\}^n$ is prefix-free {suffix-free} over C if no codeword $c \in C$ is a prefix {suffix} of w . Further by $M(C)$ { $\hat{M}(C)$ } we denote the set all binary vectors of length n that are prefix-free {suffix-free} over C .

Definition: We say that fix-free code $C(k_1, \dots, k_n)$ is a π -system if $M(C)$ is right regular, $\hat{M}(C)$ is left regular and $\chi(k_1, \dots, k_n) = \frac{1}{2}$.

We split the proof into two sections. Firstly we study the properties and develop explicit constructions of π -systems. The main result of this section is formulated by

Lemma 2: If $k_1 = \dots = k_{n-2} = 0$, $\frac{k_{n-1}}{2^{n-1}} + \frac{k_n}{2^n} = \frac{1}{2}$, then there exists a π -system $C(k_1, k_2, \dots, k_n)$.

Secondly we study the relationship between π -systems and fix-free codes and prove that an arbitrary π -system $C_1(k_1, \dots, k_p)$ can be extended to fix-free code $C_2(k_1, \dots, k_p + b, \dots, k_n)$, where $\chi(k_1, \dots, k_p + b, \dots, k_n) \leq \frac{3}{4}$.

REFERENCES

- [1] R. Ahlswede, B. Balkenhol, L. Kharchatrian, Some Properties of Fix-Free Codes. In *Proc. First INTAS International Seminar on Coding Theory and Combinatorics*, pp. 20-33, Thazkhadzor, Armenia, 1996.
- [2] Chunxuan Ye, Raymond W. Yeung, On Fix-Free Codes. In *Proc. ISIT 2000*, p. 426, Sorrento, Italy, June 25-30, 2000.

¹This work was supported by the Russian Fond for Fundamental Researches (project 01-01-00495)