

PERMUTATIONS AVOIDING A PAIR OF PATTERNS OF TYPE (1, 2) OR (2, 1)

ANDERS CLAESSION AND TOUFIK MANSOUR

ABSTRACT. In [1] Babson and Steingrímsson introduced generalized permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. Claesson [2] presented a complete solution for the number of permutations avoiding any single (generalized) pattern of type (1, 2) or (2, 1). For eight of these twelve patterns the answer is given by the Bell numbers. For the remaining four the answer is given by the Catalan numbers.

In the present paper we give a complete solution for the number of permutations avoiding a pair of patterns of type (1, 2) or (2, 1). We also conjecture the number of permutations avoiding the patterns in any set of three or more such patterns.

1. INTRODUCTION

Classically, a pattern is a permutation $\sigma \in \mathcal{S}_k$, and a permutation $\pi \in \mathcal{S}_n$ avoids σ if there is no subword of π that is order equivalent to σ . For example, $\pi \in \mathcal{S}_n$ avoids 132 if there is no $1 \leq i < j < k \leq n$ such that $\pi(i) < \pi(k) < \pi(j)$. We denote by $\mathcal{S}_n(\sigma)$ the set permutations in \mathcal{S}_n that avoids σ .

The first case to be examined was the case of permutations avoiding one pattern of length 3. Knuth [6] found that, for any $\tau \in \mathcal{S}_3$, $|\mathcal{S}_n(\tau)| = C_n$, where $C_n = \frac{1}{n+1} \binom{2n}{n}$ is the n th Catalan number. Later Simion and Schmidt [7] found the cardinality of $\mathcal{S}_n(P)$ for all $P \subseteq \mathcal{S}_3$.

In [1] Babson and Steingrímsson introduced generalized permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. The motivation for Babson and Steingrímsson in introducing these patterns was the study of Mahonian statistics. Two examples of generalized patterns are 1-32 and 13-2 (1-32 and 13-2 are of type (1, 2) and (2, 1) respectively). A permutation $\pi = a_1 a_2 \cdots a_n$ avoids 1-32 if there are no subwords $a_i a_j a_{j+1}$ of π such that $a_i < a_{j+1} < a_j$. Similarly π avoids 13-2 if there are no subwords $a_i a_{i+1} a_j$ of π such that $a_i < a_j < a_{i+1}$.

Claesson [2] presented a complete solution for the number of permutations avoiding any single (generalized) pattern of type (1, 2) or (2, 1) as follows.

Proposition 1 (Claesson [2]). *Let $n \in \mathbb{N}$. We have*

$$|\mathcal{S}_n(p)| = \begin{cases} B_n & \text{if } p \in \{1-23, 3-21, 12-3, 32-1, 1-32, 3-12, 21-3, 23-1\}, \\ C_n & \text{if } p \in \{2-13, 2-31, 13-2, 31-2\}, \end{cases}$$

where B_n and C_n are the n th Bell and Catalan numbers, respectively.

In addition, Claesson gave some results for the number of permutations avoiding a pair of patterns.

Proposition 2 (Claesson [2]). *Let $n \in \mathbb{N}$. We have*

$$\mathcal{S}_n(1-23, 12-3) = B_n^*, \quad \mathcal{S}_n(1-23, 1-32) = I_n, \quad \text{and} \quad \mathcal{S}_n(1-23, 13-2) = M_n,$$

where B_n^* is the n th Bessel number ($\#$ non-overlapping partitions of $[n]$ (see [4])), I_n is the number of involutions in \mathcal{S}_n , and M_n is the n th Motzkin number.

This paper is organized as follows. In Section 2 we define the notion of a pattern and some other useful concepts. For a proof of Proposition 1 we could refer the reader to [2]. We will however prove Proposition 1 in Section 3 in the context of binary trees. The idea being that this will be a useful aid to understanding of the proofs of Section 4. In Section 4 we give a complete solution for the number of permutations avoiding a pair of patterns of type $(1, 2)$ or $(2, 1)$. These results are summarized in the following table.

# pairs	$ \mathcal{S}_n(p, q) $	
2	$0, n > 5$	Here
2	$2(n-1)$	
4	$\binom{n}{2} + 1$	$\sum_{n \geq 0} a_n x^n = \frac{1}{1 - x - x^2 \sum_{n \geq 0} B_n^* x^n}$
34	2^{n-1}	
8	M_n	and
2	a_n	$b_{n+2} = b_{n+1} + \sum_{k=0}^n \binom{n}{k} b_k.$
4	b_n	
4	I_n	
4	C_n	
2	B_n^*	

Finally, in Section 5 we conjecture the sequences $|\mathcal{S}_n(P)|$ for sets P of three or more patterns of type $(1, 2)$ or $(2, 1)$.

2. PRELIMINARIES

By an *alphabet* X we mean a non-empty set. An element of X is called a *letter*. A *word* over X is a finite sequence of letters from X . We consider also the *empty word*, that is, the word with no letters; it is denoted by ϵ . Let $w = x_1 x_2 \cdots x_n$ be a word over X . We call $|w| := n$ the *length* of w . A *subword* of w is a word $v = x_{i_1} x_{i_2} \cdots x_{i_k}$, where $1 \leq i_1 < i_2 < \cdots < i_k \leq n$.

Let $[n] := \{1, 2, \dots, n\}$ (so $[0] = \emptyset$). A *permutation* of $[n]$ is bijection from $[n]$ to $[n]$. Let \mathcal{S}_n be the set of permutations of $[n]$, and $\mathcal{S} = \cup_{n \geq 0} \mathcal{S}_n$. We shall usually think of a permutation π as the word $\pi(1)\pi(2)\cdots\pi(n)$ over the alphabet $[n]$.

Define the *reverse* of π as $\pi^r(i) = \pi(n+1-i)$, and define the *complement* of π by $\pi^c(i) = n+1-\pi(i)$, where $i \in [n]$.

For each word $w = x_1 x_2 \cdots x_n$ over the alphabet $\{1, 2, 3, 4, \dots\}$ without repeated letters, we define the *projection* of w onto \mathcal{S}_n , which we denote $\text{proj}(w)$, by

$$\text{proj}(w) = a_1 a_2 \cdots a_n, \quad \text{where } a_i = |\{j \in [n] : x_j \leq x_i\}|.$$

Equivalently, $\text{proj}(w)$ is the permutation in \mathcal{S}_n which is order equivalent to w . For example, $\text{proj}(2659) = 1324$.

We may regard a *pattern* as a function from \mathcal{S}_n to the set \mathbb{N} of natural numbers. The patterns of main interest to us are defined as follows. Let $xyz \in \mathcal{S}_3$ and $\pi = a_1 a_2 \cdots a_n \in \mathcal{S}_n$, then

$$(x-yz)\pi = |\{a_i a_j a_{j+1} : \text{proj}(a_i a_j a_{j+1}) = xyz, 1 \leq i < j < n\}|$$

and similarly $(xy-z)\pi = (z-yx)\pi^r$. For instance

$$(1-23)491273865 = |\{127, 138, 238\}| = 3.$$

A pattern $p = p_1 - p_2 - \cdots - p_k$ containing exactly $k-1$ dashes is said to be of type $(|p_1|, |p_2|, \dots, |p_k|)$. For example, the pattern $142-5-367$ is of type $(3, 1, 3)$, and any classical pattern of length k is of type $(\underbrace{1, 1, \dots, 1}_{k-1})$.

We say that a permutation π *avoids* a pattern p if $p\pi = 0$. The set of all permutations in \mathcal{S}_n that avoids p is denoted $\mathcal{S}_n(p)$ and, more generally, $\mathcal{S}_n(P) = \bigcap_{p \in P} \mathcal{S}_n(p)$ and $\mathcal{S}(P) = \bigcup_{n \geq 0} \mathcal{S}_n(P)$.

We extend the definition of reverse and complement to patterns the following way. Let us call π the *underlying permutation* of the pattern p if π is obtained from p by deleting all the dashes in p . If p is a pattern with underlying permutation π , then p^c is the pattern with underlying permutation π^c and with dashes at precisely the same positions as there are dashes in p . We define p^r as the pattern we get from regarding p as a word and reading it backwards. For example, $(1-23)^c = 3-21$ and $(1-23)^r = 32-1$. Observe that

$$\begin{aligned} \sigma \in \mathcal{S}_n(p) &\iff \sigma^r \in \mathcal{S}_n(p^r) \\ \sigma \in \mathcal{S}_n(p) &\iff \sigma^c \in \mathcal{S}_n(p^c). \end{aligned}$$

If p is a classical pattern, that is, of type $(1, 1, \dots, 1)$, then we also have

$$\sigma \in \mathcal{S}_n(p) \iff \sigma^{-1} \in \mathcal{S}_n(p^{-1}).$$

These observations of course generalize to $\mathcal{S}_n(P)$ for any set of patterns P .

The operations reverse, complement, and inverse generates the dihedral group D_4 (the symmetry group of a square). The operations reverse and complement alone generates the group D_2 (the symmetry group of a rectangle). Since we exclusively will be concerned with patterns of type $(1, 2)$ or $(2, 1)$ the appropriate symmetry group for us will be D_2 . The orbits of D_2 in the set of patterns of type $(1, 2)$ or $(2, 1)$ will be called *symmetry classes*. For instance, the symmetry class of 1-23 is

$$\{1-23, 3-21, 12-3, 32-1\}.$$

We also talk about symmetry classes of sets of patterns (defined in the obvious way). For example, the symmetry class of $\{1-23, 3-21\}$ is $\{\{1-23, 3-21\}, \{32-1, 12-3\}\}$.

A set of patterns P such that if $p, p' \in P$ then, for each n , $|\mathcal{S}_n(p)| = |\mathcal{S}_n(p')|$ is called a *Wilf-class*. For instance, by Proposition 1, the Wilf-class of 1-23 is

$$\{1-23, 3-21, 12-3, 32-1, 1-32, 3-12, 21-3, 23-1\}.$$

We also talk about Wilf-classes of sets of patterns (defined in the obvious way). It is clear that symmetry classes are Wilf-classes, but as we have seen the converse does not hold in general.

In what follows we will frequently use the following well known bijection between increasing binary trees and permutations (see [8, p. 24]). Let π be any word on the alphabet $\{1, 2, 3, 4, \dots\}$ with no repeated letters. If $\pi \neq \epsilon$ then we can factor π as $\pi = \sigma \tilde{1} \tau$, where $\tilde{1}$ is the minimal element of π . Define $T(\epsilon) = \bullet$ (a leaf) and

$$T(\pi) = \begin{array}{c} \tilde{1} \\ \swarrow \quad \searrow \\ T(\sigma) \quad T(\tau) \end{array}$$

In addition, we define $U(t)$ as the unlabelled counterpart of the labelled tree t . For instance

$$T(316452) = \begin{array}{c} 1 \\ \swarrow \quad \searrow \\ 3 \quad 2 \\ \swarrow \quad \searrow \\ 4 \\ \swarrow \quad \searrow \\ 6 \quad 5 \end{array} \quad U \circ T(316452) = \begin{array}{c} \circ \\ \swarrow \quad \searrow \\ \circ \quad \circ \\ \swarrow \quad \searrow \\ \circ \\ \swarrow \quad \searrow \\ \circ \quad \circ \end{array}$$

Note that we, for ease of presentation, do not display the leafs (\bullet).

3. SINGLE PATTERNS

There are 3 symmetry classes and 2 Wilf-classes of single patterns. The details are as follows.

Proposition 3 (Claesson [2]). *Let $n \in \mathbb{N}$. We have*

$$|\mathcal{S}(p)| = \begin{cases} B_n & \text{if } p \in \{1-23, 3-21, 12-3, 32-1\}, \\ B_n & \text{if } p \in \{1-32, 3-12, 21-3, 23-1\}, \\ C_n & \text{if } p \in \{2-13, 2-31, 13-2, 31-2\}, \end{cases}$$

where B_n and C_n are the n th Bell and Catalan numbers, respectively.


Proof of the first case. Note that

$$\sigma 1\tau \in \mathcal{S}(1-23) \iff \begin{cases} \text{proj}(\sigma) \in \mathcal{S}(1-23) \\ \text{proj}(\tau) \in \mathcal{S}(12) \\ \sigma 1\tau \in \mathcal{S} \end{cases}$$

where of course $\mathcal{S}(12) = \{\epsilon, 1, 21, 321, 4321, \dots\}$. This enable us to give a bijection Φ between $\mathcal{S}_n(1-23)$ and the set of partitions of $[n]$, by induction. Let the elements of 1τ form the first block of $\Phi(\sigma 1\tau)$ and let the rest of the blocks be as in $\Phi(\sigma)$. \square

The most transparent way to see the above correspondence is perhaps to view the permutation as an increasing binary tree.

Example 4. The tree




corresponds to the partition $\{\{1, 3, 8\}, \{2\}, \{4, 5, 7, 9\}, \{6\}\}$.

Proof of the second case. This case is analogous to the previous one. We have

$$\sigma 1\tau \in \mathcal{S}(1-32) \iff \begin{cases} \text{proj}(\sigma) \in \mathcal{S}(1-32) \\ \text{proj}(\tau) \in \mathcal{S}(21) \\ \sigma 1\tau \in \mathcal{S} \end{cases}$$

We give a bijection Φ between $\mathcal{S}_n(1-23)$ and the set of partitions of $[n]$, by induction. Let the elements of 1τ form the first block of $\Phi(\sigma 1\tau)$ and let the rest of the blocks be as in $\Phi(\sigma)$. \square

Example 5. The tree



corresponds to the partition $\{\{1, 3, 8\}, \{2\}, \{4, 5, 7, 9\}, \{6\}\}$.

Now that we have seen the structure of $\mathcal{S}(1-23)$ and $\mathcal{S}(1-32)$, it is trivial to give a bijection between the two sets. Indeed, if $\Theta : \mathcal{S}(1-23) \rightarrow \mathcal{S}(1-32)$ is given by $\Theta(\epsilon) = \epsilon$ and $\Theta(\sigma 1\tau) = \Theta(\sigma) 1\tau^r$ then Θ is such a bijection. Actually Θ is its own inverse.

Proof of the third case. It is plain that a permutation avoids 2-13 if and only if it avoids 2-1-3 (see [2]). Note that

$$\sigma 1\tau \in \mathcal{S}(2-1-3) \iff \begin{cases} \text{proj}(\sigma), \text{proj}(\tau) \in \mathcal{S}(2-1-3) \\ \tau > \sigma \\ \sigma 1\tau \in \mathcal{S} \end{cases}$$

where $\tau > \sigma$ means that any letter of τ is greater than any letter of σ . Hence we get a unique labelling of the binary tree corresponding to $\sigma 1\tau$, that is, if $\pi_1, \pi_2 \in \mathcal{S}(2-1-3)$ and $U \circ T(\pi_1) = U \circ T(\pi_2)$ then $\pi_1 = \pi_2$. It is well known that there are exactly C_n (unlabelled) binary trees with n (internal) nodes. The validity of the last statement is for example seen from the following simple bijection between Dyck words and binary trees. Fixing notation, we let the set of Dyck words be the smallest set of words over $\{u, d\}$ that contains the empty word and is closed under $(\alpha, \beta) \mapsto u\alpha d\beta$. Now the promised bijection is given by $\Psi(\bullet) = \epsilon$ and

$$\Psi\left(\begin{array}{c} \circ \\ / \quad \backslash \\ L \quad R \end{array}\right) = u\Psi(L)d\Psi(R).$$

□

4. PAIRS OF PATTERNS

There are $\binom{12}{2} = 66$ pairs of patterns altogether. It turns out that there are 21 symmetry classes and 10 Wilf-classes. The details are as follows.

4.1. The Wilf-class corresponding to $\{0\}_n$.

Proposition 6. *Let $n \in \mathbb{N}$ with $n > 5$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 32-1\}, \\ \{3-21, 12-3\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 0.$$

Proof. We have

$$\sigma 1\tau \in \mathcal{S}(1-23, 32-1) \iff \begin{cases} \text{proj}(\sigma) \in \mathcal{S}(21, 1-23) \\ \text{proj}(\tau) \in \mathcal{S}(12, 32-1) \\ \sigma 1\tau \in \mathcal{S} \end{cases}$$

The result now follows from $\mathcal{S}(21, 1-23) = \{\epsilon, 1, 12\}$ and $\mathcal{S}(12, 32-1) = \{\epsilon, 1, 21\}$.

□

4.2. The Wilf-class corresponding to $\{2(n-1)\}_n$.


Proposition 7. *Let $n \in \mathbb{N}$ with $n > 1$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 3-21\}, \\ \{32-1, 12-3\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2(n-1).$$

Proof. Since 3-21 is the complement of 1-23, the number of permutations in $\mathcal{S}_n(1-23, 3-21)$ is twice the number of permutations in $\mathcal{S}_n(1-23, 3-21)$ in which 1 precedes n . In addition, 1 and n must be adjacent letters in a permutation avoiding 1-23 and 3-21. Let $\sigma 1n\tau$ be such a permutation. Note that τ must be both increasing and decreasing, that is, $\tau \in \{\epsilon, 2, 3, 4, \dots, n-1\}$, so there are $n-1$ choices for τ . Furthermore, there is exactly one permutation in $\mathcal{S}_n(1-23, 3-21)$ of the form $\sigma 1n$, namely $(\lceil \frac{n+1}{2} \rceil, \dots, n-2, 3, n-1, 2, n, 1)$, and similarly there is exactly one of the form $\sigma 1nk$ for each $k \in \{2, 3, \dots, n-1\}$. This completes our argument. □

Hence a permutation in $\mathcal{S}(1-23, 23-1)$ is given by the following procedure. Choose a subset $S \subseteq \{2, 3, 4, \dots, n\}$, let σ be the word obtained by writing the elements of S in decreasing order, and let τ be the word obtained by writing the elements of $\{2, 3, 4, \dots, n\} \setminus S$ in decreasing order. \square

Example 12. The tree



corresponds to the subset $\{2, 4\}$ of $\{2, 3, 4, 5, 6\}$.

Proposition 13. Let $n \in \mathbb{N}$ with $n > 0$.

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 31-2\}, \\ \{3-21, 13-2\}, \\ \{12-3, 2-31\}, \\ \{32-1, 2-13\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. This case is essentially identical to the case dealt with in Proposition 9. \square

Proposition 14. Let $n \in \mathbb{N}$ with $n > 0$.

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 2-13\}, \\ \{3-12, 2-31\}, \\ \{13-2, 21-3\}, \\ \{23-1, 31-2\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. The bijection Θ between $\mathcal{S}(1-23)$ and $\mathcal{S}(1-32)$ (see page 3) provides a one-to-one correspondence between $\mathcal{S}_n(1-32, 2-13)$ and $\mathcal{S}_n(1-23, 2-13)$. Consequently the result follows from Proposition 9. \square

Proposition 15. Let $n \in \mathbb{N}$ with $n > 0$.

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 2-31\}, \\ \{3-12, 2-13\}, \\ \{31-2, 21-3\}, \\ \{23-1, 13-2\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. We have


$$\sigma 1 \tau \in \mathcal{S}(3-12, 2-13) \iff \begin{cases} \text{proj}(\sigma), \text{proj}(\tau) \in \mathcal{S}(3-12, 2-13) \\ \sigma = \epsilon \text{ or } \tau = \epsilon \\ \sigma 1 \tau \in \mathcal{S} \end{cases}$$

Thus a bijection between $\mathcal{S}_n(3-12, 2-13)$ and $\{0, 1\}^{n-1}$ is given by $\Psi(\epsilon) = \epsilon$ and

$$\Psi(\sigma 1 \tau) = x \Psi(\sigma \tau) \text{ where } x = \begin{cases} 1 & \text{if } \sigma \neq \epsilon, \\ 0 & \text{if } \tau \neq \epsilon, \\ \epsilon & \text{otherwise.} \end{cases}$$

\square

Example 16. The tree



corresponds to $01011 \in \{0,1\}^5$.

Proposition 17. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 3-12\}, \\ \{23-1, 21-3\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. Since 3-12 is the complement of 1-32, the number of permutations in $\mathcal{S}_n(1-32, 3-12)$ is twice the number of permutations in $\mathcal{S}_n(1-32, 3-12)$ in which 1 precedes n . In addition, n must be the last letter in such a permutation or else a hit of 1-32 would be formed. We have

$$\begin{aligned} \sigma 1 \tau n \in \mathcal{S}(1-32, 3-12) &\iff \begin{cases} \text{proj}(\sigma 1 \tau) \in \mathcal{S}(1-32, 3-12) \\ \text{proj}(\tau) \in \mathcal{S}(21) \\ \sigma 1 \tau \in \mathcal{S} \end{cases} \\ &\iff \begin{cases} \text{proj}(\sigma) \in \mathcal{S}(1-32, 3-12) \\ \text{proj}(\tau) \in \mathcal{S}(21) \\ \sigma < \tau \\ \sigma 1 \tau \in \mathcal{S} \end{cases} \end{aligned}$$

The rest of the proof follows the same lines as the proof of Proposition 9. \square

Proposition 18. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 23-1\}, \\ \{3-12, 21-3\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. We can copy almost verbatim the proof of Proposition 15, indeed, it is not hard to see that $\mathcal{S}_n(1-32, 23-1) = \mathcal{S}_n(1-32, 2-31)$. \square

Proposition 19. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 31-2\}, \\ \{3-12, 13-2\}, \\ \{21-3, 2-31\}, \\ \{23-1, 2-13\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. We can copy almost verbatim the proof of Proposition 17, indeed, it is not hard to see that $\mathcal{S}_n(1-32, 31-2) = \mathcal{S}_n(1-32, 3-12)$. \square

Proposition 20. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{2-13, 2-31\}, \\ \{31-2, 13-2\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. $|\mathcal{S}_n(2-13, 2-31)| = |\mathcal{S}_n(2-1-3, 2-3-1)| = 2^{n-1}$. (See [7, Lemma 5(d)].) \square

Proposition 21. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{2-13, 13-2\}, \\ \{2-31, 31-2\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. $|\mathcal{S}_n(2-13, 13-2)| = |\mathcal{S}_n(1-3-2, 2-1-3)| = 2^{n-1}$. (See [7, Lemma 5(b)].) \square

Proposition 22. *Let $n \in \mathbb{N}$ with $n > 0$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{2-13, 31-2\}, \\ \{2-31, 13-2\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = 2^{n-1}.$$

Proof. $|\mathcal{S}_n(2-13, 31-2)| = |\mathcal{S}_n(2-1-3, 3-1-2)| = 2^{n-1}$. (See [7, Lemma 5(c)].) \square

4.5. The Wilf-class corresponding to $\{M_n\}_n$.

Proposition 23. Let $n \in \mathbb{N}$.

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 13-2\}, \\ \{3-21, 31-2\}, \\ \{12-3, 2-13\}, \\ \{32-1, 2-31\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = M_n,$$

where M_n is the n th Motzkin number.

Proof. See Proposition 2. □

Proposition 24. Let $n \in \mathbb{N}$.

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 21-3\}, \\ \{3-21, 23-1\}, \\ \{12-3, 1-32\}, \\ \{32-1, 3-12\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = M_n,$$

where M_n is the n th Motzkin number.

Proof. We give a bijection $\Lambda : \mathcal{S}_n(1-23, 21-3) \rightarrow \mathcal{S}_n(1-23, 13-2)$ by means of induction. Let $\pi \in \mathcal{S}_n(1-23, 21-3)$. Define $\Lambda(\pi) = \pi$ for $n \leq 1$. Assume $n \geq 2$ and $\pi = a_1 a_2 \cdots a_n$. It is plain that either $a_1 = n$ or $a_2 = n$, so we can define

$$\Lambda(\pi) = \begin{cases} (a'_1 + 1, \dots, a'_{n-1} + 1, a'_{n-2} + 1, 1) & \text{if } \begin{cases} a_1 = n & \& \\ a'_1 \cdots a'_{n-1} = \Lambda(a_2 a_3 a_4 \cdots a_n), \end{cases} \\ (a'_1 + 1, \dots, a'_{n-1} + 1, 1, a'_{n-2} + 1) & \text{if } \begin{cases} a_2 = n & \& \\ a'_1 \cdots a'_{n-1} = \Lambda(a_1 a_3 a_4 \cdots a_n). \end{cases} \end{cases}$$

Observing that if $\sigma \in \mathcal{S}_n(1-23, 13-2)$ then $\sigma(n-1) = 1$ or $\sigma(n) = 1$, it easy to find the inverse of Λ . □

4.6. The Wilf-class corresponding to $\{1, 1, 2, 4, 9, 22, 58, 164, 496, 1601, \dots\}$.

In [2] Claesson introduced the notion of a monotone partition. A partition is *monotone* if its non-singleton blocks can be written in increasing order of their least element and increasing order of their greatest element, simultaneously. He then proved that monotone partitions and non-overlapping partitions are in one-to-one correspondence. Non-overlapping partitions were first studied by Flajolet and Schot in [4]. A partition π is *non-overlapping* if for no two blocks A and B of π we have $\min A < \min B < \max A < \max B$. Let B_n^* be the number of non-overlapping partitions of $[n]$; this number is called the n th *Bessel number*. Proposition 2 tells us that there is a bijection between non-overlapping partitions and permutations avoiding 1-23 and 12-3. Below we define a new class of partitions called strongly monotone partitions and then show that there is a bijection between strongly monotone partitions and permutations avoiding 1-32 and 21-3.

Definition 25. Let π be an arbitrary partition whose blocks $\{A_1, \dots, A_k\}$ are ordered so that for all $i \in [k-1]$, $\min A_i > \min A_{i+1}$. If $\max A_i > \max A_{i+1}$ for all $i \in [k-1]$, then we call π a *strongly monotone partition*.

In other words a partition is strongly monotone if its blocks can be written in increasing order of their least element and increasing order of their greatest element, simultaneously. Let us denote by a_n the number of strongly monotone partitions of $[n]$. The sequence $\{a_n\}_0^\infty$ starts with

$$1, 1, 2, 4, 9, 22, 58, 164, 496, 1601, 5502, 20075, 77531, 315947, 1354279.$$

It is routine to derive the continued fraction expansion

$$\sum_{n \geq 0} a_n x^n = \frac{1}{1 - 1 \cdot x - \frac{x^2}{1 - 1 \cdot x - \frac{x^2}{1 - 2 \cdot x - \frac{x^2}{1 - 3 \cdot x - \frac{x^2}{1 - 4 \cdot x - \frac{x^2}{\ddots}}}}}}$$

using the standard machinery of Flajolet [3] and Françon and Viennot [5]. One can also note that there is a one-to-one correspondence between strongly monotone partitions and non-overlapping partition, π , such that if $\{x\}$ and B are blocks of π then either $x < \min B$ or $\max B < x$. In addition, we observe that

$$\sum_{n \geq 0} a_n x^n = \frac{1}{1 - x - x^2 B^*(x)},$$

where $B^*(x) = \sum_{n \geq 0} B_n^* x^n$ is the ordinary generating function for the Bessel numbers.

Proposition 26. *Let $n \in \mathbb{N}$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1\text{-}32, 21\text{-}3\}, \\ \{3\text{-}12, 23\text{-}1\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = a_n,$$

where a_n is the number of strongly monotone partitions of $[n]$ (see Definition 25).

Proof. Suppose $\pi \in \mathcal{S}_n$ has $k+1$ left-to-right minima $1, 1', 1'', \dots, 1^{(k)}$ such that

$$1 < 1' < 1'' < \dots < 1^{(k)}, \text{ and } \pi = 1^{(k)} \tau^{(k)} \dots 1' \tau' 1 \tau.$$

Then π avoids 1-32 if and only if, for each i , $\tau^{(i)} \in \mathcal{S}(21)$. If π avoids 1-32 and $x_i = \max 1^{(i)} \tau^{(i)}$ then π avoids 21-3 precisely when $x_0 < x_1 < \dots < x_k$. This follows from observing that the only potential (21-3)-subwords of π are $x_{i+1} 1^{(k)} x_j$ with $j \leq i$.

Mapping π to the partition $\{1\sigma, 1'\sigma', \dots, 1^{(k)} \tau^{(k)}\}$ we thus get a one-to-one correspondence between permutations in $\mathcal{S}_n(1\text{-}32, 21\text{-}3)$ and strongly monotone partitions of $[n]$. \square

4.7. The Wilf-class corresponding to $\{1, 1, 2, 4, 9, 23, 65, 199, 654, 2296, \dots\}$.

Proposition 27. *Let $n \in \mathbb{N}$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1\text{-}23, 3\text{-}12\}, \\ \{3\text{-}21, 1\text{-}32\}, \\ \{23\text{-}1, 12\text{-}3\}, \\ \{32\text{-}1, 21\text{-}3\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = b_n,$$

where the sequence $\{b_n\}$ satisfies $b_0 = 1$ and, for $n \geq -2$,

$$b_{n+2} = b_{n+1} + \sum_{k=0}^n \binom{n}{k} b_k.$$

Proof. Suppose $\pi \in \mathcal{S}_n$ has $k+1$ left-to-right minima $1, 1', 1'', \dots, 1^{(k)}$ such that

$$1 < 1' < 1'' < \dots < 1^{(k)}, \text{ and } \pi = 1^{(k)} \tau^{(k)} \dots 1' \tau' 1 \tau.$$

Then π avoids 1-23 if and only if, for each i , $\tau^{(i)} \in \mathcal{S}(12)$. If π avoids 1-23 and $x_i = \max 1^{(i)} \tau^{(i)}$ then π avoids 3-12 precisely when

$$j > i \ \& \ x_i \neq 1^{(i)} \implies x_j < x_i.$$

This follows from observing that the only potential (3-12)-subwords of π are $x_j 1^{(k)} x_i$ with $j \leq i$. Thus we have established

$$\sigma 1\tau \in \mathcal{S}_n(1-23, 3-12) \iff \begin{cases} \text{proj}(\sigma) \in \mathcal{S}(1-23, 3-12) \\ \tau \neq \epsilon \Rightarrow \tau = \tau'n \ \& \ \text{proj}(\tau') \in \mathcal{S}(12) \\ \sigma 1\tau \in \mathcal{S}_n \end{cases}$$

If we know that $\sigma 1\tau'n \in \mathcal{S}_n(1-23, 3-12)$ and $\text{proj}(\tau') \in \mathcal{S}_k(12)$ then there are $\binom{n-2}{k}$ candidates for τ' . In this way the recursion follows. \square

4.8. The Wilf-class corresponding to I_n .

Proposition 28. *Let $n \in \mathbb{N}$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 1-32\}, \\ \{3-21, 3-12\}, \\ \{21-3, 12-3\}, \\ \{32-1, 23-1\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = I_n,$$

where I_n is the number of involutions in \mathcal{S}_n .

Proof. See Proposition 2. \square

4.9. The Wilf-class corresponding to C_n .

Proposition 29. *Let $n \in \mathbb{N}$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-32, 13-2\}, \\ \{3-12, 31-2\}, \\ \{21-3, 2-13\}, \\ \{23-1, 2-31\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = C_n,$$

where C_n is the n th Catalan number.

Proof. $\mathcal{S}_n(1-32, 13-2) = \mathcal{S}_n(1-3-2)$. \square

4.10. The Wilf-class corresponding to B_n^* .

Proposition 30. *Let $n \in \mathbb{N}$.*

$$\text{If } \{p, q\} \in \left\{ \begin{array}{l} \{1-23, 12-3\}, \\ \{3-21, 32-1\} \end{array} \right\} \text{ then } |\mathcal{S}_n(p, q)| = B_n^*,$$

where B_n^* is the n th Bessel number.

Proof. See Proposition 2. \square

5. MORE THAN TWO PATTERNS

Let P be a set of patterns of type (1, 2) or (2, 1). With the aid of a computer we have calculated the cardinality of $\mathcal{S}_n(P)$ for sets P of three or more patterns. From these results we arrived at the plausible conjectures of table 1 (some of which are trivially true). We use the notation $m \times n$ to express that there are m symmetric classes each of which contains n sets. Moreover, we denote by F_n the n th Fibonacci number ($F_0 = F_1 = 1, F_{n+1} = F_n + F_{n-1}$).

For $ P = 3$ there are 220 sets, 55 symmetry classes and 9 Wilf-classes.	For $ P = 4$ there are 495 sets, 135 symmetry classes, and 9 Wilf-classes.																																								
<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>7×4</td> </tr> <tr> <td>3</td> <td>1×4</td> </tr> <tr> <td>n</td> <td>24×4</td> </tr> <tr> <td>$1 + \binom{n}{2}$</td> <td>2×4</td> </tr> <tr> <td>F_n</td> <td>7×4</td> </tr> <tr> <td>$\binom{n}{\lfloor n/2 \rfloor}$</td> <td>$1 \times 4$</td> </tr> <tr> <td>$2^{n-2} + 1$</td> <td>$1 \times 4$</td> </tr> <tr> <td>$2^{n-1}$</td> <td>$10 \times 4$</td> </tr> <tr> <td>$M_n$</td> <td>$2 \times 4$</td> </tr> </tbody> </table>	cardinality	# sets	0	7×4	3	1×4	n	24×4	$1 + \binom{n}{2}$	2×4	F_n	7×4	$\binom{n}{\lfloor n/2 \rfloor}$	1×4	$2^{n-2} + 1$	1×4	2^{n-1}	10×4	M_n	2×4	<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>$1 \times 1 + 6 \times 2 + 30 \times 4$</td> </tr> <tr> <td>2</td> <td>$2 \times 1 + 5 \times 2 + 35 \times 4$</td> </tr> <tr> <td>3</td> <td>1×4</td> </tr> <tr> <td>n</td> <td>$37 \times 4 + 1 \times 2$</td> </tr> <tr> <td>$1 + \binom{n}{2}$</td> <td>1×4</td> </tr> <tr> <td>F_n</td> <td>$9 \times 4 + 1 \times 2$</td> </tr> <tr> <td>$\binom{n}{\lfloor n/2 \rfloor}$</td> <td>$1 \times 2$</td> </tr> <tr> <td>$2^{n-2} + 1$</td> <td>$1 \times 2$</td> </tr> <tr> <td>$2^{n-1}$</td> <td>$1 \times 4 + 3 \times 2$</td> </tr> </tbody> </table>	cardinality	# sets	0	$1 \times 1 + 6 \times 2 + 30 \times 4$	2	$2 \times 1 + 5 \times 2 + 35 \times 4$	3	1×4	n	$37 \times 4 + 1 \times 2$	$1 + \binom{n}{2}$	1×4	F_n	$9 \times 4 + 1 \times 2$	$\binom{n}{\lfloor n/2 \rfloor}$	1×2	$2^{n-2} + 1$	1×2	2^{n-1}	$1 \times 4 + 3 \times 2$
cardinality	# sets																																								
0	7×4																																								
3	1×4																																								
n	24×4																																								
$1 + \binom{n}{2}$	2×4																																								
F_n	7×4																																								
$\binom{n}{\lfloor n/2 \rfloor}$	1×4																																								
$2^{n-2} + 1$	1×4																																								
2^{n-1}	10×4																																								
M_n	2×4																																								
cardinality	# sets																																								
0	$1 \times 1 + 6 \times 2 + 30 \times 4$																																								
2	$2 \times 1 + 5 \times 2 + 35 \times 4$																																								
3	1×4																																								
n	$37 \times 4 + 1 \times 2$																																								
$1 + \binom{n}{2}$	1×4																																								
F_n	$9 \times 4 + 1 \times 2$																																								
$\binom{n}{\lfloor n/2 \rfloor}$	1×2																																								
$2^{n-2} + 1$	1×2																																								
2^{n-1}	$1 \times 4 + 3 \times 2$																																								
For $ P = 5$ there are 792 sets, 198 symmetry classes, and 5 Wilf-classes.	For $ P = 6$ there are 924 sets, 246 symmetry classes, and 4 Wilf-classes.																																								
<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>84×4</td> </tr> <tr> <td>1</td> <td>16×4</td> </tr> <tr> <td>2</td> <td>74×4</td> </tr> <tr> <td>n</td> <td>20×4</td> </tr> <tr> <td>F_n</td> <td>4×4</td> </tr> </tbody> </table>	cardinality	# sets	0	84×4	1	16×4	2	74×4	n	20×4	F_n	4×4	<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>$17 \times 2 + 124 \times 4$</td> </tr> <tr> <td>1</td> <td>$4 \times 2 + 38 \times 4$</td> </tr> <tr> <td>2</td> <td>$7 \times 2 + 51 \times 4$</td> </tr> <tr> <td>n</td> <td>$1 \times 2 + 3 \times 4$</td> </tr> <tr> <td>F_n</td> <td>1×2</td> </tr> </tbody> </table>	cardinality	# sets	0	$17 \times 2 + 124 \times 4$	1	$4 \times 2 + 38 \times 4$	2	$7 \times 2 + 51 \times 4$	n	$1 \times 2 + 3 \times 4$	F_n	1×2																
cardinality	# sets																																								
0	84×4																																								
1	16×4																																								
2	74×4																																								
n	20×4																																								
F_n	4×4																																								
cardinality	# sets																																								
0	$17 \times 2 + 124 \times 4$																																								
1	$4 \times 2 + 38 \times 4$																																								
2	$7 \times 2 + 51 \times 4$																																								
n	$1 \times 2 + 3 \times 4$																																								
F_n	1×2																																								
For $ P = 7$ there are 792 sets, 198 symmetry classes, and 3 Wilf-classes.	For $ P = 8$ there are 495 sets, 135 symmetry classes, and 3 Wilf-classes.																																								
<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>140×4</td> </tr> <tr> <td>1</td> <td>40×4</td> </tr> <tr> <td>2</td> <td>18×4</td> </tr> </tbody> </table>	cardinality	# sets	0	140×4	1	40×4	2	18×4	<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>$2 \times 1 + 14 \times 2 + 94 \times 4$</td> </tr> <tr> <td>1</td> <td>$4 \times 2 + 18 \times 4$</td> </tr> <tr> <td>2</td> <td>$1 \times 1 + 2 \times 4$</td> </tr> </tbody> </table>	cardinality	# sets	0	$2 \times 1 + 14 \times 2 + 94 \times 4$	1	$4 \times 2 + 18 \times 4$	2	$1 \times 1 + 2 \times 4$																								
cardinality	# sets																																								
0	140×4																																								
1	40×4																																								
2	18×4																																								
cardinality	# sets																																								
0	$2 \times 1 + 14 \times 2 + 94 \times 4$																																								
1	$4 \times 2 + 18 \times 4$																																								
2	$1 \times 1 + 2 \times 4$																																								
For $ P = 9$ there are 220 sets, 55 symmetry classes, and 2 Wilf-classes.	For $ P = 10$ there are 66 sets, 21 symmetry classes, and 2 Wilf-classes.																																								
<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>50×4</td> </tr> <tr> <td>1</td> <td>5×4</td> </tr> </tbody> </table>	cardinality	# sets	0	50×4	1	5×4	<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>$8 \times 2 + 12 \times 4$</td> </tr> <tr> <td>1</td> <td>1×2</td> </tr> </tbody> </table>	cardinality	# sets	0	$8 \times 2 + 12 \times 4$	1	1×2																												
cardinality	# sets																																								
0	50×4																																								
1	5×4																																								
cardinality	# sets																																								
0	$8 \times 2 + 12 \times 4$																																								
1	1×2																																								
For $ P = 11$ there are 12 sets, 3 symmetry classes, and 1 Wilf-class.	For $ P = 12$ there is 1 set, 1 symmetry class, and 1 Wilf-class.																																								
<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>3×4</td> </tr> </tbody> </table>	cardinality	# sets	0	3×4	<table border="1"> <thead> <tr> <th>cardinality</th> <th># sets</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>1×1</td> </tr> </tbody> </table>	cardinality	# sets	0	1×1																																
cardinality	# sets																																								
0	3×4																																								
cardinality	# sets																																								
0	1×1																																								

TABLE 1. The cardinality of $\mathcal{S}_n(P)$ for $|P| > 2$.

ACKNOWLEDGEMENTS

The first author wishes to express his gratitude towards Einar Steingrímsson, Kimmo Eriksson, and Mireille Bousquet-Mélou; Einar for his guidance and infectious enthusiasm; Kimmo for useful suggestions and a very constructive discussion on the results of this paper; Mireille for her great hospitality during a stay at LaBRI, where some of the work on this paper was done.

We would like to thank N. J. A. Sloane for his excellent web site “The On-Line Encyclopedia of Integer Sequences”

<http://www.research.att.com/~njas/sequences/>.

It is simply an indispensable tool for all studies concerned with integer sequences.

REFERENCES

- [1] E. Babson and E. Steingrímsson. Generalized permutation patterns and a classification of the Mahonian statistics. *Séminaire Lotharingien de Combinatoire*, B44b:18pp, 2000.
- [2] A. Claesson. Generalized pattern avoidance. *To appear in: European Journal of Combinatorics*, 2001.
- [3] P. Flajolet. Combinatorial aspects of continued fractions. *Annals of Discrete Mathematics*, 8:217–222, 1980.
- [4] P. Flajolet and R. Schott. Non-overlapping partitions, continued fractions, Bessel functions and a divergent series. *European Journal of Combinatorics*, 11:421–432, 1990.
- [5] J. Françon and G. Viennot. Permutations selon leurs pics, creux, doubles montées et double descentes, nombres d’Euler et nombres de Genocchi. *Discrete Math.*, 28(1):21–35, 1979.
- [6] D. E. Knuth. *The art of computer programming*, volume 3. Addison-Wesley, 1973.
- [7] R. Simion and F. W. Schmidt. Restricted permutations. *European Journal of Combinatorics*, 6:383–406, 1985.
- [8] R. P. Stanley. *Enumerative Combinatorics*, volume 1. Cambridge University Press, 1997.

MATEMATIK, CHALMERS TEKNISKA HÖGSKOLA OCH GÖTEBORGS UNIVERSITET, S-412 96 GÖTEBORG, SWEDEN

E-mail address: `claesson@math.chalmers.se`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF HAIFA, HAIFA, ISRAEL 31905

E-mail address: `tmansur@study.haifa.ac.il`