

Management of arid soils of the Badia region of Northeastern Jordan for sustainable agriculture

RAWAJFIH Zahir (1), **KHRESATS A.** (1), **BUCK B.** (2) and **MONGER H.C.** (3)

- (1) Department of Natural Resources and the Environment, Jordan University of Science and Technology, P.O.B. 3030, Irbid-22110, Jordan
- (2) Dept. of Geoscience, University of Nevada Las Vegas, Las Vegas NV 89154, USA
- (3) Dept. of Agronomy and Horticulture, New Mexico State University, LasCruces NM 88003, USA

Abstract

Arid lands have always been important to the world's human population, but their significance has increased over the past few decades because of population demographics and continued use of natural resources (Hoekstra and Shachak, 1999). Jordan is faced with increasing population pressure in its already settled areas; hence, it will have to utilize the underpopulated Badia so that it can contribute to the economy in a sustainable way without damaging the fragile desert environments. This can be done by gaining a better understanding of the natural resource base and suggesting actions which will lead to the protection of the resources and to their sustainable usage for the long-term benefit of the local population.

Azraq basin is an important part of the Jordanian Badia region. To develop soil resources in the Azraq basin, research is needed to establish the best practices for their management, improvement, and maintenance. Four different areas in the basin were investigated. The choice of these areas was based on geologic studies which showed the proximity of these areas to promising underground water resources. Most of the studied soils contain considerable amounts of carbonates. This leads to alkaline reaction of the soils with pH values mostly above 8, resulting in low availability of certain nutrients like phosphorus and micronutrients. If such soils are to be cultivated, the proper management practices should be adopted to ensure the availability of those nutrients to crops.

Analysis of the collected soil samples also showed that those soils contain soluble salts in the upper horizons in amounts enough to impede plant growth. Salts would have to be leached using good quality water before successful growing of crops can be carried out. Most of the studied soils have high erodibility and susceptibility to seal and crust formation. The soils in some of the studied areas contain high amounts of gypsum and carbonates and also very high content of soluble salts. The use of organic amendments would be highly recommended to improve the physical and chemical characteristics of the Badia soils in order to achieve a sustainable agricultural production.

Keywords: soil management, arid soils, Jordan

Introduction

From ancient times to the present, humans tried to utilize the arid lands of the world with only marginal success. People have been more successful in their attempts to utilize semiarid lands. However, heavy use of semiarid lands has often led to a reduction in available moisture for vegetation and has initiated succession from semiarid types of vegetation toward arid types of vegetation. Such human-caused succession of vegetation from semiarid to arid or from arid to extremely arid is currently referred to as *desertification* (Dick-Peddie, 1991). Le Houerou (1986) used the term “desertization” for this process in northern Africa and emphasized that the process is caused by human beings, not climate.

Physical properties that are suggested to be common to many arid and semiarid region soils include high erodibility, high runoff generation potential, susceptibility to seal and crust formation, poor water-holding capacity, profile hardening, structural instability, low clay content, low activity clay, high bulk density, and high surface temperatures in the summer months (El-Swaify *et al.*, 1984; Guthrie, 1982; Lal, 1985).

Soils developing in an arid environment tend to retain the alkaline earth and alkali cations to a great extent, and subsequently hydroxides of these cations form and alkaline pH ensues. The upward gradient of water movement, the retention of these cations by inorganic and organic colloids, and their reprecipitation as cationic constituents (like calcite, apatites, gypsum, and halides) promote their concentration (Knight, 1991).

Desert is the prevalent ecosystem in Jordan, covering over 80 percent of the country. Jordan is faced with increasing population pressure in its already settled areas; hence, it will have to utilize the underpopulated Badia so that it can contribute to the economy in a sustainable way without damaging the fragile desert environments. This can be done by gaining a better understanding of the natural resource base and suggesting actions which will lead to the protection of the resources and to their sustainable usage for the long-term benefit of the local population. Azraq basin is an important part of the Jordanian Badia region. To develop soil resources in the Azraq basin, research is needed to establish the best practices for their management, improvement, and maintenance. As a first step towards understanding the soil resource in the Azraq basin, the present work has been carried out.

Materials and Methods

Four different areas were investigated. A team studying the geology of the Azraq basin recommended several areas in the basin as potential development areas for agricultural production. Their recommendations were based on the probability of deep soils and the proximity of these areas to promising underground water resources. Profiles were dug and described in the field for physical characterization. Morphology and classification of the studied soils is reported in a separate paper (Khresat *et al.*, 2001). Soil samples were collected from each profile and taken to the laboratory for chemical and physical analysis. The bulk soil samples were air dried, crushed with a mortar and pestle, and sieved to remove coarse (>2 mm) fragments. Particle size distribution was determined by the hydrometer method (Gee and Bauder, 1986). Soil pH was measured on 1:1 soil: water suspensions (Thomas, 1996). Soluble salts were determined by measuring the electrical conductivity of 1:1 soil: water extracts (Rhoades, 1996); organic matter was determined using the Walkley-Black method (Nelson and Sommers, 1996); calcium carbonate equivalent values were obtained using

the acid neutralization method (Richards, 1954). Nitrogen was determined by the Kjeldahl method (Bremner, 1996). Cation exchange capacity was determined by the sodium saturation method (Chapman, 1965).

Results and Discussion

Physical properties

Particle size analysis for the studied profiles is shown in Table 1. The upper horizon of most of the profiles has more than 30% silt. Soil erodibility has been shown to be primarily a function of particle size distribution and various materials that bind particles together into aggregates and stabilize the aggregates. Wischmeier *et al.* (1971) developed a graphic solution to an equation for predicting soil erodibility in which increases in silt and very fine sand are predicted to increase soil erodibility. Other factors in the nomograph are soil organic matter, soil structure, and permeability of the least permeable horizon.

Table 1 Particle size analysis of the studied profiles.

Sample	% Sand	% Silt	% Clay	Classification
Muwaqqr III Ap (0-8)	30.3	30.3	30.6	Xeric Haplocalcids
Muwaqqr III B (8-20)	12.5	12.5	50.0	
Muwaqqr III Bk1 (20-40)	19.6	19.6	52.1	
Muwaqqr III Bk2 (40-120)	17.2	17.2	18.0	
Shawmari I Av (0-4)	20.8	20.8	36.6	Typic Calcigypsis
Shawmari I A (4-20)	14.6	14.6	36.9	
Shawmari I By (20-60)	15.2	15.2	53.5	
Shawmari I Bky (60-100)	10.7	10.7	61.8	
Shawmari II A (0-4)	21.6	35.4	43.0	Typic Calcigypsis
Shawmari II Bk (4-20)	21.4	16.1	62.5	
Shawmari II Bky (20-75)	31.9	16.7	16.7	
Swaeid II A (0-20)	22.0	31.0	47.0	Typic Calciargids
Swaeid II Bk (20-55)	14.0	29.0	57.0	
Swaeid II Btk (55-140)	14.0	26.0	60.0	
Shubayka II Ap (0-10)	1.3	40.3	58.4	Typic Haplogypsis
Shubayka II By1 (10-35)	2.6	35.1	62.3	
Shubayka II By2 (35-90)	1.0	34.0	65.0	
Shubayka II Bty (90-140)	1.0	37.0	61.0	

Water and wind erosion are both serious problems in these soils. Even though the region has very low rainfall, soils are subject to water erosion because they are bare and subject to the direct, although infrequent impact of raindrops. Wind erosion is a constant hazard for the reason that the dry, unprotected soil surface is easily moved by wind. Throughout the arid and semiarid climate zone, wind erosion is a real or potential problem. Throughout the Middle East, dust storms are a serious problem (Middleton, 1986).

Managing soil erosion in arid and semiarid areas requires careful water management, surface vegetation management, and control of wind speeds with wind breaks. Increasing surface roughness can decrease wind erosion as can the use of windbreaks to reduce the critical wind speed near the ground. Maintaining soil moisture

during critical parts of the year with mulches can also contribute to lower erosion rates. Gupta *et al.* (1983) have shown that a three-row windbreak consisting of *Cassia siamea*, *Albizia lebbek*, and *C. siamea*, when 4-5 m tall, was most effective in reducing wind velocity, wind erosion, and evaporative losses downwind from the shelterbelt.

The soils in this region would also tend to form slowly permeable seals and crusts that decrease infiltration and increase runoff, further contributing to soil erosion. Sealing and crusting and the low infiltration rate they produce may often be responsible for the low water availability of arid and semiarid region soils (Singer, 1991). Both forms of erosion can be controlled by careful choice of farming systems that are designed to maintain maximum cover on the soil surface during the periods of the year with the highest soil erosion potential.

Chemical properties

Relevant chemical properties of the studied profiles are shown in Table 2. The data reveal a very wide range of variation in the properties for the different soil profiles representing Azraq basin. This would require different systems of management for the different regions of the basin. Some common properties of the investigated soils include their alkalinity, content of carbonates, and low organic matter content. The ability of these soils to supply essential growth elements would be restricted not only by the availability of water directly, but by the solubility of controlling mineral phases. Stable mineral or amorphous phases can exert controlling influences on the availability of calcium, phosphorus, and micronutrients (e.g., iron, zinc, copper, and manganese). In regard to iron and phosphorus in arid region soils, nutritional limits to plant growth are common, and in both these cases organic exudates of plant and microbial origin appear important to the natural remedy of deficiencies. The evaporative gradient in arid regions can also promote the accumulation of trace elements toxic to plants and/or animals (i.e. boron, molybdenum, and selenium).

Accumulation of the organic matter in soils is a rather delicate balance between the rate of addition of biological residues and the rate of their decomposition, with both processes highly dependent on climate. In arid regions, moisture limits the amount of plant growth, the overwhelming contributor of the raw material for humus synthesis. This is clearly the case with the soils of the Azraq basin.

Some of the studied profiles contain very high electrical conductivity, reflecting high content of soluble salts. The fundamental concerns about salt-affected soils are salinity and sodicity. High salinity results in the reduction of the water potential of the soil solution, and the concentration of toxic ions that interfere with normal plant metabolism.

Specific ion toxicities are associated with chloride, sodium, and boron (James *et al.*, 1982). High sodicity manifests itself in both the physical deterioration of the soil structure and in the potential for sodium toxicity to plants.

Two profiles (Swaeid and Shubayka) represent mud flat areas formed due to erosion and deposition of soil material in low areas. These limited areas are cropped with wheat and barley where extra moisture is received from run-off from surrounding slopes.

Table 2 Relevant chemical analysis of the studied profiles.

Sample	% N	Organic C %	pH	EC (dS m ⁻¹)	CaCO ₃ %	CEC cmol(+) kg ⁻¹
Muwaqqaer III Ap (0-8)	0.062	1.92	8.59	1.005	28.0	21.7
Muwaqqaer III B (8-20)	0.037	1.30	8.46	2.319	28.3	25.2
Muwaqqaer III Bk1 (20-40)	0.035	1.28	8.02	7.36	31.4	
Muwaqqaer III Bk2 (40-120)	0.018	1.20	7.92	18.05	30.6	
Shawmari I Av (0-4)	0.0099	1.07	7.79	36.3	28.9	21.0
Shawmari I A (4-20)	0.021	0.93	8.03	11.27	30.8	19.9
Shawmari I By (20-60)	0.0015	1.06	7.72	43.2	28.1	
Shawmari I Bky (60-100)	0.0015	1.17	7.75	46.1	29.8	
Shawmari II A (0-4)	0.02	0.98	8.83	1.146	30.5	18.3
Shawmari II Bk (4-20)	0.016	0.96	9.01	3.23	30.0	22.8
Shawmari II Bky (20-75)	0.012	0.91	8.04	17.28	30.4	
Swaeid II A (0-20)	0.0295	1.95	7.42	0.280	22.7	22.8
Swaeid II Bk (20-55)	0.0280	1.81	7.20	0.287	23.8	21.7
Swaeid II Btk (55-140)	0.019	1.04	7.41	0.329	21.5	
Shubayka II Ap (0-10)	0.0545	2.05	8.06	0.366	14.7	36.3
Shubayka II By1 (10-35)	0.0560	1.86	8.17	3.04	14.7	38.2
Shubayka II By2 (35-90)	0.0465	1.05	7.91	0.913	14.3	
Shubayka II Bty (90-140)	0.0580	1.64	7.90	1.154	14.6	

Management

Although the number of the profiles is limited, their locations gave a fairly good idea about the basin. Certain general recommendations could be given to manage some areas so that sustainable agriculture could be practiced in parts of the basin. Some of the areas studied represent potential new lands provided adequate water supplies of good quality are available.

Organic matter is an essential component of intensive rehabilitation of arid lands and deserts, and therefore its application is a must, if a sustainable use of arid soils is to be accomplished. Organic matter content and composition in soils have received considerable attention because of the disproportionately favorable effect on the chemical, physical, microbiological, and morphological properties of soil compared with the mineral matter (Tucker and Fuller, 1971; Fuller and Tucker, 1977). Moreover, organic matter is the nitrogen carrier of soils. Continued soil productivity depends upon the replenishment and maintenance of some organic matter. The maintenance of a given, fixed level is often not practical, nor is it advised. The practice of incorporating readily available plant residues, manures, straw, leaves, composts, and municipal wastes and sludges into the soil is more important than the establishment of a fixed level. Some of the more prominent functions of organic matter include: direct contribution of nutrients for plants, such as nitrogen, phosphorus, and micronutrients; indirect contribution to plant nutrition by making soil elements more available; food and nutrient supply for microorganisms; soil structure development, and thereby water behavior and conservation, bulk density, infiltration, and reclamation of salt-affected soils; storehouse of elements attached to clay particles; and increased water-holding capacity.

Tillage is widely used to control weeds; incorporate soil amendments, nutrients, herbicides, and insecticides; and modify soil physical conditions, thereby improving soil conditions for crop establishment, growth, and yield. Tillage is also used to conserve soil and water. The number of tillage operations used to produce a crop varies widely and depends on the crop grown; soil and climatic conditions; and the economic level, degree of mechanization used, attitudes, and management capability of the producer. The proper tillage system must be used under the specific production system to protect the soil from water and wind erosion and to maintain the level of organic matter in the soil.

Salinity is very high in some of the profiles studied making the utilization of those soils for agricultural production not recommended at all. In the areas where soluble salts content is reasonable, leaching of the salts with a good quality of irrigation water must take place before any successful plant production can take place.

Finally, the proper types of mineral fertilizers must be used. All soils examined are calcareous with alkaline reaction. Phosphorus and micronutrients availability to crops would be a real problem if not managed well.

Conclusion

The studied profiles in the Azraq basin in northeastern Jordan represent an attempt to evaluate the potential of utilizing this large area for sustainable agricultural production. Although the number of the profiles is limited, their locations gave a fairly good idea about the areas which should be excluded from any attempt to utilize the basin for a viable agricultural production system.

References

- Bremner, J.M. 1996. Nitrogen-Total, pp. 1085-1121. *In* D.L. Sparks *et al.* (eds.). Methods of Soil Analysis. Part 3. SSSA Book Ser. 5. SSSA, Madison, WI.
- Chapman, H.D. 1965. Cation-exchange capacity. *In* C.A. Black *et al.* (eds.). Methods of Soil Analysis. Part 2. First edition. American Society of Agronomy, Madison, Wisconsin.
- Dick-Peddie, W.A. 1991. Semiarid and arid lands: A worldwide scope, pp. 3-32. *In* J. Skujins (ed.). Semiarid lands and deserts, Marcel Dekker, Inc.
- El-Swaify, S.A., T.S. Walker and S.M. Virmani. 1984. Dryland Management Alternatives and Research Needs for Alfisols in the Semiarid Tropics. International Crops Research Institute for the Semiarid Tropics, Patancheru, P.O. Andhra Pradesh, India.
- Fuller, W.H. and T.C. Tucker. 1977. Land utilization and disposal of organic wastes in arid regions. *In* L.F. Elliot and A.J. Stevenson (eds.). Soil for Management of Organic Wastes and Waste Waters, American Society of Agronomy, Madison, WI.
- Gee, G.W. and J.W. Bauder. 1986. Particle-size analysis. *In* A. Klute (ed.). Methods of Soil Analysis. Part 1. Second edition. American Society of Agronomy, Madison, Wisconsin.
- Gupta, J.P., G.G.S.N. Rao, G.N. Gupta and B.V.R. Rao. 1983. Soil drying and wind erosion as affected by different types of shelterbelts planted in the desert region of western Rajasthan, India. *J. Arid. Environ.* 6:53-58.

- Guthrie, R.L. 1982. Distribution of great groups of Aridisols in the United States, pp. 29-36. *In* D.H. Yaalon (ed.). *Aridic Soils and Geomorphic Processes*, Catena Supplement 1. Cremlingen, FRG.
- Hoekstra, T.W. and M. Shachak. 1999. *Arid Lands Management, Toward Ecological Sustainability*. University of Illinois Press.
- James, D.W., R.J. Hanks and J.J. Jurinak. 1982. *Modern Irrigated Soils*. John Wiley, New York.
- Khresat, S.A., B. Buck, Z. Rawajfih and H.C. Monger. 2001. Genesis of arid soils in the Badia region, northeastern Jordan. (Accepted for presentation at the 17WCSS, Thailand).
- Knight, W.G. 1991. Chemistry of arid region soils, pp. 111-171. *In* J. Skujins (ed.). *Semiarid Lands and Deserts*, Marcel Dekker, Inc.
- Lal, R. 1985. Soil surface management, pp. 273-300. *In* R.C. Muchow (ed.). *Agro-research for the Semiarid Tropics: North-west Australia* University of Queensland Press, St. Lucia, Queensland.
- Le Houerou, H.N. 1986. The desert and arid zones of northern Africa, pp. 101-147. *In* M. Evenari, I. Noy-Meir and D.W. Goodall (eds.). *Ecosystems of the World*, vol. 12B, Hot deserts and arid shrublands. Elsevier Science Publishers, Amsterdam.
- Middleton, N.J. 1986. Dust storms in the Middle East. *J. Arid Environ.* 10:83-97.
- Nelson D.W. and L.E. Sommers. 1996. Total carbon, organic carbon, and organic matter, pp. 961-1010. *In* D.L. Sparks *et al.* (eds.) *methods of soil analysis*. Part 3. SSSA Book Ser. 5. SSSA, Madison, WI.
- Rhoades, J. D. 1996. Salinity: Electrical conductivity and total dissolved solids, pp. 417-435. *In* D.L. Sparks *et al.* (eds.) *methods of soil analysis*. Part 3. SSSA Book Ser. 5. SSSA, Madison, WI.
- Richards, L.A. 1954. *Diagnosis and improvement of saline and alkaline soils*. U.S. Department of Agriculture, Handbook no. 60. U.S. Government. Printing Office, Washington. D.C.
- Singer, M.J. 1991. Physical properties of arid region soils, pp. 81-109. *In* J. Skujins (ed.). *Semiarid Lands and Deserts*, Marcel Dekker, Inc.
- Thomas, G.W. 1996. Soil pH and soil acidity, pp. 475-490. *In* D.L. Sparks *et al.* (eds.) *Methods of soil analysis*. Part 3. SSSA Book Ser. 5. SSSA, Madison, WI.
- Tucker, T.C. and W.H. Fuller. 1971. Soil management, humid vs. arid areas, pp. 277-286. *In* W.G. McGinnies, B.J. Godman and P. Paylore (eds.). *Food, Fiber, and the Arid Land*. University of Arizona Press, Tucson.
- Wischmeier, W.H., C.B. Johnson and B.V. Cross. 1971. A soil erodibility nomograph for farmland and construction sites. *J. Soil Water Conservation*. 26:189-193.