

Safranal, a constituent of *Crocus sativus* (saffron), attenuated cerebral ischemia induced oxidative damage in rat hippocampus

Hossein Hosseinzadeh¹ and Hamid R. Sadeghnia²

1-Pharmaceutical Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, Mashhad, I. R. Iran

2- Department of Pharmacology, Faculty of Medicine, Mashhad University of Medical Science, Mashhad, I.R. Iran

Received March 19, 2005; Revised June 15, 2005, Accepted August 15, 2005, Published August 22, 2005

Abstract Increased oxidative stress has been implicated in the mechanisms of delayed neuronal cell death following cerebral ischemic insult. In this study, we investigated whether safranal, an active constituent of *Crocus sativus* L. stigmas, may ameliorate ischemia-reperfusion injury (IRI)-induced oxidative damage in rat hippocampus. Male NMRI rats were divided into six groups, namely, sham, control, ischemia and ischemia treated with safranal (four groups). The transient global cerebral ischemia was induced using four-vessel-occlusion method for 20 min. Safranal was injected intraperitoneally (727.5 mg/kg, 363.75 mg/kg, 145.5 mg/kg, and 72.75 mg/kg body weight) 5 min. prior to reperfusion and the administration was continued every 24 hours for 72 hours after induction of ischemia. The markers of oxidative stress including thiobarbituric acid reactive substances (TBARS), total sulfhydryl (SH) groups and antioxidant capacity of hippocampus (using FRAP assay) were measured. The transient global cerebral ischemia induced a significant increase in TBARS levels ($p < 0.001$), decrement in both antioxidant power (FRAP value) ($p < 0.05$) and total sulfhydryl (SH) concentrations ($p < 0.001$) in comparison with sham-operated animals. Following safranal administration the total SH contents (3.2 vs. 0.7 $\mu\text{mol/g}$, $p < 0.001$, safranal 727.5 mg/kg) and antioxidant capacity (4.12 vs. 1.16 $\mu\text{mol/g}$, $p < 0.001$; 727.5 mg/kg) were elevated in hippocampus in comparison with ischemic group. The MDA level was declined significantly in hippocampus (52.31 vs. 159.70 nmol/g, $p < 0.001$;

727.5 mg/kg). It is concluded that safranal have some protective effects on different markers of oxidative damage in hippocampal tissue from ischemic rats.

INTRODUCTION

As the brain's high oxygen consumption, high lipid contents, especially polyunsaturated fatty acids, high concentrations of transition metals, and low antioxidants activity, the oxidative stress has been implicated as a potential contributor to the pathogenesis of acute central nervous system (CNS) injury. After brain injury, for example by ischemia, the production of reactive oxygen species (ROS) may increase, leading to tissue damage via several different cellular molecular pathways. Radicals can cause damage to cardinal cellular components such as lipids, proteins, and nucleic acids (e.g., DNA), leading to subsequent cell death by modes of necrosis or apoptosis. Therefore, the use of antioxidants, free radical scavengers or trapping agents may be rational in acute and chronic CNS injuries, especially cerebral ischemia-reperfusion injury (IRI) [1-3]. Recently, there is overwhelming attention to plant products and natural agents that can limit free radical-mediated injuries, for better therapeutic management of IRI.

Crocus sativus L., commonly known as saffron, is used in folk medicine as an antispasmodic, eupeptic, gingival sedative, anticatarrhal, nerve sedative, carminative, diaphoretic, expectorant, stimulant, stomachic, aphrodisiac and emmenagogue [4]. Furthermore, modern pharmacological studies have demonstrated that saffron extract or its active constituents have anticonvulsant [5], antidepressant [6], anti-inflammatory [7] and antitumour effects, radical scavenger as well as learning and memory improving properties [4, 8-11] and promote the diffusivity of oxygen in different tissues [4]. Saffron extract also has chemopreventive and genoprotective effects and protects from genotoxins-induced oxidative stress in mice [12-15].

The aim of present study was to assess the protective effects of safranal, the active constituent of *Crocus sativus* L. stigmas, on ischemia-reperfusion injury (IRI)-induced oxidative damage in rat hippocampus.

Correspondence author: Hossein Hosseinzadeh, Pharmaceutical Research Center, Faculty of Pharmacy, Mashhad University of Medical Sciences, P.O. Box: 1365-91775, Mashhad, I. R. Iran, hosseinzadehh@yahoo.com

MATERIALS AND METHODS

Animals

Adult male NMRI rats weighing 200-300 g were used throughout the study. All of them were kept in the same room under a constant temperature (22 ± 2 °C) and illuminated 7:00 a.m. to 7:00 p.m., with food pellets and water available *ad libitum*.

The animals were divided into six groups, each of which contained 8 rats. Group 1 was the sham group in which only surgery was done without induction of ischemia; group 2 was the control group in which saline solution was given intraperitoneally. In groups 3-6, safranal (727.5 mg/kg, 363.75 mg/kg, 145.5 mg/kg and 72.75 mg/kg, i.p., [6]) was administered 5 min prior to reperfusion and the administration was continued every 24 hours for 72 hours after the induction of ischemia.

Chemicals

DTNB (2, 2'-dinitro-5, 5'-dithiodibenzoic acid), TPTZ (2, 4, 6-tri (2-pyridyl)-1, 3, 5-triazine), TBA (2-thiobarbituric acid), n-butanol, tris, Na₂EDTA, sodium acetate, glacial acetic acid, phosphoric acid, potassium chloride, tetramethoxypropane (TMP), ferric chloride (FeCl₃.6H₂O), ferrous sulfate and hydrochloric acid was obtained from Merck. Safranal was purchased from Fluka.

Induction of global cerebral ischemia

Global cerebral ischemia was induced using the four-vessel occlusion method (4-VO), described by Pulsinelli *et al* [16]. Briefly, under intraperitoneal ketamine/xylazine anesthesia (60 mg/kg and 6 mg/kg, respectively), the alar foramina of first cervical vertebrae were exposed and the vertebral arteries were electrocauterized permanently. On the next day and under brief anesthesia, the common carotid arteries (CCAs) were dissected from surrounding tissues and temporarily ligated using the microvascular clamps for 20 min. At the end of the ischemic period, the ligature was removed and reperfusion was supplied. After maintaining of animals in suitable situation for 72 hours, the animals were decapitated and the hippocampus portion was homogenized in 1.5% cold KCl solution to give a 10% homogeneity suspension and used for biochemical assays.

Tiobarbituric acid reactive species (TBARS) measurement

The lipid peroxidation level of the hippocampus portion was measured as malondialdehyde (MDA)

which is the end product of lipid peroxidation, and reacts with thiobarbituric acid (TBA) as a thiobarbituric acid reactive substance (TBARS) to produce a red colored complex which has peak absorbance at 532 nm [17].

3 ml phosphoric acid (1%) and 1 ml TBA (0.6%) was added to 0.5 ml of homogenate in a centrifuge tube and the mixture was heated for 45 min in a boiling water bath. After cooling, 4 ml of n-butanol was added to the mixture and vortex-mixed for 1 min followed by centrifugation at 20000 rpm for 20 min. The organic layer was transferred to a fresh tube and its absorbance was measured at 532 nm. The standard curve of MDA was constructed over the concentration range of 0-40 μ M [18].

Ferric Reducing / Antioxidant Power (FRAP) assay

The FRAP assay measures the change in absorbance at 593 nm owing to the formation of a blue colored Fe^{II}-tripyridyltriazine compound from the colorless oxidized Fe^{III} form by the action of electron donating antioxidants [19].

The FRAP reagent consist of 300 mM acetate buffer (3.1 g sodium acetate + 16 ml glacial acetic acid, made up to 1 liter with distilled water; pH=3.6), 10 mM TPTZ in 40 mM HCl and 20 mM FeCl₃.6H₂O in the ratio of 10:1:1.

Briefly, 50 μ l of homogenate was added to 1.5 ml freshly prepared and prewarmed (37 °C) FRAP reagent in a test tube and incubated at 37 °C for 10 min. The absorbance of the blue colored complex was read against reagent blank (1.5 ml FRAP reagent + 50 μ l distilled water) at 593 nm. Standard solutions of Fe^{II} in the range of 100 to 1000 mM were prepared from ferrous sulphate (FeSO₄.7H₂O) in distilled water. The data was expressed as mmol ferric ions reduced to ferrous form per liter (FRAP value) [20].

Total sulfhydryl (SH) groups assay

Total SH groups were measured using DTNB (2, 2'-dinitro-5, 5'-dithiodibenzoic acid) as the reagent. This reagent reacts with the SH groups to produce a yellow colored complex which has peak absorbance at 412 nm [21].

Briefly, 1 ml Tris-EDTA buffer (pH=8.6) was added to 50 μ l homogenate in 2 ml cuvettes and sample absorbance was read at 412 nm against Tris-EDTA buffer alone (A₁). Then 20 μ l DTNB reagent (10 mM in methanol) was added to the mixture and after 15 min (stored in laboratory temperature), the sample absorbance was read again (A₂). The absorbance of DTNB reagent was also read as a

blank (B). Total thiol concentration (mM) was calculated from the following equation:

$$\text{Total thiol concentration (mM)} = (A_2 - A_1 - B) \times 1.07 / 0.05 \times 13.6$$

Statistical analysis

Data are expressed as mean \pm SEM. Statistical analysis was performed using one-way ANOVA followed by Tukey-Kramer *post-hoc* test for multiple comparisons. The p-values less than 0.05 were considered statistically significant.

RESULTS

Change of MDA levels by safranal

The degree of free radical damage following IRI was assessed using lipid peroxidation, which was measured as MDA levels. There was an increase (64.2 %) in the MDA levels following IRI as compared with sham-operated animals (159.70 ± 15.90 vs. 97.25 ± 5.18 nmol/g tissue, $p < 0.001$) (Figure 1). Safranal pretreatment resulted in a significant and dose-dependently reduction in the free radical-mediated lipid peroxidation as indicated by a decrease in the MDA levels, at various dose levels. In safranal-pretreated groups with doses 145.5 mg/kg, 363.75 mg/kg and 727.5 mg/kg, TBARS levels were 52.31, 76.85 and 98.74 nmol/g tissue, respectively (Figure 1).




Figure 1: Effect of safranal on lipid peroxidation following global cerebral ischemia. MDA levels were measured in 10% homogenates of hippocampus portion from rats subjected to 20 min of ischemia. All drugs were administered intraperitoneally 5 min prior to reperfusion. Values are mean \pm SEM (n=8). *** $p < 0.001$ as compared with vehicle (normal saline) treated animals (One-way ANOVA followed by Tukey-Kramer test)

Change of FRAP value by safranal

IRI caused a significant reduction in FRAP value (53.2 %) of homogenate samples as compared with sham-operated animals (1.16 ± 0.2 vs. 2.48 ± 0.16 $\mu\text{mol/g}$ tissue, $p < 0.001$) (Figure 2). Safranal pretreatment increased antioxidant power (FRAP value) of brain homogenate samples, in non-dose dependent manner (from 1.16 ± 0.2 to 4.12 ± 0.33 $\mu\text{mol/g}$ tissue, $p < 0.001$; 727.5 mg/kg) (Figure 2).




Figure 2: Effect of safranal on antioxidant power of hippocampus homogenate samples following global cerebral ischemia. FRAP values were measured in 10% homogenate samples from rats subjected to 20 min of ischemia. All drugs were administered intraperitoneally 5 min prior to reperfusion. Values are mean \pm SEM (n=8). * $p < 0.05$, *** $p < 0.001$ as compared with vehicle (normal saline) treated animals (One-way ANOVA followed by Tukey-Kramer test)

Effect of safranal on total thiol concentration

Following ischemia-reperfusion injury a significant reduction (77.6 %) in total SH groups (0.710 ± 0.068 vs. 3.170 ± 0.140 $\mu\text{mol/g}$ tissue, $p < 0.001$) in homogenate samples were observed (Figure 3). Safranal pretreatment induced a significant and dose dependently elevation in total thiol concentration, as compared with control group (from 0.710 ± 0.068 to 3.180 ± 0.075 $\mu\text{mol/g}$ tissue, $p < 0.001$; 727.5 mg/kg) (Figure 3).

DISCUSSION

A great deal of effort has been directed toward searching for new compounds that can be used for protection of cerebral ischemia-reperfusion injury. The results obtained in the present investigation suggest that safranal, one of the constituents of saffron stigmas with monoterpene structure, has an

overall protective effect against cerebral ischemia-reperfusion injury-induced oxidative stress in a rat model.




Figure 3: Effect of safranal on total thiol concentrations following global cerebral ischemia. Total sulfhydryl (SH) groups were measured in 10% hippocampus homogenate samples from rats subjected to 20 min of ischemia. All drugs were administrated intraperitoneally 5 min prior to reperfusion. Values are mean \pm SEM (n=8). ***p<0.001 as compared with vehicle (normal saline) treated animals (One-way ANOVA followed by Tukey-Kramer test)

A number of processes have been implicated in the pathogenesis of oxygen deprivation-induced cell injury. These include the disturbances of cell calcium homeostasis, depletion of adenine nucleotides, activation of enzymes like phospholipases with production of toxic lipid metabolites, proteases and endonucleases and generation of free radicals (ROS) that can cause oxidative damage to cellular macromolecules [22]. It is well documented that oxidative stress is a major common pathway of cellular injury following neurological and neurodegenerative disorders such as ischemia-reperfusion, seizure, Parkinson and Alzheimer's disease and antioxidant therapy have been well documented to protect against CNS injuries [3, 23, 24].

The large numbers of polyunsaturated fatty acids (PUFAs) make cell membranes particularly vulnerable to lipid peroxidation. The oxidation of PUFAs causes them to be more hydrophilic, thereby altering the structure of the membrane with resultant changes in fluidity and permeability. Lipid peroxidation can also inhibit the function of membrane bound receptors and enzymes [22, 23]. We assessed the effect of safranal on lipid peroxidation, which was measured in terms of MDA, a stable metabolite of the free radical-mediated lipid peroxidation cascade. The MDA levels increased

significantly (p<0.001) following cerebral IRI. Safranal reversed the increase of MDA levels to a considerable extent, thereby confirming its antioxidant role in IRI.

Sulfhydryl (SH) groups are highly-reactive constituents of protein molecules, and they participate in important biochemical and metabolic process such as cell division, blood coagulation, maintenance of protein systems and enzymatic activation including antioxidant enzymes (catalase, superoxide dismutase, etc.) [25]. There are also important scavengers of oxygen-derived free radicals [26]. SH groups known to be sensitive to oxidative damage and depleted following ischemic insult [27], therefore we studied the effect of this agent on the total thiol concentration during IRI. Similarly, in our studies, total sulfhydryl groups were decreased following ischemic-reperfusion injury. Safranal pretreated rats exhibited higher SH contents than their respective controls in the dose related pattern, indicating that safranal helped in replenishing the total thiol pool.

Under acute and chronic pathologic conditions such as ischemia, the balance between oxidant and antioxidant systems has been interrupted [2, 3, 28]. Therefore, we evaluate the antioxidant or reducing potential of hippocampus homogenate samples following IRI, using FRAP assay. As expected following IRI, a significant reduction in antioxidant power, as indicated by FRAP value, was observed. Safranal increased the antioxidant power of homogenate samples of hippocampus.

Saffron has chemopreventive effects and its extract inhibits tumor growth in vivo and in vitro [12, 13, 29-34]. Escribano *et al* showed that saffron extract and its constituents; crocin, safranal and picrocrocin inhibit the growth of human cancer cells (Hella cells) in vitro [9]. Abdullaev and Frenkel also showed saffron affect intracellular nucleic acid and protein synthesis [35, 36]. Another study (El Daly) demonstrated protective effects of saffron extract against cisplatin induced toxicity in rats [37]. Saffron extract also has radical scavenger properties [4] and protects from genotoxicity as well as genotoxins-induced oxidative stress in mice [14, 15]. Premkumar *et al* showed oral pretreatment with the saffron aqueous extract (40 and 80 mg/kg) for five consecutive days inhibit genotoxins-induced oxidative stress in mice liver. In this study, an increase in the levels of glutathione (GSH) concentration as well as the activities of glutathione S-transferase (GST), glutathione peroxidase (GPx), catalase and superoxide dismutase (SOD) were

observed, however, normal levels of GSH could not be attained [15].

Among the constituent of saffron stigmas, crocins and crocetin derivatives are most abundant with established antioxidant and antitumor effects [4, 8, 13]. These carotenoids scavenge free radicals, especially superoxide anions and thereby may protect cells from oxidative stress [38]. In rats, crocin dyes are known to exert protective effects against acute hepatic damage induced by aflatoxin B₁ and dimethylnitrosamine [39]. It has been shown that crocetin, the deglycosylated crocin derivative, has protective effects on aflatoxin B₁-induced hepatotoxicity and protects rat primary hepatocytes against oxidative damage [40-42]. Cancer chemopreventive as well as antitumor activities were also reported for crocins and crocetin derivatives in different assay systems [9, 43-46].

There are several reports about the antioxidant activity and anti-inflammatory effects of some monoterpenoids such as α -pinene. Moreover, there have been shown monoterpenoids such as terpineol and linalool have depressant effects on central nervous system, in vivo [47] and linalool competitively inhibits glutamate receptors [48]. There are no reports about clinical uses of safranal. Much more basic pharmacological and toxicological studies need for clinical trials to evaluate the safety, tolerability and efficacy of safranal.

Our previous studies showed that safranal has a potent depressant effect on CNS and clearly suppress pentylenetetrazole-induced seizures (unpublished data) [49]. Its may be concluded that protective effect of safranal on ischemia-reperfusion injury, at least partly, due to these mechanisms, but it needs to be further investigated.

It is concluded that safranal have some protective effects on different markers of oxidative damage in hippocampal tissue from ischemic rats.

ACKNOWLEDGEMENTS

The authors are thankful to the Research Council, Mashhad University of Medical Sciences for financial support.

REFERENCES

[1] White BC, Sullivan JM, DeGracia DJ, O'Neil BJ, Neumar RW, Grossman LI, Rafols JA, Krause GS. Brain ischemia and reperfusion: molecular mechanisms of neuronal injury. *J Neuro Sci* 2000, 179: 1-33.
[2] Streck EL, Vieira PS, Wannmacher CMD, Dutra-Filho

CS, Wajner M, Wyse ATS. In vitro effect of homocysteine on some parameters of oxidative stress in rat hippocampus. *Metab Brain Disease* 2003, 18: 147-54.

[3] Gilgun-Sherki Y, Rosenbaum Z, Melamed E, Offen D. Antioxidant Therapy in Acute Central Nervous System Injury: Current State. *Pharm Rev* 2002, 54:271-84.

[4] Rios jl, Recio MC, Ginger RM, Manz S. An update review of saffron and its active constituents. *Phytother Res* 1996, 10: 189-93.

[5] Hosseinzadeh H, Khosravan V. Anticonvulsant effects of aqueous and ethanolic extracts of *Crocus sativus* L. stigmas in mice. *Arch Imn Med* 2002, 5: 44-7.

[6] Hosseinzadeh H, Karimi Gh, Niapoor M. Antidepressant effects of *Crocus sativus* stigma extracts and its constituents, crocin and safranal, in mice. *Acta Hort (ISHS)* 2004, 650: 435-45.

[7] Hosseinzadeh H, Younesi HM. Antinociceptive and anti-inflammatory effects of *Crocus sativus* L. stigma and petal extracts in mice. *BMC Pharmacol* 2002, 2: 1-8.

[8] Abdullaev FI. Biological effects of saffron. *Biofactors* 1993, 4: 83-6.

[9] Escribano J, Alonso GL, Coca-Prados M, Fernandez JA. Crocin, safranal and picrocrocin from saffron (*Crocus sativus* L.) inhibit the growth of human cancer cells in vitro. *Cancer Lett* 1996, 100: 23-30.

[10] Zhang YX, Sugiura M, Saito H, Shoyama Y. Acute effects of *Crocus sativus* L. on passive avoidance performance in mice. *Biol Pharmacol Bull* 1994, 17: 217-21.

[11] Abe K, Sugiura M, Ymaguchi S, Shoyama Y, Saito H. Saffron extract prevents acetaldehyde-induced inhibition of long-term potentiation in the rat dentate gyrus in vivo. *Brain Res* 1999, 851: 287-89.

[12] Abdullaev FI, Caballero-Ortega H, Riveron-Nigrete L, Pereda-miranda R, Rivera-Luna R, Manuel Hernandez J, Perez-Lopez I, Espinosa-Aguirre JJ. In vitro evaluation of chemopreventive potential of saffron. *Rev Inves Clin* 2002, 54: 430-36.

[13] Nair SC, Kurumboor SK, Hasegawa JH. Saffron chemoprevention in biology and medicine: a review. *Cancer Biother* 1995, 10: 257-64.

[14] Premkumar K, Abraham SK, Santhiya ST, Gopinath PM, Ramesh A. Inhibition of genotoxicity by saffron (*Crocus sativus* L.) in mice. *Drug Chem Toxicol* 2001, 24: 421-28.

[15] Premkumar K, Abraham SK, Santhiya ST, Ramesh A. Protective effects of saffron (*Crocus sativus* L.) on genotoxins-induced oxidative stress in swiss albino mice. *Phytother Res* 2003, 17: 614-17.

[16] Pulsinelli WA, Brierley JB. A new model of bilateral hemispheric ischemia in the unanesthetized rat. *Stroke* 1979, 10: 268-72.

[17] Fernandez J, Perez-Alvarez JA, Fernandez-lopez JA. Thiobarbituric acid test for monitoring lipid oxidation in meat. *Food Chem* 1997, 99: 345-53.

[18] Uchiama M, Miahara M. Determination of malonaldehyde precursor in tissues by thiobarbituric acid

- test. Anal Biochem 1978, 86: 279-86.
- [19] Benzie IFF, Strain J. The ferric reducing ability of plasma (FRAP) as a measure of antioxidant power: The FRAP assay. Anal Biochem 1996, 239: 70-6.
- [20] Benzie IFF, Strain JJ. Ferric reducing/antioxidant power assay: direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. Methods Enzymol 1999, 299: 15-27.
- [21] Ellman G. Tissue sulfhydryl groups. Arch Biochem Biophys 1959, 82: 70-7.
- [22] Fisher M. Stroke therapy. 2nd ed. Butter worth-Heinemann 2001, 25-50.
- [23] Love S. Oxidative stress in brain ischemia. Brain Pathol 1999, 9: 119-31.
- [24] Sun AY, Chen YM. Oxidative stress and neurodegenerative disorders. J Biomed Sci 1998, 401-414.
- [25] Jansen EV. Sulfhydryl-disulfide interchanges. Science 1959, 130: 1319-1323.
- [26] Dormandy TL. An approach to free radicals in medicine and biology. Acta Physiol Scand 1980 492, 153-168.
- [27] Soszynski M, Bartosz G. Decrease in accessible thiols as an index of oxidative damage to membrane proteins. Free Rad Biol Med 1997, 23: 463-69.
- [28] Abdollahi M, Ranjbar R, Shadnia S, Nikfar S, Rezaie A. Pesticides and oxidative stress: a review. Med Sci Monit 2004, 10: 141-47.
- [29] Salomi MJ, Nair SC, Panikkar KR. Inhibitory effects of *Nigella sativa* and *Crocus sativus* on chemical carcinogenesis in mice and its non-mutagenic activity. Proc Ker Sci Congress 1990, 3: 125.
- [30] Salomi MJ, Nair SC, Panikkar KR. Inhibitory effects of *Nigella sativa* and saffron (*Crocus sativus*) on chemical carcinogenesis in mice. Nutr Cancer 1991, 16: 67-72.
- [31] Abdullaev FI. Cancer chemopreventive and tumoricidal properties of saffron (*Crocus sativus*). Exptl Biol Med 2002, 227: 20-25.
- [32] Das I, Chakrabarty RN, Das S. Saffron can prevent chemically induced skin carcinogenesis in swiss albino mice. Asian Pac J Cancer Prev 2004, 4: 70-76.
- [33] Nair SC, Pannikar B, Pannikar KR. Antitumor activity of saffron (*Crocus sativus*). Cancer Lett 1991, 57: 109-114.
- [34] Nair SC, Varghese CD, Pannikar KR, Kurumboor SK, Parathod RK. Effect of saffron on vitamin A levels and its antitumor activity on the growth of solid tumors in mice. Int J Pharmacol 1994, 32: 105-114.
- [35] Abdullaev FI, Frenkel GD. The effect of saffron on intracellular DNA, RNA and protein synthesis in malignant and non-malignant human cells. Biofactors 1992, 4: 43-45.
- [36] Abdullaev FI, Frenkel GD. Effect of saffron on cell colony formation and cellular nucleic acid and protein synthesis. Biofactors 1992, 3: 201-204.
- [37] El Daly ES. Protective effect of cysteine and vitamin E, *Crocus sativus* and *Nigella sativa* extracts on cisplatin-induced toxicity in rats. J Pharm Belg 1998, 53:87-95.
- [38] Bors W, Saran M, Michel C. Radical intermediates involved in the bleaching of the carotenoid crocin. Hydroxyl radicals, superoxide anions and hydrated electrons. Int J Radiat Biol Relat Stud Phys Chem Med 1982, 41: 493-501.
- [39] Lin JK, Wang CJ. Protection of crocin dyes on the acute hepatic damage induced by aflatoxin B₁ and dimethylnitrosamine in rats. Carcinogenesis 1986, 7:595-599.
- [40] Wang CJ, Shioh SJ, Lin JK. Effects of crocetin on the hepatotoxicity and hepatic DNA binding of aflatoxin B₁ in rats. Carcinogenesis 1991, 12: 459-462.
- [41] Wang CJ, Hsu JD, Lin JK. Suppression of aflatoxin B₁-induced hepatotoxicity lesions by crocetin. Carcinogenesis 1991, 12:1807-1810.
- [42] Tseng TH, Chu CY, Huang JM, Shioh SJ, Wang CJ. Crocetin protects against oxidative damage in rat primary hepatocytes. Cancer Lett 1995, 97: 61-67.
- [43] Garcia-Olmo DC, Riese HH, Escribano J, Ontanon J, Fernandez JA, Atienzar M, Garcia-Olmo D. Effects of long term treatment of colon adenocarcinoma with crocin, a carotenoid from saffron (*Crocus sativus*): an experimental study in the rat. Nutr Cancer 1999, 35: 120-126.
- [44] Tarantilis PA, Morjani H, Polissou M, Manafait M. Inhibition of growth and induction of differentiation of promyelocytic leukemia (HL-60) by carotenoids from *Crocus sativus* L. AntiCancer Res 1994, 14: 1913-1918.
- [45] Konoshima T, Takasaki M, Tokuda H, Morimoto S, Tanaka H, Kawata E, Xuan LJ, Saito H, Sugiura M, Molnar J, Shoyama Y. Crocin and crocetin derivatives inhibit skin tumor promotion in mice. Phytother Res 1998, 12: 400-404.
- [46] Wang CJ, Lee MJ, Chang MC, Lin JK. Inhibition of tumor promotion in benzo(a)pyrene-initiated CD-1 mouse skin by crocetin. Carcinogenesis 1995, 16: 501-506.
- [47] Perry NSL, Bollen C, Perry EK, Ballard C. Salvia for dementia therapy: review of pharmacological activity and pilot tolerability clinical trial. Pharmacol Biochem Behav 2003, 75: 651-659
- [48] Silva Brum LF, Elisabetsky E, Souza D. Effect of linalool on [³H] MK801 and [³H] muscimol binding in mouse cortical membranes. Phytother Res 2001, 15: 422-25.
- [49] Hosseinzadeh H, Talebzadeh F. Anticonvulsant evaluation of safranal and crocin from *Crocus sativus* in mice. Fitoterapia. 2005, In Press.