

# A sample of 669 ultra steep spectrum radio sources to find high redshift radio galaxies

C. De Breuck<sup>1,2</sup>, W. van Breugel<sup>2</sup>, H.J.A. Röttgering<sup>1</sup>, and G. Miley<sup>1</sup>

<sup>1</sup> Sterrewacht Leiden, Postbus 9513, 2300 RA Leiden, The Netherlands  
e-mail: debreuck,miley,rottgeri@strw.leidenuniv.nl

<sup>2</sup> Institute of Geophysics and Planetary Physics, Lawrence Livermore National Laboratory, L-413, Livermore, CA 94550, U.S.A.  
e-mail: wil@igpp.ucllnl.org

Received November 4, 1999; accepted February 4, 2000

**Abstract.** Since radio sources with Ultra Steep Spectra (USS;  $\alpha \lesssim -1.30$ ;  $S \propto \nu^\alpha$ ) are efficient tracers of high redshift radio galaxies (HzRGs), we have defined three samples of such USS sources using the recently completed WENSS, TEXAS, MRC, NVSS and PMN radio-surveys. Our combined sample contains 669 sources with  $S_{1400} > 10$  mJy and covers virtually the entire sky outside the Galactic plane ( $|b| > 15^\circ$ ). For our 2 largest samples, covering  $\delta > -35^\circ$ , we selected only sources with angular sizes  $\Theta < 1'$ . For 410 sources, we present radio-maps with  $0''.3$  to  $\sim 5''$  resolution from VLA and ATCA observations or from the FIRST survey, which allows the optical identification of these radio sources.

Comparison with spectrally unbiased samples at similar flux density levels, shows that our spectral index, flux density, and angular size selections do not affect the angular size distribution of the sample, but do avoid significant contributions by faint foreground spiral galaxies. We find that the spectral index distribution of 143 000 sources from the WENSS and NVSS consists of a steep spectrum galaxy and a flat spectrum quasar population, with the relative contribution of flat spectrum sources doubling from  $S_{1400} > 0.1$  Jy to  $S_{1400} > 2.5$  Jy. The identification fraction of our USS sources on the POSS ( $R \lesssim 20$ ) is as low as 15%, independent of spectral index  $\alpha < -1.30$ . We further show that 85% of the USS sources that can be identified with an X-ray source are probably contained in galaxy clusters, and that  $\alpha < -1.6$  sources are excellent Galactic pulsar candidates, because the percentage of these sources is four times higher in the Galactic plane.

Our sample has been constructed to start an intensive campaign to obtain a large sample of high redshift objects

( $z > 3$ ) that is selected in a way that does not suffer from dust extinction or any other optical bias<sup>1</sup>.

**Key words:** surveys — galaxies: active — radio continuum: galaxies


## 1. Introduction

Radio galaxies have now been found out to redshifts of  $z = 5.19$  (van Breugel et al. 1999b) and radio-loud quasars out to  $z = 4.72$  (Hook & McMahon 1998). Although new optical selection techniques such as color-dropouts, deep spectroscopy of blank fields, and narrow-band Ly $\alpha$  imaging have now found galaxies at similar (Steidel et al. 1999) and even higher redshifts (up to  $z \sim 5.75$ ; Dey et al. 1998; Weymann et al. 1998; Spinrad et al. 1998, Hu et al. 1999), radio sources are still the only objects that can be selected uniformly over all redshift ranges, and in a way that does not suffer from optical biases such as dust extinction, which is known to be important at these high redshifts (e.g. Hughes et al. 1997; Ivison et al. 1998; Dickinson 1998).

At low to moderate redshift ( $z \lesssim 1$ ), powerful radio sources are uniquely identified with massive ellipticals (Lilly & Longair 1984; Owen & Laing 1989; Best et al. 1998; McLure & Dunlop 2000). The strongest indications that this is also true at higher redshifts comes from the near-IR Hubble  $K - z$  diagram of radio galaxies which shows a remarkably close correlation from the

*Send offprint requests to:* C. De Breuck

<sup>1</sup> Appendices B, C and D are available in electronic form only, at <http://www.edpsciences.org>



**Fig. 1.**  $\alpha_{1400}^{325}$  against  $z$  for 2 samples without spectral index selection (3CR, Spinrad et al. 1985 and MRC, McCarthy et al. 1996), and 2 USS samples (4C, Chambers et al. 1996a, and our new WN/TN samples, as defined in this paper). Note that the correlation is present in the spectrally unbiased 3CR and MRC, and that the 4C and our new USS samples are finding three to five times more  $z > 2$  radio galaxies than the MRC. The horizontal dotted line indicates the  $\alpha_{1400}^{325} < -1.3$  cutoff used in our USS sample

present out to  $z = 5.19$  (Lilly 1989; Eales et al. 1997; van Breugel et al. 1998; van Breugel et al. 1999b). This suggests that we can use radio galaxies to study the formation and evolution of the most massive galaxies, which, by their implied star-formation history, can put important constraints on galaxy formation models, and even on cosmological parameters (e.g. Dunlop et al. 1996; Spinrad et al. 1997). Although the unification model for radio galaxies and quasars (e.g. Barthel 1989) suggests we could also use quasars as tracers, a detailed stellar population study of quasar host galaxies is almost impossible due to the extreme luminosity of the AGN. Furthermore, samples of radio sources designed to find large quantities of quasars require additional optical selections (e.g. Gregg et al. 1996; Hook & McMahon 1998; White et al. 2000).

Considerable effort has been spent over the last decade to find these high redshift radio galaxies (HzRGs), which has led to the discovery of more than 140 radio galaxies at redshifts  $z > 2$  (see e.g. De Breuck et al. 1998a for a recent summary). However by  $z > 3$ , their numbers become increasingly sparse, and using flux limited radio surveys such as the 3CR ( $S_{178} > 10$  Jy; Laing et al. 1983), or the MRC strip ( $S_{408} > 0.95$  Jy; McCarthy et al. 1996), the highest redshift radio galaxy found so far is at  $z \sim 3.2$  (Fig. 1; Rawlings et al. 1990; McCarthy et al. 1996). This redshift

limit arises because radio power is correlated with redshift in bright flux limited samples, and an upper limit exists in the radio luminosity. Lowering the flux limit would not only substantially increase the number of sources in these samples, but at the same time the fraction of luminous very high redshift radio galaxies would decrease (Blundell et al. 1998; Jarvis et al. 1999). This fractional decrease would arise even if there is no decrease in co-moving space density at  $z \gtrsim 2.5$ . Such a redshift cutoff has been suggested by Bremer et al. (1998), but recently Jarvis et al. (1999) rule out a break at  $z \lesssim 2.5$ . To efficiently find large numbers of HzRGs in acceptable observing times, it is therefore necessary to apply additional selection criteria, at the expense of completeness.

By far the most successful selection criterion has been the ultra steep spectrum criterion (e.g. Röttgering et al. 1994; Chambers et al. 1996a; Blundell et al. 1998). Selecting sources with very steep radio spectra increases dramatically the chance of finding  $z > 2$  radio galaxies (Fig. 1). This technique is based on the results of Tielens et al. (1979) and Blumenthal & Miley (1979), who found that the identification fraction on the POSS ( $R \lesssim 20$ ) decreases with steepening spectral index, consistent with the steeper sources being at higher redshifts. It is now getting clear that this correlation can be explained by a combination of a K-correction of a concave radio spectrum and an increasing spectral curvature with redshift (Krolik & Chen 1991; Carilli et al. 1998; van Breugel et al. 1999a). To further investigate the  $z - \alpha$  correlation, we have calculated spectral indices using the flux densities from the WENSS (Rengelink et al. 1997) and NVSS (Condon et al. 1998) catalogs for four different samples: the flux density limited 3CR (Spinrad et al. 1985) and MRC (McCarthy et al. 1996) surveys, and the USS samples from the 4C (Chambers et al. 1996a) and the one presented in this paper. The results (Fig. 1) show a trend for steeper spectral index sources to have higher redshifts in flux limited, spectrally unbiased samples, confirming the empirical relation out to the highest redshifts. The efficiency of the USS criterion is clearly illustrated by the fact that the 4C USS sample (Chambers et al. 1996a) contains 50%  $z > 2$  sources, and by the early spectroscopic results on the USS samples presented in this paper, which indicate that  $\sim 2/3$  of our sources have  $z > 2$ . It is even more impressive to note that 13 of the 14 radio galaxies at  $z > 3.5$  we know of have been found from samples with a steep spectral index selection<sup>2</sup>! The limitation of this technique is that the steepest spectrum sources are rare, comprising typically only 0.5% (at  $\alpha < -1.30$ ) of a complete low frequency sample; therefore, large and deep all sky surveys are needed to obtain a significant sample of USS sources.

<sup>2</sup> The only exception is VLA J123642+6213 (Waddington et al. 1999), which has been identified in the HDF, but it does have a steep spectral index ( $\alpha_{1400}^{8500} = -0.94$ ).

**Table 1.** Radio surveys

	WENSS	TEXAS	MRC
Frequency (MHz)	325	365	408
Sky region (J2000)	$\delta > +29^\circ$	$-35^\circ.7 < \delta < +71^\circ.5$	$-85^\circ < \delta < +18^\circ.5$
# of sources	229 576	67 551	12 141
Resolution	$54'' \times 54'' \text{ cosec } \delta$	$10''^a$	$2'.62 \times 2'.86 \text{ sec } (\delta - 35^\circ.5)$
Position uncertainty (strong sources)	$1''.5$	$0''.5-1''$	$8''$
RMS noise	$\sim 4 \text{ mJy}$	20 mJy	70 mJy
Flux density limit	18 mJy	150 mJy	670 mJy
Reference	Rengelink et al. 1997	Douglas et al. 1996	Large et al. 1981
	NVSS	FIRST	PMN
Frequency (MHz)	1400	1400	4850
Sky region (J2000)	$\delta > -40^\circ{}^b$	$7^{\text{h}}20^{\text{m}} < \alpha < 17^{\text{h}}20^{\text{m}}; +22^\circ.2 < \delta < +57^\circ.5$ $21^{\text{h}}20^{\text{m}} < \alpha < 3^{\text{h}}20^{\text{m}}; -2^\circ.5 < \delta < +1^\circ.6$	$-87^\circ.5 < \delta < +10^\circ$
# of sources	1 689 515	437 429	50 814
Resolution	$45'' \times 45''$	$5'' \times 5''$	$4'.2$
Position uncertainty (strong sources)	$1''$	$0'.1$	$\sim 45''$
RMS noise	0.5 mJy	0.15 mJy	$\sim 8 \text{ mJy}$
Flux density limit	2.5 mJy	1 mJy	20 mJy
Reference	Condon et al. 1998	Becker et al. 1995	Griffith & Wright 1993

<sup>a</sup> The Texas interferometer has a complicated beam. However, sources with separations between  $10''$  and  $2'$  can be successfully modeled as doubles, and will have a single entry in the catalog. See Douglas et al. (1996) for details.

<sup>b</sup> Some small gaps are not covered. They are listed on the NVSS homepage (1998 January 19 version).

With the advent of several new deep all-sky surveys (Sect. 2), it is now possible for the first time to construct a well defined all-sky USS sample with optimized selection criteria to find large numbers of  $z > 3$  radio galaxies. In this paper, we describe the construction of such a sample, and present high resolution radio observations needed to determine accurate positions and morphologies. This information is essential for the optical and near-IR identifications, and subsequent optical spectroscopy of a significant sub-set of our sample, which will be described in future papers. The organization of the paper is as follows: we describe the radio surveys we used in Sect. 2 and define our samples in Sect. 3. We present and discuss our radio observations in Sect. 4, and present our conclusions in Sect. 5.

## 2. Description of the radio surveys

During the past years, several all-sky radio-surveys have become available (Table 1), which are 1 – 2 orders of magnitude more sensitive than previous surveys at similar frequencies (Fig. 2). The combination of these new surveys allows us to define for the first time a large sample of USS sources that covers the whole sky<sup>3</sup> in both hemispheres. We list the main survey parameters in Table 1. In this section, we will briefly discuss the usefulness of these new radio surveys for the construction of USS samples.


<sup>3</sup> To facilitate optical follow-up, we will exclude the Galactic plane at  $|b| < 15^\circ$ .

### 2.1. WENSS


The Westerbork Northern Sky Survey (WENSS; Rengelink et al. 1997) at 325 MHz is the deepest low-frequency survey with a large sky coverage (3.14 sr). We used the WENSS to define the largest, and most complete USS sample to date, covering the entire sky North of declination  $+29^\circ$ . We used version 1.0 of the main and polar WENSS catalogs. A small area is covered by both these catalogs; we selected only the sources from the main catalog in this overlapping area.

### 2.2. Texas

The Texas survey, made with the Texas interferometer from 1974 to 1983 (Douglas et al. 1996), covers 9.63 steradians at a frequency of 365 MHz with a limiting flux density about ten times higher than that of the WENSS. The Texas interferometer's 3.5 km maximum baseline provides  $< 1''$  positional accuracy, but its poor  $uv$ -coverage leads to irregular beamshapes and lobe-shifts, hampering accurate modeling of extended sources. A detailed discussion of these complications can be found in Douglas et al. (1996). To minimize these problems, we have selected only the 40.9% sources that are well modeled (listed with a “+++” flag in the catalog). This selection excludes primarily  $S_{365} \lesssim 700 \text{ mJy}$  sources (Fig. 3), but even at  $S_{365} \gtrsim 700 \text{ mJy}$ , one out of three sources is excluded by this criterion. Douglas et al. (1996) have calculated the



**Fig. 2.** Limiting flux density plotted for all major radio surveys. Lines are of constant spectral indices of  $-1.3$ . Note that WENSS, NVSS and FIRST have flux density limits  $\sim 100$  times deeper than previous surveys at comparable wavelengths




**Fig. 3.** Fraction of sources with “+++” flag in the Texas catalog (see text) as a function of Texas  $S_{365}$  flux density. Note that the selection of “+++” sources excludes primarily sources with  $S_{365} \lesssim 700$  mJy

completeness above flux density  $S$  of the Texas catalog (defined as the fraction of sources with true flux density greater than  $S$  which appear in the catalog) by comparing the Texas with the MRC (Large et al. 1981) and a variety of other low-frequency catalogs. They found that the completeness varies with declination (because the survey was done in declination strips over a large time span), and an expected increase in completeness at higher flux densities. In Table 2, we reproduce their completeness table, extended with the values after the “+++” selection.

**Table 2.** Completeness of the Texas survey

Limiting flux density	all sources	“+++” sources
250 mJy	0.8	0.2
350 mJy	0.88	0.28
500 mJy	0.92	0.40
750 mJy	0.96	0.51
1 Jy	0.96	0.50



**Fig. 4.** Ratio of integrated Texas over WENSS flux density against integrated WENSS flux density. Only Texas “+++” sources are plotted. The horizontal line is the expected ratio 0.865 due to the 40 MHz difference in central frequency, and assuming a spectral index of  $\alpha_{365}^{1400} = -0.88$ . The curved line indicates a 150 mJy flux density limit in the Texas catalog. Note the increasing amount of overestimated Texas flux densities with decreasing flux density  $S_{\text{WENSS}} < 400$  mJy

To examine the reliability of the listed flux densities, and to check to what extent the “+++” selection has removed the spurious sources from the catalog, we have correlated the Texas “+++” sources with WENSS, NVSS and FIRST. In Fig. 4, we compare the Texas flux densities with those of the WENSS. At  $S_{325} \gtrsim 500$  mJy, the ratio of the flux densities is closely distributed around 0.9. This ratio is what we expect due to the 40 MHz central frequency difference between the two surveys and assuming a spectral index  $\alpha_{365}^{1400} = -0.879$  (the median of the Texas-NVSS spectral indices). At  $S_{325} \lesssim 500$  mJy, the number of sources in the Texas catalog which are brighter than in WENSS catalog increases with decreasing flux density. This can be explained by the “up-scattering” of sources near the flux limit of the Texas catalog (i.e. only sources intrinsically brighter than  $S_{365} = 150$  mJy will be detected, but no  $S_{365} < 150$  mJy sources with a large

positive flux density measurement error). The result of this on a USS sample based on the Texas survey and correlated with a higher frequency survey (such as the NVSS), will be that with lower  $S_{365}$ , we will find more sources whose spectral indices appear steeper than they really are.

We also examined the dependence of the ratio Texas/WENSS flux density on angular size, determined from the FIRST survey (see Sect. 2.1.4). We found no significant residual variation of the flux density ratio at sizes between  $5''$  and  $2'$ .

In Fig. 5b, we plot the density of NVSS sources around a Texas source (see also Sect. 2.2.5). The width of the over-density peak ( $\sim 10''$ ) is due to the positional inaccuracies in the Texas and NVSS catalogues. However, the very broad tail of sources between  $20''$  and  $110''$  and the secondary peak coinciding with the fringe separation at  $73''$  indicates that the “+++” selection did not remove all spurious sources from the catalog.

In summary, after the selection of “+++” sources, the Texas catalog still contains  $< 5\%$  spurious sources (Douglas et al. 1996), probably due to residual lobe-shifted sources. Our comparison of the Texas flux densities with those of the WENSS survey shows that the differences are consistent with the errors quoted in the catalog. The selection of the Texas catalog with only “+++” sources is thus  $>95\%$  reliable, but only  $\sim 40\%$  complete.

### 2.3. NVSS

The NRAO VLA Sky Survey (NVSS; Condon et al. 1998) covers the 10.3 steradians north of  $-40^\circ$  at 1.4 GHz, and reaches a 50 times lower limiting flux density than previous large area 1.4 GHz surveys. At the flux density levels we are using ( $S_{1400} > 10$  mJy), the catalog is virtually complete. Because the NVSS resolution is comparable to that of the WENSS and Texas surveys, and its sky coverage is large, we use the NVSS to determine the spectral indices in our USS samples based on the WENSS and Texas surveys. The final NVSS catalog was not yet completed at the time of our USS sample construction. For our final sample, presented in this paper, we use the 1998 January 19 version. This version still lacks data in a small number of regions of the sky (listed on the NVSS homepage). As a result, the sky coverage of the area listed in Table 3 is only 99.77%.

### 2.4. FIRST

The Faint Images of Radio Sky at Twenty centimeters (FIRST, Becker et al. 1995) survey is currently being made with the VLA in the B-array at 1.4 GHz, and has a limiting flux density three times deeper than the NVSS. We used the 1998 February 4 version of the catalog, covering 1.45 steradians. As noted by Becker et al. (1995), the

photometry for extended sources in FIRST might be less reliable than that of the NVSS, due to the  $9\times$  higher resolution, which could underestimate large-scale diffuse radio emission. As the FIRST area is completely covered by the NVSS, we will consistently use NVSS flux densities for our spectral index calculation. The main advantages of FIRST over NVSS for our purposes are the much better positional accuracy ( $< 0''.5$ ) and the higher ( $5''$ ) resolution. This combination allows the identification of even the very faint ( $R > 20$ ) optical counterparts of radio sources. Additionally, the fainter detection limit of the FIRST allows an extra check on the flux densities of compact sources.

### 2.5. MRC


The Molonglo Reference Catalog of radio sources (Large et al. 1981) at 408 MHz is presently the most sensitive low-frequency catalog with reasonable positional accuracy that covers the deep southern hemisphere,  $\delta < -35^\circ$ . We will use this catalog in combination with the PMN survey (see below) to define the first USS sample at  $\delta < -40^\circ$ .

### 2.6. PMN

The Parkes-MIT-NRAO (PMN) survey is a combination of 4 strips observed with the Parkes telescope at 4.85 GHz. The strips cover different parts of the sky, each with a slightly different limiting flux density. The regions are: southern ( $-87.5^\circ < \delta < -37^\circ$ , Wright et al. 1994), zenith ( $-37^\circ < \delta < -29^\circ$ , Wright et al. 1996), tropical ( $-29^\circ < \delta < -9.5^\circ$ , Griffith et al. 1994), and equatorial ( $-9.5^\circ < \delta < +10^\circ$ , Griffith et al. 1995). For our southern hemisphere sample, we have used the southern and zenith catalogs to find USS sources at  $\delta < -30^\circ$ .

## 3. USS samples

Figure 2 shows that the surveys described above have very compatible flux density limits for defining samples of USS sources. At the same time, their sky coverage is larger and more uniform than previous surveys used for USS sample construction (Wieringa & Katgert 1992; Röttgering et al. 1994; Chambers et al. 1996a; Blundell et al. 1998; Rengelink 1998; Pursimo et al. 1999; Pedani & Grueff 1999; Andernach et al. 2000). We selected the deepest low and high frequency survey available at each part of the sky. For a small region  $-35^\circ < \delta < -30^\circ$  which is covered by both Texas and MRC, we used both surveys. This resulted in a more complete samples since the lower sensitivity of the PMN survey in the zenith strip (see Sect. 2.6) is partly compensated by the (albeit incomplete) Texas survey. To avoid problems with high Galactic extinction during optical imaging and spectroscopy, all regions at



**Fig. 5.** The density of sources from the high frequency catalog used in the correlation (NVSS or PMN) around sources from the low frequency catalog (WENSS, Texas or MRC) as a function of search radius. The dotted line represents the distribution of confusion sources (see text). The apparent under-density of WENSS–NVSS sources at search radii  $\gtrsim 60''$  is due to the grouping of multiple component sources in the  $\sim 1'$  resolution WENSS. Note the plateau and secondary peak around the  $73''$  Texas fringe separation in the Texas–NVSS correlation and the much larger uncertainty in the MRC–PMN correlation

Galactic latitude  $|b| < 15^\circ$  were excluded<sup>4</sup>, as well as the area within  $7^\circ$  of the LMC and SMC. This resulted in three USS samples that cover a total of 9.4 steradians (Fig. 6).

We designate the USS samples by a two-letter name, using the first letter of their low- and high-frequency contributing surveys. Sources from these samples are named with this 2-letter prefix followed by their IAU J2000-names using the positions from the NVSS catalog (WN and TN samples) or the MRC catalog (MP sample). We did not rename the sources after a more accurate position from our radio observations or from the FIRST survey. The sample definitions are summarized in Table 3.

### 3.1. Survey combination issues

We first discuss the problems that arise when combining radio surveys with different resolutions and positional uncertainties.

#### 3.1.1. Correlation search radius

Due to the positional uncertainties and resolution differences between radio surveys, in general the same source will be listed with slightly different positions in the catalogs.

To empirically determine the search radius within which to accept sources in 2 catalogs to be the same, we compared the density of objects around the position listed in the low-frequency survey (which has lower resolution) with the expected number of random correlations in each sample ( $\equiv$  confusion sources). To determine this number

<sup>4</sup> This also reduces the number of Galactic pulsars in our sample (see Sect. 4.7.2).

as a function of distance from the position in the most accurate catalog, we created a random position catalog by shifting one of the input catalogs by  $1^\circ$  in declination, and made a correlation with this shifted catalog. The density of sources as a function of distance from the un-shifted catalog then represents the expected number of confusion sources as a function of radial distance. In Fig. 5, we plot for each of our three samples the observed density around these sources with this confusion distribution over-plotted. The correlation search radius should thus be chosen at a distance small enough for the density of confusion sources to be negligible.

We decided to adopt the radius where the density of real sources is at least ten times higher than the density of confusion sources as the search radius for our sample construction, except for the WN sample (would be  $15''$ ) where we chose the same radius as for the TN sample ( $10''$ ). The later was done for consistency between both samples. Because of the five times lower resolution and source densities in the MRC and PMN surveys, the search radius of the MP sample is eight times larger. Summarized, the search radii we used are  $10''$  for WN and TN, and  $80''$  for MP.

#### 3.1.2. Angular size

In order to minimize errors in the spectral indices due to different resolutions and missing flux on large angular scales in the composing surveys, we have only considered sources which are not resolved into different components in the composing surveys. Effectively, this imposes an angular size cutoff of  $\sim 1'$  to the WN,  $\sim 2'$  to the TN sample and  $\sim 4'$  to the MP sample. We deliberately did not choose a smaller angular size cutoff (as e.g. Blundell et al. 1998,


**Table 3.** USS samples

Sample	Sky Area	Density sr <sup>-1</sup>	Spectral Index	Search Radius	Flux Limit mJy	C <sup>a</sup>	R <sup>a</sup>	# of Sources
WN	$29^\circ < \delta < 75^\circ,  b  > 15^\circ$ <sup>b</sup>	151	$\alpha_{325}^{1400} \leq -1.30$	10''	$S_{1400} > 10$	96%	90%	343
TN	$-35^\circ < \delta < 29^\circ,  b  > 15^\circ$ <sup>b</sup>	48 <sup>c</sup>	$\alpha_{365}^{1400} \leq -1.30$	10''	$S_{1400} > 10$	97% <sup>c</sup>	93%	268
MP	$\delta < -30^\circ,  b  > 15^\circ$	26	$\alpha_{408}^{4800} \leq -1.20$	80''	$S_{408} > 700; S_{4850} > 35$	100%	100%	58

<sup>a</sup> C = completeness and R = reliability accounting only for scattering across the spectral index limit (see Sect. 3.3.1).

<sup>b</sup> Coverage is only 99.7% because some small patches of sky we not covered at the time of writing. They are listed on the NVSS homepage (1998 January 19 version).

<sup>c</sup> Because we selected only problem free sources from the Texas survey, the effective completeness of the TN sample is  $\sim 30\%$ .



**Fig. 6.** Sky coverage of our 3 USS samples. Constant declination lines denote the boundaries between our the WN and TN and between the TN and MP samples, as indicated on the right. Note the difference in source density and the exclusion of the Galactic Plane

did for the 6C\* sample), because (1) higher resolution angular size information is only available in the area covered by the FIRST survey, and (2) even a 15'' cutoff would only reduce the number of sources by 30%, while it would definitely exclude several HzRGs from the sample. For example, in the 4C USS sample (Chambers et al. 1996b), three out of eight  $z > 2$  radio galaxies have angular sizes  $> 15''$ .

We think that our  $\sim 1'$  angular size cutoff will exclude almost no HzRGs, because the largest angular size for  $z > 2$  radio galaxies in the literature is  $53''$  (4C 23.56 at  $z = 2.479$ ; Chambers et al. 1996a; Carilli et al. 1997), while all 45  $z > 2.5$  radio galaxies with good radio maps are  $< 35''$  (Carilli et al. 1997). Although the sample of known  $z > 2$  radio galaxies is affected by angular size selection effects, very few HzRGs larger than  $1'$  would be expected.

The main incompleteness of our USS sample stems from the spectral index cutoff and the flux limit (Sect. 3.2). However, our flux limit ( $S_{1400} = 10$  mJy) is low enough to break most of the redshift-radio power degeneracy at  $z > 2$ . To achieve this with flux limited samples, multiple samples are needed (e.g. Blundell et al. 1999).

## 3.2. Sample definition

### 3.2.1. WENSS-NVSS (WN) sample

A correlation of the WENSS and NVSS catalogs with a search radius of  $10''$  centered on the WENSS position (see Sect. 3.1.1) provides spectral indices for  $\sim 143\,000$  sources. Even with a very steep  $\alpha_{325}^{1400} \leq -1.30$  spectral index criterion, we would still have 768 sources in our sample. To facilitate follow-up radio observations, and to increase the accuracy of the derived spectral indices (see Sect. 3.3.1), we have selected only NVSS sources with  $S_{1400} > 10$  mJy. Because the space density of the highest redshift galaxies is low, it is important not to limit the sample area (see e.g. Rawlings et al. 1998) to further reduce the number of sources in our sample. Because the NVSS has a slightly higher resolution than the WENSS ( $45''$  compared to  $54'' \times 54'' \cos \delta$ ), some WENSS sources have more than one associated NVSS source. We have rejected the 11 WN sources that have a second NVSS source within one WENSS beam ( $54'' \times 54'' \cos \delta$ ), we have used a circular  $72''$  WENSS beam, corresponding to the major axis of the beam at  $\delta = 48^\circ$ , the position that divides the WN sample into equal numbers to the North and South. The final WN sample contains 343 sources.

### 3.2.2. Texas-NVSS (TN) sample

Because the Texas and NVSS both have a large sky-coverage, the area covered by the TN sample includes 90% of the WN area. In the region  $\delta > 29^\circ$ , we have based our sample on the WENSS, since it does not suffer from lobe-shift problems and reaches ten times lower flux densities than the Texas survey (Sect. 2.2). In the remaining 5.28 steradians South of declination  $+29^\circ$ , we have spectral indices for  $\sim 25\,200$  sources. Again, we used a  $10''$  search radius (see Sect. 3.1.1), and for the same reason as in the WN sample we selected only NVSS sources with  $S_{1400} > 10$  mJy. Combined with the  $\alpha_{365}^{1400} \leq -1.30$  criterion, the number of USS TN sources is 285. As for the WN sample, we further excluded sources with more than one  $S_{1400} > 10$  mJy NVSS source within  $60''$  around the TEXAS position, leaving 268 sources in the final TN sample. We remind (see Sect. 2.2) that the selection of the

TEXAS survey we used is only  $\sim 40\%$  complete with a strong dependence on flux density. Using the values from Table 2, we estimate that the completeness of our TN sample is  $\sim 30\%$ .

### 3.2.3. MRC-PMN (MP) sample

In the overlapping area, we preferred the TN over the MP sample for the superior positional accuracies and resolutions of both Texas and NVSS compared to MRC or PMN. Because the MRC survey has a low source density, we would have only 13 MP sources with  $\alpha_{408}^{4850} \leq -1.30$ . We therefore relaxed this selection criterion to  $\alpha_{408}^{4850} \leq -1.20$ , yielding a total sample of 58 sources in the deep South ( $\delta < -30^\circ$ ).

## 3.3. Discussion

### 3.3.1. Spectral index errors

We have listed the errors in the spectral indices due to flux density errors in the catalogs in Tables A.1 to A.3. The WN and TN samples have the most accurate spectral indices: the median spectral index errors are  $\Delta\bar{\alpha}_{325}^{1400} = 0.04$  for WN sources and  $\Delta\bar{\alpha}_{365}^{1400} = 0.04$  ( $S_{365} > 1$  Jy) to  $0.07$  ( $S_{365} > 150$  mJy) for TN sources. For the MP sample,  $\Delta\bar{\alpha}_{408}^{4850} \approx 0.1$ , with little dependence on flux density ( $S_{408} > 750$  mJy).

Because our sample selects the sources in the steep tail of the spectral index distribution (Figs. 7 and 8), there will be more sources with an intrinsic spectral index flatter than our cutoff spectral index that get scattered into our sample by measurement errors than there will be sources with intrinsic spectral index steeper than the cutoff that get scattered out of our sample.

Following the method of Rengelink (1998), we fitted the steep tail between  $-1.60 < \alpha < -1.0$  with a Gaussian function. For each of our three samples, we generated a mock sample drawn from this distribution, and added measurement errors by convolving this true spectral index distribution with a Gaussian distribution with as standard deviation the mean error of the spectral indices. The WN mock sample predicts that 13  $\alpha_{325}^{1400} < -1.30$  sources get scattered out of the sample while 36  $\alpha_{325}^{1400} > -1.30$  sources get scattered into the USS sample. Thus, the WN sample is 96% complete and 90% reliable. For the TN sample, we expect to lose 7  $\alpha_{365}^{1400} < -1.30$  sources<sup>5</sup>, and have 18 contaminating  $\alpha_{365}^{1400} > -1.30$  sources. The completeness is thus 97% and the reliability 93%. For the MP sample, this spectral index scattering is negligible, because there are too few sources in the steep spectral index tail.

<sup>5</sup> Only due to the spectral index cutoff, the sample has more important incompleteness factors; see Sect. 2.2.


Our reliability and completeness are significantly better than the values of  $\sim 75\%$  and  $\sim 50\%$  of Rengelink (1998) because (1) our spectral indices are more accurate because they were determined from a wider frequency interval than the 325 – 610 MHz used by Rengelink (1998), and (2) our sample has a steeper cutoff spectral index, where the spectral index distribution function contains fewer sources and has a shallower slope, leading to fewer sources that can scatter in or out of the sample.

### 3.3.2. Spectral index distributions

Using the 143 000 spectral indices from the WENSS-NVSS correlation, we examined the flux density dependence of the steep and flat spectrum sources. Selecting sources with  $S_{325} > 50$  mJy or  $S_{1400} > 100$  mJy assures that we will detect all sources with  $\alpha_{325}^{1400} > \frac{\ln(S_{NVSS}^{lim}/50)}{\ln(325/1400)} = -1.82$  or  $\alpha_{325}^{1400} < \frac{\ln(S_{WENSS}^{lim}/100)}{\ln(325/1400)} = 0.82$  respectively, where  $S_{NVSS}^{lim} = 3.5$  mJy and  $S_{WENSS}^{lim} = 30$  mJy are the lowest flux densities where the NVSS and WENSS are complete (Condon et al. 1998; Rengelink et al. 1997). The results shown in Fig. 7 therefore reflect only the effect of a different selection frequency. Two populations are present in both the  $S_{325}$  and  $S_{1400}$  selected distributions. The peaks of the steep and flat populations at  $\bar{\alpha}_{325}^{1400} \approx -0.8$  and  $\bar{\alpha}_{325}^{1400} \approx -0.4$  do not show significant shifts over three orders of magnitude in flux density. This is consistent with the results that have been found at 4.8 GHz (Witzel et al. 1979; Machalski & Ryś 1981; Owen et al. 1983), with the exception that their  $\bar{\alpha}_{1400}^{4800} \approx 0.0$  for the flat spectrum component is flatter than the  $\bar{\alpha}_{325}^{1400} \approx -0.4$  we found. However, we find that the relative contribution of the flat spectrum component increases from 25% at  $S_{1400} > 0.1$  Jy to 50% at  $S_{1400} > 2.5$  Jy.

Because the steep- and flat-spectrum populations are best separated in the  $S_{1400} > 2.5$  Jy bin, we have searched the literature for identifications of all 58  $S_{1400} > 2.5$  Jy sources to determine the nature of both populations. All but one (3C 399, Martel et al. 1998) of the objects outside of the Galactic plane ( $|b| > 15^\circ$ ) were optically identified. Of the 30 steep spectrum ( $\alpha_{325}^{1400} < -0.6$ ) sources, two thirds were galaxies, while the rest were quasars. Half of the flat spectrum ( $\alpha_{325}^{1400} > -0.6$ ) sources were quasars, 20% blazars, and 30% galaxies. Figure 7 therefore confirms that the steep and flat spectral index populations are dominated by radio galaxies and quasars respectively. We also find that while the relative strength between the steep and flat spectrum populations changes due to the selection frequency, the median spectral index and width of the population does not change significantly over three orders in magnitude of flux density. Even fainter studies would eventually start to get contamination from the faint blue galaxy population (see e.g. Windhorst et al. 1985).






**Fig. 7.** Spectral index distributions from the WENSS-NVSS correlation. The left and right panels show the variation with 325 MHz and 1.4 GHz flux density. A low frequency selected sample is more appropriate to study the steep-spectrum population. The parameters of a two-component Gaussian fit (dotted line = steep, dashed line = flat) are shown in each panel. The solid line is the sum of both Gaussians

**Table 4.** Radio observations

UT Date	Telescope	Config.	Frequency	Resolution	# of sources
1996 October 28	VLA	A	4.86 GHz	0''3	90 WN, 25 TN
1997 January 25	VLA	BnA	4.86 GHz	0''6	29 TN
1997 March 10	VLA	BnA	4.885 GHz	0''6	8 TN
1997 December 15	ATCA	6C	1.420 GHz	6'' × 6'' cosec δ	41 MP, 32 TN
1998 August 12+17	VLA	B	4.86 GHz	1''0	151 WN



**Fig. 8.** Logarithmic spectral index distribution for WENSS-NVSS (full line), Texas-NVSS (dot-dash line) and MRC-PMN (dotted line). The vertical line indicates the  $-1.3$  cutoff used in our spectral index selection. Note the difference in number density and the sharper fall-off on the flat-end part of the TN and MP compared to the WN

### 3.3.3. Consistency of the three USS samples

We compare the spectral index distributions of our three USS samples in logarithmic histograms (Fig. 8). The distributions are different in two ways. First, the WENSS-NVSS correlation contains nine times more sources than the Texas-NVSS, and 14 times more than the MRC-PMN correlation. Second, the shapes of the distributions are different: while the steep side of the TN sample coincides with that of the WN, its flat end part falls off much faster. The effect is so strong that it even shifts the TN peak steep-wards by  $\sim 0.15$ . For the MP sample, the same effect is less pronounced, though still present.

Both effects are due to the different flux density limits of the catalogs. The deeper WENSS catalog obviously contains more sources than the TEXAS or MRC catalogs, shifting the distributions vertically in Fig. 2. The relative “shortage” of flat spectrum sources in the Texas-NVSS and MRC-PMN correlations can be explained as follows. A source at the flux density limit in both WENSS and NVSS would have a spectral index of  $\alpha_{325}^{1400} = -1.3$ , while for Texas and NVSS this would be  $\alpha_{365}^{1400} = -1.7$  (see Fig. 2). Faint NVSS sources with spectral indices flatter than these limits will thus more often get missed in the TEXAS catalog than in the WENSS catalog. This effect is even strengthened by the lower completeness at low flux densities of the Texas catalog. However, very few USS sources will be missed in either the WENSS-NVSS or Texas-NVSS correlations<sup>6</sup>. The parallel slope also indicates that the USS sources from both the WENSS-NVSS and Texas-NVSS correlations were drawn from the same

<sup>6</sup> The TN sample will have more spurious sources at low flux density levels; see Sect. 2.2.

population of radio sources. We therefore expect a similar efficiency in finding HzRGs from both samples.

The MP sample has been defined using a spectral index with a much wider frequency difference. However, the observed ATCA 1.420 GHz flux densities can be used to construct  $\alpha_{408}^{1420}$ . An “a posteriori” selection using  $\alpha_{408}^{1420} \leq -1.30$  from our ATCA observations (see Sect. 4.2) would keep  $\sim 60\%$  of the MP sources in a WN/TN USS sample.

## 4. Radio observations

Of all the major radio surveys described in Sect. 2, only FIRST has sufficient positional accuracy and resolution for the optical identification of  $R > 20$  objects. We present FIRST maps of 139 WN and 8 TN sources in Appendices B.2 and B.4.

Outside the area covered by FIRST, we have observed all the remaining WN sources, 30% of the TN sample, and 71% of the MP sources at  $0''.3$  to  $5''$  resolution using the Very Large Array (VLA; Napier et al. 1983) and Australia Telescope Compact Array (ATCA; Frater et al. 1992) telescopes. A log of the radio observations is given in Table 4. We observed targets for our VLA runs on the basis of declination (A-array for  $\delta > 0^\circ$  and BnA-array for  $\delta < 0^\circ$ ) and sky coverage of the WN and TN samples, which were still incomplete at the time of the 1996 observations. We observed all WN, and most TN sources with the VLA, and all MP sources with the ATCA. We observed TN sources between  $-31^\circ < \delta < -10^\circ$  with either VLA or ATCA, depending on the progress of the NVSS at the time of the observations.

### 4.1. VLA observations and data reduction

We observed all sources in the standard 4.86 GHz C-band with a 50 MHz bandwidth, resulting in a resolution of  $\sim 0''.3$  in the A-array and  $\sim 1''$  in the BnA-array. We spent 5 minutes on each source, implying a theoretical rms level of  $75 \mu\text{Jy}$ , or a ratio of total integrated signal over map noise of 110 for the weakest sources, assuming no spectral curvature beyond 1.4 GHz. We performed calibration and data editing in *AIPS*, the Astronomical Image Processing System from NRAO. We used 3C 286 as the primary flux calibrator in all runs. Comparison of the flux density of 3C 48 with the predicted values indicated the absolute flux density scale was accurate up to 2%. We observed nearby (within  $15^\circ$ ) secondary flux calibrators every 15 to 20 minutes to calibrate the phases. After flagging of bad data, we split the *uv*-data up into separate data sets for imaging and self-calibration in DIFMAP, the Caltech difference mapping program (Sheppard et al. 1997). We used field sizes of  $164''$  (A-array) or  $256''$  (BnA-array) with pixel scales of  $0''.08/\text{pixel}$  (A-array) or  $0''.25/\text{pixel}$

(BnA–array). Even the smallest field of view is still four times larger than the resolution of the NVSS, so all components of an unresolved NVSS source will be covered.

We cleaned each source brighter than the  $5\sigma$  level, followed by a phase-only self-calibration. We repeated the latter for all sources in the field of a source. Next, we made a new model from the (self-calibrated)  $uv$ -data, and subsequently cleaned to the level reached before. The last stage in the mapping routine was a deep clean with a 1% gain factor over the entire field. Most of the resulting maps have noise levels in the range 75 to 100  $\mu\text{Jy}$ , as expected.

#### 4.2. ATCA observations and data reduction

We used the ATCA in the 6C configuration, which has a largest baseline of 6 km. We observed at a central frequency of 1.384 GHz, which was selected to avoid local interference. We used 21 of the 30 frequency channels that had high enough signal, which resulted in an effective central frequency of 1.420 GHz, with a 84 MHz bandwidth. In order to obtain a good  $uv$ -coverage, we observed each source eight to ten times for three minutes, spread in hour angle. The primary flux calibrator was the source 1934-648; we used secondary flux calibrators within  $20^\circ$  of the sources to calibrate the phases. We performed editing and calibration in *ATPS*, following standard procedures. We made maps using the automated mapping/self-calibration procedure MAPIT in *ATPS*. The resulting 1.420 GHz maps (Fig. B.6.) have noise levels of  $\sim 5$  mJy.

#### 4.3. Results

Of all 343 WN sources, 139 have FIRST maps (Appendix B.2). All remaining 204 sources were observed, and 141 were detected. The remaining 30% were too faint at 4.86 GHz to be detected in 5 min snapshots, because their high frequency spectral index steepens more than expected, or they were over-resolved. Because they are significantly brighter, all the observed 89 TN and 41 MP sources were detected. We present contour maps of all the detections in Appendices B.1, B.3, B.5, and B.6 and list the source parameters in Tables A.1 to A.3.

We have subdivided our sources into 5 morphological classes, using a classification similar to that used by Röttgering et al. (1994). Note that this classification is inevitably a strong function of the resolution, which varies by a factor of 20 between the VLA A–array and the ATCA observations.

We have determined the source parameters by fitting two-dimensional Gaussian profile to all the components of a source. The results are listed in Tables A.1 to A.3 which contain:

*Column 1:* Name of the source in IAU J2000 format. The 2-letter prefix indicates the sample: WN: WENSS–NVSS, TN: Texas–NVSS, MP: MRC–PMN.

*Column 2:* The integrated flux density from the low-frequency catalog.

*Column 3:* The integrated flux density at the intermediate frequency, determined from the NVSS for WN and TN, or from the 1.420 GHz ATCA observations for the MP sample.

*Column 4:* The integrated flux density at 4.86 GHz, determined from the VLA observations for WN and TN, and from the PMN survey for the MP sample.

*Column 5:* The lower frequency two-point spectral index. This is the spectral index used to define the WN and TN samples.

*Column 6:* The higher frequency two-point spectral index. This is the spectral index used to define the MP sample.

*Column 7:* Morphological classification code: single (S), double (D), triple (T) and multiple (M) component sources, and irregularly shaped diffuse (DF) sources.

*Column 8:* Largest angular size. For single component sources, this is the de-convolved major axis of the elliptical Gaussian, or, for unresolved sources (preceded with <), an upper limit is given by the resolution. For double, triple and multiple component sources, this is the largest separation between their components. For diffuse sources this is the maximum distance between the source boundaries defined by three times the map rms noise.

*Column 9:* De-convolved position angle of the radio structure, measured North through East.

*Columns 10–11:* J2000 coordinates, determined from the map with position code listed in Col. 12. The positions in the VLA and ATCA maps have been fitted with a single two-dimensional elliptical Gaussian. For double (D) sources, the geometric midpoint is given; for triples (T) and multiples (M), the core position is listed. For diffuse (DF) sources we list the center as determined by eye.

*Column 12:* Position code, indicating the origin of the morphological and positional data in Cols. 7 to 11: A = ATCA, F = FIRST, M = MRC, N = NVSS, and V = VLA.

#### 4.4. Notes on individual sources

**WN J0043+4719:** The source  $18''$  north of the NVSS position is not detected in the NVSS. This is therefore not a real USS source because the NVSS flux density was underestimated.

**WN J0048+4137:** Our VLA map probably doesn't go deep enough to detect all the flux of this source.

**WN J0727+3020:** The higher resolution FIRST map shows that both components of this object are indeed identified on the POSS, even though the NVSS position is too far off to satisfy our identification criterion.

**WN J0717+4611:** Optical and near-IR spectroscopy revealed this object as a red quasar at  $z = 1.462$  (De Breuck et al. 1998b).

**WN J0725+4123:** The extended POSS identification suggests this source is located in a galaxy cluster.

**WN J0829+3834:** The NVSS position of this unresolved source is  $7''$  ( $3\sigma$ ) from the FIRST position, which itself is only at  $2''$  from the WENSS position.

**WN J0850+4830:** The difference with the NVSS position indicates that our VLA observations are not deep enough to detect a probable north-eastern component.

**WN J0901+6547:** This  $38''$  large source is over-resolved in our VLA observations, and probably even misses flux in the NVSS, and is therefore not a real USS source.

**WN J1012+3334:** The bend morphology and bright optical sources to the east indicate this object is probably located in a galaxy cluster.

**WN J1101+3520:** The faint FIRST component  $20''$  north of the brighter Southern component is not listed in the FIRST catalog, but is within  $1''$  of a faint optical object. This might be the core of a  $70''$  triple source.

**WN J1152+3732:** The distorted radio morphology and bright, extended POSS identification suggest this source is located in a galaxy cluster.

**WN J1232+4621:** This optically identified and diffuse radio source suggest this source is located in a galaxy cluster.

**WN J1314+3515:** The diffuse radio source appears marginally detected on the POSS.

**WN J1329+3046A,B, WN J1330+3037, WN J1332+3009 & WN J1333+3037:** The noise in the FIRST image is almost ten times higher than average due to the proximity of the  $S_{1400}=15$  Jy source 3C 286.

**WN J1330+5344:** The difference with the NVSS position indicates that our VLA observations are not deep enough to detect a probable south-eastern component.

**WN J1335+3222:** Although the source appears much like the hotspot of a larger source with the core  $90''$  to the east, no other hotspot is detected in the FIRST within  $5'$ .

**WN J1359+7446:** The extended POSS identification suggests this source is located in a galaxy cluster.

**WN J1440+3707:** The equally bright galaxy  $30''$  south of the POSS identification suggests that this source is located in a galaxy cluster.

**WN J1509+5905:** The difference with the NVSS position indicates that our VLA observations are not deep enough to detect a probable western component.

**WN J1628+3932:** This is the well studied galaxy NGC 6166 in the galaxy cluster Abell 2199 (e.g. Zabludoff et al. 1993).

**WN J1509+5905:** The difference with the NVSS position indicates that our VLA observations are not deep enough to detect a probable west-south-western component.

**WN J1821+3601:** The source  $35''$  south-west of the NVSS position is not detected in the NVSS. This is therefore not a real USS source because the NVSS flux density was underestimated.

**WN J1832+5354:** The source  $19''$  north-east of the NVSS position is not detected in the NVSS. This is therefore not a real USS source because the NVSS flux density was underestimated.

**WN J1852+5711:** The extended POSS identification suggests this source is located in a galaxy cluster.

**WN J2313+3842:** The extended POSS identification suggests this source is located in a galaxy cluster.

**TN J0233+2349:** This is probably the north-western hotspot of a  $35''$  source, with the south-eastern component barely detected in our VLA map.

**TN J0309–2425:** We have classified this source as a  $13''$  double, but the western component might also be the core of a  $45''$  source, with the other hotspot around  $\alpha = 3^{\text{h}}9^{\text{m}}10^{\text{s}}, \delta = -24^{\circ}25'50''$ .

**TN J0349–1207:** The core-dominated structure is reminiscent of the red quasar WN J0717+4611.

**TN J0352–0355:** This is probably the south-western hotspot of a  $30''$  source.

**TN J0837–1053:** Given the  $10''$  difference between the positions of the NVSS and diffuse VLA source, this is probably the northern component of a larger source.

**TN J0408–2418:** This is the  $z = 2.44$  source MRC 0406–244 (McCarthy et al. 1996). The bright object on the POSS is a foreground star to the north-east of the  $R = 22.7$  galaxy.


**TN J0443–1212:** Using the higher resolution VLA image, we can identify this radio source with a faint object on the POSS.

**TN J2106–2405:** This is the  $z = 2.491$  source MRC 2104–242 (McCarthy et al. 1996). The identification is an  $R = 22.7$  object, not the star to the north-north-west of the NVSS position.

#### 4.5. Radio spectra and spectral curvature

We have used the CATS database at the Special Astronomical Observatory (Verkhodanov et al. 1997) to search for all published radio measurements of the sources in our samples. In Appendix C, we show the radio spectra for all sources with flux density information for more than two frequencies (the  $S_{4860}$  points from our VLA observations are also included). These figures show that most radio spectra have curved spectra, with flatter spectral indices below our selection frequencies, as has been seen in previous USS studies (see e.g. Röttgering et al. 1994; Blundell et al. 1998).

This low frequency flattening and high frequency steepening is obvious in the radio “color-color diagrams” of the WN sample (Fig. 9). The median spectral index at low frequencies ( $\nu < 325$  MHz) is  $-1.16$ , while the median  $\bar{\alpha}_{325}^{1400} = -1.38$ . At higher frequencies ( $\nu > 1400$  MHz), the steepening continues to a median  $\bar{\alpha}_{1400}^{4850} = -1.44$ . Note that the real value of the latter is probably even



**Fig. 9.** Radio “color-color” plots for the WN sample. The abscissa is the  $\alpha_{325}^{1400}$  spectral index used to construct the sample. The ordinates are the low-frequency spectral indices determined from the 8C (38 MHz, Rees 1990) or 6C (151 MHz, Hales et al. 1993) and the 325 MHz WENSS (left panel), and the high frequency spectral index determined between the 1.4 GHz NVSS and our 4.86 GHz VLA observations (right panel). The line in each panel indicates a straight power law spectrum. Note the unequal number of points on either side of these lines, indicating substantial spectral curvature

**Table 5.** Radio structure distribution

Morphology	USS Samples			
	WN	TN	MP	Combined
Single	157 (56 ± 4%)	43 (48 ± 7%)	23 (56 ± 12%)	223 (54 ± 4%)
Double	81 (29 ± 3%)	28 (31 ± 6%)	16 (39 ± 10%)	125 (31 ± 3%)
Triple	22 (8 ± 2%)	9 (10 ± 3%)	0 (0 ± 0%)	31 (8 ± 1%)
Multiple	2 (1 ± 1%)	4 (5 ± 2%)	0 (0 ± 0%)	6 (1 ± 1%)
Diffuse	18 (6 ± 2%)	5 (6 ± 3%)	2 (5 ± 3%)	25 (6 ± 1%)
# Observed	280	89	41	410


steeper, as 30% of the WN sources were not detected in our 4.86 GHz VLA observations, and may therefore have even more steepened high-frequency spectral indices.

#### 4.6. Radio source properties

##### 4.6.1. Radio source structure and angular size

In Table 5, we give the distribution of the radio structures of the 410 USS sources for which we have good radio-maps. At first sight, all three our samples have basically the same percentage of resolved sources, but the similar value for the MP sample is misleading, as it was observed at much lower resolution.

Our results are different from the USS sample of Röttgering et al. (1994), which contains only 18% unresolved sources at comparable resolution (1''). To check if this effect is due to the fainter sources in our sample, we compared our sample with the deep high resolution



**Fig. 10.** Median angular size for the flux density limited, spectrally unbiased WSRT samples of Oort 1988 (open triangles), and for our combined USS samples (filled squares). The sources have been binned in equal number bins, and errors represent the 35% and 65% levels of the distribution. Note that our USS selection does not affect the value of the median, and that our USS samples also exclude sources that fall below the break at  $S_{1400} \lesssim 10$  mJy

VLA observations of spectrally unbiased sources (Oort 1988; Coleman & Condon 1985). The resolution of our observations is significantly better than the median angular size for  $S_{1400} > 1$  mJy sources, allowing us to accurately determine the median angular sizes in our samples. We find that our USS sources have a constant median angular size of  $\sim 6''$  between 10 mJy and 1 Jy (Fig. 10). This is indistinguishable from the results from samples without spectral index selection. It indicates that our USS selection of sources with  $\alpha < -1.3$  and  $\Theta < 1'$  does not bias the angular size distribution in the resulting sample. The “downturn” in angular sizes that occurs at  $\sim 1$  mJy is probably due to a different radio source population, which consist of lower redshift sources in spiral galaxies (see e.g., Coleman & Condon 1985; Oort et al. 1987; Benn et al. 1993). By selecting only sources with  $S_{1400} > 10$  mJy, we have avoided “contamination” of our sample by these foreground sources.

We have searched for further correlations between spectral index or spectral curvatures and angular size or flux density, but found no significant results, except for a trend for more extended sources to have lower than expected 4.86 GHz flux densities, but this effect can be explained by missing flux at large scales in our VLA observations.

## 4.7. Identifications

### 4.7.1. POSS


We have searched for optical identifications of our USS sources on the digitized POSS-I. We used the likelihood ratio identification criterion as described by e.g. de Ruiter et al. (1977). In short, this criterion compares the probability that a radio and optical source with a certain positional difference are really associated with the probability that this positional difference is due to confusion with a field object (mostly a foreground star), thereby incorporating positional uncertainties in both radio and optical positions. The ratio of these probabilities is expressed as the likelihood ratio  $LR$ . In the calculation, we have assumed a density of POSS objects  $\rho = 4 \cdot 10^{-4''^{-2}}$ , independent of galactic latitude  $b$ . We have adopted a likelihood ratio cutoff  $\mathcal{L} = 1.0$ , slightly lower than the values used by de Ruiter et al. (1977) and Röttgering et al. (1994). We list sources with  $LR > 1.0$  for our USS samples in Tables A.4 to A.6. We have included four WN sources (WN J0704+6318, WN J1259+3121, WN J1628+3932 and WN J2313+3842), two TN sources (TN J0510–1838 and TN J1521 + 0741) and four MP sources (MP J0003–3556, MP J1921–5431, MP J1943–4030 and MP J2357–3445) as identifications because both their optical and radio morphologies are diffuse and overlapping, making it impossible to measure a common radio and optical component, while they are very likely to be associated.

Figure 11 shows the identification fraction of USS sources on the POSS ( $R \lesssim 20$ ). Because the distributions for the WN and TN are very similar, we have combined both samples to calculate the identification fraction. Unlike the results for 4C USS (Tielens et al. 1979; Blumenthal & Miley 1979), we do not detect a decrease of the identification fraction with steepening spectral index<sup>7</sup>. We interpret the constant  $\sim 15\%$  identification fraction from our sample as a combined population of foreground objects, primarily consisting of clusters (see next section). Our extremely steep spectral index criterion would then select only radio galaxies too distant to be detected on the POSS ( $R \gtrsim 20.0$ ).

### 4.7.2. Literature

Using the NASA Extragalactic Database (NED), the SIMBAD database and the W<sup>3</sup>Browse at the High Energy Astrophysics Science Archive Research Center, we have searched for known optical and X-ray identifications

<sup>7</sup> In the Westerbork faint USS (Wieringa & Katgert 1992) or the USS sample from Röttgering et al. (1994), there is also a decrease in the identification fraction, even at limiting magnitudes of  $R = 22.5$  and  $R = 23.7$ , indicating that this trend continues out to fainter magnitudes and radio fluxes.




**Fig. 11.** Identification fraction on the POSS as a function of spectral index for the combined WN and TN sample. Note the absence of a further decrease in the identification fraction with steepening spectral index

of sources in our samples (see Appendices A.7 to A.9). Of the bright optical ( $R \lesssim 20$ ) identifications, only one source is known as a K0-star, three (TN J0055+2624, TN J0102–2152, and TN J1521+0742) are “Relic radio galaxies” (Komissarov & Gubanov 1994; Giovannini et al. 1999), while all others are known galaxy clusters.

All optical cluster identifications, except MP J1943–4030, are also detected in the ROSAT All-Sky survey Bright Source Catalogue (RASS-BSC; Voges et al. 1999). Conversely, of the 23 X-ray sources, seven are known galaxy clusters, and three known galaxies. The remaining 13 sources are good galaxy cluster candidates because they either show a clear over-density of galaxies on the POSS (eight sources), or they have low X-ray count rates ( $< 0.02$  counts  $s^{-1}$ ), suggesting that these might be more distant galaxy clusters too faint to be detected on the POSS. We conclude that probably  $>3\%$  of our USS sources are associated with galaxy clusters, and that the combined USS + X-ray selection is an efficient (up to 85%) selection technique to find galaxy clusters<sup>8</sup>.

Three of our USS sources (WN J2313+4253, TN J0630–2834 and TN J1136+1551) are previously known pulsars (Kaplan et al. 1998). It is worth noting that two out of nine sources in our USS samples with  $\alpha < -2$  are known pulsars. Because Lorimer et al. (1995) found the median spectral index of pulsars to

<sup>8</sup> In the RASS-BSC, only 14% of the extra-galactic sources are identified with galaxy clusters (Voges et al. 1999).



**Fig. 12.** Percentage of  $\alpha_{325}^{1400} < -1.60$  radio sources from a WENSS–NVSS correlation as a function of Galactic latitude. Note the clear peak near the Galactic plane, indicating that these  $\alpha_{325}^{1400} < -1.60$  objects might well be Galactic pulsars

be  $\sim -1.6$ , we examined the distribution of spectral indices as a function of Galactic latitude. In Fig. 12, we plot the percentage of  $\alpha_{325}^{1400} < -1.60$  pulsar candidates as a function of Galactic latitude. The four times higher density near the Galactic plane strongly suggests that the majority of these  $\alpha_{325}^{1400} < -1.60$  sources are indeed pulsars, which are confined to our Galaxy. A sample of such  $\alpha_{325}^{1400} < -1.60$  sources at  $|b| < 15^\circ$  would be an efficient pulsar search method.

We also note that no known quasars are present in our sample. Preliminary results from our optical spectroscopy campaign (De Breuck et al. 1998b, 2000) indicate that  $\sim 10\%$  of our sample are quasars. We interpret this lack of previously known quasars as a selection bias in quasar samples against USS sources.

At  $R \gtrsim 20$ , all five USS sources with known redshift are HzRGs, indicating a selection of sources without detections on the POSS strongly increases our chances of finding HzRGs.

## 5. Conclusions

We have constructed three spatially separated samples of USS sources containing a total of 669 objects. High-resolution radio observations of more than half of these show that the median size is  $\sim 6''$ , independent of 1.4 GHz flux density, which is consistent with results

of similar resolution surveys of samples without spectral index selection. The absence of a downturn in angular size at the lowest fluxes indicates that we do not include significant numbers of spiral galaxies in our sample. A USS sample fainter than ours would therefore include more of these foreground sources, and be less efficient to find HzRGs.

The identification fraction on the POSS is  $\sim 15\%$ , with no clear dependence on spectral index, indicating that the HzRGs in the sample are all too distant to be detected, and the POSS detections consist of different classes of objects. A correlation of our USS samples with X-ray catalogs showed that at least 85% of the X-ray identifications seem to be galaxy clusters known from the literature or by inspection of the POSS. We conclude that (1) the majority of the “non HzRG” USS sources in our sample are clusters, and (2) the combined selection of USS and X-ray sources is an extremely efficient technique to select galaxy clusters.

The above results indicate that up to 85% of our USS sources might be HzRGs. To identify these objects, we have started an intensive program of  $R$ - and  $K$ -band imaging on 3 – 10 m class telescopes. Initial results from optical spectroscopy indicate that 2/3 are indeed  $z > 2$  radio galaxies (De Breuck et al. 1998a), and  $K$ -band imaging of optically undetected ( $R > 25$ ) sources (see e.g., van Breugel et al. 1999a) has already lead to the discovery of the first radio galaxy at  $z > 5$  (van Breugel et al. 1999b).

*Acknowledgements.* We are grateful for the excellent help provided by the staff of the VLA and ATCA observatories, with special thanks to Chris Carilli and Greg Taylor (NRAO), and Ray Norris and Kate Brooks (ATNF) for help in with observation planning and data reduction. We thank Hien “Napkin” Tran for his comments on the manuscript. The VLA is a facility of the National Radio Astronomy Observatory, which is operated by Associated Universities Inc. under cooperative agreement with the National Science Foundation. The Australia Telescope is funded by the Commonwealth of Australia for operation as a National Facility managed by CSIRO. The authors made use of the database CATS (Verkhodanov et al. 1997) of the Special Astrophysical Observatory, the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration, and the High Energy Astrophysics Science Archive Research Center Online Service, provided by the NASA/Goddard Space Flight Center. The work by C.D.B. and W.V.B. was performed under auspices of the U.S. Department of Energy by University of California Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

## Appendix A: Source lists

Table A.1. WN sample

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J0000+4654	204 ± 9	21.1 ± 1.4	2.97 ± 0.13	-1.55 ± 0.06	-1.58 ± 0.06	D	3.6	173	00 00 28.88	+46 54 40.7	V
WN J0007+3641	447 ± 18	28.8 ± 1.7	2.51 ± 0.11	-1.88 ± 0.05	-1.96 ± 0.06	S	1.7	67	00 07 02.92	+36 41 55.9	V
WN J0007+4615	144 ± 7	16.9 ± 0.5	...	-1.47 ± 0.04	...	...	...	...	00 07 15.05	+46 15 40.8	N
WN J0016+4311	105 ± 6	15.7 ± 1.5	...	-1.30 ± 0.08	...	...	...	...	00 16 40.63	+43 11 03.5	N
WN J0029+3439	288 ± 12	40.9 ± 2.0	44.76 ± 1.05	-1.34 ± 0.04	0.07 ± 0.04	DF	12.8	80	00 29 01.61	+34 39 34.7	V
WN J0034+4142	84 ± 6	11.0 ± 0.5	2.50 ± 0.12	-1.39 ± 0.05	-1.19 ± 0.05	D	23.7	154	00 34 29.21	+41 42 09.5	V
WN J0034+3238	120 ± 9	16.4 ± 0.5	...	-1.36 ± 0.04	...	...	...	...	00 34 56.36	+32 38 36.4	N
WN J0038+7859	109 ± 6	15.1 ± 0.5	1.30 ± 0.09	-1.35 ± 0.04	-1.97 ± 0.06	S	2.5	159	00 38 01.96	+78 59 51.5	V
WN J0040+3857	127 ± 7	17.5 ± 0.5	3.13 ± 0.14	-1.36 ± 0.04	-1.38 ± 0.04	S	1.4	31	00 40 56.23	+38 57 30.0	V
WN J0043+4719 <sup>†</sup>	99 ± 6	14.2 ± 1.3	41.81 ± 1.01	-1.33 ± 0.08	0.87 ± 0.08	T	18.5	157	00 43 53.13	+47 19 48.3	V
WN J0048+4137 <sup>*</sup>	277 ± 11	34.5 ± 1.9	1.75 ± 0.10	-1.43 ± 0.05	-2.40 ± 0.06	S	2.5	17	00 48 46.34	+41 37 20.8	V
WN J0059+3958	152 ± 7	11.4 ± 1.4	...	-1.77 ± 0.09	...	...	...	...	00 59 51.07	+39 58 46.5	N
WN J0112+4039	111 ± 7	15.4 ± 0.5	...	-1.35 ± 0.05	...	...	...	...	01 12 26.31	+40 39 30.1	N
WN J0117+3715	165 ± 8	22.5 ± 0.6	6.62 ± 0.21	-1.26 ± 0.04	-0.98 ± 0.03	S	1.4	71	01 17 10.02	+37 15 16.3	V
WN J0121+4305	389 ± 16	50.8 ± 2.5	...	-1.39 ± 0.04	...	...	...	...	01 21 10.71	+43 05 18.3	N
WN J0135+3303	94 ± 6	12.1 ± 0.5	10.22 ± 0.28	-1.40 ± 0.06	-0.14 ± 0.04	T	1.8	18	01 35 07.24	+33 03 47.4	V
WN J0137+3250	177 ± 9	23.8 ± 1.6	16.06 ± 0.40	-1.37 ± 0.06	-0.32 ± 0.06	T	27.9	104	01 36 59.85	+32 50 40.7	V
WN J0155+8036	338 ± 14	38.0 ± 0.9	6.87 ± 0.21	-1.50 ± 0.03	-1.38 ± 0.03	D	4.6	159	01 55 43.77	+80 36 48.1	V
WN J0207+3655	118 ± 6	17.5 ± 1.4	...	-1.31 ± 0.07	...	...	...	...	02 07 22.59	+36 55 16.0	N
WN J0231+3600	303 ± 13	45.2 ± 2.2	7.66 ± 0.22	-1.30 ± 0.04	-1.43 ± 0.05	D	14.8	59	02 31 11.48	+36 00 26.6	V
WN J0240+3526	79 ± 5	10.2 ± 0.5	...	-1.40 ± 0.06	...	...	...	...	02 40 31.10	+35 26 55.1	N
WN J0303+3733	1071 ± 43	127.9 ± 5.7	17.32 ± 0.43	-1.46 ± 0.04	-1.61 ± 0.04	D	4.4	3	03 03 26.01	+37 33 41.6	V
WN J0303+3629	117 ± 6	17.1 ± 0.5	...	-1.32 ± 0.05	...	...	...	...	03 03 29.87	+36 29 56.1	N
WN J0305+3525	110 ± 6	15.8 ± 1.3	3.74 ± 0.15	-1.33 ± 0.07	-1.16 ± 0.07	S	1.9	64	03 05 47.42	+35 25 13.4	V
WN J0310+3644	289 ± 11	24.2 ± 0.6	2.45 ± 0.12	-1.70 ± 0.03	-1.84 ± 0.05	S	2.0	118	03 10 54.80	+36 44 02.5	V
WN J0315+3757	95 ± 5	13.3 ± 0.5	4.74 ± 0.17	-1.35 ± 0.05	-0.83 ± 0.04	S	1.7	53	03 15 17.29	+37 57 05.0	V
WN J0323+3738	205 ± 9	22.6 ± 1.5	8.11 ± 0.25	-1.51 ± 0.06	-0.83 ± 0.06	DF	9.7	100	03 23 38.24	+37 38 39.3	V
WN J0343+7540	118 ± 6	12.5 ± 1.6	...	-1.54 ± 0.09	...	...	...	...	03 43 39.77	+75 40 43.2	N
WN J0346+3039	286 ± 12	35.6 ± 1.9	4.95 ± 0.18	-1.43 ± 0.05	-1.59 ± 0.05	D	0.4	89	03 46 42.60	+30 39 51.0	V
WN J0352+3143	154 ± 7	21.8 ± 1.5	...	-1.34 ± 0.06	...	...	...	...	03 52 24.49	+31 43 03.1	N
WN J0359+3000	88 ± 6	12.1 ± 0.5	...	-1.36 ± 0.06	...	...	...	...	03 59 12.86	+30 00 51.7	N
WN J0528+6549	96 ± 6	10.6 ± 2.4	1.08 ± 0.09	-1.51 ± 0.16	-1.84 ± 0.19	S	1.9	108	05 28 46.07	+65 49 57.3	V
WN J0533+7205	117 ± 7	17.4 ± 1.6	2.42 ± 0.12	-1.30 ± 0.08	-1.59 ± 0.08	D	2.9	140	05 33 38.64	+72 05 52.7	V
WN J0538+7348	132 ± 7	18.1 ± 0.5	3.20 ± 0.14	-1.36 ± 0.04	-1.39 ± 0.04	S	2.1	113	05 38 25.59	+73 48 39.6	V
WN J0559+6926	87 ± 5	12.9 ± 0.5	1.88 ± 0.13	-1.31 ± 0.06	-1.55 ± 0.06	S	0.3	85	05 59 06.04	+69 26 37.8	V
WN J0610+6611	98 ± 6	12.5 ± 1.9	1.19 ± 0.09	-1.41 ± 0.11	-1.89 ± 0.14	S	2.2	114	06 10 08.59	+66 11 43.1	V
WN J0617+5012	196 ± 9	26.5 ± 0.7	6.80 ± 0.23	-1.37 ± 0.04	-1.09 ± 0.03	D	3.4	11	06 17 39.37	+50 12 54.7	V
WN J0625+5708	71 ± 4	10.3 ± 0.5	1.54 ± 0.10	-1.32 ± 0.07	-1.53 ± 0.06	D	7.3	18	06 25 01.67	+57 08 26.1	V
WN J0627+7311	470 ± 19	61.9 ± 2.9	4.00 ± 0.17	-1.39 ± 0.04	-2.20 ± 0.05	S	< 0.3	0	06 27 03.64	+73 11 54.2	V
WN J0633+6024	115 ± 6	13.4 ± 2.8	2.07 ± 0.11	-1.47 ± 0.15	-1.50 ± 0.17	S	2.2	118	06 33 19.70	+60 24 37.1	V
WN J0633+4653	3661 ± 146	520.1 ± 22.8	80.88 ± 1.74	-1.34 ± 0.04	-1.50 ± 0.04	T	4.0	19	06 33 52.18	+46 53 40.5	V
WN J0641+4325	154 ± 7	15.6 ± 0.5	...	-1.57 ± 0.04	...	...	...	...	06 41 08.23	+43 25 00.3	N
WN J0646+3912	166 ± 8	20.5 ± 2.4	...	-1.43 ± 0.09	...	...	...	...	06 46 34.97	+39 12 00.3	N
WN J0648+4309	215 ± 9	30.4 ± 1.7	5.00 ± 0.19	-1.34 ± 0.05	-1.45 ± 0.05	DF	1.5	151	06 48 01.10	+43 09 47.8	V
WN J0648+4137	91 ± 5	10.9 ± 1.5	1.69 ± 0.11	-1.45 ± 0.11	-1.50 ± 0.12	S	13.2	94	06 48 14.58	+41 37 18.8	V
WN J0650+4106	393 ± 16	53.5 ± 1.2	15.04 ± 0.41	-1.37 ± 0.03	-1.02 ± 0.03	DF	3.0	26	06 50 41.66	+41 06 39.1	V
WN J0653+4434	115 ± 6	13.1 ± 0.5	...	-1.49 ± 0.05	...	...	...	...	06 53 23.84	+44 34 55.4	N
WN J0656+5301	137 ± 7	18.7 ± 1.5	...	-1.36 ± 0.07	...	D	56.1	88	06 56 38.40	+53 01 18.1	F
WN J0658+4444	407 ± 17	50.6 ± 2.4	13.91 ± 0.35	-1.43 ± 0.04	-1.04 ± 0.04	S	1.7	73	06 58 27.43	+44 44 16.4	V
WN J0702+4448	147 ± 7	17.0 ± 0.5	...	-1.48 ± 0.04	...	S	7.8	119	07 02 16.79	+44 48 59.7	F
WN J0711+4242	140 ± 6	16.0 ± 0.5	...	-1.49 ± 0.04	...	S	3.3	17	07 11 40.25	+42 42 09.6	F
WN J0715+4655	89 ± 6	10.4 ± 0.5	1.86 ± 0.13	-1.47 ± 0.06	-1.38 ± 0.07	S	0.2	114	07 15 02.03	+46 55 26.3	V
WN J0716+5107	204 ± 9	28.4 ± 0.7	4.54 ± 0.18	-1.35 ± 0.04	-1.47 ± 0.04	D	1.2	8	07 16 40.50	+51 07 04.7	V
WN J0717+4611 <sup>*</sup>	724 ± 29	107.8 ± 4.8	45.30 ± 1.00	-1.30 ± 0.04	-0.70 ± 0.04	T	6.2	16	07 17 58.49	+46 11 39.1	V
WN J0720+5140	95 ± 6	11.6 ± 0.5	...	-1.44 ± 0.05	...	S	3.2	101	07 20 00.20	+51 40 28.5	F
WN J0720+5758	93 ± 7	13.9 ± 0.5	2.37 ± 0.12	-1.30 ± 0.05	-1.42 ± 0.05	S	2.0	109	07 20 09.56	+57 58 32.9	V
WN J0721+3324	313 ± 13	36.8 ± 1.9	2.35 ± 0.13	-1.47 ± 0.05	-2.21 ± 0.06	DF	2.0	143	07 21 56.85	+33 24 53.8	V
WN J0725+4123 <sup>*</sup>	272 ± 12	35.5 ± 1.9	...	-1.39 ± 0.05	...	S	8.3	156	07 25 57.08	+41 23 05.1	F
WN J0727+3020 <sup>*</sup>	93 ± 6	12.8 ± 1.9	...	-1.36 ± 0.11	...	D	34.8	103	07 27 48.50	+30 21 00.7	F
WN J0736+6845	84 ± 5	12.4 ± 0.5	...	-1.31 ± 0.06	...	...	...	...	07 36 36.42	+68 45 56.8	N
WN J0737+7146	79 ± 5	10.3 ± 0.5	...	-1.40 ± 0.07	...	...	...	...	07 37 09.93	+71 46 29.3	N
WN J0740+6319	254 ± 11	26.1 ± 1.7	...	-1.56 ± 0.05	...	...	...	...	07 40 27.01	+63 19 55.2	N
WN J0741+5611	313 ± 13	44.4 ± 1.0	7.08 ± 0.24	-1.34 ± 0.03	-1.48 ± 0.03	S	0.5	20	07 41 15.38	+56 11 35.9	V
WN J0747+3654	289 ± 12	36.8 ± 0.8	...	-1.41 ± 0.03	...	S	2.1	2	07 47 29.38	+36 54 38.1	F
WN J0747+4527	138 ± 7	19.1 ± 2.2	...	-1.35 ± 0.09	...	S	3.5	109	07 47 29.56	+45 27 16.3	F



Table A.1. continued

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J0751+3300	98 ± 7	11.5 ± 0.5	1.90 ± 0.12	-1.47 ± 0.05	-1.44 ± 0.06	S	0.4	139	07 51 48.04	+33 00 07.1	V
WN J0756+5010	204 ± 9	29.9 ± 0.7	...	-1.31 ± 0.04	...	S	1.7	93	07 56 24.63	+50 10 12.5	F
WN J0801+7134	120 ± 6	13.5 ± 2.3	...	-1.50 ± 0.13	...	...	...	...	08 01 18.68	+71 34 01.6	N
WN J0809+6624	257 ± 11	32.2 ± 2.3	...	-1.42 ± 0.06	...	...	...	...	08 09 19.28	+66 24 09.5	N
WN J0810+4948	146 ± 7	18.1 ± 1.4	2.16 ± 0.13	-1.43 ± 0.07	-1.74 ± 0.08	S	< 0.3	0	08 10 34.92	+49 48 37.5	V
WN J0812+3424	160 ± 8	19.9 ± 1.4	...	-1.43 ± 0.06	...	D	6.6	58	08 12 14.43	+34 24 09.3	F
WN J0813+4828	184 ± 9	23.3 ± 1.7	15.20 ± 0.40	-1.42 ± 0.06	-0.32 ± 0.06	D	0.9	173	08 13 38.10	+48 28 41.5	V
WN J0819+5628	95 ± 6	11.7 ± 0.5	...	-1.43 ± 0.06	...	...	...	...	08 19 33.65	+56 28 38.1	N
WN J0829+3834*	112 ± 6	12.7 ± 1.3	...	-1.49 ± 0.08	...	S	1.6	0	08 29 17.35	+38 34 52.6	F
WN J0830+3018	109 ± 6	13.9 ± 1.2	...	-1.41 ± 0.07	...	S	3.5	89	08 30 36.91	+30 18 11.8	F
WN J0834+4820	128 ± 7	17.9 ± 0.5	3.65 ± 0.18	-1.35 ± 0.04	-1.27 ± 0.04	S	0.4	99	08 34 25.85	+48 20 00.8	V
WN J0835+3439	143 ± 7	19.1 ± 1.3	...	-1.38 ± 0.06	...	D	58.0	168	08 35 22.92	+34 39 49.8	F
WN J0842+3540	110 ± 6	15.0 ± 0.5	...	-1.36 ± 0.04	...	S	1.1	0	08 42 10.82	+35 40 42.2	F
WN J0842+3101	76 ± 5	10.5 ± 3.6	...	-1.36 ± 0.24	...	...	...	...	08 42 16.35	+31 01 11.6	N
WN J0843+3723	290 ± 13	35.8 ± 1.9	...	-1.43 ± 0.05	...	S	1.1	5	08 43 02.09	+37 23 18.1	F
WN J0850+4830*	192 ± 9	18.9 ± 1.4	1.53 ± 0.13	-1.59 ± 0.06	-2.02 ± 0.09	S	0.3	52	08 50 56.50	+48 30 46.2	V
WN J0852+3311	179 ± 8	24.6 ± 1.5	...	-1.36 ± 0.05	...	D	6.1	13	08 52 50.55	+33 11 41.0	F
WN J0901+6547†	140 ± 7	19.0 ± 0.5	71.60 ± 1.57	-1.37 ± 0.04	1.07 ± 0.03	D	37.6	117	09 01 27.69	+65 47 25.8	V
WN J0901+3151	135 ± 8	20.1 ± 1.4	...	-1.30 ± 0.06	...	D	20.1	109	09 01 48.50	+31 51 59.4	F
WN J0903+8246	112 ± 6	15.5 ± 0.5	...	-1.35 ± 0.04	...	...	...	...	09 03 44.81	+82 46 02.0	N
WN J0911+6306	118 ± 6	12.9 ± 1.3	...	-1.52 ± 0.08	...	...	...	...	09 11 34.24	+63 06 21.7	N
WN J0913+6104	297 ± 12	33.8 ± 1.8	...	-1.49 ± 0.05	...	...	...	...	09 13 05.21	+61 04 11.7	N
WN J0915+8133	127 ± 5	18.6 ± 0.6	2.99 ± 0.13	-1.32 ± 0.04	-1.47 ± 0.04	S	2.3	127	09 15 09.29	+81 33 34.1	V
WN J0920+5544	142 ± 7	13.5 ± 1.2	...	-1.61 ± 0.07	...	D	8.9	129	09 20 46.78	+55 44 21.8	F
WN J0923+4602	107 ± 5	12.1 ± 0.5	...	-1.49 ± 0.06	...	D	21.6	172	09 23 10.44	+46 02 59.0	F
WN J0928+6003	119 ± 6	11.0 ± 0.5	2.14 ± 0.11	-1.63 ± 0.05	-1.32 ± 0.06	S	3.4	128	09 28 44.58	+60 03 01.7	V
WN J0930+4358	118 ± 7	12.0 ± 0.5	...	-1.57 ± 0.05	...	S	3.1	88	09 30 12.14	+43 58 27.2	F
WN J0930+3207	218 ± 9	31.1 ± 1.7	...	-1.33 ± 0.05	...	S	11.7	58	09 30 53.32	+32 07 20.8	F
WN J0939+6323	268 ± 11	39.7 ± 2.0	10.86 ± 0.29	-1.31 ± 0.05	-1.04 ± 0.05	T	30.3	27	09 39 54.89	+63 23 31.9	V
WN J0940+3838	108 ± 6	11.8 ± 1.6	...	-1.52 ± 0.10	...	...	...	...	09 40 26.15	+38 38 35.0	N
WN J0946+6726	632 ± 25	84.7 ± 3.7	40.19 ± 0.88	-1.38 ± 0.04	-0.60 ± 0.04	T	44.8	114	09 46 44.25	+67 26 41.8	V
WN J0948+6305	731 ± 29	90.5 ± 1.9	14.46 ± 0.38	-1.43 ± 0.03	-1.48 ± 0.03	D	3.1	134	09 48 40.83	+63 05 42.0	V
WN J0952+5153	211 ± 10	16.1 ± 1.3	...	-1.76 ± 0.06	...	S	2.7	166	09 52 49.14	+51 53 05.0	F
WN J0955+6023	103 ± 6	10.9 ± 1.6	4.32 ± 0.15	-1.54 ± 0.11	-0.75 ± 0.12	DF	3.1	129	09 55 30.01	+60 23 17.1	V
WN J1002+5512	119 ± 5	17.7 ± 0.5	...	-1.30 ± 0.05	...	S	10.4	58	10 02 39.24	+55 12 55.5	F
WN J1003+4448	94 ± 5	13.9 ± 2.0	...	-1.31 ± 0.11	...	DF	17.2	93	10 03 30.22	+44 48 16.2	F
WN J1012+3334*	168 ± 8	16.4 ± 1.6	...	-1.59 ± 0.07	...	D	22.1	6	10 12 00.43	+33 35 00.9	F
WN J1013+3254	102 ± 7	14.2 ± 0.5	...	-1.35 ± 0.05	...	S	< 0.3	89	10 13 06.54	+32 54 33.2	F
WN J1014+7407	170 ± 8	21.8 ± 0.6	3.76 ± 0.14	-1.41 ± 0.04	-1.41 ± 0.04	S	2.2	150	10 14 15.75	+74 07 56.2	V
WN J1015+3038	285 ± 12	38.0 ± 1.9	...	-1.38 ± 0.04	...	D	12.8	151	10 15 08.92	+30 38 02.0	F
WN J1015+7432	146 ± 7	21.4 ± 0.6	...	-1.31 ± 0.05	...	...	...	...	10 15 52.49	+74 32 20.0	N
WN J1018+3634	137 ± 7	20.0 ± 1.4	...	-1.32 ± 0.06	...	S	< 0.3	180	10 18 03.13	+36 34 43.2	F
WN J1019+5244	85 ± 5	10.5 ± 0.4	...	-1.43 ± 0.06	...	D	16.5	145	10 19 20.59	+52 44 33.2	F
WN J1022+3308	90 ± 5	10.1 ± 0.4	...	-1.50 ± 0.05	...	S	7.8	136	10 22 37.44	+33 08 43.6	F
WN J1026+2943	81 ± 6	12.0 ± 1.2	...	-1.31 ± 0.08	...	S	11.9	20	10 26 12.04	+29 43 48.0	F
WN J1029+4838	118 ± 8	17.2 ± 1.4	2.80 ± 0.16	-1.32 ± 0.07	-1.46 ± 0.08	S	2.0	131	10 29 32.99	+48 38 09.0	V
WN J1030+5415	223 ± 10	29.2 ± 1.6	...	-1.39 ± 0.05	...	S	6.1	44	10 30 42.94	+54 15 35.9	F
WN J1036+3936	153 ± 7	16.7 ± 0.5	...	-1.52 ± 0.04	...	T	17.5	107	10 36 21.29	+39 36 59.4	F
WN J1038+4229	816 ± 33	114.0 ± 5.1	...	-1.35 ± 0.04	...	S	1.2	3	10 38 41.04	+42 29 51.5	F
WN J1041+5859	193 ± 9	25.2 ± 0.6	6.96 ± 0.22	-1.39 ± 0.04	-1.04 ± 0.03	D	11.2	119	10 41 29.64	+58 59 23.5	V
WN J1043+3953	95 ± 8	13.9 ± 0.5	...	-1.32 ± 0.05	...	S	< 0.3	79	10 43 32.24	+39 53 46.8	F
WN J1052+4826	1039 ± 41	154.2 ± 6.8	...	-1.31 ± 0.04	...	D	10.0	108	10 52 53.05	+48 26 33.8	F
WN J1053+5424	467 ± 20	66.4 ± 1.4	7.94 ± 0.25	-1.34 ± 0.03	-1.70 ± 0.03	D	1.1	34	10 53 36.31	+54 24 42.1	V
WN J1055+3047	342 ± 14	39.0 ± 2.0	...	-1.49 ± 0.05	...	S	7.1	147	10 55 18.52	+30 47 23.2	F
WN J1057+3156	221 ± 10	22.2 ± 1.5	...	-1.57 ± 0.06	...	T	28.7	20	10 57 56.56	+31 56 21.0	F
WN J1057+3007	276 ± 12	39.5 ± 2.0	...	-1.33 ± 0.05	...	D	24.6	90	10 57 57.45	+30 07 02.9	F
WN J1058+3506	205 ± 10	24.0 ± 1.5	...	-1.47 ± 0.05	...	S	1.3	126	10 58 14.74	+35 06 40.9	F
WN J1058+7003	82 ± 4	10.5 ± 0.5	...	-1.41 ± 0.06	...	...	...	...	10 58 34.95	+70 03 17.6	N
WN J1059+4341	99 ± 8	12.3 ± 1.6	...	-1.43 ± 0.10	...	D	29.1	66	10 59 43.88	+43 41 05.2	F
WN J1101+3520*	120 ± 7	17.5 ± 2.1	...	-1.32 ± 0.09	...	S	8.1	1	11 01 14.35	+35 20 12.7	F
WN J1111+3311	92 ± 6	13.3 ± 0.5	...	-1.32 ± 0.05	...	S	6.7	16	11 11 40.79	+33 12 00.9	F
WN J1115+5016	232 ± 10	33.9 ± 0.8	5.47 ± 0.22	-1.32 ± 0.03	-1.46 ± 0.04	D	0.2	117	11 15 06.87	+50 16 23.9	V
WN J1117+5251	1439 ± 57	198.7 ± 8.7	...	-1.36 ± 0.04	...	D	9.5	6	11 17 24.37	+52 51 54.6	F
WN J1122+3239	143 ± 7	19.5 ± 1.4	...	-1.36 ± 0.06	...	S	4.2	89	11 22 33.53	+32 39 40.9	F
WN J1123+3141	623 ± 25	74.0 ± 3.3	...	-1.46 ± 0.04	...	T	25.8	83	11 23 55.85	+31 41 26.1	F

Table A.1. continued

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J1124+3228	263 ± 11	38.9 ± 0.9	...	-1.31 ± 0.03	...	S	5.7	165	11 24 34.17	+32 28 19.6	F
WN J1126+8318	175 ± 8	25.8 ± 1.6	...	-1.31 ± 0.05	...	...	...	...	11 26 26.48	+83 18 23.8	N
WN J1128+4822	280 ± 12	29.9 ± 0.7	...	-1.53 ± 0.03	...	S	1.2	171	11 28 05.19	+48 22 56.6	F
WN J1128+6416	257 ± 11	31.0 ± 1.7	3.20 ± 0.17	-1.45 ± 0.05	-1.82 ± 0.06	S	0.8	116	11 28 12.10	+64 16 25.3	V
WN J1130+4911	134 ± 6	18.9 ± 1.5	...	-1.34 ± 0.07	...	T	37.0	145	11 30 17.53	+49 11 17.7	F
WN J1139+3048	819 ± 33	119.6 ± 5.3	...	-1.32 ± 0.04	...	D	9.0	40	11 39 14.34	+30 48 50.7	F
WN J1139+3706	139 ± 8	16.7 ± 1.4	...	-1.45 ± 0.07	...	S	3.5	32	11 39 43.04	+37 06 55.7	F
WN J1141+6924	81 ± 5	11.8 ± 0.5	...	-1.32 ± 0.07	...	...	...	...	11 41 09.71	+69 24 16.3	N
WN J1148+5116	148 ± 7	20.5 ± 1.7	...	-1.35 ± 0.07	...	D	6.7	50	11 48 01.92	+51 16 29.6	F
WN J1148+6233	98 ± 5	14.1 ± 0.5	1.37 ± 0.09	-1.33 ± 0.05	-1.88 ± 0.06	S	2.4	144	11 48 44.42	+62 33 24.6	V
WN J1148+3519	114 ± 7	13.9 ± 1.7	...	-1.44 ± 0.09	...	D	20.0	7	11 48 54.29	+35 19 10.1	F
WN J1151+4436	78 ± 5	11.1 ± 0.5	...	-1.34 ± 0.06	...	D	15.1	20	11 51 16.80	+44 36 10.0	F
WN J1152+3732*	414 ± 17	17.1 ± 1.4	...	-2.18 ± 0.06	...	S	15.0	119	11 52 36.34	+37 32 43.9	F
WN J1154+5415	153 ± 7	19.1 ± 0.5	1.05 ± 0.10	-1.42 ± 0.04	-2.33 ± 0.08	S	< 0.3	0	11 54 18.97	+54 15 11.4	V
WN J1203+8350	132 ± 6	18.5 ± 0.5	...	-1.35 ± 0.04	...	...	...	...	12 03 30.63	+83 50 37.1	N
WN J1204+5014	94 ± 5	11.3 ± 1.7	...	-1.45 ± 0.11	...	...	...	...	12 04 31.11	+50 14 49.8	N
WN J1206+3136	167 ± 9	22.2 ± 1.5	...	-1.38 ± 0.06	...	D	16.6	53	12 06 48.10	+31 36 50.9	F
WN J1208+4301	162 ± 8	13.9 ± 1.4	...	-1.68 ± 0.08	...	T	8.0	27	12 08 42.77	+43 01 42.3	F
WN J1216+4446	191 ± 8	13.6 ± 0.5	...	-1.81 ± 0.04	...	S	6.1	98	12 16 46.84	+44 46 51.3	F
WN J1218+3143	1906 ± 76	281.5 ± 12.4	...	-1.31 ± 0.04	...	S	5.6	65	12 18 31.47	+31 43 41.7	F
WN J1219+4644	140 ± 8	20.9 ± 0.6	2.84 ± 0.14	-1.30 ± 0.04	-1.59 ± 0.05	S	0.2	157	12 19 31.79	+46 44 49.6	V
WN J1223+4256	116 ± 6	17.2 ± 0.5	...	-1.31 ± 0.04	...	S	4.3	179	12 23 11.32	+42 56 57.3	F
WN J1224+4956	145 ± 7	21.4 ± 1.6	...	-1.31 ± 0.06	...	T	39.6	66	12 24 21.03	+49 56 49.2	F
WN J1224+5436	790 ± 31	69.1 ± 1.5	6.30 ± 0.20	-1.67 ± 0.03	-1.92 ± 0.03	D	0.9	16	12 24 52.35	+54 36 39.9	V
WN J1226+4836	155 ± 8	17.0 ± 0.5	...	-1.51 ± 0.04	...	S	4.8	115	12 26 22.45	+48 36 39.6	F
WN J1232+4621*	133 ± 6	19.8 ± 1.8	...	-1.30 ± 0.07	...	DF	12.5	156	12 32 39.79	+46 21 48.1	F
WN J1242+3915	549 ± 22	78.4 ± 1.7	...	-1.33 ± 0.03	...	D	11.2	51	12 42 53.09	+39 15 48.6	F
WN J1247+6121	103 ± 5	15.4 ± 2.4	...	-1.30 ± 0.11	...	...	...	...	12 47 45.31	+61 21 23.3	N
WN J1249+4043	160 ± 7	22.8 ± 1.4	...	-1.33 ± 0.05	...	D	37.0	80	12 49 27.43	+40 43 49.3	F
WN J1258+3212	182 ± 12	19.9 ± 1.5	...	-1.52 ± 0.06	...	S	6.7	131	12 58 23.62	+32 12 42.4	F
WN J1258+5041	230 ± 11	28.8 ± 2.0	...	-1.42 ± 0.06	...	T	44.9	58	12 58 50.96	+50 41 42.3	F
WN J1259+3121	268 ± 11	34.0 ± 1.8	...	-1.41 ± 0.05	...	DF	20.6	1	12 59 51.96	+31 21 05.6	F
WN J1300+5311	157 ± 7	19.2 ± 0.5	2.70 ± 0.16	-1.44 ± 0.04	-1.58 ± 0.05	S	0.3	7	13 00 36.25	+53 11 52.4	V
WN J1302+3206	383 ± 16	54.0 ± 1.2	...	-1.34 ± 0.03	...	S	4.7	102	13 02 03.07	+32 06 35.6	F
WN J1303+8024	165 ± 7	22.2 ± 0.6	5.05 ± 0.18	-1.37 ± 0.03	-1.19 ± 0.04	D	4.1	145	13 03 33.92	+80 24 42.2	V
WN J1303+5437	145 ± 7	15.4 ± 1.4	...	-1.54 ± 0.07	...	D	16.5	88	13 03 45.48	+54 37 28.9	F
WN J1306+4726	100 ± 7	13.1 ± 0.5	...	-1.39 ± 0.05	...	S	10.2	40	13 06 35.99	+47 26 10.1	F
WN J1310+5706	112 ± 6	15.9 ± 0.5	...	-1.34 ± 0.04	...	S	4.2	99	13 10 22.96	+57 06 57.9	F
WN J1310+3820	69 ± 4	10.3 ± 1.5	...	-1.30 ± 0.12	...	S	4.4	18	13 10 24.08	+38 20 44.1	F
WN J1312+6646	291 ± 12	41.6 ± 0.9	8.56 ± 0.24	-1.33 ± 0.03	-1.27 ± 0.03	D	6.3	118	13 12 45.59	+66 46 36.1	V
WN J1314+3649	286 ± 11	36.4 ± 0.8	...	-1.41 ± 0.03	...	S	1.3	90	13 14 17.86	+36 49 14.6	F
WN J1314+3515*	169 ± 7	20.2 ± 1.7	...	-1.45 ± 0.07	...	DF	34.2	47	13 14 25.00	+35 15 53.9	F
WN J1315+4337	149 ± 8	16.3 ± 0.5	...	-1.52 ± 0.04	...	S	1.3	152	13 15 19.05	+43 37 57.1	F
WN J1321+3311	141 ± 8	19.0 ± 0.6	...	-1.37 ± 0.04	...	D	14.7	63	13 21 13.55	+33 11 31.2	F
WN J1323+4713	106 ± 5	14.9 ± 0.5	...	-1.34 ± 0.04	...	S	2.5	139	13 23 08.31	+47 13 10.7	F
WN J1327+5341	196 ± 9	27.4 ± 0.7	...	-1.35 ± 0.04	...	S	0.6	162	13 27 12.77	+53 41 20.6	F
WN J1327+5332	163 ± 7	20.5 ± 0.6	2.70 ± 0.17	-1.42 ± 0.04	-1.61 ± 0.06	D	0.3	89	13 27 37.86	+53 32 10.8	V
WN J1329+3046*	86 ± 5	11.7 ± 1.7	...	-1.37 ± 0.11	...	S	1.6	6	13 29 31.75	+30 46 13.6	F
WN J1329+3046*	88 ± 5	12.1 ± 2.0	...	-1.36 ± 0.13	...	S	3.6	129	13 29 55.12	+30 46 51.2	F
WN J1330+6505	74 ± 6	10.5 ± 0.5	2.34 ± 0.12	-1.34 ± 0.06	-1.21 ± 0.06	S	1.8	5	13 30 02.77	+65 04 59.3	V
WN J1330+3037*	496 ± 20	64.5 ± 2.9	...	-1.40 ± 0.04	...	D	25.5	140	13 30 53.67	+30 37 59.1	F
WN J1330+3604	103 ± 5	15.1 ± 0.5	...	-1.31 ± 0.05	...	D	9.6	0	13 30 56.83	+36 04 01.6	F
WN J1330+5344*	525 ± 21	71.0 ± 3.2	8.10 ± 0.29	-1.37 ± 0.04	-1.74 ± 0.05	DF	2.5	173	13 30 58.90	+53 44 07.8	V
WN J1331+2937	93 ± 5	12.5 ± 1.3	3.44 ± 0.13	-1.37 ± 0.09	-1.04 ± 0.09	S	2.7	40	13 31 21.58	+29 37 12.6	V
WN J1332+3009*	239 ± 10	35.3 ± 2.0	...	-1.31 ± 0.05	...	D	39.2	161	13 32 45.73	+30 09 59.9	F
WN J1333+3037*	423 ± 17	21.5 ± 0.6	...	-2.04 ± 0.03	...	S	0.4	179	13 33 21.20	+30 37 35.1	F
WN J1333+4913	149 ± 7	22.2 ± 0.6	4.78 ± 0.23	-1.30 ± 0.04	-1.22 ± 0.04	T	5.4	40	13 33 56.42	+49 13 27.1	V
WN J1335+3222*	150 ± 7	18.3 ± 1.4	...	-1.44 ± 0.06	...	D	8.8	85	13 35 35.44	+32 22 48.7	F
WN J1336+3820	116 ± 6	15.6 ± 0.5	...	-1.37 ± 0.04	...	S	3.4	61	13 36 40.88	+38 20 02.2	F
WN J1337+3149	142 ± 7	19.6 ± 1.9	...	-1.36 ± 0.08	...	D	35.2	21	13 37 02.46	+31 49 49.2	F
WN J1337+3401	109 ± 8	14.3 ± 0.5	...	-1.39 ± 0.04	...	S	3.5	56	13 37 10.50	+34 01 26.8	F
WN J1339+5320	104 ± 5	13.9 ± 0.5	1.80 ± 0.15	-1.38 ± 0.05	-1.63 ± 0.07	D	0.1	138	13 39 59.25	+53 20 07.8	V
WN J1341+4953	153 ± 8	16.4 ± 1.4	...	-1.53 ± 0.07	...	D	18.6	160	13 41 25.70	+49 53 41.8	F
WN J1346+6736	107 ± 6	14.5 ± 0.5	2.82 ± 0.16	-1.37 ± 0.04	-1.32 ± 0.05	S	< 0.3	0	13 46 37.02	+67 36 21.7	V
WN J1347+3033	341 ± 14	42.4 ± 2.2	...	-1.43 ± 0.05	...	D	37.3	151	13 47 16.09	+30 33 07.6	F

Table A.1. continued

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J1352+4259	1213 ± 48	173.9 ± 7.7	...	-1.33 ± 0.04	...	D	12.1	19	13 52 28.39	+42 59 18.1	F
WN J1353+3336	107 ± 6	15.9 ± 1.9	...	-1.31 ± 0.09	...	...	...	...	13 53 07.46	+33 36 24.6	N
WN J1355+3848	204 ± 8	25.6 ± 0.6	...	-1.42 ± 0.03	...	S	2.3	59	13 55 29.50	+38 48 11.1	F
WN J1359+7446*	214 ± 9	12.9 ± 0.5	1.26 ± 0.10	-1.92 ± 0.05	-1.87 ± 0.07	S	2.5	12	13 59 16.90	+74 46 42.6	V
WN J1400+4348	178 ± 7	20.2 ± 1.5	...	-1.49 ± 0.06	...	S	6.4	93	14 00 51.84	+43 48 07.2	F
WN J1403+3109	328 ± 13	45.7 ± 2.2	...	-1.35 ± 0.04	...	S	11.0	25	14 03 39.47	+31 09 11.4	F
WN J1410+4615	103 ± 5	14.2 ± 1.4	...	-1.36 ± 0.08	...	S	4.4	79	14 10 48.17	+46 15 57.5	F
WN J1416+3821	94 ± 6	13.1 ± 0.5	...	-1.35 ± 0.05	...	D	7.6	14	14 16 13.21	+38 21 56.7	F
WN J1418+4546	186 ± 8	17.9 ± 0.5	2.70 ± 0.14	-1.60 ± 0.04	-1.52 ± 0.05	S	0.5	18	14 18 38.40	+45 46 38.4	V
WN J1420+6735	382 ± 15	33.1 ± 1.7	1.22 ± 0.09	-1.67 ± 0.05	-2.65 ± 0.07	S	2.0	173	14 20 03.34	+67 35 07.5	V
WN J1420+4126	128 ± 6	18.7 ± 0.6	...	-1.32 ± 0.04	...	S	5.3	65	14 20 50.23	+41 26 26.1	F
WN J1421+3103	110 ± 5	13.6 ± 0.5	...	-1.43 ± 0.05	...	S	0.4	127	14 21 07.69	+31 03 03.9	F
WN J1422+3452	163 ± 7	10.7 ± 1.3	...	-1.86 ± 0.09	...	S	13.3	151	14 22 05.17	+34 52 14.4	F
WN J1422+4212	258 ± 10	35.4 ± 1.8	...	-1.36 ± 0.05	...	S	5.1	164	14 22 44.38	+42 12 00.2	F
WN J1426+3522	89 ± 5	11.5 ± 1.6	...	-1.40 ± 0.11	...	...	...	...	14 26 56.32	+35 22 34.7	N
WN J1431+3015	101 ± 5	14.3 ± 1.5	...	-1.34 ± 0.09	...	D	28.9	158	14 31 17.77	+30 15 00.2	F
WN J1431+7317	129 ± 6	17.9 ± 1.3	4.65 ± 0.19	-1.35 ± 0.06	-1.08 ± 0.07	D	11.2	47	14 31 28.01	+73 17 29.7	V
WN J1433+3044	144 ± 6	16.7 ± 1.3	...	-1.48 ± 0.06	...	D	14.5	100	14 33 56.51	+30 44 25.5	F
WN J1435+3523	160 ± 7	22.3 ± 1.6	...	-1.35 ± 0.06	...	S	8.0	64	14 35 53.99	+35 24 06.8	F
WN J1436+6319	3507 ± 140	514.0 ± 22.4	64.84 ± 1.39	-1.31 ± 0.04	-1.67 ± 0.04	D	3.8	154	14 36 37.28	+63 19 13.9	V
WN J1437+7409	111 ± 5	16.4 ± 0.5	1.32 ± 0.11	-1.31 ± 0.05	-2.02 ± 0.07	S	< 0.3	0	14 37 35.89	+74 09 22.2	V
WN J1439+3729	142 ± 7	17.1 ± 0.5	...	-1.45 ± 0.04	...	D	7.2	157	14 39 49.99	+37 29 03.4	F
WN J1440+3707*	219 ± 9	16.7 ± 1.3	...	-1.76 ± 0.06	...	S	7.4	61	14 40 03.63	+37 07 27.4	F
WN J1444+4114	97 ± 5	13.2 ± 1.4	...	-1.37 ± 0.09	...	D	30.1	151	14 44 27.65	+41 14 38.3	F
WN J1444+4112	109 ± 6	13.1 ± 0.5	...	-1.45 ± 0.05	...	D	12.6	96	14 44 48.82	+41 12 25.5	F
WN J1447+5423	97 ± 6	13.3 ± 0.5	...	-1.36 ± 0.05	...	S	11.0	174	14 47 54.36	+54 23 25.2	F
WN J1450+3534	95 ± 5	10.2 ± 0.4	...	-1.53 ± 0.05	...	S	3.3	62	14 50 20.69	+35 34 46.5	F
WN J1451+3649	254 ± 10	29.8 ± 2.3	...	-1.47 ± 0.06	...	T	50.7	101	14 51 11.62	+47 56 34.4	F
WN J1454+3210	264 ± 11	37.6 ± 0.9	...	-1.33 ± 0.03	...	S	3.0	22	14 54 11.64	+32 10 16.8	F
WN J1459+6405	339 ± 14	40.0 ± 3.6	...	-1.46 ± 0.07	...	...	...	...	14 59 33.34	+64 05 58.0	N
WN J1459+4947	175 ± 8	18.7 ± 0.5	...	-1.53 ± 0.04	...	S	3.0	109	14 59 43.19	+49 47 15.9	F
WN J1500+3613	390 ± 16	52.9 ± 2.5	...	-1.37 ± 0.04	...	S	2.2	76	15 00 09.01	+36 13 25.8	F
WN J1502+4756	221 ± 9	19.1 ± 1.5	...	-1.68 ± 0.06	...	D	12.3	37	15 02 03.58	+47 56 34.7	F
WN J1508+5839	122 ± 6	12.5 ± 0.5	1.37 ± 0.09	-1.56 ± 0.05	-1.78 ± 0.06	D	35.2	128	15 08 08.17	+58 39 13.3	V
WN J1509+5905*	752 ± 30	107.3 ± 4.6	8.60 ± 0.26	-1.33 ± 0.04	-2.03 ± 0.04	DF	1.8	74	15 09 32.34	+59 05 24.5	V
WN J1525+3010	145 ± 7	17.2 ± 0.5	...	-1.46 ± 0.04	...	S	1.2	123	15 25 01.21	+30 10 30.2	F
WN J1525+5130	94 ± 5	10.7 ± 0.5	2.02 ± 0.13	-1.49 ± 0.06	-1.34 ± 0.06	S	0.8	7	15 25 12.93	+51 30 06.1	V
WN J1528+6317	332 ± 14	40.0 ± 2.0	20.80 ± 0.52	-1.45 ± 0.04	-0.53 ± 0.05	DF	6.0	121	15 28 06.47	+63 17 40.2	V
WN J1529+3454	124 ± 6	13.0 ± 0.5	...	-1.54 ± 0.05	...	S	2.3	35	15 29 41.52	+34 54 31.3	F
WN J1537+3402	83 ± 5	11.6 ± 0.5	...	-1.35 ± 0.05	...	S	1.7	31	15 37 50.57	+34 02 33.3	F
WN J1543+3512	231 ± 10	23.6 ± 1.5	...	-1.56 ± 0.05	...	D	9.2	0	15 43 29.49	+35 12 29.7	F
WN J1546+3935	155 ± 7	13.7 ± 0.5	...	-1.66 ± 0.04	...	D	8.5	102	15 46 41.95	+49 35 52.7	F
WN J1546+3005	92 ± 6	13.0 ± 0.5	...	-1.34 ± 0.05	...	S	2.3	144	15 46 57.65	+30 05 38.1	F
WN J1550+3830	229 ± 10	32.9 ± 1.7	...	-1.33 ± 0.05	...	S	4.2	178	15 50 19.42	+38 30 14.8	F
WN J1552+3715	546 ± 22	27.3 ± 1.6	...	-2.05 ± 0.05	...	D	18.6	120	15 52 07.07	+37 15 07.1	F
WN J1555+4011	359 ± 14	50.9 ± 2.4	...	-1.34 ± 0.04	...	S	1.1	90	15 55 02.50	+40 11 58.2	F
WN J1558+7028	114 ± 6	15.7 ± 2.1	1.68 ± 0.11	-1.36 ± 0.11	-1.80 ± 0.12	S	2.1	38	15 58 37.92	+70 28 11.2	V
WN J1559+6110	130 ± 6	18.3 ± 1.3	3.47 ± 0.14	-1.34 ± 0.06	-1.34 ± 0.07	S	0.3	20	15 59 50.65	+61 10 57.8	V
WN J1559+5926	116 ± 6	15.3 ± 1.3	41.70 ± 0.92	-1.39 ± 0.07	...	D	31.0	60	15 59 54.42	+59 26 35.3	V
WN J1604+5505	142 ± 7	20.7 ± 0.6	...	-1.32 ± 0.04	...	S	1.1	15	16 04 21.91	+55 05 45.4	F
WN J1606+6346	130 ± 6	19.0 ± 0.6	5.60 ± 0.19	-1.32 ± 0.04	-0.98 ± 0.04	D	5.6	14	16 06 12.44	+63 47 06.1	V
WN J1606+4142	83 ± 5	12.3 ± 0.5	...	-1.31 ± 0.05	...	S	3.2	145	16 06 30.58	+41 42 10.8	F
WN J1609+5725	70 ± 5	10.2 ± 0.5	2.13 ± 0.11	-1.32 ± 0.07	-1.26 ± 0.06	S	1.7	60	16 09 28.14	+57 25 02.8	V
WN J1618+5736	233 ± 10	33.5 ± 1.8	6.38 ± 0.20	-1.33 ± 0.05	-1.34 ± 0.05	S	2.1	102	16 18 07.35	+57 36 11.7	V
WN J1622+3447	156 ± 7	21.1 ± 0.6	...	-1.37 ± 0.04	...	S	1.6	0	16 22 10.93	+34 47 44.6	F
WN J1623+8213	166 ± 7	19.8 ± 0.6	...	-1.46 ± 0.04	...	...	...	...	16 23 01.71	+82 13 17.4	N
WN J1624+4202	75 ± 5	11.0 ± 1.7	...	-1.31 ± 0.12	...	D	24.4	19	16 24 37.86	+42 02 10.5	F
WN J1627+5430	146 ± 7	20.9 ± 0.6	...	-1.33 ± 0.04	...	S	1.6	176	16 27 24.09	+54 30 55.2	F
WN J1628+3932	26737 ± 1069	3680.7 ± 35.1	...	-1.36 ± 0.03	...	M	48.8	77	16 28 38.56	+39 33 00.1	F
WN J1633+7351	75 ± 5	10.1 ± 1.3	23.88 ± 0.55	-1.37 ± 0.11	0.69 ± 0.11	DF	21.8	90	16 33 07.45	+73 51 38.1	V
WN J1645+4413	135 ± 6	18.9 ± 0.6	...	-1.35 ± 0.04	...	S	1.1	56	16 45 35.69	+44 13 42.1	F
WN J1704+3839	335 ± 14	34.5 ± 1.8	...	-1.56 ± 0.05	...	D	23.8	145	17 04 00.50	+38 39 56.7	F
WN J1713+3656	116 ± 6	14.8 ± 0.5	...	-1.41 ± 0.04	...	S	1.1	169	17 13 55.63	+36 56 34.5	F
WN J1714+7031	132 ± 6	14.2 ± 1.4	0.95 ± 0.09	-1.53 ± 0.08	-2.18 ± 0.11	S	2.4	46	17 14 35.18	+70 31 30.7	V
WN J1714+5251	262 ± 11	37.9 ± 2.0	...	-1.32 ± 0.05	...	D	26.6	80	17 14 37.93	+52 51 30.1	F

Table A.1. continued

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J1717+3828	368 ± 15	35.8 ± 2.0	...	-1.60 ± 0.05	...	M	33.6	156	17 17 16.22	+38 28 16.5	F
WN J1718+5823	1098 ± 44	133.2 ± 2.7	11.26 ± 0.32	-1.44 ± 0.03	-1.98 ± 0.03	S	0.3	107	17 18 18.51	+58 23 21.8	V
WN J1723+5822	88 ± 6	10.4 ± 0.5	2.27 ± 0.11	-1.46 ± 0.06	-1.23 ± 0.06	S	4.7	103	17 23 01.98	+58 22 44.0	V
WN J1723+6844	105 ± 6	15.7 ± 1.3	...	-1.30 ± 0.07	...	...	...	...	17 23 05.64	+68 44 17.7	N
WN J1731+4640	200 ± 9	15.7 ± 0.5	...	-1.74 ± 0.04	...	D	11.6	106	17 31 46.10	+46 40 03.1	F
WN J1731+4654	323 ± 13	42.0 ± 2.1	...	-1.40 ± 0.05	...	S	1.9	176	17 31 59.63	+46 54 00.2	F
WN J1732+6757	85 ± 5	11.9 ± 0.5	2.31 ± 0.11	-1.35 ± 0.05	-1.32 ± 0.05	D	1.5	178	17 32 30.61	+67 57 02.5	V
WN J1733+4037	80 ± 5	11.2 ± 2.0	...	-1.35 ± 0.13	...	...	...	...	17 33 31.13	+40 37 00.6	N
WN J1734+4527	87 ± 5	11.0 ± 0.5	...	-1.42 ± 0.06	...	D	10.4	174	17 34 07.58	+45 27 25.6	F
WN J1734+3606	69 ± 3	10.2 ± 3.2	3.12 ± 0.13	-1.31 ± 0.22	-0.95 ± 0.25	S	2.1	137	17 34 24.13	+36 06 38.8	V
WN J1736+6502	193 ± 8	27.4 ± 1.6	4.40 ± 0.16	-1.34 ± 0.05	-1.47 ± 0.06	D	16.0	159	17 36 37.50	+65 02 28.7	V
WN J1739+5309	78 ± 5	11.3 ± 0.5	...	-1.32 ± 0.06	...	S	7.3	114	17 39 28.39	+53 09 41.8	F
WN J1749+5659	124 ± 7	15.9 ± 1.6	...	-1.41 ± 0.08	...	...	...	...	17 49 21.19	+56 59 59.6	N
WN J1749+6248	104 ± 6	13.3 ± 0.5	2.93 ± 0.12	-1.41 ± 0.05	-1.22 ± 0.05	S	1.7	48	17 49 27.04	+62 48 54.0	V
WN J1752+2949	135 ± 6	16.9 ± 0.5	2.29 ± 0.14	-1.42 ± 0.04	-1.61 ± 0.05	S	3.2	102	17 52 56.67	+29 49 28.3	V
WN J1801+3336	136 ± 6	11.6 ± 0.5	1.26 ± 0.12	-1.69 ± 0.05	-1.78 ± 0.08	S	2.5	34	18 01 07.26	+33 36 40.4	V
WN J1802+3948	1102 ± 44	157.3 ± 6.8	27.67 ± 0.62	-1.33 ± 0.04	-1.40 ± 0.04	T	19.2	13	18 02 31.65	+39 48 59.6	V
WN J1804+5547	173 ± 8	22.9 ± 1.7	...	-1.38 ± 0.06	...	...	...	...	18 04 15.35	+55 47 16.3	N
WN J1804+3048	192 ± 8	27.7 ± 0.7	5.88 ± 0.18	-1.33 ± 0.04	-1.25 ± 0.03	S	2.0	84	18 04 42.01	+30 48 45.0	V
WN J1806+6332	138 ± 7	17.7 ± 0.5	...	-1.41 ± 0.04	...	...	...	...	18 06 23.66	+63 32 07.9	N
WN J1807+5027	483 ± 19	71.4 ± 3.3	11.67 ± 0.31	-1.31 ± 0.04	-1.46 ± 0.04	D	3.7	173	18 07 07.39	+50 27 25.8	V
WN J1807+6628	76 ± 5	10.9 ± 1.4	1.20 ± 0.11	-1.33 ± 0.10	-1.78 ± 0.13	S	0.3	103	18 07 29.75	+66 28 33.4	V
WN J1810+6635	115 ± 8	14.1 ± 0.5	2.62 ± 0.12	-1.44 ± 0.04	-1.36 ± 0.05	S	1.6	15	18 10 28.41	+66 35 15.1	V
WN J1811+6008	160 ± 7	17.8 ± 0.5	7.87 ± 0.24	-1.50 ± 0.04	-0.66 ± 0.03	DF	3.5	31	18 11 18.38	+60 08 42.9	V
WN J1813+4847	92 ± 5	12.1 ± 0.5	...	-1.39 ± 0.06	...	...	...	...	18 13 16.68	+48 47 50.1	N
WN J1814+5009	120 ± 7	16.7 ± 0.5	...	-1.35 ± 0.05	...	...	...	...	18 14 01.40	+50 09 58.0	N
WN J1815+3656	124 ± 6	17.6 ± 0.6	2.96 ± 0.14	-1.34 ± 0.04	-1.43 ± 0.05	S	1.6	48	18 15 23.11	+36 56 00.9	V
WN J1816+3840	186 ± 8	27.3 ± 1.6	...	-1.31 ± 0.05	...	...	...	...	18 16 26.11	+38 40 53.0	N
WN J1818+7042	1624 ± 65	199.6 ± 8.8	30.50 ± 0.74	-1.44 ± 0.04	-1.51 ± 0.04	D	1.4	10	18 18 04.24	+70 42 59.5	V
WN J1818+3852	105 ± 6	12.6 ± 0.5	1.29 ± 0.09	-1.45 ± 0.05	-1.84 ± 0.07	S	2.6	64	18 18 42.17	+38 52 20.4	V
WN J1818+3428	167 ± 7	23.7 ± 0.6	5.68 ± 0.19	-1.34 ± 0.04	-1.15 ± 0.03	T	7.5	103	18 18 47.90	+34 28 24.0	V
WN J1818+6144	149 ± 6	21.9 ± 1.5	1.11 ± 0.10	-1.31 ± 0.06	-2.40 ± 0.09	S	1.6	11	18 18 50.55	+61 44 18.4	V
WN J1819+3122	77 ± 5	10.5 ± 0.5	...	-1.36 ± 0.06	...	...	...	...	18 19 28.60	+31 22 19.3	N
WN J1819+6213	106 ± 6	12.7 ± 0.5	0.47 ± 0.08	-1.45 ± 0.05	-2.66 ± 0.14	S	1.6	41	18 19 52.26	+62 13 58.0	V
WN J1820+5711	136 ± 7	20.3 ± 1.5	...	-1.30 ± 0.06	...	...	...	...	18 20 51.14	+57 11 06.5	N
WN J1821+3602	3398 ± 136	99.0 ± 4.2	233.33 ± 4.90	-2.42 ± 0.04	0.69 ± 0.04	T	35.0	55	18 21 02.20	+36 02 15.7	V
WN J1821+3601 <sup>†</sup>	151 ± 6	10.2 ± 1.5	105.26 ± 2.53	-1.85 ± 0.11	1.88 ± 0.12	D	34.9	55	18 21 22.59	+36 01 03.3	V
WN J1829+6914	207 ± 9	13.0 ± 1.3	2.25 ± 0.12	-1.90 ± 0.08	-1.41 ± 0.09	S	1.8	22	18 29 05.68	+69 14 06.1	V
WN J1829+4919	98 ± 6	14.2 ± 0.5	7.95 ± 0.24	-1.32 ± 0.06	-0.47 ± 0.04	D	12.5	173	18 29 33.74	+49 19 55.6	V
WN J1829+5945	346 ± 14	50.4 ± 2.4	8.82 ± 0.25	-1.32 ± 0.04	-1.40 ± 0.04	S	1.7	26	18 29 57.26	+59 45 03.2	V
WN J1830+6422	70 ± 3	10.3 ± 0.5	...	-1.31 ± 0.07	...	...	...	...	18 30 45.82	+64 22 26.3	N
WN J1832+5354 <sup>†</sup>	226 ± 10	28.7 ± 1.6	23.82 ± 0.59	-1.41 ± 0.05	-0.15 ± 0.05	D	19.4	56	18 32 41.16	+53 54 34.0	V
WN J1836+5210	191 ± 9	24.2 ± 1.5	3.54 ± 0.15	-1.41 ± 0.05	-1.55 ± 0.06	S	1.4	10	18 36 23.22	+52 10 28.4	V
WN J1839+4710	176 ± 8	22.8 ± 1.6	...	-1.40 ± 0.06	...	...	...	...	18 39 14.54	+47 10 59.4	N
WN J1843+5932	998 ± 40	148.2 ± 6.6	30.87 ± 0.70	-1.31 ± 0.04	-1.26 ± 0.04	T	7.7	64	18 43 31.70	+59 32 59.3	V
WN J1847+5423	108 ± 6	15.8 ± 1.3	6.67 ± 0.23	-1.32 ± 0.07	-0.69 ± 0.07	DF	2.9	67	18 47 04.03	+54 23 06.6	V
WN J1852+5711 <sup>*</sup>	387 ± 16	53.4 ± 2.5	...	-1.36 ± 0.04	...	...	...	...	18 52 08.35	+57 11 42.7	N
WN J1857+7411	160 ± 7	19.9 ± 0.6	...	-1.43 ± 0.04	...	...	...	...	18 57 03.60	+74 11 31.2	N
WN J1859+5900	98 ± 6	11.7 ± 0.5	1.64 ± 0.12	-1.46 ± 0.06	-1.56 ± 0.07	S	0.5	171	18 59 40.34	+59 00 36.8	V
WN J1859+5416	81 ± 4	10.1 ± 1.6	2.70 ± 0.14	-1.43 ± 0.12	-1.06 ± 0.13	S	1.7	17	18 59 57.65	+54 16 21.8	V
WN J1907+8532	69 ± 4	10.2 ± 0.4	...	-1.31 ± 0.05	...	...	...	...	19 07 26.54	+85 32 35.2	N
WN J1911+6342	185 ± 10	23.3 ± 1.5	3.37 ± 0.14	-1.42 ± 0.05	-1.56 ± 0.06	S	1.8	7	19 11 49.54	+63 42 09.6	V
WN J1912+8627	126 ± 7	18.8 ± 1.2	2.55 ± 0.13	-1.30 ± 0.05	-1.61 ± 0.07	S	2.4	178	19 12 06.26	+86 27 10.1	V
WN J1917+6635	75 ± 5	11.2 ± 0.5	2.37 ± 0.12	-1.30 ± 0.06	-1.25 ± 0.06	S	1.9	152	19 17 35.50	+66 35 38.5	V
WN J1917+7149	280 ± 12	40.4 ± 2.1	...	-1.33 ± 0.05	...	...	...	...	19 17 56.11	+71 49 19.7	N
WN J1923+6047	83 ± 5	12.4 ± 0.5	2.56 ± 0.13	-1.30 ± 0.06	-1.27 ± 0.05	S	2.4	91	19 23 33.18	+60 47 56.8	V
WN J1925+5203	199 ± 9	27.5 ± 1.7	4.30 ± 0.16	-1.36 ± 0.05	-1.49 ± 0.06	D	8.6	52	19 25 16.97	+52 03 34.4	V
WN J1925+5742	100 ± 6	14.5 ± 0.5	1.58 ± 0.11	-1.32 ± 0.06	-1.78 ± 0.06	S	2.0	9	19 25 22.22	+57 42 27.3	V
WN J1926+5710	136 ± 7	20.2 ± 0.6	4.55 ± 0.18	-1.31 ± 0.04	-1.20 ± 0.04	S	1.8	161	19 26 43.85	+57 10 00.5	V
WN J1927+6436	661 ± 26	80.3 ± 3.5	6.62 ± 0.21	-1.44 ± 0.04	-2.01 ± 0.04	D	20.1	120	19 27 22.83	+64 36 02.6	V
WN J1944+6552	1367 ± 55	147.2 ± 6.5	19.45 ± 0.47	-1.53 ± 0.04	-1.63 ± 0.04	S	1.8	165	19 44 23.99	+65 52 23.8	V
WN J1953+7052	75 ± 7	10.3 ± 0.5	3.22 ± 0.15	-1.36 ± 0.07	-0.94 ± 0.05	D	14.4	150	19 53 21.10	+70 52 19.8	V
WN J1954+7011	460 ± 19	63.4 ± 2.9	...	-1.36 ± 0.04	...	...	...	...	19 54 30.53	+70 11 33.5	N
WN J2044+7044	311 ± 13	41.7 ± 0.9	7.18 ± 0.22	-1.38 ± 0.03	-1.42 ± 0.03	D	1.3	83	20 44 57.80	+70 44 03.8	V
WN J2052+6925	113 ± 6	16.1 ± 1.3	...	-1.33 ± 0.07	...	...	...	...	20 52 33.78	+69 25 09.3	N

Table A.1. continued

Name	$S_{325}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{325}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
WN J2053+6849	109 ± 5	15.8 ± 0.5	2.59 ± 0.13	-1.32 ± 0.05	-1.45 ± 0.05	S	2.8	13	20 53 38.68	+68 48 54.7	V
WN J2139+3125	249 ± 10	28.0 ± 0.7	...	-1.50 ± 0.04	...	...	...	...	21 39 32.31	+31 25 18.7	N
WN J2146+3330	105 ± 6	10.8 ± 0.5	...	-1.56 ± 0.05	...	...	...	...	21 46 55.47	+33 30 01.7	N
WN J2147+3137	215 ± 9	25.4 ± 1.6	...	-1.46 ± 0.05	...	...	...	...	21 47 35.21	+31 37 58.9	N
WN J2158+3424	182 ± 8	25.3 ± 0.7	4.48 ± 0.17	-1.35 ± 0.04	-1.39 ± 0.04	T	7.9	163	21 58 54.14	+34 24 47.8	V
WN J2213+3411	320 ± 13	29.8 ± 2.2	...	-1.63 ± 0.06	...	...	...	...	22 13 12.48	+34 11 42.8	N
WN J2219+2951	151 ± 10	13.2 ± 0.5	...	-1.67 ± 0.04	...	...	...	...	22 19 28.65	+29 51 57.4	N
WN J2221+3800	99 ± 6	14.4 ± 0.5	2.18 ± 0.11	-1.32 ± 0.05	-1.52 ± 0.05	S	1.4	5	22 21 49.53	+38 00 38.4	V
WN J2222+3305	232 ± 10	30.4 ± 1.7	3.53 ± 0.14	-1.39 ± 0.05	-1.73 ± 0.05	S	1.4	13	22 22 15.19	+33 05 44.1	V
WN J2245+3937	204 ± 8	15.5 ± 1.3	5.65 ± 0.21	-1.76 ± 0.07	-0.81 ± 0.07	DF	4.0	168	22 45 02.74	+39 37 27.9	V
WN J2250+4131	128 ± 7	15.2 ± 1.4	2.03 ± 0.11	-1.46 ± 0.07	-1.62 ± 0.09	S	1.5	25	22 50 51.16	+41 31 16.4	V
WN J2313+4053	101 ± 6	11.3 ± 1.4	...	-1.50 ± 0.09	...	...	...	...	23 13 06.34	+40 53 40.6	N
WN J2313+4253	94 ± 7	13.4 ± 1.5	3.41 ± 0.14	-1.33 ± 0.09	-1.10 ± 0.10	S	1.4	13	23 13 08.62	+42 53 13.0	V
WN J2313+3842 <sup>†</sup>	304 ± 13	13.2 ± 2.6	...	-2.15 ± 0.14	...	...	...	...	23 13 47.75	+38 42 26.2	N
WN J2319+4251	321 ± 14	40.2 ± 2.9	3.88 ± 0.14	-1.42 ± 0.06	-1.88 ± 0.07	S	1.5	20	23 19 47.25	+42 51 09.2	V
WN J2337+3421	84 ± 4	12.4 ± 1.3	...	-1.31 ± 0.09	...	...	...	...	23 37 23.43	+34 21 51.5	N
WN J2338+4047	115 ± 6	16.9 ± 0.5	5.06 ± 0.18	-1.31 ± 0.04	-0.97 ± 0.04	D	15.7	8	23 38 11.10	+40 47 19.6	V
WN J2350+3631	196 ± 8	22.8 ± 0.6	2.20 ± 0.11	-1.47 ± 0.04	-1.88 ± 0.04	DF	2.7	165	23 50 25.69	+36 31 27.6	V

<sup>†</sup> Not a real USS source; see notes.

\* See notes.

Table A.2. TN sample

Name	$S_{365}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{365}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
TN J0008-0912	638 ± 64	100.2 ± 4.4	...	-1.38 ± 0.08	...	...	...	...	00 08 52.43	-09 12 01.4	N
TN J0018+1904	498 ± 30	84.5 ± 3.8	...	-1.32 ± 0.06	...	...	...	...	00 18 07.35	+19 04 50.0	N
TN J0026+1501	241 ± 26	39.4 ± 2.0	...	-1.35 ± 0.09	...	...	...	...	00 26 03.31	+15 01 29.2	N
TN J0027+0059	735 ± 55	98.0 ± 4.4	...	-1.50 ± 0.07	...	...	...	...	00 27 39.36	+00 59 53.6	N
TN J0033-2731	512 ± 41	66.1 ± 3.0	...	-1.52 ± 0.07	...	...	...	...	00 33 24.28	-27 31 48.1	N
TN J0037+2629	442 ± 35	70.8 ± 3.2	...	-1.36 ± 0.07	...	...	...	...	00 37 10.05	+26 29 42.3	N
TN J0038-1540	471 ± 33	63.9 ± 1.4	10.8 ± 0.6	-1.49 ± 0.06	-1.43 ± 0.05	D	3.2	18	00 38 47.19	-15 40 06.6	V
TN J0040+1417	435 ± 41	71.1 ± 1.5	...	-1.35 ± 0.07	...	...	...	...	00 40 37.04	+14 17 14.6	N
TN J0041+1250	451 ± 65	71.0 ± 3.3	21.0 ± 1.1	-1.38 ± 0.11	-0.98 ± 0.06	D	10.7	103	00 41 31.27	+12 50 32.3	V
TN J0042+2649	484 ± 57	69.6 ± 3.2	...	-1.44 ± 0.09	...	...	...	...	00 42 32.46	+26 49 06.2	N
TN J0055+2624	8670 ± 457	1375.5 ± 56.2	...	-1.37 ± 0.05	...	...	...	...	00 55 50.38	+26 24 35.8	N
TN J0102-1055	775 ± 41	110.4 ± 4.9	16.4 ± 0.9	-1.45 ± 0.05	-1.53 ± 0.06	D	1.5	136	01 01 59.99	-10 55 56.0	V
TN J0102-2152	1900 ± 88	168.3 ± 6.7	...	-1.80 ± 0.05	...	...	...	...	01 02 41.09	-21 52 30.2	N
TN J0114-0333	367 ± 36	54.7 ± 2.6	12.9 ± 0.7	-1.42 ± 0.08	-1.16 ± 0.06	T	2.4	146	01 14 42.50	-03 33 58.8	N
TN J0119-2054	522 ± 60	87.8 ± 4.0	...	-1.33 ± 0.09	...	...	...	...	01 19 47.65	-20 54 46.8	N
TN J0121+1320	348 ± 33	57.3 ± 2.7	8.4 ± 0.5	-1.34 ± 0.08	-1.55 ± 0.06	S	0.3	89	01 21 42.74	+13 20 58.3	V
TN J0154+0044	465 ± 29	78.6 ± 3.6	...	-1.32 ± 0.06	...	...	...	...	01 54 38.48	+00 44 37.7	N
TN J0156+1619	437 ± 45	62.9 ± 2.9	...	-1.44 ± 0.09	...	...	...	...	01 56 18.01	+16 19 52.5	N
TN J0201-1302	360 ± 33	57.0 ± 2.7	11.0 ± 0.6	-1.37 ± 0.08	-1.32 ± 0.06	S	< 1.3	0	02 01 15.80	-13 02 19.4	V
TN J0205+2242	381 ± 26	60.4 ± 2.8	8.8 ± 0.5	-1.37 ± 0.06	-1.55 ± 0.06	D	2.7	148	02 05 10.69	+22 42 50.2	V
TN J0208+1305	403 ± 61	68.3 ± 3.1	...	-1.32 ± 0.12	...	...	...	...	02 08 32.45	+13 05 53.0	N
TN J0215+2651	431 ± 49	69.3 ± 3.2	...	-1.36 ± 0.09	...	...	...	...	02 15 45.31	+26 51 12.3	N
TN J0218+0844	395 ± 32	61.8 ± 1.3	8.8 ± 0.5	-1.38 ± 0.06	-1.56 ± 0.05	S	0.4	169	02 18 25.56	+08 44 31.3	V
TN J0230-0255	573 ± 56	90.1 ± 4.0	22.5 ± 1.2	-1.38 ± 0.08	-1.12 ± 0.06	M	6.7	156	02 30 21.97	-02 55 05.1	V
TN J0230-2001	380 ± 33	57.8 ± 2.7	...	-1.40 ± 0.07	...	...	...	...	02 30 45.73	-20 01 19.0	N
TN J0233+2349*	1075 ± 86	176.8 ± 7.8	43.5 ± 2.3	-1.34 ± 0.07	-1.13 ± 0.06	DF	4.0	116	02 33 10.32	+23 49 52.2	V
TN J0234-1215	451 ± 46	64.2 ± 2.9	...	-1.45 ± 0.08	...	...	...	...	02 34 57.31	-12 15 21.7	N
TN J0243+1405	395 ± 51	58.4 ± 1.3	26.2 ± 1.4	-1.42 ± 0.10	-0.64 ± 0.05	T	3.1	126	02 43 10.52	+14 05 36.9	V
TN J0244+0327	542 ± 62	84.8 ± 3.8	...	-1.38 ± 0.09	...	...	...	...	02 44 53.97	+03 27 46.8	N
TN J0245+2700	388 ± 53	54.0 ± 2.5	...	-1.47 ± 0.11	...	...	...	...	02 45 22.27	+27 00 06.3	N
TN J0250+0130	275 ± 25	39.1 ± 2.0	...	-1.45 ± 0.08	...	...	...	...	02 50 20.94	+01 30 24.9	N
TN J0254-1039	421 ± 60	68.1 ± 3.1	25.5 ± 1.4	-1.36 ± 0.11	-0.79 ± 0.06	D	11.7	129	02 54 59.89	-10 39 49.2	V
TN J0256-2717	1045 ± 70	177.9 ± 7.4	...	-1.32 ± 0.06	...	...	...	...	02 56 51.44	-27 17 56.5	N
TN J0301+0155	3950 ± 180	402.7 ± 17.3	...	-1.70 ± 0.05	...	...	...	...	03 01 38.49	+01 55 14.9	N
TN J0306+0524	909 ± 76	155.1 ± 6.3	...	-1.32 ± 0.07	...	...	...	...	03 06 08.10	+05 24 23.8	N
TN J0309-2425*	483 ± 53	62.6 ± 2.9	14.8 ± 0.9	-1.52 ± 0.09	-1.16 ± 0.06	D	12.9	23	03 09 09.26	-24 25 11.7	V
TN J0310+0814	593 ± 67	96.8 ± 4.3	...	-1.35 ± 0.09	...	...	...	...	03 10 37.37	+08 14 59.2	N
TN J0311-2553	1102 ± 66	174.1 ± 7.5	...	-1.37 ± 0.06	...	...	...	...	03 11 47.40	-25 53 21.1	N
TN J0312-2622	442 ± 41	46.4 ± 2.3	...	-1.68 ± 0.08	...	...	...	...	03 12 34.19	-26 22 23.9	N
TN J0316-0815	1950 ± 92	305.1 ± 13.4	42.2 ± 2.2	-1.38 ± 0.05	-1.59 ± 0.05	T	3.2	34	03 16 18.59	-08 15 54.8	V

Table A.2. continued

Name	$S_{365}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{365}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
TN J0321+1631	1084 ± 85	175.7 ± 7.7	...	-1.35 ± 0.07	...	...	...	...	03 21 36.86	+16 31 55.8	N
TN J0327-0948	281 ± 30	39.0 ± 2.0	...	-1.47 ± 0.09	...	...	...	...	03 27 00.55	-09 48 05.2	N
TN J0344+1903	297 ± 22	50.7 ± 2.4	...	-1.32 ± 0.07	...	...	...	...	03 44 00.12	+19 03 53.1	N
TN J0344-2029	538 ± 71	74.0 ± 1.6	19.2 ± 1.1	-1.48 ± 0.10	-1.08 ± 0.05	D	6.5	42	03 44 37.88	-20 29 03.8	V
TN J0349-1207*	460 ± 61	77.5 ± 3.5	23.3 ± 1.3	-1.32 ± 0.11	-0.97 ± 0.06	T	9.4	101	03 49 49.22	-12 07 10.6	V
TN J0351-1947	642 ± 86	111.4 ± 4.9	...	-1.30 ± 0.11	...	...	...	...	03 51 33.13	-19 47 10.5	N
TN J0352-0355*	681 ± 63	114.9 ± 4.8	13.5 ± 0.8	-1.32 ± 0.08	-1.65 ± 0.06	DF	...	0	03 52 32.04	-03 55 47.2	V
TN J0355+0440	2015 ± 119	262.0 ± 11.5	...	-1.52 ± 0.05	...	...	...	...	03 55 12.79	+04 40 41.0	N
TN J0356-3028	1030 ± 65	167.5 ± 7.3	34.1 ± 1.8	-1.35 ± 0.06	-1.28 ± 0.06	D	3.1	86	03 56 45.54	-30 28 36.9	V
TN J0401-0156	1001 ± 65	172.8 ± 7.1	...	-1.31 ± 0.06	...	...	...	...	04 01 08.43	-01 56 08.9	N
TN J0402+1007	479 ± 54	74.6 ± 3.4	13.4 ± 0.8	-1.38 ± 0.09	-1.38 ± 0.06	D	0.8	67	04 02 34.44	+10 07 14.9	V
TN J0404-0541	666 ± 69	114.1 ± 5.1	...	-1.31 ± 0.08	...	...	...	...	04 04 09.66	-05 41 07.3	N
TN J0408-2418*	3881 ± 172	647.7 ± 28.1	...	-1.33 ± 0.05	...	...	...	...	04 08 51.38	-24 18 15.5	N
TN J0410+1019	1185 ± 88	166.9 ± 3.4	26.5 ± 1.4	-1.46 ± 0.06	-1.26 ± 0.05	S	0.2	109	04 10 40.52	+10 19 13.6	V
TN J0429-2118	561 ± 58	95.5 ± 4.3	...	-1.32 ± 0.08	...	...	...	...	04 29 39.79	-21 18 01.7	N
TN J0431+0725	175 ± 24	30.4 ± 0.7	...	-1.30 ± 0.10	...	...	...	...	04 31 54.21	+07 25 13.8	N
TN J0433+0717	229 ± 27	33.1 ± 1.8	...	-1.44 ± 0.10	...	...	...	...	04 33 05.61	+07 17 55.2	N
TN J0435-0736	358 ± 71	47.0 ± 2.3	16.6 ± 0.9	-1.51 ± 0.15	-0.84 ± 0.06	T	11.9	59	04 35 32.20	-07 36 12.3	V
TN J0437+0259	410 ± 54	71.1 ± 3.3	...	-1.30 ± 0.10	...	...	...	...	04 37 33.04	+02 59 06.2	N
TN J0443-1212*	412 ± 59	56.8 ± 2.7	14.6 ± 0.8	-1.47 ± 0.11	-1.09 ± 0.06	M	20.2	25	04 43 53.72	-12 12 46.7	V
TN J0452-1737	747 ± 73	118.1 ± 2.5	23.7 ± 1.3	-1.37 ± 0.07	-1.29 ± 0.05	D	2.4	176	04 52 26.66	-17 37 54.0	V
TN J0510-1838	6801 ± 325	641.3 ± 25.6	...	-1.76 ± 0.05	...	D	36.1	63	05 10 32.30	-18 38 41.4	A
TN J0515-3410	589 ± 52	97.4 ± 4.4	...	-1.34 ± 0.07	...	...	...	...	05 15 18.86	-34 10 09.5	N
TN J0517-0641	285 ± 30	40.0 ± 0.9	6.8 ± 0.4	-1.46 ± 0.08	-1.43 ± 0.05	S	1.0	0	05 17 36.89	-06 41 14.6	V
TN J0525-1832	549 ± 73	86.4 ± 3.9	24.1 ± 1.3	-1.38 ± 0.11	-1.02 ± 0.06	T	8.1	172	05 25 54.85	-18 32 32.5	V
TN J0547-0706	2552 ± 156	362.6 ± 16.0	...	-1.45 ± 0.06	...	...	...	...	05 47 27.57	-07 06 35.0	N
TN J0549-2459	427 ± 55	58.0 ± 1.3	...	-1.49 ± 0.10	...	S	23.5	8	05 49 28.61	-24 59 15.2	A
TN J0551-0756	473 ± 71	68.5 ± 3.2	...	-1.44 ± 0.12	...	...	...	...	05 51 30.37	-07 56 17.5	N
TN J0552-0433	1618 ± 83	269.7 ± 11.8	37.7 ± 2.0	-1.33 ± 0.05	-1.56 ± 0.06	D	4.3	152	05 52 14.38	-04 33 28.8	V
TN J0553-2033	887 ± 80	148.0 ± 6.2	...	-1.33 ± 0.07	...	...	...	...	05 53 37.24	-20 33 07.6	N
TN J0558-1358	425 ± 54	73.4 ± 3.4	...	-1.31 ± 0.10	...	...	...	...	05 58 39.43	-13 58 45.5	N
TN J0559-3406	773 ± 69	123.5 ± 5.4	...	-1.36 ± 0.07	...	S	4.6	8	05 59 14.30	-34 06 51.0	A
TN J0602-2741	810 ± 54	117.2 ± 5.2	20.7 ± 1.2	-1.44 ± 0.06	-1.39 ± 0.06	T	4.3	88	06 02 41.32	-27 41 10.1	V
TN J0630-2834	343 ± 29	14.3 ± 0.5	...	-2.36 ± 0.07	...	S	13.5	2	06 30 49.49	-28 34 40.4	A
TN J0634-3232	394 ± 63	67.7 ± 3.1	...	-1.31 ± 0.12	...	...	...	...	06 34 17.10	-32 32 34.1	N
TN J0729+2436	1062 ± 87	68.9 ± 3.1	...	-2.03 ± 0.07	...	D	19.4	72	07 29 28.65	+24 36 26.2	F
TN J0733+1753	291 ± 45	46.1 ± 2.3	...	-1.37 ± 0.12	...	...	...	...	07 33 17.53	+17 53 28.4	N
TN J0805+2738	987 ± 79	153.0 ± 6.7	...	-1.39 ± 0.07	...	T	71.3	119	08 05 15.50	+27 38 01.2	F
TN J0806+0401	560 ± 53	92.1 ± 2.0	...	-1.34 ± 0.07	...	...	...	...	08 06 32.25	+04 01 37.7	N
TN J0812+0915	277 ± 46	43.4 ± 2.2	...	-1.38 ± 0.13	...	...	...	...	08 12 16.84	+09 15 43.7	N
TN J0818-0741	381 ± 48	59.3 ± 2.8	...	-1.38 ± 0.10	...	...	...	...	08 18 51.89	-07 41 48.2	N
TN J0831+0046	451 ± 55	72.8 ± 3.3	...	-1.36 ± 0.10	...	...	...	...	08 31 39.92	+00 46 56.5	N
TN J0831+0851	567 ± 53	83.8 ± 3.8	98.3 ± 5.2	-1.42 ± 0.08	0.13 ± 0.06	DF	3.0	2	08 31 52.74	+08 51 14.1	V
TN J0837-1053*	450 ± 81	69.2 ± 3.1	11.6 ± 0.7	-1.39 ± 0.14	-1.43 ± 0.06	D	23.5	11	08 37 41.62	-10 53 46.0	V
TN J0855-0000	416 ± 57	65.9 ± 3.0	13.8 ± 0.8	-1.37 ± 0.11	-1.26 ± 0.06	D	3.8	85	08 55 57.26	-00 00 58.2	V
TN J0856-1510	469 ± 68	70.0 ± 3.2	3.7 ± 0.3	-1.41 ± 0.11	-2.01 ± 0.07	S	1.0	14	08 56 12.44	-15 10 35.7	V
TN J0902+1809	491 ± 37	55.5 ± 2.6	...	-1.62 ± 0.07	...	...	...	...	09 02 00.00	+18 09 03.4	N
TN J0906-0701	939 ± 101	142.5 ± 5.8	...	-1.40 ± 0.09	...	...	...	...	09 06 40.23	-07 01 12.4	N
TN J0910-2228	465 ± 32	55.3 ± 2.6	4.0 ± 0.3	-1.58 ± 0.06	-2.11 ± 0.07	S	2.0	0	09 10 34.15	-22 28 47.4	V
TN J0914+1119	343 ± 53	49.5 ± 1.1	...	-1.44 ± 0.12	...	...	...	...	09 14 30.81	+11 19 08.0	N
TN J0919+1845	522 ± 50	90.0 ± 4.0	...	-1.31 ± 0.08	...	...	...	...	09 19 33.25	+18 45 51.2	N
TN J0920-0712	760 ± 53	100.4 ± 4.5	16.3 ± 0.9	-1.51 ± 0.06	-1.46 ± 0.06	S	1.4	135	09 20 22.43	-07 12 17.6	V
TN J0920-1405	842 ± 88	133.6 ± 5.5	...	-1.37 ± 0.08	...	...	...	...	09 20 37.77	-14 05 56.3	N
TN J0924-2201	656 ± 72	73.3 ± 1.5	8.6 ± 0.5	-1.63 ± 0.08	-1.72 ± 0.05	D	1.2	74	09 24 19.92	-22 01 41.5	V
TN J0936+0422	6025 ± 362	992.0 ± 42.5	...	-1.34 ± 0.05	...	...	...	...	09 36 32.02	+04 22 10.7	N
TN J0936-2243	488 ± 63	69.1 ± 3.2	18.5 ± 1.0	-1.45 ± 0.10	-1.06 ± 0.06	D	8.1	77	09 36 32.53	-22 43 04.9	V
TN J0941-1628	1810 ± 115	304.4 ± 13.4	64.7 ± 3.8	-1.33 ± 0.06	-1.24 ± 0.06	D	1.9	175	09 41 07.43	-16 28 02.5	V
TN J0946-0306	238 ± 24	38.2 ± 2.0	...	-1.36 ± 0.09	...	...	...	...	09 46 00.88	-03 06 26.2	N
TN J0954-1635	1531 ± 98	260.0 ± 11.4	...	-1.32 ± 0.06	...	...	...	...	09 54 52.30	-16 35 35.1	N
TN J0956-1500	505 ± 49	68.9 ± 1.5	...	-1.48 ± 0.07	...	...	...	...	09 56 01.46	-15 00 21.6	N
TN J0958-1103	431 ± 63	70.6 ± 3.2	...	-1.35 ± 0.11	...	...	...	...	09 58 21.84	-11 03 47.5	N
TN J1000+0549	410 ± 55	67.3 ± 3.1	...	-1.34 ± 0.11	...	...	...	...	10 00 44.58	+05 49 31.1	N
TN J1002+2225	343 ± 41	54.2 ± 2.6	12.2 ± 0.7	-1.37 ± 0.10	-1.20 ± 0.06	D	11.2	10	10 02 54.08	+22 25 21.9	V
TN J1026-2116	399 ± 29	61.5 ± 1.3	8.5 ± 0.5	-1.39 ± 0.06	-1.59 ± 0.05	S	1.0	0	10 26 22.37	-21 16 07.7	V
TN J1028+1114	459 ± 56	79.7 ± 3.6	...	-1.30 ± 0.10	...	...	...	...	10 28 24.31	+11 14 52.4	N

Table A.2. continued

Name	$S_{365}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{365}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
TN J1031+0259	416 ± 62	60.0 ± 1.3	...	-1.44 ± 0.11	...	...	...	...	10 31 13.36	+02 59 05.3	N
TN J1033-1339	981 ± 75	153.8 ± 6.8	26.0 ± 1.4	-1.38 ± 0.07	-1.43 ± 0.06	S	2.0	107	10 33 10.70	-13 39 52.0	V
TN J1043-1718	578 ± 70	91.0 ± 3.9	18.9 ± 1.0	-1.38 ± 0.10	-1.26 ± 0.06	D	34.7	63	10 43 19.42	-17 18 53.5	V
TN J1043+2404	439 ± 34	52.0 ± 2.5	...	-1.59 ± 0.07	...	S	3.7	90	10 43 43.28	+24 04 47.3	F
TN J1045+1832	348 ± 27	53.6 ± 1.2	...	-1.39 ± 0.06	...	...	...	...	10 45 42.93	+18 32 37.5	N
TN J1049-1258	818 ± 56	112.2 ± 5.0	17.2 ± 0.9	-1.48 ± 0.06	-1.50 ± 0.06	M	10.1	104	10 49 06.22	-12 58 18.3	V
TN J1053-1518	383 ± 62	63.7 ± 2.9	...	-1.33 ± 0.13	...	...	...	...	10 53 44.52	-15 18 29.8	N
TN J1056-0400	449 ± 57	57.2 ± 2.7	...	-1.53 ± 0.10	...	...	...	...	10 56 52.64	-04 00 15.9	N
TN J1102+1029	3713 ± 224	633.6 ± 27.8	...	-1.32 ± 0.06	...	...	...	...	11 02 17.47	+10 29 08.0	N
TN J1102-1651	688 ± 68	112.5 ± 5.0	20.2 ± 1.1	-1.35 ± 0.08	-1.38 ± 0.06	D	3.0	70	11 02 47.13	-16 51 34.4	V
TN J1112-2948	663 ± 51	101.0 ± 4.5	16.6 ± 0.9	-1.40 ± 0.07	-1.45 ± 0.06	D	9.1	119	11 12 23.86	-29 48 06.4	V
TN J1114-2221	386 ± 53	64.9 ± 3.0	...	-1.33 ± 0.11	...	...	...	...	11 14 04.82	-22 21 26.6	N
TN J1117-1409	785 ± 69	134.5 ± 5.9	...	-1.31 ± 0.07	...	...	...	...	11 17 23.00	-14 09 09.5	N
TN J1118-2612	514 ± 44	62.3 ± 2.9	...	-1.57 ± 0.07	...	...	...	...	11 18 54.08	-26 12 20.6	N
TN J1121-1155	332 ± 65	50.1 ± 2.4	...	-1.41 ± 0.15	...	...	...	...	11 21 32.75	-11 55 03.2	N
TN J1121+1706	1156 ± 83	196.8 ± 8.4	...	-1.32 ± 0.06	...	...	...	...	11 21 39.89	+17 06 14.8	N
TN J1123-2154	407 ± 64	50.6 ± 2.4	8.5 ± 0.5	-1.55 ± 0.12	-1.43 ± 0.06	S	0.8	29	11 23 10.15	-21 54 05.3	V
TN J1125-0342	354 ± 52	61.4 ± 2.9	...	-1.30 ± 0.11	...	...	...	...	11 25 57.41	-03 42 04.0	N
TN J1133-2715	2938 ± 125	447.9 ± 9.2	...	-1.40 ± 0.04	...	...	...	...	11 33 31.53	-27 15 22.9	N
TN J1136+1551	384 ± 33	21.7 ± 0.6	...	-2.14 ± 0.07	...	...	...	...	11 36 03.28	+15 51 06.7	N
TN J1136+0610	698 ± 50	118.8 ± 5.3	...	-1.32 ± 0.06	...	...	...	...	11 36 38.91	+06 10 26.8	N
TN J1139+0935	508 ± 58	87.3 ± 3.9	...	-1.31 ± 0.09	...	...	...	...	11 39 22.75	+09 35 26.1	N
TN J1146-1052	843 ± 79	120.9 ± 5.4	...	-1.44 ± 0.08	...	S	8.8	1	11 46 07.21	-10 52 08.6	A
TN J1148-0901	571 ± 67	80.4 ± 3.7	...	-1.46 ± 0.09	...	S	< 6.0	5	11 48 39.90	-09 01 48.8	A
TN J1149+1844	457 ± 39	77.6 ± 1.6	...	-1.32 ± 0.07	...	...	...	...	11 49 36.78	+18 44 21.5	N
TN J1151-3013	435 ± 28	52.7 ± 1.2	7.0 ± 0.4	-1.57 ± 0.05	-1.62 ± 0.05	S	1.0	0	11 51 59.42	-30 13 40.8	V
TN J1156-3105	317 ± 48	48.3 ± 2.3	...	-1.40 ± 0.12	...	S	10.7	167	11 56 25.60	-31 05 40.0	A
TN J1159-1629	324 ± 34	52.9 ± 1.2	10.4 ± 0.6	-1.35 ± 0.08	-1.30 ± 0.05	D	1.4	38	11 59 53.25	-16 29 48.2	V
TN J1204+1630	804 ± 56	129.7 ± 5.8	...	-1.36 ± 0.06	...	...	...	...	12 04 20.49	+16 30 49.5	N
TN J1210+1738	397 ± 50	68.1 ± 3.2	...	-1.31 ± 0.10	...	...	...	...	12 10 36.00	+17 38 27.8	N
TN J1216+1944	546 ± 44	91.7 ± 4.1	...	-1.33 ± 0.07	...	...	...	...	12 16 01.76	+19 44 43.3	N
TN J1216-2850	870 ± 43	145.2 ± 6.4	...	-1.33 ± 0.05	...	S	< 7.0	163	12 16 17.38	-28 50 48.7	A
TN J1220+0604	491 ± 61	65.1 ± 3.0	...	-1.50 ± 0.10	...	...	...	...	12 20 07.82	+06 04 16.5	N
TN J1221-2646	1403 ± 82	217.1 ± 9.5	...	-1.39 ± 0.05	...	S	8.3	162	12 21 42.91	-26 46 37.7	A
TN J1227-2255	271 ± 27	44.0 ± 2.2	...	-1.35 ± 0.08	...	S	< 7.0	158	12 27 47.23	-22 55 35.1	A
TN J1239+1005	830 ± 89	79.3 ± 1.7	...	-1.75 ± 0.08	...	...	...	...	12 39 02.08	+10 05 49.2	N
TN J1239+1043	569 ± 63	87.1 ± 3.8	...	-1.40 ± 0.09	...	...	...	...	12 39 31.67	+10 43 22.9	N
TN J1239-0319	1629 ± 74	277.3 ± 12.2	...	-1.32 ± 0.05	...	...	...	...	12 39 38.91	-03 19 02.2	N
TN J1245-1127	518 ± 62	88.0 ± 4.0	...	-1.32 ± 0.10	...	...	...	...	12 45 02.47	-11 27 25.9	N
TN J1247+1547	405 ± 44	70.1 ± 3.2	...	-1.30 ± 0.09	...	...	...	...	12 47 04.84	+15 47 38.9	N
TN J1251-2714	609 ± 53	96.5 ± 4.3	...	-1.37 ± 0.07	...	S	< 7.0	158	12 51 14.86	-27 14 19.9	A
TN J1256-0911	457 ± 38	76.4 ± 3.5	...	-1.33 ± 0.07	...	...	...	...	12 56 19.48	-09 11 24.8	N
TN J1303-3349	2431 ± 108	360.8 ± 7.7	...	-1.42 ± 0.04	...	...	...	...	13 03 02.67	-33 49 15.5	N
TN J1306-0436	331 ± 57	55.2 ± 2.6	...	-1.33 ± 0.13	...	...	...	...	13 06 36.11	-04 36 17.4	N
TN J1313-0459	369 ± 34	58.1 ± 2.7	...	-1.38 ± 0.08	...	...	...	...	13 13 34.37	-04 59 30.0	N
TN J1317+0339	399 ± 62	39.8 ± 2.0	...	-1.71 ± 0.12	...	...	...	...	13 17 48.12	+03 39 12.3	N
TN J1323-2604	564 ± 60	97.3 ± 2.0	...	-1.31 ± 0.08	...	...	...	...	13 23 32.43	-26 04 04.7	N
TN J1326-2330	523 ± 33	83.2 ± 3.8	...	-1.37 ± 0.06	...	...	...	...	13 26 25.24	-23 30 23.5	N
TN J1327+1437	788 ± 68	115.0 ± 5.1	...	-1.43 ± 0.07	...	...	...	...	13 27 23.93	+14 37 00.4	N
TN J1336+2450	358 ± 71	51.3 ± 2.5	...	-1.45 ± 0.15	...	M	54.4	100	13 36 15.12	+24 50 05.4	F
TN J1338-1942	718 ± 63	122.9 ± 5.4	22.1 ± 1.2	-1.31 ± 0.07	-1.34 ± 0.06	S	1.4	152	13 38 26.06	-19 42 30.1	V
TN J1339-0114	349 ± 51	59.5 ± 1.3	...	-1.32 ± 0.11	...	...	...	...	13 39 55.26	-01 14 10.8	N
TN J1340+2723	827 ± 93	137.8 ± 6.1	24.8 ± 1.4	-1.33 ± 0.09	-1.38 ± 0.06	DF	1.7	0	13 40 29.38	+27 23 25.0	V
TN J1346-1004	363 ± 59	49.6 ± 2.4	...	-1.48 ± 0.13	...	...	...	...	13 46 16.84	-10 04 57.2	N
TN J1351+1328	662 ± 54	93.8 ± 4.2	...	-1.45 ± 0.07	...	...	...	...	13 51 54.72	+13 28 38.1	N

Table A.2. continued

Name	$S_{365}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{365}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
TN J1352-0800	554 ± 88	95.6 ± 4.3	...	-1.31 ± 0.12	...	...	...	...	13 52 55.32	-08 00 14.1	N
TN J1353+2302	403 ± 38	59.5 ± 1.3	23.3 ± 1.3	-1.42 ± 0.07	-0.75 ± 0.05	DF	2.5	44	13 53 42.26	+23 02 50.9	V
TN J1358+2003	227 ± 42	35.0 ± 1.8	...	-1.39 ± 0.14	...	...	...	...	13 58 29.63	+20 03 13.4	N
TN J1403-1223	396 ± 69	65.8 ± 3.0	...	-1.34 ± 0.14	...	S	30.6	169	14 03 37.33	-12 23 57.3	A
TN J1408-0855	694 ± 73	114.3 ± 5.1	...	-1.34 ± 0.09	...	...	...	...	14 08 18.46	-08 55 38.5	N
TN J1411-0824	542 ± 71	76.1 ± 3.5	...	-1.46 ± 0.10	...	...	...	...	14 11 53.33	-08 24 02.9	N
TN J1412-2548	452 ± 59	78.0 ± 3.6	...	-1.31 ± 0.10	...	...	...	...	14 12 51.27	-25 48 05.5	N
TN J1418-2256	527 ± 59	82.0 ± 3.7	...	-1.38 ± 0.09	...	S	7.6	148	14 18 16.65	-22 56 58.6	A
TN J1428+2425	411 ± 34	69.1 ± 1.5	...	-1.33 ± 0.07	...	S	1.0	0	14 28 10.20	+24 25 11.3	F
TN J1438+2334	564 ± 69	96.5 ± 4.2	...	-1.31 ± 0.10	...	D	21.1	157	14 38 56.52	+23 34 25.6	F
TN J1439-3226	622 ± 65	90.7 ± 1.9	...	-1.43 ± 0.08	...	...	...	...	14 39 02.87	-32 26 46.6	N
TN J1449+1440	464 ± 50	80.3 ± 3.7	...	-1.30 ± 0.09	...	...	...	...	14 49 02.50	+14 40 42.2	N
TN J1451+2351	595 ± 44	93.2 ± 4.2	...	-1.38 ± 0.07	...	D	12.6	109	14 51 10.71	+23 51 38.8	F
TN J1452-1113	344 ± 55	53.9 ± 2.6	...	-1.38 ± 0.13	...	S	< 7.5	2	14 52 04.88	-11 13 35.7	A
TN J1452+2013	291 ± 40	43.4 ± 1.0	...	-1.42 ± 0.11	...	...	...	...	14 52 22.93	+20 13 09.8	N
TN J1453+1106	833 ± 56	144.9 ± 6.4	...	-1.30 ± 0.06	...	...	...	...	14 53 21.97	+11 06 04.9	N
TN J1454+0017	1309 ± 88	214.8 ± 9.2	...	-1.34 ± 0.06	...	...	...	...	14 54 40.87	+00 17 41.9	N
TN J1459-2730	321 ± 40	52.5 ± 2.5	...	-1.35 ± 0.10	...	S	6.5	143	14 59 34.96	-27 30 08.3	A
TN J1506+2728	360 ± 58	60.7 ± 1.3	15.9 ± 0.9	-1.32 ± 0.12	-1.07 ± 0.05	D	1.1	131	15 06 35.41	+27 28 55.6	V
TN J1513-2417	504 ± 33	83.6 ± 3.7	...	-1.34 ± 0.06	...	...	...	...	15 13 31.67	-24 17 55.3	N
TN J1513-1801	534 ± 36	85.3 ± 3.9	...	-1.36 ± 0.10	...	S	< 7.5	4	15 13 55.31	-18 01 07.9	A
TN J1515-2651	352 ± 44	58.0 ± 2.7	...	-1.34 ± 0.06	...	S	6.5	9	15 15 59.29	-26 51 14.9	A
TN J1520+1410	478 ± 55	66.0 ± 1.4	...	-1.47 ± 0.09	...	...	...	...	15 20 53.36	+14 10 45.5	N
TN J1521+0741	1777 ± 123	105.6 ± 4.5	...	-2.10 ± 0.06	...	...	...	...	15 21 50.75	+07 41 41.1	N
TN J1522-2540	464 ± 63	54.1 ± 2.6	...	-1.60 ± 0.11	...	S	8.2	10	15 22 23.28	-25 40 07.1	A
TN J1524+0642	479 ± 55	82.9 ± 1.7	...	-1.30 ± 0.09	...	...	...	...	15 24 06.39	+06 42 15.9	N
TN J1531-3234	522 ± 59	80.6 ± 3.6	...	-1.39 ± 0.09	...	S	4.8	27	15 31 33.59	-32 34 01.7	A
TN J1547-2218	427 ± 62	72.4 ± 3.3	...	-1.32 ± 0.11	...	...	...	...	15 47 49.95	-22 18 25.6	N
TN J1556-2759	612 ± 56	66.4 ± 3.1	...	-1.65 ± 0.08	...	...	...	...	15 56 43.63	-27 59 43.1	N
TN J1559-1727	575 ± 71	97.4 ± 4.3	...	-1.32 ± 0.10	...	...	...	...	15 59 34.08	-17 27 41.6	N
TN J1604-0248	345 ± 33	46.8 ± 1.0	...	-1.49 ± 0.07	...	...	...	...	16 04 59.59	-02 48 47.9	N
TN J1621+2043	539 ± 48	61.4 ± 2.9	...	-1.62 ± 0.08	...	...	...	...	16 21 06.18	+20 43 36.4	N
TN J1628+1604	602 ± 44	55.5 ± 2.6	...	-1.77 ± 0.07	...	...	...	...	16 28 19.16	+16 04 00.0	N
TN J1634-2222	4714 ± 214	809.2 ± 35.4	...	-1.31 ± 0.05	...	...	...	...	16 34 49.79	-22 22 12.7	N
TN J1637-1931	432 ± 40	36.8 ± 0.9	...	-1.83 ± 0.07	...	S	10.8	6	16 37 44.80	-19 31 24.8	A
TN J1646-1328	608 ± 63	102.5 ± 4.6	...	-1.32 ± 0.09	...	...	...	...	16 46 51.67	-13 28 49.1	N
TN J1652-1339	328 ± 33	56.5 ± 1.3	...	-1.31 ± 0.08	...	...	...	...	16 52 24.29	-13 39 04.3	N
TN J1653-1155	1131 ± 60	197.0 ± 8.7	...	-1.30 ± 0.05	...	...	...	...	16 53 52.81	-11 55 59.2	N
TN J1656-1521	587 ± 79	93.9 ± 4.2	...	-1.36 ± 0.11	...	...	...	...	16 56 03.77	-15 21 23.3	N
TN J1701-0101	999 ± 55	172.2 ± 7.6	...	-1.31 ± 0.05	...	...	...	...	17 01 37.58	-01 01 13.6	N
TN J1701+0252	356 ± 38	57.1 ± 2.7	...	-1.36 ± 0.09	...	...	...	...	17 01 54.89	+02 52 13.4	N
TN J1702-0811	436 ± 38	69.9 ± 3.2	...	-1.36 ± 0.07	...	...	...	...	17 02 04.71	-08 11 07.5	N
TN J1702+2015	322 ± 48	43.4 ± 2.1	...	-1.49 ± 0.12	...	...	...	...	17 02 23.02	+20 15 34.2	N
TN J1702-0145	808 ± 74	135.6 ± 6.0	...	-1.33 ± 0.08	...	...	...	...	17 02 43.11	-01 45 54.0	N
TN J1707-1120	425 ± 72	54.2 ± 2.6	...	-1.53 ± 0.13	...	...	...	...	17 07 51.37	-11 20 21.0	N
TN J1710-0826	445 ± 74	69.2 ± 3.2	...	-1.38 ± 0.13	...	...	...	...	17 10 20.36	-08 26 40.3	N
TN J1714+2226	285 ± 48	43.3 ± 1.0	...	-1.40 ± 0.13	...	S	1.6	38	17 14 53.42	+22 26 18.5	F
TN J1821+2433	1416 ± 95	227.5 ± 4.7	...	-1.36 ± 0.05	...	...	...	...	18 21 37.07	+24 33 13.8	N
TN J1855-3430	654 ± 85	87.8 ± 3.8	...	-1.49 ± 0.10	...	...	...	...	18 55 24.12	-34 30 27.7	N
TN J1905-3520	799 ± 75	131.3 ± 5.4	...	-1.34 ± 0.08	...	...	...	...	19 05 22.67	-35 20 50.9	N
TN J1932-1931	5678 ± 335	813.6 ± 35.5	...	-1.45 ± 0.05	...	...	...	...	19 32 07.22	-19 31 49.7	N
TN J1941-1952	1194 ± 103	183.0 ± 3.8	...	-1.40 ± 0.07	...	S	7.9	164	19 41 00.07	-19 52 14.0	A
TN J1953-0541	478 ± 83	59.0 ± 2.8	...	-1.56 ± 0.13	...	...	...	...	19 53 27.04	-05 41 28.5	N
TN J1954-1207	657 ± 67	102.8 ± 4.6	...	-1.38 ± 0.08	...	S	< 7.0	177	19 54 24.15	-12 07 48.7	A
TN J2007-1316	777 ± 46	115.4 ± 5.2	...	-1.42 ± 0.06	...	S	7.2	174	20 07 53.23	-13 16 43.6	A
TN J2008-1344	798 ± 71	132.3 ± 2.7	...	-1.34 ± 0.07	...	S	8.2	176	20 08 07.48	-13 44 17.8	A
TN J2009-3040	409 ± 65	65.3 ± 3.0	...	-1.36 ± 0.12	...	S	< 7.0	144	20 09 48.13	-30 40 07.0	A
TN J2014-2115	348 ± 33	48.0 ± 1.1	...	-1.47 ± 0.07	...	D	56.4	11	20 14 31.96	-21 14 36.7	A
TN J2021+0839	880 ± 72	145.4 ± 6.4	...	-1.34 ± 0.07	...	...	...	...	20 21 33.11	+08 39 09.2	N
TN J2028+0811	455 ± 58	74.8 ± 1.6	...	-1.34 ± 0.10	...	...	...	...	20 28 37.12	+08 11 21.4	N



Table A.2. continued

Name	$S_{365}$ mJy	$S_{1400}$ mJy	$S_{4850}$ mJy	$\alpha_{365}^{1400}$	$\alpha_{1400}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
TN J2028−1934	814 ± 88	134.4 ± 5.8	...	−1.34 ± 0.09	...	S	19.2	169	20 28 48.57	−19 34 03.2	A
TN J2029−0858	396 ± 34	47.2 ± 2.3	...	−1.58 ± 0.07	...	...	...	...	20 29 49.60	−08 58 48.5	N
TN J2034−2735	510 ± 70	67.4 ± 1.5	...	−1.51 ± 0.10	...	S	< 8.0	163	20 34 15.74	−27 35 45.6	A
TN J2042+1354	281 ± 45	42.0 ± 1.0	...	−1.41 ± 0.12	...	...	...	...	20 42 07.25	+13 54 16.6	N
TN J2048−0618	881 ± 99	150.7 ± 6.6	...	−1.31 ± 0.09	...	...	...	...	20 48 06.24	−06 18 40.5	N
TN J2049−2223	823 ± 86	126.1 ± 5.6	...	−1.40 ± 0.08	...	D	95.0	97	20 49 43.04	−22 22 56.3	A
TN J2051−0527	377 ± 79	54.7 ± 1.2	...	−1.44 ± 0.16	...	...	...	...	20 51 13.39	−05 27 06.1	N
TN J2051+0133	391 ± 33	66.6 ± 3.1	...	−1.32 ± 0.07	...	...	...	...	20 51 32.31	+01 33 34.9	N
TN J2052−0555	321 ± 35	41.5 ± 0.9	...	−1.52 ± 0.08	...	...	...	...	20 52 49.93	−05 55 49.1	N
TN J2102+2158	483 ± 52	73.5 ± 1.6	...	−1.40 ± 0.08	...	...	...	...	21 02 52.80	+21 58 35.1	N
TN J2103−1917	1410 ± 109	228.6 ± 10.1	...	−1.35 ± 0.07	...	...	...	...	21 03 42.91	−19 17 47.7	N
TN J2106−2405*	2232 ± 150	366.3 ± 15.9	...	−1.34 ± 0.06	...	...	...	...	21 06 58.42	−24 05 04.2	N
TN J2107+0933	476 ± 56	68.0 ± 3.1	...	−1.45 ± 0.09	...	...	...	...	21 07 10.61	+09 33 32.5	N
TN J2108+0249	573 ± 41	65.8 ± 3.1	...	−1.61 ± 0.06	...	...	...	...	21 08 26.75	+02 49 46.3	N
TN J2113+0111	553 ± 68	90.8 ± 4.1	...	−1.34 ± 0.10	...	...	...	...	21 13 50.43	+01 11 22.3	N
TN J2125+2013	352 ± 45	51.9 ± 2.5	...	−1.42 ± 0.10	...	...	...	...	21 25 51.74	+20 13 39.8	N
TN J2133+1629	1737 ± 123	283.4 ± 12.4	...	−1.35 ± 0.06	...	...	...	...	21 33 29.32	+16 29 19.4	N
TN J2134+0946	351 ± 47	48.0 ± 2.3	...	−1.48 ± 0.11	...	...	...	...	21 34 42.85	+09 46 56.6	N
TN J2136+0947	426 ± 60	66.9 ± 3.1	...	−1.38 ± 0.11	...	...	...	...	21 36 36.29	+09 47 22.8	N
TN J2147−0645	382 ± 63	57.3 ± 2.7	...	−1.41 ± 0.13	...	...	...	...	21 47 57.13	−06 45 48.8	N
TN J2155−2109	449 ± 87	77.2 ± 1.6	...	−1.31 ± 0.15	...	...	...	...	21 55 50.90	−21 09 30.0	N
TN J2158−2516	1225 ± 59	200.9 ± 8.9	...	−1.34 ± 0.05	...	...	...	...	21 58 11.14	−25 16 01.5	N
TN J2201−2822	703 ± 54	120.5 ± 5.1	...	−1.31 ± 0.07	...	...	...	...	22 01 22.94	−28 22 14.0	N
TN J2204+1621	381 ± 42	57.3 ± 2.7	...	−1.41 ± 0.09	...	...	...	...	22 04 14.18	+16 21 43.7	N
TN J2205−3201	430 ± 60	65.7 ± 3.1	...	−1.40 ± 0.11	...	...	...	...	22 05 27.56	−32 01 23.9	N
TN J2217−1913	321 ± 34	40.5 ± 2.0	...	−1.54 ± 0.09	...	...	...	...	22 17 28.22	−19 13 21.1	N
TN J2219−1648	301 ± 53	32.0 ± 0.8	...	−1.67 ± 0.13	...	...	...	...	22 19 40.06	−16 48 03.1	N
TN J2232−2917	528 ± 59	91.7 ± 1.9	...	−1.30 ± 0.08	...	...	...	...	22 32 57.25	−29 17 56.5	N
TN J2237−0242	727 ± 48	125.0 ± 2.6	...	−1.31 ± 0.05	...	...	...	...	22 37 22.60	−02 42 23.0	N
TN J2239−0429	3807 ± 177	571.4 ± 24.9	...	−1.41 ± 0.05	...	...	...	...	22 39 32.85	−04 29 33.9	N
TN J2245+0007	297 ± 53	51.6 ± 1.1	...	−1.30 ± 0.13	...	...	...	...	22 45 44.99	+00 07 11.3	N
TN J2251−1634	551 ± 64	60.6 ± 1.3	...	−1.64 ± 0.09	...	...	...	...	22 51 14.06	−16 34 03.2	N
TN J2251−1034	862 ± 72	138.8 ± 5.9	...	−1.36 ± 0.07	...	...	...	...	22 51 22.01	−10 34 18.0	N
TN J2252−1618	397 ± 60	51.6 ± 2.5	...	−1.52 ± 0.12	...	...	...	...	22 52 20.15	−16 18 14.9	N
TN J2253−2156	425 ± 72	59.3 ± 2.8	...	−1.47 ± 0.13	...	...	...	...	22 53 17.03	−21 56 38.0	N
TN J2301+0555	484 ± 62	74.1 ± 3.3	...	−1.40 ± 0.10	...	...	...	...	23 01 18.11	+05 55 01.0	N
TN J2303−0251	405 ± 56	66.5 ± 3.1	...	−1.34 ± 0.11	...	...	...	...	23 03 55.67	−02 51 08.6	N
TN J2310+1358	1497 ± 102	251.9 ± 5.2	...	−1.33 ± 0.05	...	...	...	...	23 10 52.32	+13 58 20.5	N
TN J2312−0040	293 ± 25	45.5 ± 1.0	...	−1.39 ± 0.07	...	...	...	...	23 12 46.90	−00 40 53.0	N
TN J2312−0208	1295 ± 80	194.8 ± 8.6	...	−1.41 ± 0.06	...	...	...	...	23 12 55.71	−02 08 39.9	N
TN J2314+2053	386 ± 27	46.8 ± 2.3	5.9 ± 0.4	−1.57 ± 0.06	−1.67 ± 0.07	S	0.3	129	23 14 56.04	+20 53 37.4	V
TN J2320+1222	1566 ± 93	264.3 ± 5.4	...	−1.32 ± 0.05	...	...	...	...	23 20 07.79	+12 22 06.6	N
TN J2333−1508	504 ± 68	70.2 ± 3.3	...	−1.47 ± 0.11	...	...	...	...	23 33 06.99	−15 08 12.8	N
TN J2335+0634	507 ± 58	84.3 ± 3.8	...	−1.33 ± 0.09	...	...	...	...	23 35 06.42	+06 34 38.8	N
TN J2350−1321	419 ± 35	56.1 ± 1.2	...	−1.50 ± 0.06	...	...	...	...	23 50 15.92	−13 21 16.0	N
TN J2352+0814	498 ± 50	71.0 ± 3.2	...	−1.45 ± 0.08	...	...	...	...	23 52 10.44	+08 14 50.0	N
TN J2352+0450	398 ± 59	60.5 ± 2.8	...	−1.40 ± 0.12	...	...	...	...	23 52 32.20	+04 50 30.1	N
TN J2352−3139	573 ± 87	85.8 ± 3.8	29.0 ± 1.5	−1.41 ± 0.12	−0.87 ± 0.06	T	21.1	170	23 52 46.79	−31 39 36.9	V

† Not a real USS source; see notes.

\* See notes.

Table A.3. MP sample

Name	$S_{408}$ mJy	$S_{1420}$ mJy	$S_{4850}$ mJy	$\alpha_{408}^{1420}$	$\alpha_{408}^{4850}$	Str	LAS "	PA °	$\alpha_{J2000}$ h m s	$\delta_{J2000}$ ° ' "	Pos
MP J0003-3556	2440 ± 80	...	103.0 ± 13.0	...	-1.28 ± 0.05	...	...	...	00 03 13.11	-35 56 33.8	M
MP J0028-5523	1910 ± 80	389 ± 38	86.0 ± 9.0	-1.28 ± 0.10	-1.25 ± 0.05	S	3.7	57	00 28 53.76	-55 23 28.5	A
MP J0100-6403	1410 ± 150	247 ± 24	56.0 ± 7.0	-1.40 ± 0.11	-1.30 ± 0.07	S	4.2	100	01 00 32.42	-64 03 13.6	A
MP J0103-3225	2290 ± 80	...	93.0 ± 13.0	...	-1.29 ± 0.06	...	...	...	01 03 17.10	-32 25 53.3	M
MP J0114-3302	1590 ± 90	396 ± 39	76.0 ± 12.0	-1.11 ± 0.14	-1.23 ± 0.07	D	22.0	119	01 14 37.06	-33 02 10.2	A
MP J0118-6331	1040 ± 90	264 ± 26	50.0 ± 7.0	-1.10 ± 0.09	-1.23 ± 0.07	S	4.1	48	01 18 07.54	-63 31 43.2	A
MP J0130-8352	950 ± 50	159 ± 15	23.0 ± 5.0	-1.43 ± 0.11	-1.50 ± 0.09	S	3.2	66	01 30 07.45	-83 52 29.6	A
MP J0141-4421	1400 ± 60	295 ± 29	56.0 ± 9.0	-1.25 ± 0.10	-1.30 ± 0.07	D	29.9	95	01 41 13.57	-44 21 10.8	A
MP J0141-6941	3980 ± 320	755 ± 75	179.0 ± 11.0	-1.33 ± 0.09	-1.25 ± 0.04	D	16.1	153	01 41 55.18	-69 41 33.1	A
MP J0155-8521	830 ± 70	...	38.0 ± 5.0	...	-1.25 ± 0.06	...	...	...	01 55 16.98	-85 21 07.4	M
MP J0202-5425	820 ± 70	186 ± 18	41.0 ± 8.0	-1.19 ± 0.09	-1.21 ± 0.09	D	14.5	5	02 02 56.25	-54 25 13.3	A
MP J0211-5146	900 ± 40	160 ± 16	43.0 ± 8.0	-1.38 ± 0.08	-1.23 ± 0.08	S	11.1	50	02 11 44.92	-51 46 47.0	A
MP J0249-4145	1100 ± 50	242 ± 24	48.0 ± 9.0	-1.21 ± 0.08	-1.27 ± 0.08	S	8.2	49	02 49 10.08	-41 45 36.3	A
MP J0339-4018	3610 ± 140	759 ± 75	174.0 ± 13.0	-1.25 ± 0.09	-1.22 ± 0.03	S	5.6	33	03 39 48.18	-40 18 22.1	A
MP J0340-6507	1640 ± 50	383 ± 38	81.0 ± 8.0	-1.17 ± 0.10	-1.22 ± 0.04	D	14.0	18	03 40 44.92	-65 07 07.3	A
MP J0446-3305	1910 ± 100	448 ± 44	82.0 ± 12.0	-1.16 ± 0.10	-1.27 ± 0.06	D	58.8	6	04 46 00.21	-33 05 18.3	A
MP J0449-5449	900 ± 70	200 ± 20	46.0 ± 8.0	-1.20 ± 0.09	-1.20 ± 0.08	S	4.3	56	04 49 04.66	-54 49 10.1	A
MP J0601-3926	2270 ± 320	451 ± 45	97.0 ± 10.0	-1.29 ± 0.09	-1.27 ± 0.07	S	6.2	43	06 01 00.77	-39 26 18.1	A
MP J0605-5036	1040 ± 70	358 ± 35	43.0 ± 8.0	-0.85 ± 0.10	-1.29 ± 0.08	D	9.4	105	06 05 47.25	-50 36 41.7	A
MP J0618-7340	750 ± 50	...	25.0 ± 6.0	...	-1.37 ± 0.10	...	...	...	06 18 47.54	-73 40 40.8	M
MP J0648-7118	1000 ± 50	300 ± 30	42.0 ± 7.0	-0.96 ± 0.09	-1.28 ± 0.07	S	2.5	45	06 48 37.38	-71 18 26.4	A
MP J0731-6621	870 ± 40	...	40.0 ± 7.0	...	-1.24 ± 0.07	...	...	...	07 31 47.00	-66 21 30.3	M
MP J0839-7359	1450 ± 60	...	69.0 ± 7.0	...	-1.23 ± 0.04	DF	...	...	08 39 33.28	-73 59 37.4	M
MP J1033-3418	5590 ± 250	...	268.0 ± 22.0	...	-1.23 ± 0.04	...	...	...	10 33 13.04	-34 18 48.4	M
MP J1119-3631	2440 ± 70	542 ± 54	113.0 ± 14.0	-1.21 ± 0.10	-1.24 ± 0.05	S	4.8	166	11 19 21.81	-36 31 39.5	A
MP J1208-3403	4650 ± 210	...	198.0 ± 18.0	...	-1.28 ± 0.04	...	...	...	12 08 39.17	-34 03 10.7	M
MP J1250-4026	6030 ± 180	...	299.0 ± 18.0	...	-1.21 ± 0.03	...	...	...	12 50 05.68	-40 26 28.7	M
MP J1250-4119	1350 ± 60	167 ± 16	63.0 ± 9.0	-1.67 ± 0.09	-1.24 ± 0.06	S	7.4	123	12 50 46.49	-41 19 35.7	A
MP J1355-7806	1690 ± 90	...	44.0 ± 6.0	...	-1.47 ± 0.06	DF	...	...	13 55 35.57	-78 06 43.7	M
MP J1657-7423	1080 ± 50	...	50.0 ± 7.0	...	-1.24 ± 0.06	...	...	...	16 57 44.73	-74 23 09.3	M
MP J1710-7343	910 ± 70	47 ± 4	26.0 ± 6.0	-2.37 ± 0.10	-1.44 ± 0.10	S	...	149	17 10 17.73	-73 43 19.1	A
MP J1733-8342	4630 ± 210	...	198.0 ± 12.0	...	-1.27 ± 0.03	...	...	...	17 33 55.38	-83 42 48.0	M
MP J1755-6916	1580 ± 90	486 ± 48	71.0 ± 7.0	-0.94 ± 0.09	-1.25 ± 0.05	D	0.5	90	17 55 30.23	-69 16 49.8	A
MP J1758-6738	3000 ± 140	478 ± 47	112.0 ± 9.0	-1.47 ± 0.09	-1.33 ± 0.04	D	31.2	119	17 58 51.25	-67 38 30.5	A
MP J1821-7457	2190 ± 110	...	77.0 ± 7.0	...	-1.35 ± 0.04	...	...	...	18 21 59.83	-74 57 47.5	M
MP J1909-7755	900 ± 70	182 ± 18	41.0 ± 6.0	-1.28 ± 0.09	-1.25 ± 0.07	D	27.9	81	19 09 57.19	-77 56 01.5	A
MP J1912-5349	1110 ± 70	441 ± 44	56.0 ± 8.0	-0.74 ± 0.09	-1.21 ± 0.06	D	17.0	45	19 12 20.77	-53 49 01.3	A
MP J1921-6217	2110 ± 40	400 ± 40	79.0 ± 8.0	-1.33 ± 0.08	-1.33 ± 0.04	S	3.4	113	19 21 03.06	-62 17 25.2	A
MP J1921-5431	4590 ± 210	...	66.0 ± 8.0	...	-1.71 ± 0.05	...	...	...	19 21 52.92	-54 31 51.8	M
MP J1929-3732	3260 ± 110	806 ± 80	135.0 ± 12.0	-1.12 ± 0.09	-1.29 ± 0.04	S	3.7	14	19 29 08.51	-37 32 48.8	A
MP J1940-4236	1340 ± 70	...	64.0 ± 9.0	...	-1.23 ± 0.06	...	...	...	19 40 58.04	-42 36 49.5	M
MP J1942-5553	2120 ± 110	385 ± 38	74.0 ± 8.0	-1.37 ± 0.08	-1.36 ± 0.05	S	6.7	176	19 42 06.97	-55 53 32.6	A
MP J1943-4030	4400 ± 140	...	153.0 ± 12.0	...	-1.36 ± 0.03	DF	...	...	19 43 52.33	-40 30 10.9	M
MP J1953-4852	860 ± 30	143 ± 14	42.0 ± 8.0	-1.44 ± 0.09	-1.22 ± 0.08	S	4.7	115	19 53 39.88	-48 52 13.8	A
MP J2002-5252	2730 ± 130	411 ± 41	76.0 ± 9.0	-1.52 ± 0.09	-1.45 ± 0.05	S	4.5	29	20 02 22.89	-52 52 51.8	A
MP J2003-8340	1600 ± 80	253 ± 25	67.0 ± 6.0	-1.48 ± 0.09	-1.28 ± 0.04	S	3.5	106	20 03 30.73	-83 41 00.0	A
MP J2017-5747	1860 ± 160	287 ± 28	79.0 ± 8.0	-1.50 ± 0.08	-1.28 ± 0.05	S	11.8	28	20 17 48.11	-57 47 11.9	A
MP J2045-6018	11160 ± 340	2105 ± 210	447.0 ± 24.0	-1.34 ± 0.09	-1.30 ± 0.02	D	41.6	44	20 45 21.82	-60 18 51.5	A
MP J2048-5750	2810 ± 130	494 ± 49	137.0 ± 10.0	-1.39 ± 0.09	-1.22 ± 0.03	S	5.7	18	20 48 36.64	-57 50 47.6	A
MP J2126-5439	1180 ± 80	282 ± 28	57.0 ± 8.0	-1.15 ± 0.12	-1.22 ± 0.06	D	17.1	147	21 26 53.65	-54 39 36.0	A
MP J2204-5831	900 ± 70	148 ± 14	44.0 ± 8.0	-1.45 ± 0.08	-1.22 ± 0.08	S	10.5	11	22 04 05.48	-58 31 38.3	A
MP J2222-4723	910 ± 70	509 ± 50	45.0 ± 9.0	-0.47 ± 0.09	-1.21 ± 0.09	D	9.1	148	22 22 53.71	-47 23 12.1	A
MP J2226-7654	900 ± 60	203 ± 20	35.0 ± 6.0	-1.19 ± 0.09	-1.31 ± 0.07	S	4.1	107	22 26 56.90	-76 54 48.0	A
MP J2229-3824	8210 ± 210	...	412.0 ± 23.0	...	-1.21 ± 0.02	...	...	...	22 29 46.94	-38 24 02.9	M
MP J2308-6423	1750 ± 90	301 ± 30	39.0 ± 7.0	-1.41 ± 0.09	-1.54 ± 0.08	D	14.3	7	23 08 45.21	-64 23 33.3	A
MP J2313-4243	1390 ± 60	...	58.0 ± 9.0	...	-1.28 ± 0.07	...	...	...	23 13 58.76	-42 43 38.1	M
MP J2352-6154	1450 ± 40	311 ± 31	69.0 ± 8.0	-1.23 ± 0.10	-1.23 ± 0.05	D	56.1	148	23 52 55.52	-61 54 06.9	A
MP J2357-3445	8700 ± 350	...	117.0 ± 14.0	...	-1.74 ± 0.05	...	...	...	23 57 01.09	-34 45 38.9	M

**Table A.4.** WN POSS identifications

Name	$S_{1400}$ mJy	$\alpha_{325}^{1400}$	LAS "	LR <sup>a</sup>
WN J0029+3439	40.9	-1.34	12.8	4.2
WN J0121+4305	50.8	-1.39	...	7.1
WN J0315+3757	13.3	-1.35	1.7	14.8
WN J0559+6926	12.9	-1.31	0.3	4.6
WN J0641+4325	15.6	-1.57	...	195.6
WN J0725+4123	35.5	-1.39	8.3	183.0
WN J0756+5010	29.9	-1.31	1.7	127.3
WN J0830+3018	13.9	-1.41	3.5	27.3
WN J0835+3439	19.1	-1.38	58.0	24.6
WN J0923+4602	12.1	-1.49	21.6	17.0
WN J0952+5153	16.1	-1.76	2.7	194.1
WN J0955+6023	10.9	-1.54	3.1	1.0
WN J1014+7407	21.8	-1.41	2.2	147.7
WN J1026+2943	12.0	-1.31	11.9	136.5
WN J1030+5415	29.2	-1.39	6.1	16.4
WN J1052+4826	154.2	-1.31	10.0	38.1
WN J1124+3228	38.9	-1.31	5.7	101.2
WN J1130+4911	18.9	-1.34	37.0	18.9
WN J1141+6924	11.8	-1.32	...	1.4
WN J1148+5116	20.5	-1.35	6.7	1.0
WN J1152+3732	17.1	-2.18	15.0	159.9
WN J1232+4621	19.8	-1.30	12.5	42.9
WN J1258+5041	28.8	-1.42	44.9	3.7
WN J1259+3121	34.0	-1.41	20.6	0.0
WN J1330+6505	10.5	-1.34	1.8	21.5
WN J1353+3336	15.9	-1.31	...	3.3
WN J1400+4348	20.2	-1.49	6.4	29.2
WN J1403+3109	45.7	-1.35	11.0	148.5
WN J1410+4615	14.2	-1.36	4.4	42.5
WN J1440+3707	16.7	-1.76	7.4	182.0
WN J1459+4947	18.7	-1.53	3.0	47.2
WN J1558+7028	15.7	-1.36	2.1	2.8
WN J1624+4202	11.0	-1.31	24.4	182.2
WN J1628+3932	3680.7	-1.36	48.8	0.0
WN J1717+3828	35.8	-1.60	33.6	141.4
WN J1752+2949	16.9	-1.42	3.2	3.4
WN J1801+3336	11.6	-1.69	2.5	1.5
WN J1815+3656	17.6	-1.34	1.6	195.0
WN J1819+6213	12.7	-1.45	1.6	122.9
WN J1852+5711	53.4	-1.36	...	145.4
WN J1927+6436	80.3	-1.44	20.1	6.2
WN J1944+6552	147.2	-1.53	1.8	147.5
WN J2146+3330	10.8	-1.56	...	4.6
WN J2147+3137	25.4	-1.46	...	31.9
WN J2313+3842	11.3	-1.50	...	0.0
WN J2313+4053	40.2	-1.42	1.5	36.4

<sup>a</sup> Likelihood Ratio, see text.**Table A.5.** TN POSS identifications

Name	$S_{1400}$ mJy	$\alpha_{365}^{1400}$	LAS "	LR <sup>a</sup>
TN J0244+0327	84.8	-1.38	...	6.9
TN J0245+2700	54.0	-1.47	...	30.5
TN J0250+0130	39.1	-1.45	...	14.7
TN J0256-2717	177.9	-1.32	...	126.6
TN J0301+0155	402.7	-1.70	...	21.3
TN J0433+0717	33.1	-1.44	...	29.9
TN J0443-1212	56.8	-1.47	20.2	162.1
TN J0510-1838	641.3	-1.76	36.1	0.0
TN J0729+2436	68.9	-2.03	19.4	129.0
TN J0812+0915	43.4	-1.38	...	37.7
TN J0818-0741	59.3	-1.38	...	96.5
TN J0958-1103	70.6	-1.35	...	181.8
TN J1043+2404	52.0	-1.59	3.7	167.8
TN J1053-1518	63.7	-1.33	...	1.8
TN J1117-1409	134.5	-1.31	...	128.0
TN J1220+0604	65.1	-1.50	...	189.4
TN J1239+1005	79.3	-1.75	...	180.7
TN J1245-1127	88.0	-1.32	...	146.1
TN J1326-2330	83.2	-1.37	...	190.1
TN J1408-0855	114.3	-1.34	...	160.9
TN J1418-2256	82.0	-1.38	7.6	85.4
TN J1513-2417	83.6	-1.34	...	2.5
TN J1521+0741	105.6	-2.10	...	0.0
TN J1531-3234	80.6	-1.39	4.8	172.5
TN J1547-2218	72.4	-1.32	...	7.1
TN J1556-2759	66.4	-1.65	...	9.2
TN J1628+1604	55.5	-1.77	...	153.7
TN J1646-1328	102.5	-1.32	...	119.7
TN J1701+0252	57.1	-1.36	...	19.4
TN J1707-1120	54.2	-1.53	...	85.5
TN J1710-0826	69.2	-1.38	...	8.8
TN J1953-0541	59.0	-1.56	...	71.3
TN J2028+0811	74.8	-1.34	...	33.1
TN J2155-2109	77.2	-1.31	...	184.6
TN J2335+0634	84.3	-1.33	...	189.2

<sup>a</sup> Likelihood Ratio, see text.**Table A.6.** MP UKST identifications

Name	$S_{408}$ mJy	$\alpha_{408}^{4850}$	LAS "	LR <sup>a</sup>
MP J0003-3556	2440	-1.28	...	0.0
MP J0103-3225	2290	-1.29	...	88.1
MP J0618-7340	750	-1.37	...	7.5
MP J1033-3418	5590	-1.23	...	81.7
MP J1250-4026	1350	-1.24	7.4	131.0
MP J1921-5431	4590	-1.71	...	0.0
MP J1943-4030	4400	-1.36	...	0.0
MP J2313-4243	1390	-1.28	...	95.5
MP J2357-3445	8700	-1.74	...	0.0

<sup>a</sup> Likelihood Ratio, see text.

**Table A.7.** WN identifications from the literature

Name	$z$	$R$ [mag]	$F_X^a$	Identification	Reference
WN J0633+4653	...	23.6	...	4C +46.12	Chambers et al. 1996a
WN J0648+4309	...	23.0	...	B3 0644+432	Wieringa & Katgert 1992
WN J0658+4444	...	23.4	...	B3 0654+448	Wieringa & Katgert 1992
WN J0717+4611	1.462	21.6	...	B3 0714+462	De Breuck et al. 1998b
WN J0725+4123	...	...	0.1137	1RXS J072600.0+4	Voges et al. 1999
WN J0923+4602	...	...	0.0242	1WGA J0923.1+460	White et al. 1994
WN J0952+5153	0.214	...	0.2413	ZwCl 0949.6+5207	Voges et al. 1999
WN J1148+5116	...	...	0.0508	1RXS J114802.7+5	Voges et al. 1999
WN J1152+3732	...	...	0.0192	1WGA J1152.5+373	White et al. 1994
WN J1259+3121	...	16.2	...	NGP9 F323-0140639	Odewahn & Aldering 1996
WN J1332+3009	...	19.8	...	NGP9 F324-0235590	Odewahn & Aldering 1996
WN J1352+4259	...	21.7	...	4C +43.31	Vigotti et al. 1989
WN J1359+7447	...	...	0.07485	1RXS J135916.0+7	Voges et al. 1999
WN J1400+4348	...	8.2	...	HD122441	SIMBAD
WN J1410+4615	...	...	...	SHK 010	Shakhbazian 1973
WN J1436+6319	4.261	24.1	...	4C +63.20	Lacy et al. 1994
WN J1440+3707	...	...	0.07944	1RXS J144005.4+3	Voges et al. 1999
WN J1628+3932	0.031	13.1	4.5	NGC 6166	Zabludoff et al. 1993
WN J1736+6502	...	22.2	...	8C 1736+650	Lacy et al. 1993
WN J1829+6913	...	...	0.05765	1RXS J182903.8+6	Voges et al. 1999
WN J1852+5711	...	...	0.1466	1RXS J185209.4+5	Voges et al. 1999
WN J1944+6552	...	...	0.1118	1RXS J194423.1+6	Voges et al. 1999
WN J2319+4251	...	...	0.1255	1RXS J231947.4+4	Voges et al. 1999

<sup>a</sup>  $F_X$  is the number of X-ray counts  $s^{-1}$  as listed in the cited catalogs.

**Table A.8.** TN identifications from the literature

Name	$z$	$R$ [mag]	$F_X^a$	Identification	Reference
TN J0055+2624	0.1971	...	...	ABELL 85	Giovanini et al. 1999
TN J0102-2152	0.0604	...	...	ABELL 133	Komissarov & Gubanov 1994
TN J0245+2700	...	...	0.0782	1RXS J024521.0+2	Bade et al. 1998
TN J0256-2717	0.480	19.2	...	MRC 0254-27	McCarthy et al. 1996
TN J0301+0155	0.170	...	0.2053	ZwCl 0258.9+0142	Crawford et al. 1995
TN J0408-2418	2.440	22.7	...	MRC 0406-244	McCarthy et al. 1996
TN J0510-1838	...	19.5	0.00314	1RXS J051032.4-1	Voges et al. 1999
TN J0630-2834	...	...	0.00314	1WGA J0630.8-283	White et al. 1994
TN J0729+2436	...	...	0.1752	1RXS J072927.4+2	Voges et al. 1999
TN J0936-2243	1.339	23.0	...	3C 222	Heckman et al. 1994
TN J0958-1103	0.153	...	1.64	ABELL 0907	McCarthy et al. 1996
TN J1521+0742	0.045	14.5	1.64	NGC 5920	Komissarov & Gubanov 1994
TN J2106-2405	2.491	22.7	...	MRC 2104-242	McCarthy et al. 1996
TN J2239-0429	...	18.8	...	4C -04.85	Röttgering et al. 1995
TN J2320+1222	...	22.9	...	MRC 2317+120	Röttgering et al. 1995

<sup>a</sup>  $F_X$  is the number of X-ray counts  $s^{-1}$  as listed in the cited catalogs.

**Table A.9.** MP identifications from the literature

Name	$z$	$R$ [mag]	$F_X^a$	Identification	Reference
MP J0003-3556	0.0497	13.2	0.5206	ABELL 2717	Collins et al. 1995
MP J0103-3225	...	19.0	...	IRAS F01009-3241	Condon et al. 1995
MP J1250-4026	...	...	0.000595	1RXP J125006-402	Voges et al. 1999
MP J1943-4030	...	17.3	...	ABELL 3646	Abell et al. 1989
MP J2313-4243	0.0564	14.6	2.024	ABELL S1101	Stocke et al. 1991
MP J2357-3445	0.0490	13.7	2.27	ABELL 4059	Postman & Lauer 1995

<sup>a</sup>  $F_X$  is the number of X-ray counts  $s^{-1}$  as listed in the cited catalogs.

## Appendix B: Radio maps

### B.1. VLA maps of the WN sample

VLA maps of the WN sample. The contour scheme is a geometric progression in  $\sqrt{2}$ , which implies a factor 2 change in surface brightness every 2 contours. The first contour level, indicated above each plot, is at  $3\sigma_{\text{rms}}$ , where  $\sigma_{\text{rms}}$  is the rms noise determined around the sources. The restoring beams are indicated in the lower left corner of the plots. Two maps are given for each source, one showing a 6' field of view to show possible related components, and a smaller blow-up of the source to show its morphology. The open cross indicates the NVSS position. Sources identified on the POSS have been marked in the top right corner.

### B.2. FIRST maps of the WN sample

FIRST maps of the WN sample. Contours are as in Sect. B.1.

### B.3. VLA maps of the TN sample

VLA maps of the TN sample. Contours are as in Sect. B.1.

### B.4. FIRST maps of the TN sample

FIRST maps of the TN sample. Contours are as in Sect. B.1.

### B.5. ATCA maps of the TN sample

ATCA maps of the TN sample. Contours are as in Sect. B.1.

### B.6. ATCA maps of the MP sample

ATCA maps of the MP sample. Contours are as in Sect. B.1.

## Appendix C: Radio spectra

### C.1. Radio spectra for the WN sample

Radio spectra of the WN sample using data from the literature. The two connected flux points indicate the spectral index used to select the source in the USS sample. Note the steeper spectra with higher frequency in most objects.

### C.2. Radio spectra for the TN sample

Radio spectra of the TN sample using data from the literature. The two connected flux points indicate the spectral index used to select the source in the USS sample. Note the steeper spectra with higher frequency in most objects.

### C.3. Radio spectra for the MP sample

Radio spectra of the MP sample using data from the literature and our ATCA observations (diamonds). The two connected flux points indicate the spectral index used to select the source in the USS sample. Note the steeper spectra with higher frequency in most objects.

## Appendix D: POSS finding charts

### D.1. POSS finding charts for the WN sample

POSS finding charts of the WN sample. The open cross indicates the NVSS position.

### D.2. POSS finding charts for the TN sample

POSS finding charts of the TN sample. The open cross indicates the NVSS position.

### D.3. POSS finding charts for the MP sample

POSS finding charts of the MP sample. The open cross indicates the MRC position.

## References

- Abell G.O., Corwin H.G. Jr., Olowin R.P., 1989, ApJS 70, 1  
 Andernach H., Verkhodanov O., Verkhodanova N., 2000, IAU Symp. 199 (in press) astro-ph/0001473  
 Bade N., et al., 1998, A&AS 127, 145  
 Barthel P.D., 1989, ApJ 336, 606  
 Becker R.H., White R.L., Helfand D.J., 1995, ApJ 450, 559  
 Benn C.R., Rowan-Robinson M., McMahon R., Broadhurst T., Lawrence A., 1993, MNRAS 263, 98  
 Best P., Longair M., Röttgering H., 1998, MNRAS 295, 549  
 Blumenthal G., Miley G. 1979, A&A 80, 13  
 Blundell K., Rawlings S., Eales S., Taylor G., Bradley A.D., 1998, MNRAS 295, 265  
 Blundell K., Rawlings S., Willott C., 1999, AJ 117, 677  
 Bremer M.N., Rengelink R., Saunders R., Röttgering H.J.A., Miley G.K., Snellen I.A.G., 1998, in Observational Cosmology with the New Radio Surveys, Bremer M.N., Jackson N., Pérez-Fournon I. (eds.). Dordrecht: Kluwer, p. 165

- Carilli C.L., Röttgering H.J.A., van Ojik R., Miley G.K., van Breugel W.J.M., 1997, *ApJS* 109, 1
- Carilli C.L., Röttgering H.J.A., Miley G.K., Pentericci L.H., Harris D.E., 1999, in *Proc. The most distant radio galaxies*, Röttgering H.J.A., Best P.N., Lehnert M.D. (eds.). Amsterdam: KNAW, p. 123
- Chambers K.C., Miley G.K., van Breugel W.J.M., Huang J.-S., 1996, *ApJS* 106, 215
- Chambers K.C., Miley G.K., van Breugel W.J.M., Bremer M.A.R., Huang J.-S., Trentham N.A., 1996, *ApJS* 106, 247
- Coleman P.H., Condon J.J., 1985, *AJ* 90, 8
- Collins C.A., Guzzo L., Nichol R.C., Lumsden S.L., 1995, *MNRAS* 274, 1071
- Condon J.J., Anderson E., Broderick J.J., 1995, *AJ* 109, 2318
- Condon J., et al., 1998, *AJ* 115, 1693
- Crawford C., et al., 1995, *MNRAS* 274, 75
- De Breuck C., van Breugel W., Röttgering H., Miley G., 1998, in *Observational Cosmology with the New Radio Surveys*, Bremer M.N., Jackson N., Pérez-Fournon I. (eds.). Dordrecht: Kluwer, p. 185
- De Breuck C., Brotherton M.S., Tran H.D., van Breugel W., Röttgering H., 1998, *AJ* 116, 13
- De Breuck C., van Breugel W., Röttgering H., Miley G., Stern D., 2000 (in preparation)
- de Ruiter H.R., Willis A.G., Arp H.C., 1977, *A&A* 28, 211
- Dey A., Spinrad H., Stern D., Graham J.R., Chaffee F.H., 1998, *ApJ* 498, L93
- Dickinson M.E., in *Proc. STScI May 1997 Symposium "The Hubble Deep Field"*, Livio M., Fall S.M., Madau P. (eds.). Cambridge: Cambridge Univ. Press (in press) astro-ph/9802064
- Douglas J.N., Bash F.N., Bozyan F.A., Torrence G.W., Wolfe C., 1996, *AJ* 111, 1945
- Dunlop J., Peacock J., Spinrad H., Stern D., Windhorst R.A., 1996, *Nat* 381, 581
- Eales S., Rawlings S., Law-Green D., Cotter G., Lacy M., 1997, *MNRAS* 291, 593
- Faranoff B.L., Riley J.M., 1974, *MNRAS* 167, 31
- Ficarra A., Gruett G., Tomassetti G., 1985, *A&AS* 59, 255
- Frater R.H., Brooks J.W., Whiteoak J.B., 1992, *JEEE, Australia* 12 (2), 103
- Giovannini G., Tordi M., Feretti L., 1999, *New Astron.* 4, 141
- Gregg M.D., Becker R.H., White R.L., Helfand D.J., McMahon R.G., Hook I.M., 1996, *AJ* 112, 407
- Gregory P.C., Condon J.J., 1991, *ApJS* 75, 1011
- Griffith M., Wright A.E., 1993, *AJ* 105, 1666
- Griffith M., Wright A.E., Burke B.F., Ekers R.D., 1994, *ApJS* 90, 179
- Griffith M., Wright A.E., Burke B.F., Ekers R.D., 1995, *ApJS* 97, 347
- Hales S.E.G., Baldwin J.E., Warner P.G., 1993, *MNRAS* 263, 25
- Heckman T.M., O'Dea C.P., Baum S.A., Laurikainen E., 1994, *ApJ* 428, 65
- Hook I.M., McMahon R.G., 1998, *MNRAS* 294, 7L
- Hu E.M., McMahon R.G., Cowie L.L., 1999, *ApJ* 522, L9
- Hughes D.H., Dunlop J.S., Rawlings S., 1997, *MNRAS* 289, 766
- Ivison R., et al., 1998, *ApJ* 494, 211
- Jarvis M., Rawlings S., Willot C., Blundell K., Eales S., Lacy M., 1999, in *Proc. "The Hy-redshift universe"* (in press) astro-ph/9908106
- Kaplan D.L., Condon J.J., Arzoumanian Z., Cordes J.M., 1998, *ApJS* 119, 75
- Komissarov S.S., Gubanov A.G., 1994, *A&A* 285, 27
- Krolik J.H., Chen W., 1991, *AJ* 102, 1659
- Lacy M., Hill G., Kaiser M.-E., Rawlings S., 1993, *MNRAS* 263, 707
- Lacy M., et al., 1994, *MNRAS* 271, 504
- Large M.I., Mills B.Y., Little A.G., Crawford D.F., Sutton J.M., 1981, *MNRAS* 194, 693
- Laing R.A., Riley J., Longair M., 1983, *MNRAS* 204, 151
- Lilly S.J., Longair M.S., 1984, *MNRAS* 211, 833
- Lilly S.J., 1989, *ApJ* 340, 77
- Lorimer D.R., Yates J.A., Lyne A.G., Gould D.M., 1995, *MNRAS* 273, 411
- Machalski J., Rys S., 1981, *A&A* 99, 388
- Martel A., et al., 1998, *AJ* 115, 1348
- McCarthy P., 1993, *ARA&A* 693
- McCarthy P., Kapahi V.K., van Breugel W., Persson S.E., Atheya R.M., Subrahmanya C.R., 1996, *ApJS* 107, 19
- McLure R., Dunlop J., 2000, *MNRAS* (in press) astro-ph/9908214
- Napier P.J., Thompson A.R., Ekers R.D., 1983, *Proc. IEEE* 71, 1295
- Rawlings S., Lacy M., Blundell K., Eales S., Bunker A., Garrington S., 1996, *Nat* 383, 502
- Odehahn S.C., Aldering G., 1996 (private communication to NED)
- Oort M.J.A., Katgert P., Steeman F.W.M., Windhorst R.A., 1987, *A&A* 179, 41
- Oort M.J.A., 1988, *A&A* 193, 5
- Owen F.N., Condon J.J., Ledden J.E., 1983, *AJ* 88, 1
- Owen F.N., Laing R.A., 1989, *MNRAS* 238, 357
- Pedani M., Gruett G., 1999, *A&A* 350, 368
- Postman M., Lauer T.R., 1995, *ApJ* 440, 28
- Pursimo T., et al., 1999, *A&AS* 134, 50
- Rawlings S., Eales S., Warren, 1990, *MNRAS* 243, 14P
- Rawlings S., Blundell K.M., Lacy M., Willot C.J., 1998, in *Observational Cosmology with the New Radio Surveys*, Bremer M.N., Jackson N., Pérez-Fournon I. (eds.). Dordrecht: Kluwer, p. 171
- Rees N., 1990, *MNRAS* 244, 233
- Rengelink R., et al., 1997, *A&A* 124, 259
- Rengelink R., 1998, Ph.D. Thesis, Rijksuniversiteit Leiden
- Röttgering H.J.A., Lacy M., Miley G.K., Chambers K.C., Saunders R., *A&AS* 108, 79
- Röttgering H.J.A., Miley G.K., Chambers K.C., Machetto F., *A&AS* 114, 51
- Röttgering H.J.A., van Ojik R., Miley G.K., Chambers K.C., van Breugel W.J.M., de Koff S., 1997, *A&A* 326, 505
- Shakhbazian R.K., 1973, *Astrofizika* 9, 495
- Sheppard M.C., 1997, in *ASP Conf. Ser. 125, Astronomical Data Analysis Software and Systems VI*, Hunt G., Payne H.E. (eds.). San Francisco: Astron. Soc. of the Pacific, p. 77
- Spinrad H., Djorgovski S., Marr J., Aguilar L., 1985, *PASP* 97, 932
- Spinrad H., et al., 1997, *ApJ* 484, 581
- Spinrad H., et al., 1998, *AJ* 116, 2617
- Steidel C.C., Adelberger K.L., Giavalisco M., Dickinson M., Pettini M., 1999, *ApJ* 519, 1
- Stocke J., et al., 1991, *ApJS* 76, 813
- Tielsens A.G.G.M., Miley G.K., Willis A.G., 1979, *A&AS* 35, 153

- van Breugel W., Stanford S.A., Spinrad H., Stern D., Graham J.R., 1998, *ApJ* 502, 614
- van Breugel W., De Breuck C., Röttgering H., Miley G., Stanford A., 1999, in “Looking Deep in the Southern Sky”, Morganti R. (ed.). Sydney, p. 236
- van Breugel W., De Breuck C., Stanford S.A., Stern D., Röttgering H., Miley G., 1999, *ApJ* 518, 61
- Verkhodanov O.V., Trushkin S.A., Andernach H., Chernenkov V.N., 1997, in “Astronomical Data Analysis Software and Systems VI”, Hunt G., Payne H.E. (eds.). ASP Conf. Ser. 125, 322-325
- Vigotti M., Gruett G., Perley R., Clark B.G., Bridle A., 1989, *AJ* 98, 419
- Voges W., et al., 1999, *A&A* 349, 389
- Waddington I., Windhorst R., Cohen S., Partridge R., Spinrad H., Stern D., 1999, *ApJ* 526, L77
- Weymann R., et al., 1998, *ApJ* 505, L95
- White N.E., Giommi P., Angelini L., 1994, *IAU Circ.* 6100, 1
- White R.L., Becker R.H., Helfand D.J., Gregg M.D., 1997, *ApJ* 475, 479
- White R.L., et al., 2000, *ApJS* (in press) astro-ph/9912215
- Wieringa M.H., Ph.D. Thesis, Rijksuniversiteit Leiden
- Wieringa M.H., Katgert P., 1992, *A&AS* 93, 399
- Windhorst R.A., Miley G.K., Owen F.N., Kron R.G., Koo D.C., 1985, *ApJ* 289, 494
- Witzel A., Schmidt J., Pauliny-Toth I.I.K., Nauber U., 1979, *AJ* 84, 942
- Wright A.E., Griffith M.R., Burke B.F., Ekers R.D., 1994, *ApJS* 91, 111
- Wright A.E., Griffith M.R., Troup E., Hunt A., Burke B.F., Ekers R.D., 1996, *ApJS* 103, 145
- Zabuldoff A.I., Geller M.J., Huchra J.P., Vogeley M.S., 1993, *AJ* 106, 1273